



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07H 3/06, C12N 15/79, A61K 31/70, C12N 15/56, C07K 14/705, A61K 38/14		A2	(11) International Publication Number: WO 97/06176 (43) International Publication Date: 20 February 1997 (20.02.97)
(21) International Application Number: PCT/US96/12820		Edmond, OK 73034 (US). MOORE, Kevin, L. [US/US]; 612 N.W. 42nd Street, Oklahoma City, OK 73118 (US).	
(22) International Filing Date: 2 August 1996 (02.08.96)		(74) Agent: PABST, Patrea, L.; Arnall Golden & Gregory, 2800 One Atlantic Center, 1201 West Peachtree Street, Atlanta, GA 30309-3450 (US).	
(30) Priority Data: 08/510,920 3 August 1995 (03.08.95) US 60/017,794 15 May 1996 (15.05.96) US 08/649,802 17 May 1996 (17.05.96) US		(81) Designated States: AU, BG, BR, CA, CN, CZ, EE, GE, HU, IL, JP, KR, LT, LV, MX, NO, NZ, PL, RO, SG, SK, UA, US, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(60) Parent Applications or Grants (63) Related by Continuation US Filed on 08/510,920 (CIP) US Filed on 3 August 1995 (03.08.95) US Filed on 08/649,802 (CIP) US Filed on 17 May 1996 (17.05.96)		Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(71) Applicant (for all designated States except US): BOARD OF REGENTS OF THE UNIVERSITY OF OKLAHOMA [US/US]; 600 Parrington Oval, Norman, OK 73109 (US).		(72) Inventors; and (75) Inventors/Applicants (for US only): MCEVER, Rodger, P. [US/US]; 1716 Guilford Lane, Oklahoma City, OK 73120 (US). CUMMINGS, Richard, D. [US/US]; 5215 Santa Fe,	
(54) Title: PEPTIDE AND O-GLYCAN INHIBITORS OF SELECTIN MEDIATED INFLAMMATION			
(57) Abstract			
<p>Tyrosine sulfate on PSGL-1, particularly at least one of residues (46, 48) and (51), functions in conjunction with sialylated and fucosylated glycans, most preferably Thr-57, to mediate high affinity binding to P-selectin. PSGL-1 O-glycans have been determined to consist of disialylated or neutral forms of the core-2 tetrasaccharide $\text{Gal}\beta 1\rightarrow 4\text{GlcNAc}\beta 1\rightarrow 6(\text{Gal}\beta 1\rightarrow 3)\text{GalNAcOH}$. A minority of the O-glycans are $\alpha 1,3$ fucosylated that occur as two major species containing the sialyl Lewis x antigen - one species is a disialylated, monofucosylated glycan of formula (I), and the other is a monosialylated, trifucosylated glycan having a polylactosamine backbone of formula (II), wherein R = H, OH, another sugar or an aglycone such as an amino acid.</p>			
$ \begin{array}{c} \text{Fucosyl} \\ \downarrow \\ 3 \\ \text{NeuAc}\alpha 2\rightarrow 3\text{Gal}\beta 1\rightarrow 4\text{GlcNAc}\beta 1 \\ \downarrow \\ 6 \\ \text{NeuAc}\alpha 2\rightarrow 3\text{Gal}\beta 1\rightarrow 3\text{GalNAc-R,} \\ \text{(I)} \end{array} $			
$ \begin{array}{ccc} \text{Fucosyl} & \text{Fucosyl} & \text{Fucosyl} \\ \downarrow & \downarrow & \downarrow \\ 3 & 3 & 3 \\ \text{NeuAc}\alpha 2\rightarrow 3\text{Gal}\beta 1\rightarrow 4\text{GlcNAc}\beta 1\rightarrow 3\text{Gal}\beta 1\rightarrow 4\text{GlcNAc}\beta 1\rightarrow 3\text{Gal}\beta 1\rightarrow 4\text{GlcNAc}\beta 1 \\ \downarrow \\ 6 \\ \text{Gal}\beta 1\rightarrow 3\text{GalNAcR} \\ \text{(II)} \end{array} $			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

PEPTIDE AND O-GLYCAN INHIBITORS OF SELECTIN MEDIATED INFLAMMATION

Background of the Invention

5 The present invention is directed to inhibitors of selectin mediated inflammation and is particularly directed to inhibitors derived from PSGL-1, the ligand for P-selectin, and novel oligosaccharides derived from PSGL-1.

10 The United States government has rights in this invention by virtue of National Institutes of Health Grant PO1 HL54804 and RO1 HL 43510 to Richard D. Cummings and Rodger P. McEver.

15 The selectins are a family of three Ca^{2+} -dependent membrane-bound lectins that initiate adhesion of leukocytes to platelets or endothelial cells under the shear forces found in the venular circulation (Laskey, 1992) *Science* **258**, 964-969; Bevilacqua, et al., (1993) *J. Clin. Invest.* **91**, 379-387; McEver (1994) *Curr. Opin. Immunol.* **6**, 75-84). L-selectin, expressed on leukocytes, binds to constitutive or inducible ligands on endothelial cells. E-selectin, expressed by cytokine-activated endothelial cells, and P-selectin, expressed by thrombin-activated platelets 20 and endothelial cells, bind to ligands on myeloid cells and subsets of lymphocytes.

25 P-selectin is a calcium-dependent carbohydrate-binding protein that is expressed on the surfaces of activated platelets and endothelium in response to thrombin and other agonists (McEver, et al. (1995) *J. Biol. Chem.*, **270**, 11025-11028; Varki, A. (1994) *Proc. Natl. Acad. Sci.*, U.S.A. **91**, 7390-7397; Springer, T.A. (1995) *Annu. Rev. Physiol.* **57**, 827-872). Through its binding to glycoconjugate-based counterreceptors on leukocytes, P-selectin mediates rolling adhesion of these cells on activated platelets and endothelium (Lawrence and Springer (1991) *Cell* **65**, 852-873; Moore, et al. (1995) *J. Cell. Biol.* **128**, 661-671). Both sialic acid and fucose are components of the P-selectin counterreceptors

on leukocytes (Corral, et al. (1990) *Biochem. Biophys. Res. Comm.* **172**, 1349-1356; Moore, et al. (1992) *J. Cell. Biol.* **118**, 445-456; Sako, et al. (1993) *Cell* **75**, 1179-1186).

Although the selectins interact weakly with small sialylated, 5 fucosylated oligosaccharides such as sialyl Lewis x (sLe^x; NeuAc α 2-3Gal β 1-4[Fuc α 1-3]GlcNAc) (Foxall, et al., (1992) *J. Cell Biol.* **117**, 895-902; Varki, (1992) *Curr. Opin. Cell Biol.* **257**, 257-266), they bind with higher affinity to glycans displayed on a limited number of glycoproteins (Moore, et al., (1992) *J. Cell Biol.* **118**, 445-456; Lasky, et al., (1992) 10 *Cell* **69**, 927-938; Levinovitz, et al. (1993) *J. Cell Biol.* **121**, 449-459; Walcheck, et al., (1993) *J. Exp. Med.* **178**, 853-863; Baumhueter, et al., (1993) *Science* **262**, 436-438; Berg, et al., (1993) *Nature* **366**, 695-698) or proteoglycans (Norgard-Sumnicht, et al., (1993) *Science* **261**, 480-483). Oligosaccharides containing sialyl Lewis x (sLe^x), NeuAc α 2-15 3Gal β 1-4[Fuc α 1-3]GlcNAc β 1-R, a determinant present on leukocyte surfaces, inhibit adhesion of leukocytes to P-selectin (Polley, et al., (1991) *Proc. Natl. Acad. Sci. USA* **88**, 6224-6228; Foxall, et al. (1992) *J. Cell Biol.* **117**, 895-902). However, expression of sLe^x on cell surfaces is not sufficient for high affinity binding of cells to P-selectin, 20 since non-myeloid cells that express high levels of sLe^x bind poorly to P-selectin compared to myeloid cells (Zhou, et al. (1991) *J. Cell Biol.* **115**, 557-564). The high affinity ligands are potentially important as physiologic mediators of selectin-mediated leukocyte adhesion during inflammation. Thus, understanding the structural basis for high affinity 25 recognition of these glycoconjugates by the selectins has attracted increasing interest.

A subset of the high affinity selectin ligands consists of mucin-like glycoproteins (McEver, et al. (1995)). One sialomucin ligand for P-selectin is expressed by human neutrophils and the human promyelocytic 30 HL-60 cell line (Moore, et al., (1992); (Moore, et al., (1994) *J. Biol. Chem.* **269**, 23318-23327). Leukocytes express a single high affinity ligand for P-selectin, termed P-selectin glycoprotein ligand-1 (PSGL-1)

-3-

(Moore, et al., (1995); Moore, et al., (1992); Sako, et al., (1993); Norgaard, et al. (1993) *J. Biol. Chem.* **268**, 12764-127748; Moore, et al. (1994) *J. Biol. Chem.* **269**, 23318-23327). Binding of P-selectin to the ligand is Ca^{2+} dependent and is abolished by treatment of the ligand with 5 sialidase.

PSGL is a homodimer with two disulfide-linked subunits with relative molecular masses of approximately 120,000 as assessed by SDS-PAGE (Moore, et al., 1992). Each subunit has no more than three N-glycans, but has many clustered, sialylated O-glycans (Moore, et al. 10 (1992) ; Norgard, et al. (1993) *J. Biol. Chem.* **268**, 12764-12774). The extracellular domain of PSGL-1 is highly extended, a characteristic feature of mucin-like proteins. PSGL-1 is a type 1 membrane protein with an extracellular domain containing many serines, threonines, and prolines, including a series of decameric repeats (15 in human 15 promyelocytic HL-60 cells and 16 in human leukocytes) (Sako, et al. (1993) *Cell* **75**, 1179-1186; Moore, et al. (1995) *J. Cell Biol.* **128**, 661-671; Veldman, et al. (1995) *J. Biol. Chem.* **270**, 16470-16475). Following an 18-residue signal peptide, there is a propeptide spanning residues 19-41 that is removed from PSGL-1 following its synthesis in 20 leukocytes (Sako, et al. (1993); Vachino, et al. (1995) *J. Biol. Chem.* **270**, 21966-21974). The extracellular domain of the processed mature protein extends from residues 42 to 318, and is followed by a 25-residue transmembrane domain at a 69-residue cytoplasmic tail. It carries many unmodified sialic residues as well as the sLe^x antigen. The ligand 25 contains at least one PNGaseF-sensitive N-linked glycan that P-selectin does not require for recognition. In contrast, it has clustered, sialylated, O-linked oligosaccharides that render the polypeptide backbone sensitive to cleavage by the enzyme O-sialoglycoprotease. Treatment of intact HL- 30 60 cells with this enzyme eliminates the high affinity binding sites of P-selectin (Ushiyama, et al., (1993) *J. Biol. Chem.* **268**, 15229-15237) and prevents cell adhesion to immobilized P-selection (Norgard, et al., (1993) *J. Biol. Chem.* **268**, 12764-12774; Steininger, et al., *Biochem. Biophys.*

-4-

5 *Res. Commun.* (1992) **188**, 760-766), without affecting the overall surface expression of sLe^x. These data suggest that this sialomucin, which carries only a small portion of the cell surface sLe^x, corresponds to functionally important, high affinity binding sites for P-selectin on human myeloid cells.

Sako *et al.* (1993) isolated a cDNA derived from human HL-60 cells that encodes PSGL-1. The cDNA-derived sequence for each subunit of PSGL-1 predicts a type 1 transmembrane protein of 402 amino acids. The extracellular domain has an N-terminal signal peptide from residues 1 to 18 and a putative propeptide from residues 19-41. Assuming cleavage of the propeptide, the extracellular domain of the mature protein begins at residue 42 and extends to residue 308. The sequence concludes with a 25-residue transmembrane domain and a 69 residue cytoplasmic tail. The extracellular domain is rich in serines and threonines that are potential sites of O-glycosylation. The extracellular domain contains three potential sites for addition of *N*-linked oligosaccharides as well as a single cysteine that might promote dimerization and three potential tyrosine sulfation sites at residues 46, 48 and 51.

20 Although previous studies have shown that sialylation and fucosylation of PSGL-1 are required for its binding to P-selectin, other post-translational modifications of PSGL-1 may also be important. PSGL-1 is highly O-glycosylated and contains sialylated and fucosylated O-linked poly-N-acetyllactosamine, including some glycans that terminate in sLe^x (Moore, *et al.* (1994) *J. Biol. Chem.* **269**, 23318-23327). sLe^x or related glycans are not sufficient for high affinity binding of PSGL-1 to P-selectin. For example, sulfated compounds lacking either sialic acid or fucose can inhibit adhesion of leukocytes to P-selectin (Norgaard-Sumnicht, *et al.* (1993) *Science* **261**, 480-483; Nelson, *et al.* (1993) *Blood* **82**, 3253-3258; Cecconi, *et al.* (1994) *J. Biol. Chem.* **269**, 15060-15066; 25 Skinner, *et al.* (1989) *Biochem. Biophys. Res. Comm.* **164**, 1373-1379). These data suggest that sulfation of PSGL-1 may be required for its high affinity binding to P-selectin. Recombinant PSGL-1 expressed in COS

-5-

cells with a fucosyltransferase interacts with P-selectin in cell adhesion experiments (Sako, et al., 1993). However, the degree to which the oligosaccharides on recombinant PSGL-1 resemble those on the native glycoprotein ligand is unknown.

5 Although carbohydrates on PSGL-1 are critical for binding to selectins, no detailed chemical structures of the glycans are available. Much of the information about the glycosylation of the molecule has been obtained by enzymatic treatments of the native ligand and by studies on recombinant forms of PSGL-1 expressed in various cell types. While 10 these indirect methods can provide valuable information about critical determinants on the ligand, detailed structural information on O-glycans from native PSGL-1 is essential to identify glycans that are important for ligand function and to provide a clearer understanding of why PSGL-1 is a ligand for P-and E-selectin, whereas other mucins such as CD43 are 15 not.

It is therefore an object of the present invention to provide a method for making PSGL-1 which is normally glycosylated and sulfated.

It is another object of the present invention to provide a method and reagents for inhibiting binding of PSGL-1 to P-selectin and to other 20 selectins.

It is a further object of the present invention to provide novel O-glycans, which are useful for modifying binding to P-selectin.

Summary of the Invention

25 Sulfated, glycosylated peptides and the O-glycan structures uniquely present on PSGL-1 which bind to P-selectin have been developed using studies which identified the critical amino acid residues in the N-terminal region of PSGL-1, the importance of sulfation of the tyrosine residues, and the structure and role of the O-glycan carbohydrate structures coupled to threonine in the N-terminal region.


30 The examples demonstrate that PSGL-1 is sulfated, primarily on one or more of the tyrosine at residues 46, 48 and 51 which are required

for high affinity binding to P-selectin. Studies demonstrate that PSGL-1 synthesized in human HL-60 cells can be metabolically-labeled with [³⁵S]sulfate that is incorporated primarily into tyrosine sulfate and that treatment of PSGL-1 with a bacterial arylsulfatase releases sulfate from 5 tyrosine, resulting in a concordant decrease in binding to P-selectin.


Moreover, the binding of PSGL-1 to P-selectin is blocked by antiserum directed against a synthetic peptide encompassing the three putative tyrosine sulfation sites near the N-terminus of PSGL-1.

Studies also demonstrate that the expression of PSGL-1 in a 10 recombinantly engineered mammalian cell expressing two enzymes is required for proper glycosylation of PSGL-1: a fucosyltransferase (FTIII) and a core 2 β 1-6-N-acetylglucosaminyltransferase. Comparative tests demonstrate that the glycosylation of PSGL-1 expressed in cells expressing these enzymes is more similar to the glycosylation of native 15 PSGL-1 than the glycosylation of recombinant PSGL-1 expressed in COS or CHO cells not expressing the core 2 β 1-6-N-acetylglucosaminyltransferase. The structures of the Ser/Thr-linked O-glycans of PSGL-1 synthesized by HL-60 cells were then metabolically-radiolabeled with ³H-sugar precursors. In control studies, the O-glycans 20 on CD43 (leukosialin), a mucin-like glycoprotein also expressed by HL-60 cells, were analyzed and compared to those of PSGL-1. O-glycans were released from Ser/Thr residues by mild base/borohydride treatment of purified glycoproteins, and glycan structures were determined by a combination of techniques. In contrast to expectations, PSGL-1 is not 25 heavily fucosylated; a majority of the O-glycans are disialylated or neutral forms of the core-2 tetrasaccharide Gal β 1 \rightarrow 4GlcNAc β 1 \rightarrow 6(Ba β 1 \rightarrow 3)GalNAcOH. A minority of the O-glycans are α 1,3 fucosylated that occur as two major species containing the sialyl Lewis x antigen - one species is a disialylated, monofucosylated glycan:

-7-



and the other is a monosialylated, trifucosylated glycan having a polylactosamine backbone:



wherein R = H, OH, another sugar or an aglycone such as an amino acid.

These results demonstrate that PSGL-1 contains unique fucosylated O-glycans that are involved in high affinity interactions between PSGL-1 and selectins.

Brief Description of the Drawings

Figures 1A, 1B, and 1C are graphs of the retention time for anion exchange chromatography using a NaH_2PO_4 , pH 3.0 gradient (dashed line) for enzymatically digested PSGL-1 demonstrating that PSGL-1 contains tyrosine sulfate that is sensitive to arylsulfatase. ^{35}S -PSGL-1 hydrolyzed with strong base is shown in Figure 1A; ^{35}S -PSGL-1 hydrolyzed with strong base and sham-treated with 1000 mU of boiled arylsulfatase is shown in Figure 1B; and ^{35}S -PSGL-1 hydrolyzed with strong base and treated with 1000 mU of active enzyme is shown in Figure 1C. The retention times of tyrosine, tyrosine sulfate, and free sulfate are indicated. Free sulfate eluted at 40.0 min. Sulfated monosaccharides (Gal-6-sulfate, GlcNAc-6-sulfate, GalNAc-6-sulfate, and GalNAc-4-sulfate) eluted with similar retention times between 14 and 15 min.

-8-

Figure 2 is a graph of arylsulfatase release of [³⁵S]sulfate from intact ³⁵S-PSGL-1 and percent decrease in Ca²⁺-dependent binding of ¹²⁵I-PSGL-1 to P-selectin, showing that tyrosine sulfate is important for P-selectin binding.

5 Figures 3A and 3B are graphs showing that a specific anti-peptide serum to PSGL-1 blocks binding of PSGL-1 to P-selectin. ¹²⁵I-PSGL-1 was incubated in the presence of normal rabbit serum (NRS) or the antiserum directed against residues 42-56 of PSGL-1 (anti-42-56 amino acids) in microtiter plates containing immobilized recombinant soluble P-
10 selectin or human serum albumin (HSA), and binding measured as a function of serum concentration, as shown in Figure 3A. Platelet-derived P-selectin was incubated with HL-60 cells in the presence of 20% NRS or anti-42-56 amino acids serum. Binding of P-selectin was assayed by indirect immunofluorescence and flow cytometry, and cell number plotted
15 against P-selectin binding, as shown in Figure 3B.

Figures 4A-E, show that PSGL-1 fraction P-10-2b contains a minimum-sized glycan carrying the sLe^x determinant. Fraction P-10-2b derived from ³H-GlcN-PSGL-1 was treated with exoglycosidases and analyzed by descending paper chromatography for 24h. The β -eliminated
20 glycans were first desialylated with neuraminidase and the neutral glycans were separated from released sialic acid by ion-exchange chromatography.
(A) Chromatography of desialylated glycans; (B) desialylated glycans treated with β -galactosidase; (C) desialylated glycans treated with *Streptomyces* α 1,3/4-fucosidase; (D) desialylated glycans treated with a
25 combination of β -galactosidase, β -N-acetylhexosaminidase and *Streptomyces* α 1,3/4-fucosidase. (E) β -eliminated glycans derived from ³H-Fuc-PSGL-1 fraction P-10-2b were enzymatically desialylated and co-chromatographed in the same experiment. Migration of authentic standards is indicated: 5,
30 Gal β 1 \rightarrow 4(Fuc α 1 \rightarrow 3)GlcNAc β 1 \rightarrow 6(Gal β 1 \rightarrow 3)GalNAcOH; 4, Gal β 1 \rightarrow 4 GlcNA β 1 \rightarrow 6(Gal β 1 \rightarrow 3)GalNAcOH; 3*,

-9-

GlcNAc β 1 \rightarrow 6(Gal β 1 \rightarrow 3)GalNAcOH; 2, Gal β 1 \rightarrow 3 GalNAcOH; 1, GlcNAc.

Figures 5A and B, show that PSGL-1 fraction P-10-2a consists of a sialylated, polylactosamine-containing O-glycan that is trifucosylated.

5 Fraction P-10-2a derived from 3 H-GlcN-PSGL-1 was treated with exoglycosidase and analyzed by descending paper chromatography for 24 h. The β -eliminated glycans were first desialylated with neuraminidase, and the released sialic acid was separated from neutral glycans before and after treatments with either endo- β -galactosidase, or a combination of β -galactosidase and β -N-acetylhexosaminidase. (B) Desialylated glycans were chemically defucosylated and analyzed before and after treatments with either endo- β -galactosidase or a combination of β -galactosidase plus β -N-acetylhexosaminidase. Migration of authentic standards is indicated:
10 3, Gal β 1 \rightarrow 4GlcNAc β 1 \rightarrow 3Gal β 1; 3*, GlcNAc β 1 \rightarrow 6(Gal β 1 \rightarrow 3)GalNAcOH;
15 2, Gal β 1 \rightarrow 4GlcNAc; 1, GlcNAc.

Figure 6 shows that P-10-3 glycans contain the neutral core-2 tetrasaccharide. 3 H-GlcN-P-10-3 was treated with neuraminidase and the released sialic acid separated from neutral glycans by QAE-Sephadex chromatography. The desialylated glycans were analyzed by descending paper chromatography for 18 h before and after treatment with either β -galactosidase alone, or β -galactosidase plus β -N-acetylhexosaminidase.
20 Migration of authentic standards is indicated:
4, Gal β 1 \rightarrow 4GlcNAc β 1 \rightarrow 6(Gal β 1 \rightarrow 3)GalNAcOH; e*,
GlcNAc β 1 \rightarrow 6(Gal β 1 \rightarrow 3)
25 GalNAcOH; 2, Gal β 1 \rightarrow 4GlcNAc; 1, GlcNAc.

Figure 7 are structures and relative percentages of the O-glycans in PSGL-1.

Detailed Description of the Invention

Pharmaceutical and Diagnostic Peptides

30 Based on the information obtained from the examples described below, one can synthesize using recombinant techniques or peptide

-10-

synthesizers followed by chemical or enzymatic sulfation and glycosylation, peptides which are useful for inhibiting binding of P-selectin and other selectins to PSGL-1, as described by the foregoing general formula and reference to the figures and examples below. The 5 peptides preferably include at a minimum one of the three tyrosines at the amino terminus of the protein (residues 46, 48, and 51 of Sequence Listing ID No. 1), at least one of which is sulfated (-SO₄), and preferably include at least one threonine or serine which is suitable for O-linked glycosylation, for example, the threonine residue at amino acid residue 57 10 of Sequence Listing ID No. 1. An example of a minimum useful peptide corresponds to residues 48 to 57 of Sequence ID No. 1, as well as peptides having conservative substitutions which do not alter binding of the peptide to P-selectin. As demonstrated by the examples, it is important that the peptide include a core-2, sialylated, fucosylated O-glycan, as demonstrated in the following examples. 15

Amino acid sequence modifications fall into one or more of three classes: substitutional, insertional or deletional variants. Insertions include amino and/or carboxyl terminal fusions as well as intrasequence insertions of single or multiple amino acid residues. Insertions ordinarily 20 will be smaller insertions than those of amino or carboxyl terminal fusions, for example, on the order of one to four residues. Immunogenic fusion protein derivatives, such as those described in the examples, are made by fusing a polypeptide sufficiently large to confer immunogenicity to the target sequence by cross-linking *in vitro* or by recombinant cell 25 culture transformed with DNA encoding the fusion. Deletions are characterized by the removal of one or more amino acid residues from the protein sequence. Typically, no more than about from 2 to 6 residues are deleted at any one site within the protein molecule. These variants ordinarily are prepared by site specific mutagenesis of nucleotides in the 30 DNA encoding the protein, thereby producing DNA encoding the variant, and thereafter expressing the DNA in recombinant cell culture. Techniques for making substitution mutations at predetermined sites in

DNA having a known sequence are well known, for example M13 primer mutagenesis. Amino acid substitutions are typically of single residues; insertions usually will be on the order of about from 1 to 10 amino acid residues; and deletions will range about from 1 to 30 residues. Deletions or insertions preferably are made in adjacent pairs, i.e. a deletion of 2 residues or insertion of 2 residues. Substitutions, deletions, insertions or any combination thereof may be combined to arrive at a final construct. The mutations must not place the sequence out of reading frame and preferably will not create complementary regions that could produce secondary mRNA structure. Substitutional variants are those in which at least one residue has been removed and a different residue inserted in its place. Such substitutions generally are made in accordance with the following Tables 1 and 2 and are referred to as conservative substitutions.

15 TABLE 1: Amino Acid Abbreviations

	<u>Amino Acid</u>	<u>Abbreviations</u>
	alanine	Ala
	allosoleucine	Alle
	arginine	Arg
	asparagine	Asn
20	aspartic acid	Asp
	cysteine	Cys
	glutamic acid	Glu
	glutamine	Gln
	glycine	Gly
25	histidine	His
	isolelucine	Ile
	leucine	Leu
	lysine	Lys
	phenylalanine	Phe
30	proline	Pro
	pyroglutamic acid	pGlu
	serine	Ser
	threonine	Thr
	tyrosine	Tyr
35	tryptophan	Trp
	valine	Val

-12-

TABLE 2: Amino Acid Substitutions

	<u>Original Residue</u>	<u>Exemplary Substitutions</u>
--	-------------------------	--------------------------------

	Ala	ser
	Arg	lys
5	Asn	gln; his
	Asp	glu
	Cys	ser
	Gln	asn
	Glu	asp
10	Gly	pro
	His	asn;gln
	Ile	leu; val
	Leu	ile; val
	Lys	arg; gln; glu
15	Met	Leu; ile
	Phe	met; leu; tyr
	Ser	thr
	Thr	ser
	Trp	tyr
20	Tyr	trp; phe
	Val	ile; leu

Substantial changes in function or immunological identity are made by selecting substitutions that are less conservative than those in Table 2, 25 i.e., selecting residues that differ more significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site or (c) the bulk of the side chain. The substitutions which in general are expected to 30 produce the greatest changes in the protein properties will be those in which (a) a hydrophilic residue, e.g. seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g. leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g., lysyl, 35 arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g., glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain,

-13-

e.g., glycine, in this case, (e) by increasing the number of sites for sulfation and/or glycosylation

Substitutional or deletional mutagenesis can be employed to insert sites for N-glycosylation (Asn-X-Thr/Ser) or O-glycosylation (Ser or

5 Thr). Deletions of cysteine or other labile residues also may be desirable. Deletions or substitutions of potential proteolysis sites, e.g. Arg, is accomplished for example by deleting one of the basic residues or substituting one by glutaminyl or histidyl residues.

10 Certain post-translational derivatizations are the result of the action of recombinant host cells on the expressed polypeptide. Glutaminyl and asparaginyl residues are frequently post-translationally deamidated to the corresponding glutamyl and asparyl residues. Alternatively, these residues are deamidated under mildly acidic conditions. Other post-translational modifications include hydroxylation of proline and lysine, 15 phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the o-amino groups of lysine, arginine, and histidine side chains (T.E. Creighton, Proteins: Structure and Molecular Properties, W. H. Freeman & Co., San Francisco pp 79-86 [1983]), acetylation of the N-terminal amine and, in some instances, amidation of the C-terminal 20 carboxyl.

Methods of Preparation of Peptides

The peptides can generally be prepared following known techniques, as described for example in the cited publications, the teachings of which are specifically incorporated herein. Since the 25 preferred peptides are both glycosylated and sulfated, it is preferable to express the peptides in appropriate mammalian host cells as described below.

Chemical Synthesis

In a preferred synthetic method, the peptides are prepared 30 following the solid-phase synthetic technique initially described by Merrifield in

-14-

J. Amer. Chem. Soc., 85, 2149-2154 (1963). Other techniques may be found, for example, in M. Bodanszky, et al., Peptide Synthesis, second edition, (John Wiley & Sons, 1976), as well as in other reference works known to those skilled in the art.

General synthetic procedures for the synthesis of peptides by solid phase peptide synthesis using N^α-Boc protection.

			<u>REPETITIONS</u>	<u>TIME</u>
20	1.	25% TFA in CH_2Cl_2	1	3 min
	2.	50% TFA in CH_2Cl_2	1	16 min
	3.	CH_2Cl_2	5	3 min
	4.	5% DIEA in NMP	2	4 min
	5.	NMP	6	5 min
25	6.	Coupling step	1	57 min
	a.	Preformed BOC-Amino Acid- HOBT active ester in NMP		36 min
	b.	DMSO		16 min
	c.	DIEA		5 min
30	7.	10% Ac_2O , 5% DIEA in NMP	1	9 min
	8.	CH_2Cl_2	5	3 min

-15-

General synthetic procedures for solid phase peptide synthesis
using N^α- FMOC protection.

		<u>REPETITIONS</u>	<u>TIME</u>
5	1.	20% piperidine in NMP	1
	2.	20% piperidine in NMP	1
	3.	NMP	6
	4.	Coupling	1
Preformed FMOC-Amino Acid- HCBT active ester in NMP			
10	5.	NMP	6
N-terminal acetylation on the deprotected N ^α -amino group of peptides synthesized using either Boc or FMOC strategies is accomplished with 10% Ac ₂ O and 5% DIEA in NMP, followed by washing of the peptide resin with NMP and/or CH ₂ Cl ₂ .			

15 Expression Systems

As demonstrated by the following examples, the sulfated, glycosylated PSGL-1 peptides described herein which bind to P-selectin are those having at least one sulfate bound to a tyrosine residue and appropriate glycosylation. Comparative studies of recombinant PSGL-1 expressed in mammalian cells (COS cells) which express a fucosyltransferase with purified native PSGL-1 demonstrated that the glycosylation was substantially different. A new expression system has therefore been developed which results in more natural glycosylation of PSGL-1. The expression system is based on a mammalian expression system which expresses at least two glycosyltransferases: fucosyltransferase III (FTIII) and core 2 β 1-6-N-acetylglucosaminyltransferase. In a preferred embodiment, the cell line is based on a well characterized expression line such as COS or CHO cells which are transfected with the cDNA encoding the two enzymes. The cDNAs are shown below as Sequence Listing ID Nos. 2 and 3, respectively. Sequence Listing ID No. 2 encodes a human α (1,3/1,4)fucosyltransferase (FTIII) described by Kukowska-Latallo, et

-16-

al., *Genes & Devel.* **4**, 1288-1303 (1990). Sequence Listing ID No. 3 encodes Core 2 β 1-6-N-acetylglucosaminyltransferase (EC 2.4.1.102), as reported by Bierhuizen and Fukuda, *Proc. Natl. Acad. Sci. USA* **89**, 9326-9330 (1992). Prior studies had failed to recognize the importance of 5 the core 2 6-N-acetylglucosaminyltransferase for modification of PSGL-1 to confer binding to P-selectin. Alternatively, the expression system can be selected based on expression of either or both enzymes naturally, with the second enzyme being provided by transfection if necessary.

10 The cDNAs are introduced into the cells under the control of appropriate promoters and, optionally, enhancers and selection sequences.

Preferred promoters controlling transcription from vectors in mammalian host cells may be obtained from various sources, for example, the genomes of viruses such as: Simian Virus 40 (SV40), adenovirus, retroviruses, hepatitis-B virus and most preferably cytomegalovirus, or 15 from heterologous mammalian promoters, e.g. beta actin promoter. The early and late promoters of the SV40 virus are conveniently obtained as an SV40 restriction fragment which also contains the SV40 viral origin of replication. Fiers et al., *Nature*, **273**: 113 (1978). The immediate early promoter of the human-cytomegalovirus is conveniently obtained as a 20 HindIII restriction fragment. Greenaway, P. J. et al., *Gene* **18** 355-360 (1982). Of course, promoters from the host cell or related species also are useful.

Transcription of a DNA encoding a desired protein by higher 25 eukaryotes is increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually from about 10 to 300 bp, that act on a promoter to increase its transcription initiation capability. Enhancers are relatively orientation and position independent having been found 5' (Laimins, L. et al., *Proc. Natl. Acad. Sci.* **78**: 993) and 3' (Lusky, M. L., et al., *Mol. Cell Bio.* **3**: 1108 (1983)) to the transcription 30 unit, within an intron (Banerji, J. L. et al., *Cell* **33**: 729 (1983)) as well as within the coding sequence itself (Osborne, T.F., et al., *Mol. Cell Bio.* **4** 1293 (1984)). Many enhancer sequences are now known from

-17-

mammalian genes (globin, elastase, albumin, α -fetoprotein and insulin).

Typically, however, one will use an enhancer from an eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter

5 enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human or nucleated cells) may also contain sequences necessary for the termination of transcription which may affect 10 mRNA expression. These regions are transcribed as polyadenylated segments in the untranslated portion of the mRNA encoding the desired protein. The 3' untranslated regions also include transcription termination sites.

Expression vectors may contain a selection gene or selectable

15 marker. Examples of suitable selectable markers for mammalian cells are dihydrofolate reductase (DHFR), thymidine kinase or neomycin. When the selectable markers are successfully transferred into a mammalian host cell, the transformed mammalian host cell survives when placed under selective pressure. There are two widely used distinct categories of 20 selective regimes. The first category is based on a cell's metabolism and the use of a mutant cell line which lacks the ability to grow independent of a supplemented media. Two examples are: DHFR-minus CHO cells and mouse LTK cells. These cells lack the ability to grow without the addition of such nutrients as thymidine or hypoxanthine. Since these cells 25 lack certain genes necessary for a complete nucleotide synthesis pathway, they cannot survive unless the missing nucleotides are provided in a supplemented media. An alternative to supplementing the media is to introduce an intact DHFR or TK gene into cells lacking the respective genes, thus altering their growth requirements. Individual cells which 30 were not transformed with the DHFR or TK gene will not be capable of survival in non supplemented media.

-18-

The second category is dominant selection which refers to a selection scheme used in any cell type and does not require the use of a mutant cell line. These schemes typically use a drug to arrest growth of a host cell. Those cells which have a novel gene would express a protein 5 conveying drug resistance and would survive the selection. Examples of such dominant selection use the drugs neomycin, Southern P. and Berg, P., *J. Molec. Appl. Genet.* **1**: 327 (1982), mycophenolic acid, Mulligan, R. C. and Berg, P. *Science* **209**: 1422 (1980) or hygromycin, Sugden, B. et al., *Mol. Cell. Biol.*, **5** 410-413 (1985). The three examples given 10 above employ bacterial genes under eukaryotic control to convey resistance to the appropriate drug G418 or neomycin (geneticin), xgpt (mycophenolic acid) or hygromycin, respectively.

"Amplification" refers to the increase or replication of an isolated region within a cell's chromosomal DNA. Amplification is achieved 15 using a selection agent e.g. methotrexate (MTX) which inactivates DHFR. Amplification or the making of successive copies of the DHFR gene results in greater amounts of DHFR being produced in the face of greater amounts of MTX. Amplification pressure is applied notwithstanding the presence of endogenous DHFR, by adding ever greater amounts of MTX 20 to the media. Amplification of a desired gene can be achieved by cotransfected a mammalian host cell with a plasmid having a DNA encoding a desired protein and the DHFR or amplification gene permitting cointegration. One ensures that the cell requires more DHFR, which requirement is met by replication of the selection gene, by selecting only 25 for cells that can grow in the presence of ever-greater MTX concentration. So long as the gene encoding a desired heterologous protein has cointegrated with the selection gene replication of this gene gives rise to replication of the gene encoding the desired protein. The result is that increased copies of the gene, i.e. an amplified gene, 30 encoding the desired heterologous protein express more of the desired heterologous protein.

-19-

Preferred suitable host cells for expressing the vectors in higher eukaryotes include: monkey kidney CVI line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293, Graham, F.L. et al. *J. Gen Virol.* **36**: 59 (1977)); baby hamster kidney cells (BHK, 5 ATCC CCL 10); chinese hamster ovary-cells-DHFR (CHO, Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA* **77** 4216, (1980)); mouse sertoli cells (TM4, Mather, J. P., *Biol. Reprod.* **23**: 243-251 (1980)); monkey kidney cells (CVI ATCC CCL 70); african green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC 10 CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); and, TRI cells (Mather, J. P. et al., *Annals N. Y. Acad. Sci* **383**: 44-68 (1982)). Host cells can be 15 transformed with the expression vectors and cultured in conventional nutrient media modified as is appropriate for inducing promoters, selecting transformants or amplifying genes. The culture conditions, such as temperature and pH, are those previously used with the host cell selected for expression.

20 **Preparation of Diagnostic and Therapeutic Agents
Derived from the Protein or Carbohydrate Components
of the Glycoprotein Ligand for P-selectin.**

The glycoprotein ligand for P-selectin described above has a 25 variety of applications as diagnostic reagents and, potentially, in the treatment of numerous inflammatory and thrombotic disorders.

Diagnostic Reagents.

Antibodies to the ligand can be used for the detection of human 30 disorders in which P-selectin ligands might be defective. Such disorders would most likely be seen in patients with increased susceptibility to infections in which leukocytes might not be able to bind to activated platelets or endothelium. Cells to be tested, usually leukocytes, are collected by standard medically approved techniques and screened.

Detection systems include ELISA procedures, binding of radiolabeled

-20-

antibody to immobilized activated cells, flow cytometry, immunoperoxidase or immunogold analysis, or other methods known to those skilled in the arts.

Antibodies directed specifically to protein or carbohydrate

5 components of the ligand can be used to distinguish defects in expression of the core protein or in glycosyltransferases and/or modifying enzymes that construct the proper oligosaccharide chains on the protein. The antibodies can also be used to screen cells and tissues other than leukocytes for expression of the protein or carbohydrate components of

10 the ligand for P-selectin.

Complementary DNA clones encoding the protein component of the ligand can be isolated and sequenced. These cDNA probes can be used as diagnostic reagents to examine expression of RNA transcripts for the ligand in leukocytes and other tissues by standard procedures such as

15 Northern blotting of RNA isolated from cells and *in situ* hybridization of tissue sections.

A similar approach can be used to determine qualitative or quantitative disorders of P-selectin itself. The glycoprotein ligand, carbohydrates, or appropriate derivatives thereof, is labeled and tested for

20 its ability to bind to P-selectin on activated platelets from patients with disorders in which P-selectin might be defective.

The ligand, or components thereof, can also be used in assays of P-selectin binding to screen for compounds that block interactions of P-selectin with the ligand.

25 **Clinical Applications.**

Since P-selectin has several functions related to leukocyte adherence, inflammation, tumor metastases, and coagulation, clinically, compounds which interfere with binding of P-selectin and/or the other selectins, including E-selectin and L-selectin, such as the carbohydrates,

30 can be used to modulate these responses. These compounds include PSGL-1 or fragments thereof, antibodies to PSGL-1 or fragments thereof. For example, the glycoprotein ligand, or components thereof, particularly

-21-

the carbohydrate moieties, can be used to inhibit leukocyte adhesion by competitively binding to P-selectin expressed on the surface of activated platelets or endothelial cells. Similarly, antibodies to the ligand can be used to block cell adhesion mediated by P-selectin by competitively

5 binding to the P-selectin ligand on leukocytes or other cells. These therapies are useful in acute situations where effective, but transient, inhibition of leukocyte-mediated inflammation is desirable. In addition, treatment of chronic disorders may be attained by sustained administration of agents, for example, by subcutaneous or oral administration.

10 An inflammatory response may cause damage to the host if unchecked, because leukocytes release many toxic molecules that can damage normal tissues. These molecules include proteolytic enzymes and free radicals. Examples of pathological situations in which leukocytes can cause tissue damage include injury from ischemia and reperfusion, 15 bacterial sepsis and disseminated intravascular coagulation, adult respiratory distress syndrome, tumor metastasis, rheumatoid arthritis and atherosclerosis.

Reperfusion injury is a major problem in clinical cardiology. Therapeutic agents that reduce leukocyte adherence in ischemic 20 myocardium can significantly enhance the therapeutic efficacy of thrombolytic agents. Thrombolytic therapy with agents such as tissue plasminogen activator or streptokinase can relieve coronary artery obstruction in many patients with severe myocardial ischemia prior to irreversible myocardial cell death. However, many such patients still 25 suffer myocardial neurosis despite restoration of blood flow. This "reperfusion injury" is known to be associated with adherence of leukocytes to vascular endothelium in the ischemic zone, presumably in part because of activation of platelets and endothelium by thrombin and cytokines that makes them adhesive for leukocytes (Romson et al., 30 *Circulation* 67: 1016-1023, 1983). These adherent leukocytes can migrate through the endothelium and destroy ischemic myocardium just as it is being rescued by restoration of blood flow.

-22-

There are a number of other common clinical disorders in which ischemia and reperfusion results in organ injury mediated by adherence of leukocytes to vascular surfaces, including strokes; mesenteric and peripheral vascular disease; organ transplantation; and circulatory shock

5 (in this case many organs might be damaged following restoration of blood flow).

Bacterial sepsis and disseminated intravascular coagulation often exist concurrently in critically ill patients. They are associated with generation of thrombin, cytokines, and other inflammatory mediators, 10 activation of platelets and endothelium, and adherence of leukocytes and aggregation of platelets throughout the vascular system. Leukocyte-dependent organ damage is an important feature of these conditions.

Adult respiratory distress syndrome is a devastating pulmonary disorder occurring in patients with sepsis or following trauma, which is 15 associated with widespread adherence and aggregation of leukocytes in the pulmonary circulation. This leads to extravasation of large amounts of plasma into the lungs and destruction of lung tissue, both mediated in large part by leukocyte products.

Two related pulmonary disorders that are often fatal are in 20 immunosuppressed patients undergoing allogeneic bone marrow transplantation and in cancer patients suffering from complications that arise from generalized vascular leakage resulting from treatment with interleukin-2 treated LAK cells (lymphokine-activated lymphocytes).

LAK cells are known to adhere to vascular walls and release products that 25 are presumably toxic to endothelium. Although the mechanism by which LAK cells adhere to endothelium is not known, such cells could potentially release molecules that activate endothelium and then bind to endothelium by mechanisms similar to those operative in neutrophils.

Tumor cells from many malignancies (including carcinomas, 30 lymphomas, and sarcomas) can metastasize to distant sites through the vasculature. The mechanisms for adhesion of tumor cells to endothelium and their subsequent migration are not well understood, but may be

-23-

similar to those of leukocytes in at least some cases. Specifically, certain carcinoma cells have been demonstrated to bind to both E-selectin, as reported by Rice and Bevilacqua. *Science* **246**:1303-1306 (1991), and P-selectin, as reported by Aruffo, et al., *Proc. Natl. Acad. Sci. USA* **89**:2292-2296 (1992), Stone and Wagner, *I. Clin. Invst.* **92**:804-813 (1992). The association of platelets with metastasizing tumor cells has been well described, suggesting a role for platelets in the spread of some cancers. Since P-selectin is expressed on activated platelets, it is believed to be involved in association of platelets with at least some malignant tumors.

Platelet-leukocyte interactions are believed to be important in atherosclerosis. Platelets might have a role in recruitment of monocytes into atherosclerotic plaques; the accumulation of monocytes is known to be one of the earliest detectable events during atherogenesis. Rupture of a fully developed plaque may not only lead to platelet deposition and activation and the promotion of thrombus formation, but also the early recruitment of neutrophils to an area of ischemia. P-selectin is also inappropriately expressed on the surface of endothelial cells overlying atheromas, where it may mediate the initial attachment of monocytes that then migrate into the atheroma (Johnson-Tidey, et al., *Am. J. Path.* **144**:952-961, 1994).

Another area of potential application is in the treatment of rheumatoid arthritis and other autoimmune or inflammatory disorders.

In these clinical applications, the peptides or isolated O-glycan, or fragments thereof, described herein, can be administered to block selectin-dependent interactions by binding competitively to P-selectin expressed on activated cells. In addition, antibodies to the protein and/or carbohydrate components of the ligand, or fragments thereof, can be administered. The antibodies are preferably of human origin or modified to delete those portions most likely to cause an immunogenic reaction.

Peptides as described herein can also be administered as a pharmaceutically acceptable acid- or base- addition salt, formed by

reaction with inorganic acids such as hydrochloric acid, hydrobromic acid, perchloric acid, nitric acid, thiocyanic acid, sulfuric acid, and phosphoric acid, and organic acids such as formic acid, acetic acid, propionic acid, glycolic acid, lactic acid, pyruvic acid, oxalic acid, 5 malonic acid, succinic acid, maleic acid, and fumaric acid, or by reaction with an inorganic base such as sodium hydroxide, ammonium hydroxide, potassium hydroxide, and organic bases such as mono-, di-, trialkyl and aryl amines and substituted ethanolamines.

Carbohydrate components (the O-glycan structures or components 10 thereof) of the ligand or the antibodies, in an appropriate pharmaceutical carrier, are preferably administered intravenously where immediate relief is required. The carbohydrate(s) can also be administered intramuscularly, intraperitoneally, subcutaneously, orally, as the carbohydrate, conjugated to a carrier molecule, or in a drug delivery 15 device. The carbohydrate can be modified chemically to increase its *in vivo* half-life.

The carbohydrate can be isolated from cells expressing the carbohydrate, either naturally or as a result of genetic engineering as described in the transfected mammalian cell examples, or, preferably, by 20 synthetic means. These methods are known to those skilled in the art. In addition, a large number of additional glycosyltransferases have been cloned (J.C. Paulson and K.J. Colley, *J. Biol. Chem.* **264**:17615-17618, 1989). Accordingly, workers skilled in the art can use a combination of synthetic chemistry and enzymatic synthesis to make pharmaceuticals or 25 diagnostic reagents.

Carbohydrates that are biologically active are those which inhibit binding of leukocytes to P-selectin. Suitable pharmaceutical vehicles for administration to a patient are known to those skilled in the art. For parenteral administration, the carbohydrate will usually be dissolved or 30 suspended in sterile water or saline. For enteral administration, the carbohydrate will be incorporated into an inert carrier in tablet, liquid, or capsular form. Suitable carriers may be starches or sugars and include

-25-

lubricants, flavorings, binders, and other materials of the same nature.

The carbohydrate can also be administered locally at a wound or inflammatory site by topical application of a solution or cream.

Alternatively, the carbohydrate may be administered in, on or as part of, liposomes or microspheres (or microparticles). Methods for preparing liposomes and microspheres for administration to a patient are known to those skilled in the art. U.S. Patent No. 4,789,734 describe methods for encapsulating biological materials in liposomes. Essentially, the material is dissolved in an aqueous solution, the appropriate phospholipids and lipids added, along with surfactants if required, and the material dialyzed or sonicated, as necessary. A good review of known methods is by G. Gregoriadis, Chapter 14. "Liposomes", Drug Carriers in Biology and Medicine pp. 287-341 (Academic Press, 1979).

Microspheres formed of polymers or proteins are well known to those skilled in the art, and can be tailored for passage through the gastrointestinal tract directly into the bloodstream. Alternatively, the carbohydrate can be incorporated and the microspheres, or composite of microspheres, implanted for slow release over a period of time, ranging from days to months. See, for example, U.S. Patent No. 4,906,474, 4,925,673, and 3,625,214.

The carbohydrates should be active when administered parenterally in amounts above about 1 $\mu\text{g}/\text{kg}$ of body weight. For treatment of most inflammatory disorders, the dosage range will be between 0.1 to 30 mg/kg of body weight. A dosage of 70 mg/kg may be required for some of the carbohydrates characterized in the examples.

The criteria for assessing response to therapeutic modalities employing antibodies or carbohydrate is dictated by the specific condition and will generally follow standard medical practices. For example, the criteria for the effective dosage to prevent extension of myocardial infarction would be determined by one skilled in the art by looking at marker enzymes of myocardial necrosis in the plasma, by monitoring the electrocardiogram, vital signs, and clinical response. For treatment of

acute respiratory distress syndrome, one would examine improvements in arterial oxygen, resolution of pulmonary infiltrates, and clinical improvement as measured by lessened dyspnea and tachypnea. For treatment of patients in shock (low blood pressure), the effective dosage 5 would be based on the clinical response and specific measurements of function of vital organs such as the liver and kidney following restoration of blood pressure. Neurologic function would be monitored in patients with stroke. Specific tests are used to monitor the functioning of transplanted organs; for example, serum creatinine, urine flow, and serum 10 electrolytes in patients undergoing kidney transplantation.

The present invention will be further understood by reference to the following non-limiting examples.

Example 1: Determination of the structure and function of the oligosaccharides attached to PSGL-1.

15 The effects of various glycosidases on the structure and function of PSGL-1 from human neutrophils was determined. This molecule, unlike recombinant PSGL-1 expressed in COS cells with FTIII, is largely resistant to treatment with endo- α -N-acetylgalactosaminidase, suggesting that it has few simple core 1 Gal β 1-3GalNAc disaccharides linked to serine or threonine. Human neutrophil PSGL-1 displays sLe x and its 20 nonsialylated counterpart, Le x , primarily on *O*-linked poly-*N*-acetyllactosamine. These data suggest that PSGL-1 on human neutrophils presents an array of sialylated, fucosylated, *O*-linked poly-*N*-acetyllactosamine that P-selectin preferentially recognizes as compared to 25 recombinant PSGL-1 expressed in COS cells with FTIII.

30 The effects of glycosidases on purified, radioiodinated PSGL-1 from human neutrophils were examined to further characterize the structure and function of the attached oligosaccharides. PSGL-1 was found to have poly-*N*-acetyllactosamine, only some of which could be removed with endo- β -galactosidase. The majority of the Le x and sLe x structures were on endo- β -galactosidase-sensitive chains. Peptide: *N*-glycosidase F (PNGaseF) treatment removed at least two of the three

-27-

possible *N*-linked oligosaccharides from PSGL-1. Expression of Le^x and sLe^x was not detectably altered by PNGaseF digestion, indicating that these structures were primarily on *O*-linked poly-*N*-acetyllactosamine. Endo- β -galactosidase-treated PSGL-1 retained the ability to bind to P-selectin, suggesting that some of the oligosaccharides recognized by P-selectin were either on enzyme-resistant poly-*N*-acetyllactosamine or on chains which lack poly-*N*-acetyllactosamine. PNGaseF treatment did not affect the ability of PSGL-1 to bind to P-selectin, demonstrating that the oligosaccharides required for P-selectin recognition are *O*-linked. These data demonstrate that PSGL-1 from human neutrophils displays complex, sialylated, and fucosylated *O*-linked poly-*N*-acetyllactosamine that promote high affinity binding to P-selectin.

Materials and Methods

The abbreviations used are: CHO, Chinese hamster ovary; Con A, Concanavalin A; HSA, human serum albumin; Le^x , Lewis x; PBS, phosphate-buffered saline; PAGE, polyacrylamide gel electrophoresis; PNGaseF, peptide:*N*-glycosidase F; PSGL-1, P-Selectin Glycoprotein Ligand 1; sLe^x , sialyl Lewis x; tES, truncated soluble E-selectin; tPS, truncated soluble P-selectin; WGA, wheat germ agglutinin.

Materials--Triton X-100, Brij-58, and sialidase from *Arthrobacter ureafaciens* (75 U/mg, EC 3.2.1.18) were obtained from CalBioChem-Behring Corp. (La Jolla, CA). Iodobeads®, 3M Emphaze® Biosupport Medium, and Protein A Sepharose CL4B were from Pierce Chemical Co. (Rockford, IL). Endo- β -galactosidase from *Escherichia freundii* (5 U/mg, EC 3.2.1.103) was obtained from V-Labs, Inc. (Covington, LA). *Streptomyces* sp.142 α 1,3/4-L-fucosidase (EC 3.2.1.51) was obtained from PanVera Corp. (Madison, WI) and β -*N*-acetylglucosaminidase from Jack Bean (EC 3.2.1.30, 53 U/mg) was from Sigma Chemical Co. (St. Louis, MO). Recombinant peptide:*N*-glycosidase F (EC 3.2.2.18, 25,000 U/mg, PNGase F), and endo- α -*N*-acetylgalactosaminidase from *Diplococcus pneumoniae* (EC 3.2.1.97, O-glycanase®) were purchased from Genzyme Corp. (Cambridge, MA). Concanavalin A Sepharose

-28-

(Con A, 10 mg/ml resin) was from Pharmacia/LKB (Uppsala, Sweden). Wheat germ agglutinin (WGA) agarose (7.6 mg/ml resin) and tomato lectin were from Vector Laboratories, Inc. (Burlingame, VT). Tomato lectin was coupled to cyanogen bromide-activated Sepharose to a density of 2 mg/ml resin in the presence of 7.5 mg/ml chitotriose.

5 *Antibodies*--The anti-human P-selectin mAbs S12 and G1 were prepared and characterized as previously described by McEver and Martin (1984), Geng, et al., (1990). The anti-human E-selectin mAbs CL2 and CL37 (Mulligan, et al., *J. Clin. Invest.* **88**, 1396-1406 (1991)) were 10 provided by C. Wayne Smith (Baylor College of Medicine, Houston, TX). LeuM1 (CD15, IgM) was purchased from Becton-Dickinson & Co. (San Jose, CA) . The CSLEX-1 hybridoma (HB 8580) was purchased from the American Type Culture Collection and the IgM mAb was purified from ascites fluid by boric acid precipitation and gel filtration on 15 Sepharose CL4B in PBS, pH 7.4 as described by Shattil, et al., *J. Biol. Chem.* **260**, 11107-11114 (1985). Goat anti-mouse μ -chain specific IgG and MOPC104E (IgM) were purchased from Cappel-Organon Technika (Durham, NC).

20 *Peptide Antisera Production and Immunoprecipitations.* Peptides corresponding to residues 42-56 (QATEYEYLDYDFLPEC) (Sequence ID No. 1) and 354-376 (CISSLLPDGGEGPSATANGGLSK) (Sequence ID No. 1) of the cDNA-derived amino acid sequence of PSGL-1 were synthesized on an Applied Biosystems Model 431 peptide synthesizer. Peptide 42-56 was coupled to maleimide-activated keyhole limpet 25 hemocyanin (Pierce Chemical Co.) through the added cysteine and injected into New Zealand white rabbits (Montana State University Animal Resources Center). Immune sera were collected and used to immunoprecipitate 125 I-PSGL-1. Protein A beads (25 μ l, 50% suspension) were preincubated with 1 μ l of a 1:4 dilution of either normal or immune 30 rabbit serum for 90 min at 37°C. The beads were washed once with 0.1 M NaCl, 20 mM MOPS, pH 7.5, 0.1% Triton X-100 (MBS/0.1% Triton). Purified 125 I-PSGL-1 or radioiodinated-WGA eluate was then

-29-

incubated with the beads in the presence or absence of 0.2 mg/ml of either the immunizing peptide or PSGL-1 peptide 354-376 which served as a negative control. After a 90 min incubation at 37°C the beads were washed five times with MBS/0.1% Triton and the eluted by boiling for 5 min with SDS sample buffer (Laemmli, *Nature* 227, 680-685 (1970)). Immunoprecipitates were analyzed by SDS-PAGE, followed by autoradiography.

Generation of CHO D⁻ cells Stably Expressing a Soluble Form of E-selectin. An expression plasmid (ELAM-1:BBG 57, obtained from 10 British BioTechnology Products, Ltd.) containing the full-length cDNA for human E-selectin was used as a template in the polymerase chain reaction to modify the 5' and 3' ends of E-selectin for high level production of recombinant soluble E-selectin in CHO D⁻ cells. The oligonucleotide pair (5' GTC CCT CTA GAC CAC CAT GAT TGC 15 TTC ACA GTT T + 5' TCC CGG TCG ACT TAG GGA GCT TCA CAG GT) (Sequence Listing ID Nos. 4 and 5, respectively) used in the polymerase chain reaction was designed to introduce an *Xba* I site and a perfect Kozak sequence (Kozak, *Nucleic Acids Res.* 9, 5233-5252 (1981)) preceding the E-selectin initiator codon, and to introduce a *Sal* I site 20 following the stop codon introduced after amino acid 550 located between the sixth consensus repeat and the transmembrane domain of full-length E-selectin. DNA sequence analysis verified the changes generated by the polymerase chain reaction as well as the integrity of the remainder of the E-selectin cDNA. The truncated E-selectin (tES) gene was then 25 introduced as an *Xba* I - *Sal* I fragment into the mammalian expression vector pDSR α 2 (European patent application A20398753) that was modified to include unique *Xba* I and *Sal* I restriction sites. The tES expression vector was transfected into a CHO cell line deficient in dihydrofolate reductase (DHFR-minus CHO) (Urlaub, et al., *Proc. Natl. 30 Acad. Sci. USA* 77, 4216-4220 (1980)), and transfecants were selected in medium lacking hypoxanthine and thymidine (Bourdrel, et al., *Protein Exp. Purif.* 4, 130-140 (1993)). An RNase protection assay was used to

-30-

screen for transfectants that had high levels of E-selectin-specific mRNA (Turner, et al., *Blood* 80, 374-381 (1992)).

Protein Purification. PSGL-1 was purified from human neutrophil membranes prepared as described by Moore, et al., (1992), 5 with the following modifications. Neutrophil membranes extracted with 5% Triton X-100 in 0.1 M NaCl, 20 mM MOPS, pH 7.5, 0.02% NaN₃, were applied to WGA-agarose, and bound proteins were eluted with 0.5 M N-acetylglucosamine (WGA eluate). After extensive dialysis against 0.1 M NaCl, 20 mM MOPS, pH 7.5, 0.02% NaN₃, 0.02% Brij-58, 2 10 mM CaCl₂, 2 mM MgCl₂ (equilibration buffer), the WGA eluate was loaded on a truncated P-selectin (tPS)-Emphaze column (10 mg lectin/ml resin). The column was extensively washed with equilibration buffer and eluted with 5 mM EDTA. Ligand-containing fractions were pooled and loaded on a Mono Q PC 1.6/5 column equilibrated with 0.1 M NaCl, 20 15 mM MOPS, pH 7.5, 2 mM EDTA, 0.02% NaN₃, 0.02% Brij-58 using a SMART® Micro Separation System (Pharmacia/LKB). The column was developed with a 2-ml linear gradient (0.1-1.0 M NaCl) at 50 µml/min. Aliquots of the fractions were iodinated with Na¹²⁵I using Iodobeads® 20 according to the instructions of the manufacturer. The purity of the fractions was assessed by SDS-PAGE and autoradiography of the iodinated fractions.

A recombinant soluble form of P-selectin (tPS) truncated after the ninth consensus repeat was immunoaffinity purified from conditioned medium of permanently transfected 293 cells as described by Ushiyama, et al., (1993). As determined by sedimentation velocity and equilibrium analysis, tPS is an asymmetric monomer with a molecular weight of 25 103,600 Da. The E₂₈₀^{1%} for tPS is 12.3.

Recombinant soluble truncated E-selectin (tES) was immunoaffinity purified from conditioned medium of CHO D⁻ cells stably 30 secreting tES. Cells were cultured in equal parts of Dulbecco's modified Eagle's medium (DMEM) and Ham's F-12 nutrient mixture, supplemented with 10% fetal bovine serum, 10 U/ml penicillin, and 10

mg/ml streptomycin. At confluence, the cells were shifted to serum-free medium, and conditioned medium was harvested every 7 days. Conditioned medium was clarified, sterile-filtered, using two Sartorius filters (10 μ m and 0.2 μ m) connected in series, and then concentrated 50-fold (Filtron Maximate, 10 kDa MW_{CO}). The anti-E-selectin mAb H18/7 (Bevilacqua, et al., *Proc. Natl. Acad. Sci. USA* **84**, 9238-9242 (1987)) was coupled to cyanogen bromide-activated Sepharose to a density of 6 mg/ml resin. The H18/7-Sepharose column was equilibrated at 4°C with PBS, pH 7.5, containing 1 mM CaCl₂ and 1 mM MgCl₂, and 500 ml of concentrated medium was loaded on the column (V_T = 200 ml) at a flow rate of 10 cm/h. The column was eluted with 0.1 M glycine, pH 2.7 and the fractions were rapidly neutralized by addition of Tris-HCl, pH 7.5 to a final concentration of 0.1 M. The eluate was dialyzed against 25 mM sodium phosphate, pH 7.5, and loaded on a Q-High Performance anion-exchange column (V_T = 200 ml, Pharmacia) equilibrated in the same buffer. The column was developed with a linear gradient of NaCl (0.17 - 0.22 M). Purity was assessed by SDS-PAGE and by N-terminal sequence analysis. Protein concentration was determined using an E₂₈₀^{1%} = 12.5 that was calculated using the method of Gill, and von Hippel, *Anal. Biochem.* **182**, 319-326 (1989). The yield was 250 mg of tES per approximately 500 ml of concentrated conditioned medium.

Sedimentation Equilibrium Analysis of Soluble tES. Sedimentation equilibrium experiments were conducted using a 4-hole titanium rotor, short column kel-f centerpieces, and sapphire windows in a Beckman Optima XLA analytical ultracentrifuge equipped with on-line Rayleigh interference optics. Data were acquired at rotor speeds of 10,000, 15,000, 20,000, and 25,000 rpm at 20°C using a television-based camera and an on-line acquisition and analysis system (Laue, T. M. (1981) *Ph.D. Dissertation*, University of Connecticut, Storres, CT; Laue, et al., (1992) in *Analytical Ultracentrifugation in Biochemistry and Polymer Science* (Harding, S., Rowe, A., and Horton, J. C., eds) pp. 90-125, Royal Society of Chemistry, London). Samples were loaded at concentrations of

-32-

approximately 1.0, 0.5, 0.25, and 0.13 mg/ml in PBS, pH 7.4. Data were collected at intervals after the estimated time to equilibrium and tested for equilibrium by subtracting successive scans (Yphantis, D. A. (1964) *Biochemistry* 3, 297-317)). The data within the optical window 5 were selected using the program REEDIT (provided by David Yphantis). Data were analyzed to estimate the monomer molecular weight of tES using the NONLIN program (Yphantis, D. A. (1964) *Biochemistry* 3, 297-317). Three or more channels of sedimentation equilibrium data obtained at different loading concentration of tES, radial positions, and 10 angular velocities were fit by simultaneous nonlinear least-squares analysis. For molecular weight calculations, the partial specific volume of tES was estimated to be 0.688 ml/g with an inaccuracy of \pm 0.03 ml/g to account for uncertainty in the carbohydrate composition of tES (Laue, et al. (1992)).

15 *Sedimentation Velocity Analysis of Soluble tES.* Sedimentation velocity experiments were conducted in the Beckman Optima XLA analytical ultracentrifuge using Rayleigh interference optics. Experiments were conducted at 60,000 rpm and 20°C, using a 4-hole titanium rotor, two-channel, charcoal-filled, epon centerpieces, and sapphire windows. 20 Sample was loaded at a concentration of 0.5 mg/ml in PBS, pH 7.4. Sedimentation coefficients were determined from the time derivative of the concentration profile as described by. Stafford, W. (1992) *Anal. Biochem.* 203, 295-301.

25 *Protein Iodinations.* The mAbs G1 and CL2 (200 μ g) were iodinated with 400 μ Ci carrier-free Na¹²⁵I (ICN Biomedical, Inc., Costa Mesa, CA) using Iodobeads®. Free ¹²⁵I was removed by gel filtration through a Sephadex G-25 column (PD-10, Pharmacia) equilibrated in 0.1 M NaCl, 20 mM MOPS, pH 7.5, 0.1% Triton X-100. The labeled antibodies were then centrifuged for 30 min at 90,000 $\times g$ in a TL-100 30 ultracentrifuge (Beckman, Palo Alto, CA). The concentration of labeled proteins was determined with a Micro BCA protein assay kit (Pierce)

-33-

using the respective unlabeled proteins as standards. The specific activity of the labeled antibodies ranged from 0.5-1.0 μ Ci/ μ g protein.

PSGL-1 (approximately 0.5 μ g) or WGA eluate (approximately 100 μ g) was iodinated with 400 μ Ci Na¹²⁵I and desalted as described above. Radioabeled PSGL-1 was concentrated to approximately 200 μ l using a Centicon-30® device (Pharmacia/LKB), and gel filtered on a Superose 6 PC 3.2/30 column equilibrated with 0.15 M NaCl, 20 mM MOPS, pH 7.5, 0.02% NaN₃, 0.1% Brij-58 using a SMART® Micro Separation System (Pharmacia/LKB). The specific activity of the radiolabeled ligand ranged from 14-30 μ Ci/ μ g protein. Labeled proteins were routinely greater than 95% trichloroacetic acid precipitable.

Binding of ¹²⁵I-PSGL-1 to Immobilized tPS or tES. tPS or tES (50 μ l, 10 μ g/ml) in HBSS, 0.02% NaN₃ (HBSS/Az) were incubated overnight at 4°C in microtiter plates (Immulon I Removawell® Strips, Dynatech Laboratories, Inc., Chantilly, VA). After three washes with HBSS/Az the wells were blocked with HBSS/Az/1% human serum albumin (HSA) for 2 h at 22°C. The wells were then washed once with HBSS/Az and ¹²⁵I-PSGL-1 diluted in HBSS/Az/0.1% HSA was added (50 μ l, 5,000-10,000 cpm/well). After 1 h at 22°C the wells were rapidly washed five times with HBSS/Az using a 12-channel plate washer (Nunc-Immuno Wash 12, Nunc Inc., Naperville, IL). The individual wells were then counted in a γ counter. All assays were performed in quadruplicate. In certain assays binding of ¹²⁵I-PSGL-1 to immobilized tPS was measured in the presence of increasing concentrations of fluid-phase tPS or tES.

Binding of ¹²⁵I-PSGL-1 to Immobilized Anti-Carbohydrate Antibodies. Goat anti-mouse μ -chain specific IgG (50 μ l, 10 μ g/ml) in HBSS/Az was incubated overnight at 4°C in Immulon I Removawell® strips. The wells were washed and blocked as described above. Then, either CSLEX-1, LeuM1, or MOPC104E was added (50 μ l, 5 μ g/ml). After incubation for 1 h at 22°C, the wells were washed three times with HBSS/Az, and ¹²⁵I-PSGL-1 in HBSS/Az/0.1% HSA (50 μ l, 5,000-10,000 cpm/well) was added. After 1 h the wells were washed as described

-34-

above, and individual wells were counted in a γ counter. All assays were performed in quadruplicate.

Enzyme Digestion of ^{125}I -PSGL-1. Digestions with sialidase (0.2 U/ml, 15 h), endo- β -galactosidase (2 U/ml, 15 h), and α 1,3/4-L-fucosidase (0.32 mU/ml, 15 h) were performed in 0.1 M NaCl, 50 mM sodium acetate, pH 5.5. Digestions with PNGaseF (40 U/ml, 15 h) were done in 0.1 M sodium phosphate, pH 8.6. PSGL-1 was generally not pretreated with SDS because PNGase F had similar effects with or without SDS pretreatment. Sequential digestions with sialidase (0.2 U/ml, 2 h) followed by endo- α -N-acetylgalactosaminidase (0.1 U/ml, 15 h) were performed in 0.1 M NaCl, 50 mM sodium acetate, pH 6.0. All glycosidase digestions were performed at 37°C in the presence of 20 μM leupeptin, 30 μM antipain, 1 mM benzamidine, and 0.02% NaN_3 .

Lectin Affinity Chromatography. ^{125}I -PSGL-1 was applied to a 15 Concanavalin A (Con A) Sepharose column (V_T = 1 ml) equilibrated in 0.1 M NaCl, 20 mM Tris, pH 7.5, 2 mM CaCl₂, 0.1% Brij-58, 0.02% NaN_3 . The column was washed with five column volumes of equilibration buffer and eluted with 0.5 M α -methylmannoside in equilibration buffer preheated to 65°C. Tomato lectin chromatography 20 was similarly performed except that the column (V_T = 1 ml) was equilibrated with 0.1 M NaCl, 20 mM MOPS, pH 7.5, 0.1% Brij-58, 0.02% NaN_3 , and step eluted with 20 mg/ml chitotriose in equilibration buffer. Greater than 85% of the counts loaded on the lectin columns were recovered.

Site Density Determinations. mAbs to P-selectin (G1) or E-selectin (CL2) that inhibit interactions with myeloid cells were used for site density measurements. Wells were coated with tPS or tES (50 μl , 10 $\mu\text{g}/\text{ml}$) and blocked as described above. Saturating concentrations of ^{125}I -labeled mAb (2 $\mu\text{g}/\text{ml}$) in HBSS/Az/0.1% HSA were added to the wells 30 and incubated for 1 h at 22°C. The wells were washed 5 times and then counted in a γ counter. Specific binding was defined as cpm bound to selectin-coated wells minus cpm bound to HSA-coated wells. Site

densities were calculated assuming monovalent binding of antibody at saturation. All assays were performed in quadruplicate.

Neutrophil Adhesion Assay. Human neutrophils were isolated from heparinized blood from consenting volunteer donors by dextran sedimentation, hypotonic lysis, and Ficoll-Hypaque density gradient centrifugation as described by Zimmerman, et al. (1985) *J. Clin. Invest.* **76**, 2235-2246. Quantitation of neutrophil adhesion to tPS or tES (50 μ l, 10 μ g/ml), immobilized on Immulon I Removawell® microtiter plates, was performed as described by Geng, et al., (1990).

10 Results

An Antiserum to a PSGL-1 Peptide Recognizes the P-selectin Glycoprotein Ligand from Human Neutrophils. To determine whether the polypeptide core of the sialomucin ligand for P-selectin characterized in human neutrophils and HL-60 cells was immunologically related to PSGL-1, a glycoprotein ligand for P-selectin identified by expression cloning from a HL-60 cDNA library, antiserum prepared in rabbits against a peptide corresponding to residues 42-56 of the cDNA-derived amino acid sequence of PSGL-1, was tested to see if this antiserum could recognize the glycoprotein ligand that we isolated by P-selectin affinity chromatography. The radiochemical purity of the 125 I-P-selectin glycoprotein ligand used in this study showed that it exhibited a relative molecular mass of approximately 250,000 under nonreducing conditions and approximately 120,000 under reducing conditions. Thus, the radioiodinated glycoprotein contained two disulfide-linked subunits of M_r 120,000, consistent with previous results using protein blotting or metabolic labeling. As noted previously, a small portion of the dimeric glycoprotein was resistant to reduction under the conditions used. The immune serum, but not normal rabbit serum, precipitated the purified 125 I-P-selectin ligand. Precipitation by the antiserum was specific for the 42-56 peptide sequence, because it was completely inhibited by inclusion of this peptide, but not by a control peptide corresponding to amino acids 354-376 of PSGL-1. The specificity of the antiserum was further

-36-

demonstrated by its selective immunoprecipitation of the ligand from a crude mixture of radiolabeled neutrophil membrane proteins that were eluted from a WGA column.

Two tryptic peptides were derived from the P-selectin glycoprotein ligand purified from human neutrophils. The sequences of these peptides, His-Met-Tyr-Pro-Val-Arg and Pro-Gly-Lys-Thr-Pro-Glu-Pro, are identical to amino acids 340-345 and 380-386 in the cDNA-derived sequence of PSGL-1 (Sequence ID No. 1). These data and the results from the immunoprecipitation experiments establish the identity of the polypeptide backbones of the P-selectin ligands studied by both groups.

Purified ¹²⁵I-PSGL-1 Binds Specifically to P-selectin. ¹²⁵I-PSGL-1 isolated from human neutrophils bound in a time-dependent fashion to tPS immobilized on microtiter wells. Binding increased as a function of the amount of tPS coated on the wells. Binding was Ca^{2+} -dependent and was abolished by G1, a mAb to P-selectin that inhibits leukocyte adhesion to P-selectin, but not by S12, a mAb that does not block adhesion. In addition, $84.6 \pm 2.4\%$ (mean \pm SD, n = 4) of the radiolabeled glycoprotein rebound to a column of recombinant, soluble P-selectin (tPS) and was eluted with buffer containing EDTA. This result demonstrates that the function of ¹²⁵I-PSGL-1 was not substantially altered by iodination.

PSGL-1 Contains Substituted Poly-N-acetyllactosamine. PSGL-1 from human myeloid cells contains clustered, sialylated O-linked oligosaccharides that are released from the polypeptide backbone by β -elimination (Norgard, *et al.* 1993). To determine whether these O-linked glycans had simple structures, ¹²⁵I-PSGL-1 was treated with sialidase, then with endo- α -N-acetylgalactosaminidase, an enzyme that releases non-sialylated Gal β 1-3GalNAc core 1 disaccharides, but not more complex O-linked glycans, from serine or threonine residues (Umemoto, *et al.*, (1977) *J. Biol. Chem.* **252**, 8609-8614). Treatment with sialidase slowed the electrophoretic mobility of PSGL-1, confirming previous results.

Subsequent addition of endo- β -N-acetylgalactosaminidase increased the mobility of PSGL-1 only slightly more than sialidase-treated PSGL-1, suggesting that very few of the O-linked oligosaccharides were simple structures that were susceptible to this enzyme.

PSGL-1 is fucosylated, because it contains the sialylated, fucosylated tetrasaccharide antigen, sLe^x (Norgard, et al., (1993)). Treatment of the glycoprotein with α 1,3/4 fucosidase, which removes fucose attached to a penultimate GlcNAc, had no effect on electrophoretic mobility, although it did remove fucose residues that were part of the Le^x epitopes on the molecule. The ligand was then treated with endo- β -galactosidase, which hydrolyzes internal β 1-4 linkages between galactose and N-acetylglucosamine (Gal β 1-4GlcNAc) in extended unbranched poly-N-acetyllactosamine chains (Scudder, et al., (1983) *Biochem. J.* **213**, 485-494; Scudder, et al., (1984) *J. Biol. Chem.* **259**, 6586-6592). This enzyme slightly accelerated the mobility and lessened the electrophoretic heterogeneity of PSGL-1, suggesting that PSGL-1 contained some poly-N-acetyllactosamine.

To confirm the presence of poly-N-acetyllactosamine on PSGL-1, the ability of ¹²⁵I-PSGL-1 to bind to a column containing immobilized tomato lectin, which avidly binds to poly-N-acetyllactosamine, was tested. A single poly-N-acetyllactosamine chain is sufficient to confer binding of a glycoprotein to tomato lectin in this system (Merkle, et al. (1987) *J. Biol. Chem.* **262**, 8179-8189). Greater than 95% of the PSGL-1 bound to the column, indicating that essentially every molecule contained at least one poly-N-acetyllactosamine (Table 3). Treatment of PSGL-1 with endo- β -galactosidase modestly reduced to 84% the amount of material binding to the column. This suggests that most, if not all, of the poly-N-acetyllactosamine could be removed from a portion of the PSGL-1 molecules by endo- β -galactosidase. Because substitutions of poly-N-acetyllactosamine with sialic acid and/or fucose inhibit the efficiency of this enzyme, PSGL-1 was pretreated with sialidase and fucosidase before addition of endo- β -galactosidase. Treatment with sialidase and fucosidase

had no effect on binding of PSGL-1 to the column, consistent with the fact that substitutions with sialic acid or fucose do not affect binding of poly-*N*-acetyllactosamine to tomato lectin. However, subsequent addition of endo- β -galactosidase reduced to 69% the portion of the glycoprotein 5 binding to the column. Because tomato lectin binds weakly to terminal GlcNAc residues that are exposed by endo- β -galactosidase treatment (Cummings, (1994) *Methods Enzymol.* **230**, 66-86), β -*N*-acetylglucosaminidase was added to remove these terminal residues. Only 53% of this material bound to tomato lectin. These data clearly 10 demonstrated that PSGL-1 contained poly-*N*-acetyllactosamine. Furthermore, at least some of these chains were resistant to endo- β -galactosidase because of substitutions with sialic acid and/or fucose. The inability of sialidase, fucosidase, endo- β -galactosidase, and β -*N*-acetylglucosaminidase to eliminate binding of all PSGL-1 molecules to 15 tomato lectin may reflect enzyme resistance because of the clustered nature of *O*-linked glycans, the presence of internal fucose residues, and/or the presence of additional substitutions such as sulfate on the poly-*N*-acetyllactosamine.

Table 3. Effects of Glycosidases on Binding of ^{125}I -PSGL-1 to a Tomato Lectin Affinity Column

Enzyme Treatment	Percentage Bound
Sham	> 95
5 Endo- β -galactosidase	84
Sialidase + Fucosidase	> 95
Sialidase + Fucosidase + Endo- β -galactosidase	69
10 Sialidase + Fucosidase + Endo- β -galactosidase + β -N-acetylglucosaminidase	53
Peptide:N-glycosidase F	> 95

15 ^{125}I -PSGL-1 was treated with the indicated glycosidase(s) and applied to a column of immobilized Tomato lectin ($V_T = 1 \text{ ml}$, 2 mg lectin/ml resin). After washing, the column was eluted with 20 mg/ml of chitotriose. The percentage of PSGL-1 bound was defined as the radioactivity eluted by chitotriose divided by the sum of the radioactivity in the column flowthrough, wash, and eluate.

20 *PSGL-1 Contains a Limited Number of Heterogeneous N-linked Oligosaccharides.* PNGaseF treatment slightly increased the electrophoretic mobility of [^{125}I]PSGL-1, consistent with previous studies indicating that all molecules contained a limited number of N-linked glycans sensitive to this enzyme. The cDNA-derived sequence of PSGL-1 25 indicates that the molecule contains only three potential sites for attachment of N-linked oligosaccharides (Sako, et al., (1993)). To explore possible heterogeneity of the N-linked structures, sham-treated and PNGaseF-treated [^{125}I]PSGL-1 was applied to a column of

Concanavalin A (Con A), a plant lectin that binds avidly to high mannose *N*-linked glycans, less avidly to complex biantennary and hybrid *N*-linked oligosaccharides, and very weakly to complex triantennary and tetraantennary *N*-linked chains (Ogata, et al. (1975) *J. Biochem. (Tokyo)* **78**, 687-696). 41.9 \pm 4.2% of the sham-treated ligand bound to Con A, whereas only 3.7 \pm 1.0% of the PNGaseF-treated material bound (mean \pm SD, n = 4). Greater than 85% of the counts loaded on the column were recovered. This result demonstrated that PNGaseF removed *N*-linked glycans recognized by Con A from PSGL-1.

10 The fraction of PSGL-1 not bound by Con A migrated slightly slower than the fraction bound by Con A. To confirm that the Con A-unbound material also contained *N*-linked oligosaccharides, both the bound and unbound fractions were treated with PNGaseF. The enzyme decreased the apparent molecular weights of the Con A-unbound and Con A-bound fractions by 14.5 kDa and 13.7 kDa, respectively. Thus, both fractions had PNGaseF-sensitive *N*-linked glycans. These data suggested that the Con A-unbound fraction contained triantennary and/or tetraantennary *N*-linked glycans that Con A did not recognize. The largest *N*-linked glycan to be described on myeloid cells (molecular mass approximately 6,600) is a disialylated tetraantennary structure with poly-*N*-acetyllactosamine sequences on each antenna and both external and core fucose (Spooncer, et al. (1984) *J. Biol. Chem.* **259**, 4792-4801). Therefore, the observed change in electrophoretic mobility after PNGaseF treatment is consistent with the removal of at least two and possibly three complex *N*-linked glycans. Even after PNGaseF treatment, the Con A-unbound material migrated slightly slower than the Con A-bound material. This may indicate that the Con A-unbound fraction retained a single *N*-linked chain that was resistant to PNGaseF. Alternatively, there may be other structural differences between the Con-A-bound and Con-unbound fractions, of which the most likely is heterogeneity in *O*-linked glycosylation.

Treatment with sialidase or endo- β -galactosidase had similar effects on the electrophoretic mobilities of the Con A-unbound and Con A-bound fractions, probably because these enzymes primarily affected the abundant *O*-linked oligosaccharides. Endo- β -galactosidase produced 5 similar increases in the electrophoretic mobilities of PSGL-1 pretreated with PNGaseF and PSGL-1 that contained all its *N*-linked oligosaccharides, suggesting that at least some of the poly-*N*-acetyllactosamine was on the *O*-linked glycans.

PSGL-1 Expresses Le^x and sLe^x on *O*-linked Poly-*N*-acetyllactosamine. To determine whether PSGL-1 expresses Le^x or sLe^x on poly-*N*-acetyllactosamine, an assay was developed to measure the binding of ^{125}I -PSGL-1 to an immobilized monoclonal IgM antibody to sLe^x (CSLEX-1) or Le^x (LeuM1), or to an immobilized irrelevant IgM mAb (MOPC104E). PSGL-1 bound to CSLEX1 but not to MOPC104E, 15 confirming previous immunoblotting data that the glycoprotein expressed sLe^x . Greater than 85% of the ^{125}I -PSGL-1 molecules bound to a CSLEX-1 affinity column, indicating that virtually all molecules contained sLe^x . PSGL-1 also bound to LeuM1, indicating that it contained Le^x , although it is possible that desialylation generated this structure during 20 purification of the glycoprotein. Treatment of PSGL-1 with sialidase eliminated binding to CSLEX-1 and increased binding to LeuM1, consistent with the known specificities of these antibodies. Treatment of PSGL-1 with fucosidase had no effect on binding to CSLEX-1, in keeping with the inhibition of this enzyme by terminal sialic acid (Sano, et al., 25 (1992) *J. Biol. Chem.* **267**, 1522-1527; Maemura, K. and Fukuda, M. (1992) *J. Biol. Chem.* **267**, 24379-24386). In contrast, the fucosidase eliminated the binding of PSGL-1 to LeuM1.

After establishing the specificity of the assay, it was determined whether treatment of PSGL-1 with endo- β -galactosidase affected the 30 expression of Le^x or sLe^x . This enzyme abolished binding of PSGL-1 to immobilized LeuM1 and decreased binding to CSLEX-1 by $64 \pm 14\%$ (mean \pm SI, n = 4). These data clearly indicated that poly-*N*-

-42-

acetyllactosamine cleaved by endo- β -galactosidase carried some, and perhaps all, of the Le^x as well as a portion of the sLe^x on PSGL-1. The remaining sLe^x structures may be on branched or substituted poly-*N*-acetyllactosamine that was resistant to the enzyme or on chains that lacked poly-*N*-acetyllactosamine.

5 In contrast to the effects of endo- β -galactosidase, PNGaseF did not affect binding of PSGL-1 to immobilized LeuM1 or CSLEX-1, indicating that few, if any, Le^x or sLe^x structures were present on PNGaseF sensitive *N*-linked oligosaccharides. In conjunction with the 10 other data, this observation indicated that *O*-linked oligosaccharides carried most, and perhaps all, of the Le^x and sLe^x structures. Furthermore, at least some of these structures were on *O*-linked poly-*N*-acetyllactosamine.

15 *Effects of Glycosidases on Binding of PSGL-1 to P-selectin.* The effects of the glycosidases on binding of ¹²⁵I-PSGL-1 to immobilized tPS were then determined. Treatment with sialidase, which eliminated binding to the anti-sLe^x antibody but enhanced binding to the anti-Le^x antibody, completely prevented binding of PSGL-1 to tPS. In contrast, treatment with fucosidase, which eliminated binding to the anti-Le^x 20 antibody but not to the anti-sLe^x antibody, had no effect on the interaction of PSGL-1 with tPS. These data are consistent with previous results from other assays that high-affinity binding of P-selectin to myeloid cells or to PSGL-1 required sialic acid on the ligand.

25 Treatment of PSGL-1 with endo- β -galactosidase resulted in only a modest reduction in binding to immobilized tPS. In four independent experiments, endo- β -galactosidase reduced binding by 25 \pm 11% (mean \pm SD, n = 4). These data suggested that some, but not all, of the sialylated chains required for P-selectin recognition were on poly-*N*-acetyllactosamine that were sensitive to the enzyme. The remaining 30 glycans required for recognition might be on poly-*N*-acetyllactosamine that were resistant to endo- β -galactosidase or on oligosaccharides that lacked poly-*N*-acetyllactosamine. Endo- β -galactosidase inhibited binding

-43-

of PSGL-1 to the anti-sLe^x antibody more than it inhibited binding to tPS. The levels of sLe^x with the interaction with tPS could not be accurately correlated, since it is not known how many sLe^x structures were required for PSGL-1 to bind to either the antibody or to tPS in these assays.

5 Treatment of PSGL-1 with PNGaseF did not affect its ability to bind to tPS, supporting the conclusion that *N*-linked oligosaccharide played little if any role in P-selectin recognition.

10 The structure and function of PSGL-1 from human neutrophils was further characterized using assays with the purified, radioiodinated glycoprotein. The results suggest that this molecule presents an array of sialylated and fucosylated, *O*-linked poly-*N*-acetyllactosamine that creates high-affinity binding structures for P-selectin.

15 Although the great majority of the oligosaccharides on PSGL-1 are *O*-linked, there are a few *N*-linked glycans which were determined to be heterogeneous. All PSGL-1 molecules contained PNGaseF-sensitive *N*-linked oligosaccharides, but only 42% of the glycoprotein molecules bound to Con A, a lectin that binds well to high mannose *N*-linked glycans and less well to complex biantennary and hybrid *N*-linked chains. Based on the reductions in apparent molecular weight observed in SDS 20 gels, PNGaseF removed at least two of the three possible *N*-linked glycans from both the Con A-bound and Con A-bound fractions of PSGL-1. One chain on the Con A-unbound fraction of PSGL-1 may be resistant to PNGaseF, since the Con A-unbound material still migrated slightly slower in SDS gels than the Con A-bound material following treatment 25 with PNGaseF. Alternatively, PNGaseF may have removed all the *N*-linked chains from both the Con A-bound and Con A-unbound fractions, in which case the mobility difference may reflect heterogeneous *O*-linked glycosylation. The PNGaseF-sensitive *N*-linked glycans carried little or no Le^x and sLe^x and did not contribute measurably to the binding 30 interaction with P-selectin. These results strongly suggest that the *O*-linked glycans display the relevant structures that P-selectin recognizes.

It is highly improbable that a single PNGaseF-resistant *N*-linked glycan on a subpopulation of PSGL-1 is responsible for P-selectin recognition.

Because at least some of the *O*-linked oligosaccharides on PSGL-1 carry (α 2-3)sialic acid and (α 1-3)fucose, they must have relatively large, branched structures. In agreement with this prediction, PSGL-1 was found to carry few of the Ser/Thr-linked Gal β 1-3GalNAc core 1 disaccharides that endo- α -*N*-acetylgalactosaminidase cleaves following removal of terminal sialic acids. These data are consistent with previous observations that less than 10% of the 3 H-labeled *O*-linked oligosaccharides released by β -elimination from PSGL-1 were small as assessed by gel filtration.

The *O*-linked glycans on PSGL-1 include a heterogeneous group of poly-*N*-acetyllactosamine. As measured by mobility on SDS gels and by binding to tomato lectin, endo- β -galactosidase cleaved some of the poly-*N*-acetyllactosamine at the internal β 1-4 linkage between the Gal and GlcNAc. Treatment with this enzyme also reduced the amount of Le x and sLe x , indicating that some of these structures were on poly-*N*-acetyllactosamine. As measured by binding to tomato lectin, endo- β -galactosidase cleaved other poly-*N*-acetyllactosamine only after treatment with sialidase and fucosidase, demonstrating that they had sialic acid and fucose moieties at or near their terminus that prevented access to endo- β -galactosidase. Because a significant portion of the PSGL-1 molecules still bound to tomato lectin after treatment with all three enzymes, there may be other *O*-linked poly-*N*-acetyllactosamine chains that remained resistant to endo- β -galactosidase because of branching or additional internal substitutions. Such structures might be responsible, at least in part, for the ability of PSGL-1 to bind to immobilized tPS after digestion with endo- β -galactosidase.

Previous studies have shown that human myeloid cells express poly-*N*-acetyllactosamine on some *O*-linked glycans. These structures occur almost exclusively on the branch attached to the C-6 of GalNAc as part of a core 2 *O*-linked oligosaccharide (Fukuda, et al., (1986) *J. Biol.*

Chem. **261**, 12796-12806; Hanisch, et al., (1989) *J. Biol. Chem.* **264**, 872-883). Of the approximately 80 *O*-linked glycans on the major sialomucin of human myeloid cells, leukosialin (CD43), only one or two contain poly-*N*-acetyllactosamine, and approximately half of these have a 5 terminal sLe^x moiety. In contrast, PSGL-1 from neutrophils appears to contain a higher percentage of core 2, poly-*N*-acetyllactosamine. Direct structural analysis will be required to determine the specific monosaccharides, linkages, and modifications that constitute the *O*-linked glycans of PSGL-1. However, the current data suggest that some of these 10 glycans may differ from those in leukosialin, implying that human myeloid cells can differentially glycosylate the polypeptide backbones of two distinct sialomucins. This differential glycosylation is functionally important, because P-selectin binds with high affinity to PSGL-1, but not to leukosialin.

15 The selectins require sialylation and fucosylation of most carbohydrate ligands for recognition. Similarly, PSGL-1 requires sialic acid and fucose to interact with P-selectin. Although PSGL-1 expresses the terminal sLe^x tetrasaccharide, it is not clear whether it requires this structure *per se* for recognition. A simple model is that PSGL-1 presents 20 many terminal sLe^x moieties that increase the avidity with which it interacts with P-selectin. However, previous observations indicate that high concentrations of cell surface sLe^x are not sufficient to confer optimal interactions with P-selectin. It is more likely that additional modifications on each oligosaccharide chain are essential for optimal 25 recognition. Specific features of several oligosaccharides in proximity may also present a clustered saccharide patch that promotes high affinity binding to P-selectin.

Sako et al. (1993) generated a recombinant form of PSGL-1 that bound to P-selectin when expressed in COS cells co-transfected with an 30 α 1-3/4 fucosyltransferase (FTIII), but not in COS cells lacking the fucosyltransferase. The relative molecular mass of the recombinant molecule was similar to that of PSGL-1 in human myeloid HL-60 cells,

and the authors apparently concluded that the structure and function of recombinant PSGL-1 were equivalent to those of the native molecule. However, the specific oligosaccharide structures attached to native PSGL-1 and the glycosyltransferases required to construct these structures are 5 unknown. Moreover, the α 1-3/4 fucosyltransferase (FTIII) expressed in the COS cells to generate recombinant PSGL-1 differs from the fucosyltransferases in human myeloid cells (Goelz, et al. (1990) *Cell* **63**, 1349-1356; Sasaki, et al. (1994) *J. Biol. Chem.* **269**, 14730-14737). Thus, it is not clear that the glycosylation, and hence the function, of 10 recombinant PSGL-1 is identical to that of the native glycoprotein in human myeloid cells. Indeed, following sialidase digestion of recombinant PSGL-1, addition of endo- α -N-galactosaminidase caused a substantial increase in mobility of the glycoprotein, indicating that 15 recombinant PSGL-1 expressed in COS cells with FTIII, in contrast to the native glycoprotein, contains many simple *O*-linked core 1 Gal β 1-3GalNAc disaccharides that are susceptible to this enzyme.

Sako et al. found that recombinant PSGL-1 no longer bound to P-selectin after treatment with sialidase and endo- β -N-galactosaminidase, and concluded that *O*-linked oligosaccharides were important for 20 recognition by P-selectin. Based on the data presented here, this conclusion appears to be correct. However, the experiment of Sako et al. cannot be used to support a role for *O*-linked glycans in recognition. Prolonged digestion with sialidase is required to eliminate binding to native PSGL-1, and the conditions for sialidase digestion of recombinant 25 PSGL-1 used by Sako et al. reduced, but did not eliminate, binding of P-selectin. The loss of P-selectin binding after addition of endo- α -N-galactosaminidase, which releases only non-sialylated core 1 Gal β 1-3GalNAc disaccharides, was more likely due to the previously observed contamination of this enzyme with sialidase, which would remove the 30 remaining sialic acids required for recognition.

Sako et al. also suggested that the *N*-linked oligosaccharides contributed to binding to P-selectin, because recombinant P-selectin did

not reprecipitate all of the PNGaseF-treated recombinant PSGL-1. However, these authors did not demonstrate whether the reprecipitation assay was quantitative by testing whether additional precipitation steps increased recovery of PSGL-1. The same group also concluded that *N*-linked glycosylation was important for P-selectin recognition because they observed that myeloid HL-60 cells treated with tunicamycin, an inhibitor of *N*-linked glycosylation, did not adhere to P-selectin-expressing cells (Larsen, et al., (1992) *J. Biol. Chem.* **267**, 11104-11110). However, tunicamycin exerts indirect effects on cells, such as inhibiting the exit of some proteins from the endoplasmic reticulum, that make it difficult to interpret its effects on cell adhesion. The studies in this application provide no evidence that the *N*-linked glycans of PSGL-1 from human neutrophils play a role in recognition by P-selectin.

Example 2: Expression of PSGL-1 in CHO cells transfected with cDNA encoding fucosyltransferase III and a core 2 β 1-6-N-acetylglucosaminyltransferase.

A functional, recombinant form of PSGL-1 that has many of the properties of native PSGL-1 purified from human neutrophils has been expressed. The study demonstrates that PSGL-1, in order to bind P-selectin, requires sialylated and fucosylated O-linked oligosaccharides containing the branched core 2 structure.

PSGL-1 was expressed in Chinese hamster ovary (CHO) cells, which were selected because the *N*- and O-linked oligosaccharide structures on these cells are very well characterized. In particular, the Ser/Thr-linked oligosaccharides on secreted and surface glycoproteins are simple, core 1, mono- and disialylated derivatives of Ga1 β 1-3Ga1NAc α 1-Ser/Thr (Sasaki et al., *J. Biol. Chem.* **262**:12059-12076, 1987; Seguchi et al., *Arch. Biochem. Biophys.* **284**: 245-256, 1991). DHFR-minus CHO cells were stably transfected with the cDNA encoding fucosyltransferase III (FTIII) (Kukowska-Latallo, et al., 1990), with the cDNA encoding the core 2 β 1-6-N-acetylglucosaminyltransferase (Bierhuizen and Fukuda, *Proc. Natl. Acad. Sci. USA* **89**:9326-933), 1992; Bierhuizen et al., *J. Biol. Chem.* **269**:4473-4479, 1994), or with both cDNAs. The cDNAs

-48-

were ligated into the pRc/RSV or pCDNA3 expression vectors (Invitrogen). Following transfection with the calcium phosphate precipitation method, stably transfected clones were selected in medium containing G418. Stable clones expressing either or both

5 glycosyltransferases were identified using specific glycosyltransferase assays performed on cell lysates. Cells expressing FTIII also expressed high levels of the sialyl Lewis x antigen (NeuAc α 2-3Gal β 1-4[Fuca1-3]GlcNAc β 1-4-R) on their surfaces, as measured by binding of the CSLEX1 monoclonal antibody (Fukushima et al., *Cancer Res.* **44**:5279-10 5285, 1984).

A cDNA encoding a full-length form of PSGL-1 (Moore et al., *J. Cell Biol.* **128**: 661-671, 1995) was inserted into the pZeoSV expression vector (Invitrogen). CHO cells expressing the glycosyltransferases were transfected with the PSGL-1 cDNA using the Lipofectamine reagent (Gibco BRL Life Technologies). In transient expression analysis, cells were analyzed by flow cytometry 48 h after transfection. Stable clones were selected in medium containing both G418 and Zeocin (Invitrogen). Analysis of cell-bound antibodies by flow cytometry was performed as described in Moore et al, (1995). Analysis of cell-bound platelet-derived 15 P-selectin by flow cytometry was performed as described by Moore and Thompson, *Biochem. Biophys. Res. Commun.* **186**:173-181 (1992).

In transient expression studies, PSGL-1 was co-expressed on the majority of transfected cells expressing FTIII, the core 2 β 1-6-N-acetylglucosaminyltransferase, or both glycosyltransferases. Co-expression was documented by strong binding of anti-PSGL-1 monoclonal antibodies PL1 or PL2 (Moore, et al. 1995) as measured by flow cytometry. In contrast, flow cytometric analysis indicated that P-selectin bound only to CHO cells expressing FTIII, the core 2 β 1-6-N-acetylglucosaminyltransferase and PSGL-1. P-selectin did not bind to 20 CHO cells expressing either or both glycosyltransferases, in the absence of PSGL-1. P-selectin also failed to bind to CHO cells expressing either FTIII or the core 2 β 1-6-N-acetylglucosaminyltransferase, even in the

-49-

presence of PSGL-1. Thus, a functional form of PSGL-1 capable of binding P-selectin was expressed only on CHO cells that also expressed FTIII and the core 2 β 1-6-N-acetylglucosaminyltransferase.

CHO cell clones were also selected that stably expressed FTIII
5 and the core 2 β 1-6-N-acetylglucosaminyltransferase. These cells
expressed high levels of PSGL-1 as detected by binding of PL1 and PL2
by flow cytometry. P-selectin also bound well to these stably selected
cells. Thus, high affinity binding of P-selectin to CHO cells required the
simultaneous expression of FTIII, the core 2 β 1-6-N-
10 acetylglucosaminyltransferase, and PSGL-1 by the cells. For cells
expressing PSGL-1 either transiently or stably, binding of P-selectin was
specific, as it was blocked by an inhibitory monoclonal antibody to P-
selectin, G1, but not by a noninhibitory monoclonal antibody to P-
selectin, S12 (Geng et al., *Nature* 343:757-760, 1990). Binding was also
15 Ca^{2+} - dependent, consistent with the requirement for Ca^{2+} for selectins to
bind carbohydrate ligands. P-selectin also bound specifically to PSGL-1
on the cells: PL1, but not PL2, completely blocked binding of P-selectin.
Furthermore, the anti-42-56 serum, discussed below, but not control
rabit serum, blocked binding of P-selectin.

20 These data indicate that CHO cells can be genetically engineered
to express a recombinant form of PSGL-1 that binds specifically and with
high affinity to P-selectin as does native PSGL-1 from human leukocytes.
The inhibition of P-selectin binding by PL1 and by the anti-42-56 serum
indicates that P-selectin binds to the same region on recombinant PSGL-1
25 and on native PSGL-1 from human leukocytes. The requirement for the
core 2 enzyme in the CHO cells indicates that a critical component of the
P-selectin recognition site on PSGL-1 consists of branched, core 2 O-
linked oligosaccharide(s) with a $\text{Gal}\beta 1-3(\text{Gal}\beta 1-4\text{GlcNAc}\beta 1-6)\text{GalNAc}$ -
Ser/Thr backbone. Sialylation and fucosylation of PSGL-1 is also
30 required for high affinity binding to P-selectin; these modifications must
occur on the critical core 2 O-linked glycans.

Example 3: Determination of a sulfated tyrosine containing region in PSGL-1 critical for P-selectin binding.

PSGL-1 has an extracellular domain of 308 residues. The first 18 residues constitute a signal peptide. Residues 19-41 constitute a putative propeptide that is predicted to be cleaved following synthesis, probably in the trans-Golgi network. Therefore, the putative N-terminus of the mature protein begins at residue 42. The three tyrosines within the consensus sequence for tyrosine sulfation are at residues 46, 48, and 51, near the N-terminus. A rabbit polyclonal antiserum to a synthetic residue encoding residues 42-56 completely blocks binding of PSGL-1 to P-selectin, suggesting that a least part of the critical recognition site for P-selectin lies in this region. All references to residue number refer to Sequence Listing ID No. 1.

Two IgG monoclonal antibodies to PSGL-1, termed PL1 and PL2, were previously described in Moore et al, *J Cell Biol.* 128:661-671 (1995)). PL1, but not PL2, completely blocks binding of PSGL-1 to P-selectin, and abolishes adhesion of leukocytes to P-selectin under both static and shear conditions. To map the regions on PSGL-1 where the epitopes for PL1 and PL2 are located, a series of fusion proteins containing overlapping regions of the extracellular domain of PSGL-1 were expressed in bacteria. The PSGL-1 sequences in the fusion proteins were as follows: Fragment 1, residues 18-78; Fragment 2, residues 63-123; Fragment 3, residues 109-168; Fragment 4, residues 154-200; Fragment 5, residues 188-258; and Fragment 6, residues 243-308. The DNA encoding each fragment was fused in frame to the DNA encoding dihydrofolate reductase and a 6X His tag, in the expression vector pQE-40 (Qiagen). Each fragment was expressed in bacteria and purified on a nickel affinity column which bound to the 6X His tag. The reactivity of each fusion protein with the antibodies was assessed by Western blotting under reducing conditions.

PL1 bound strongly to Fragment 1, but not to any of the other fragments. This indicates that the PL1 epitope is located within residues

-51-

18-78. Assuming proteolytic removal of the propeptide from native mature PSGL-1, the epitope is located within residues 42-78. Since PL1 does not bind Fragment 2 (residues 63-123), the epitope is likely located within residues 42-70. Thus, PL1 binds to an N-terminal epitope that is 5 near the epitopes for the polyclonal anti-42-56 serum. This result further suggests that a critical component of the P-selectin recognition site resides within a small region at the N-terminus of PSGL-1. In contrast, the nonblocking monoclonal antibody PL2 bound to Fragments 5 and 6 (but not the other fragments), suggesting that it recognizes an epitope within 10 residues 243-258, which are present in both fragments.

These results demonstrate that P-selectin binds to a specific region on both native and recombinant PSGL-1. This region resides near the N-terminus of the mature molecule. High affinity binding of PSGL-1 to P-selectin requires both tyrosine sulfation and sialylated, fucosylated, core 2 15 branched O-linked oligosaccharides. Fragments of PSGL-1, or derivatives thereof, can be constructed that contain the binding site for P-selectin. Such fragments may be as short as that encoding residues 42-58; this fragment contains the three tyrosines that are capable of being sulfated as well as two threonines that may be attachment sites for core 2, 20 O-linked oligosaccharides. Longer fragments extending from residue 42 should also bind with high affinity to P-selectin.

Experimental Procedures

Chemicals and reagents. Carrier Free [³⁵S]sulfate (1100-1600 Ci/mmol) was purchased from Dupont/NEN (Wilmington, DE).

25 Emphaze® affinity support resin was purchased from Pierce Biochemicals (Rockford, IL). *Aerobacter aerogenes* arylsulfatase, *Arthrobacter ureafaciens* neuraminidase, and sulfated monosaccharides were obtained from Sigma Chemical Co. (St. Louis, MO). Recombinant peptide:N-glycosidase F was purchased from Boehringer Mannheim (Indianapolis, 30 IN). All cell culture reagents were obtained from GIBCO/BRL (Grand Island, NY). Authentic tyrosine sulfate standard was synthesized as described (Huttner, (1984) *Meth. Enzymol.* **107**, 200-224) using

concentrated H₂SO₄ and tyrosine. Other chemicals were ACS grade or better and were obtained from Fisher Scientific.

Isolation of radiolabeled PSGL-1. PSGL-1 was purified from human neutrophils and radiolabeled with Na¹²⁵I as described by Moore, et al., (1994). HL-60 cells (1-2 x 10⁶ cells/ml) were labeled for 48 h with 100 µCi/ml [³⁵S]sulfate at 37°C in sulfate-deficient medium (S-MEM) containing 10% dialyzed fetal bovine serum. ³⁵S-PSGL-1 was purified using affinity chromatography in a column containing recombinant soluble P-selectin (Ushiyama, et al., (1993) *J. Biol. Chem.* **268**, 15229-15237) coupled to Emphaze® at a density of 5 mg/ml (Moore, et al. (1994)). The enriched EDTA-eluted sample was rechromatographed on the P-selectin column after dialysis into Ca²⁺-containing buffer. Purified ³⁵S-PSGL-1 was treated with reducing or nonreducing SDS sample buffer and electrophoresed in a 7.5% polyacrylamide gel (Laemmli (1970)). Gels were dried and then exposed to Fuji RX x-ray film at -80°C.

Detection of tyrosine sulfate. Tyrosine sulfation of PSGL-1 was determined by base hydrolysis of ³⁵S-PSGL-1. Following identification of ³⁵S-PSGL-1 by autoradiography, the dried gel was aligned with the exposed x-ray film and the band from the nonreducing lane was subjected to strong base hydrolysis in 1.0 M NaOH at 110°C, for 24 h as described (Hortin, et al., *J. Biol. Chem.* **261**, 15827-15830 (1986). The hydrolysate was passed over a Dowex 50 (H⁺) column washed with water, lyophilized, and analyzed by anion exchange chromatography using a Varian AX-5 column (4 mm x 30 cm) that was eluted with a gradient of NaH₂PO₄, pH 3.0.

Enzymatic treatment of PSGL-1. All arylsulfatase digestions were performed with *Aerobacter aerogenes* arylsulfatase (EC 3.1.6.1) (Fowler and Rammier (1964) *Biochemistry* **3**, 230-237) in approximately 500 µl of 0.01 M Tris-HCl, pH 7.5, overnight at 37°C. Following digestion, the enzyme reactions were terminated by boiling for 15 in. Sham digestions were performed with 1000 mU of boiled enzyme. ³⁵S-tyrosine sulfate from ³⁵S-PSGL-1 was prepared by base hydrolysis as described above.

-53-

The base hydrolysate was treated with 50 mU or 1000 mU of active arylsulfatase, or with 1000 mU of boiled arylsulfatase, then analyzed by anion exchange chromatography.

The specificity of the *A. aerogenes* arylsulfatase was determined 5 for the sulfated monosaccharides GlcNAc-6-sulfate, Gal-6-sulfate, GalNAc-6-sulfate, and GalNAc-4-sulfate. Each of the sulfated monosaccharides (100 nmol) was separately treated with 1000 mU of boiled or active enzyme. After overnight incubation, the reaction mixtures were boiled for 15 min, lyophilized, resuspended in water and 10 analyzed by high pH anion exchange chromatography with a Dionex CarboPac PA-1 column and PAD detection as described (Shilatifard and Cummings (1995) *Glycobiology* 5, 291-297.

Intact ³⁵S-PSGL-1 or ¹²⁵I-PSGL-1 was treated with *A. aerogenes* arylsulfatase as described above. The [³⁵S]sulfate released from ³⁵S- 15 PSGL-1 by arylsulfatase was quantified by BaSO₄ precipitation, in which 100 μ l saturated Na₂SO₃ as a carrier (Shilatifard, et al., (1993) *J. Virol.* 67, 943-945). In the graphic representation of the data, the sulfate released from the sham-treated sample was adjusted setting the sham-treated sample equal to 0%. The actual percentage of radioactivity 20 released from the sham-treated samples ranged from 1.3-5.0%.

In a control experiment, ¹²⁵I-PSGL-1 was treated with 1000 mU of *Arthrobacter ureafaciens* neuraminidase in 0.05 M sodium acetate, pH 5.0 overnight at 37°C prior to application to a CSLEX-1 antibody column. Chromatography on CSLEX-1 was performed as described previously 25 (Moore, et al, 1994).

³⁵S-PSGL-1 was treated with peptide:N-glycosidase F as described Shilatifard, et al., (1993). ³⁵S-PSGL-1 was denatured by boiling 5 min in 30 50 μ l of 50 mM sodium phosphate, pH 7.5, 50 mM 2-mercaptoethanol, 0.5% SDS. The SDS was then diluted to a final concentration of 0.2% with 1.5% NP-40. 10 U of enzyme was added to the denatured ³⁵S-PSGL-1 and incubated overnight at 37°C.

-54-

Rebinding of arylsulfatase-treated PSGL-1 to P-selectin.

Arylsulfatase-treated PSGL-1 was applied to a P-selectin affinity column in Ca^{2+} -containing buffer, and the bound radioactivity was eluted with EDTA. The percentage of ^{125}I -PSGL-1 eluted from the column with 5 EDTA was corrected setting the sham-treated sample equal to 100% (actual rebinding ranged from 87-99%).

Immunoprecipitation and P-selectin binding assays.

10 Immunoprecipitation of ^{35}S -PSGL-1 was performed as described by Moore, et al., (1994), using a rabbit antiserum (Moore, et al., (1995), Moore, et al., 1994)) to a peptide corresponding to residues 42-56 (QATEYEYLDYDFLPE) of PSGL-1 (Sako, et al., (1993)) (anti-42-56) or normal rabbit serum (NRS). The immunoprecipitates were analyzed by SDS-PAGE under nonreducing conditions, followed by autoradiography. The solid phase P-selectin binding assay was performed as described by 15 Moore, et al., 1994). Briefly, ^{125}I -PSGL-1 was incubated in the presence of NRS or the antiserum directed against residues 42-56 of PSGL-1 (anti-42-56) in microtiter plates containing immobilized recombinant soluble P-selectin (50 μl , 10 $\mu\text{g}/\text{ml}$) or human serum albumin (HSA) (50 μl , 1% HSA). Binding of platelet-derived P-selectin to HL-60 cells was 20 measured by flow cytometry as described in the presence of 20% NRS or 20% anti-42-56 serum.

Results and Discussion

To determine if PSGL-1 is post-translationally sulfated, human 25 promyelocytic leukemia HL-60 cells were metabolically-labeled with $[^{35}\text{S}]$ sulfate, and PSGL-1 was purified from lysates of these cells by affinity chromatography on a P-selectin column. An $[^{35}\text{S}]$ sulfate-labeled protein was detected that migrated in SDS gels with a relative molecular mass of 120,000 under reducing conditions and 240,000 under nonreducing conditions.

30 These mobilities are consistent with the disulfide-linked homodimeric structure of PSGL-1 (Moore, et al., 1992). Furthermore, the $[^{35}\text{S}]$ sulfate-labeled protein was immunoprecipitated with a specific

rabbit antiserum generated against a synthetic peptide encoding residues 42-56 of the extracellular domain of PSGL-1. Analysis of the supernatants showed that anti-42-56 precipitated all of the ^{35}S -PSGL-1, whereas normal rabbit serum (NRS) precipitated none of the ^{35}S -PSGL-1.

5 These results demonstrate that PSGL-1 is post-translationally sulfated.

Sulfate can be incorporated into eukaryotic glycoproteins as tyrosine sulfate (Huttner, et al. (1988) *Mod. Cell Biol.* **6**, 97-140) or as sulfated carbohydrates (Fiete, et al., (1991) *Cell* **65**, 1103-1110; Kjellen and Lindahl; (1991) *Ann. Rev. Biochem.* **60**, 443-475). In preliminary 10 studies, sulfated carbohydrates were not detected on PSGL-1, using techniques that detected such structures on other glycoproteins (Shilatifard, et al. (1995, 1993)). The cDNA sequence of PSGL-1 predicts four extracytoplasmic tyrosine residues, three of which are clustered at positions 46, 48 and 51 within a predicted consensus sequence 15 for tyrosine sulfation (Huttner and Baeuerle (1988)). To determine if PSGL-1 has tyrosine sulfate, the gel slice containing the 240 kDa ^{35}S -PSGL-1 was hydrolyzed with strong base and analyzed by anion exchange chromatography. A single radioactive peak was recovered that co-eluted with tyrosine sulfate but not with sulfated sugar standards (Figure 1A).

20 Tyrosine sulfate was then tested for importance for binding of PSGL-1 to P-selectin. Although the functions of tyrosine sulfate within proteins are not clear, some proteins require this modification for optimal activity (Pittman, et al. (1992) *Biochemistry* **31**, 3315-3325; Dong, et al., (1994) *Biochemistry* **33**, 13946-13953). The usual approach for assessing 25 the importance of tyrosine sulfate is to prevent sulfation of newly synthesized proteins with chemical inhibitors or to replace tyrosine with phenylalanine by site-directed mutagenesis. An alternative approach in which sulfate was enzymatically removed from tyrosine on intact PSGL-1 was tested. The ability of an arylsulfatase from *Aerobacter aerogenes* to 30 release sulfate from ^{35}S -tyrosine sulfate was also determined. As little as 50 mU of this arylsulfatase quantitatively released [^{35}S]sulfate from ^{35}S -tyrosine sulfate derived from PSGL-1 (Figures 1B, 1C). This cleavage

-56-

was specific, since 1000 mU of the enzyme did not release any sulfate from Gal-6-sulfate, GlcNAc-6-sulfate, GalNAc-4-sulfate, and GalNAc-6-sulfate.

The ability of the arylsulfatase to release sulfate from intact ^{35}S -PSGL-1 was examined. $[^{35}\text{S}]$ sulfate released by the enzyme was quantified by precipitation as insoluble BaSO_4 . Up to 50% of the $[^{35}\text{S}]$ sulfate on PSGL-1 was released by 500 mU of arylsulfatase; increasing amounts of enzyme did not release more radioactivity (Figure 2).

10 The functional importance of the tyrosine sulfate on PSGL-1 was assessed by treating ^{125}I -PSGL-1 with arylsulfatase and measuring the rebinding of the treated ligand to a P-selectin column. Binding of arylsulfatase-treated ^{125}I -PSGL-1 to P-selectin was reduced in a dose-dependent manner; the decreased binding was inversely related to the 15 amount of sulfate released from ^{35}S -PSGL-1 (Figure 2). The reduced binding of PSGL-1 to P-selectin following arylsulfatase treatment was not due to general release of sialic acid and/or fucose by contaminating exoglycosidases, because PSGL-1 treated with 1000 mU of arylsulfatase bound quantitatively to any affinity column containing CSLEX-1, a 20 monoclonal antibody of sLe x .

Arylsulfatase released approximately 50% of the $[^{35}\text{S}]$ sulfate from ^{35}S -PSGL-1 and decreased rebinding of ^{125}I -PSGL-1 to P-selectin by the same degree. The possibility was considered that the fraction of ^{35}S -PSGL-1 that rebound to P-selectin following treatment with arylsulfatase 25 retained critical tyrosine sulfate residues, whereas the fraction that did not rebind to P-selectin had lost tyrosine sulfate. ^{35}S -PSGL-1 was treated with 1000 mU of arylsulfatase and then applied to a P-selectin column. The bound and unbound fractions were analyzed by SDS-PAGE. A band corresponding to ^{35}S -PSGL-1 was observed in the bound fractions, 30 whereas no band was seen in the unbound fractions. The radioactivity in the unbound fractions represented free sulfate. When the band of ^{35}S -PSGL-1 in the bound fractions was hydrolyzed, radioactivity was

recovered in tyrosine sulfate. In the control experiment, ^{125}I -PSGL-1 was treated with 1000 mU of arylsulfatase and then applied to the P-selectin column. SDS-PAGE analysis revealed that intact ^{125}I -PSGL-1 was recovered in both the unbound and bound fractions. Thus, arylsulfatase 5 removed sulfate quantitatively from a subset of PSGL-1, which no longer bound P-selectin, but the enzyme did not otherwise degrade PSGL-1.

The tyrosine sulfate remaining in the P-selectin- bound- subset of PSGL-1 may be resistant to arylsulfatase because of its inaccessibility or because of some other feature of PSGL-1 that blocks action of the 10 enzyme. When ^{35}S -PSGL-1 was partly deglycosylated by treatment with peptide:N-glycosidase F and *Arthrobacter ureafaciens* neuraminidase, subsequent treatment with arylsulfatase released up to 70% of the 15 radioactivity as $[^{35}\text{S}]$ sulfate. Since neuraminidase also eliminates binding of PSGL-1 to P-selectin, it could not be determined whether the increased removal of $[^{35}\text{S}]$ sulfate further reduced binding of PSGL-1 to P-selectin. These results suggest that the extensive glycosylation of PSGL-1 may account for the inaccessibility of some tyrosine sulfate sites to arylsulfatase.

To further demonstrate the importance of tyrosine sulfation for the 20 function of PSGL-1, the antiserum to the peptide encoding residues 42-56 of the extracellular domain of PSGL-1 was used. As noted above, this region contains the three potential tyrosine sulfation sites at residues 46, 48, and 51. The antiserum inhibited binding of ^{125}I -PSGL-1 to immobilized P-selectin (Figure 3A), and also abolished binding of P-selectin to HL-60 cells, as measured by flow cytometry (Figure 3B). Thus, antiserum to the 42-56 peptide binds to native PSGL-1 and effectively blocks its recognition by P-selectin.

These results demonstrate that PSGL-1 contains tyrosine sulfate 30 that is required for high affinity binding to P-selectin. It has been shown previously that PSGL-1 contains the sLe α determinant on O-linked oligosaccharides and that both sialic acid and fucose are required for binding of PSGL-1 to P-selectin. Tyrosine sulfate may be important

because it promotes appropriate presentation of the glycans that bind directly to P-selectin. Alternatively, tyrosine sulfate may directly interact with P-selectin. This latter possibility seems more likely, since sulfatide and sulfated oligosaccharides are known to bind P-selectin. Sulfate is also 5 a critical determinant for the binding of GlyCAM-1 to L-selectin, but GlyCAM-1 contains sulfate in Gal-6-sulfate and GlcNAc-6-rather than on monosaccharides.

The results suggest a model in which PSGL-1 presents both carbohydrate and tyrosine sulfate as components of a critical recognition 10 site for P-selectin. It is hypothesized that this site is located at the N-terminal, membrane-distal region of PSGL-1, near the three potentially sulfated tyrosine residues. Consistent with this hypothesis, PL1, a monoclonal antibody that recognizes a membrane-distal epitope on PSGL-1, blocks binding of fluid-phase P-selectin to leukocytes and abolished 15 adhesion of neutrophils to P-selectin under both static and shear conditions. In contrast, PL2, a monoclonal antibody that recognizes a membrane-proximal epitope on PSGL-1, does not inhibit binding to P-selectin. The concept that P-selectin recognizes a localized site on a mucin-like glycoprotein is distinct from a model in which a selectin 20 recognizes multiple, clustered O-linked glycans attached along the entire polypeptide (Lasky, et al., (1992) *Cell* **69**, 927-938; Baumheuter, et al., (1993) *Science* **262**, 436-438).

Example 4: Determination of O-glycan Structures on PSGL.

The structures of the O-glycans on PSGL-1 synthesized by HL-60 25 cells metabolically-radiolabeled with ³H-sugar precursors were determined. The glycosylation of PSGL-1 was compared with that of CD43 to determine whether two sialomucins expressed by the same cells are O-glycosylated differently. The HL-60 cell line was used in these studies because the post-translational modifications known to be important 30 for binding to P-and E-selectin on both HL-60 and neutrophil PSGL-1 are comparable.

The studies demonstrate that the majority of O-glycans of PSGL-1 are disialylated or neutral forms of the core-2 tetrasaccharide. Only a few of the O-glycans are fucosylated and these contain the structural determinant for the sLe^x antigen. These results demonstrate that PSGL-1
5 is glycosylated differently from CD43 and that PSGL-1 contains unique O-glycans that are likely to be critical for high affinity interactions with P- and E-selectin.

The abbreviations used are: sLe^x, sialyl Lewis x; NDV, Newcastle disease virus; CHO, Chinese hamster ovary; GlcN, 10 glucosamine; GalN, galactosamine; GalNOH, galactosaminitol; GalNAcOH, N-acetylgalactosaminitol; FT, fucosyltransferase; C2GnT, core 2 β 1,6 N-acetylglucosaminyltransferase; HPAEC, high pH anion exchange chromatography.

I. Experimental Procedures

15 *Materials* - Protein A-Sepharose, QAE-Sephadex, *Arthrobacter ureafaciens* neuraminidase, jack bean β -galactosidase, jack bean β -N-acetylhexosaminidase, and Gal β 1 \rightarrow 3GalNAc were obtained from Sigma. Pronase and TPCK-treated trypsin were purchased from Worthington Biochemicals. *Escherichia freundii* endo- β -galactosidase was obtained 20 from V-labs. *Streptomyces* sp.142 α 1,3/4-fucosidase was purchased from Takara. Newcastle Disease Virus neuraminidase was obtained from Oxford Glycosystems. D-[6-³H]glucosamine hydrochloride (20-45 Ci/mmol), D-[2-³H]mannose (20-30 Ci/mmol), and L-[6-³H]fucose (70-90 Ci/mmol) were purchased from Dupont/NEN and NaB[³H]₄ (40-60 Ci/mmol) was purchased from ARC. Rabbit anti-mouse IgG1 was purchased from Zymed. EmphazeTM affinity support resin was obtained 25 from Pierce Biochemicals. Bio-Gel P-4 and P-10 resins and molecular weight markers were obtained from BioRad. All cell culture reagents were obtained from GIBCO/BRL. Other chemicals were ACS grade or better and were obtained from Fisher Scientific.

30 *Isolation of Metabolically-radiolabeled PSGL-1* - ³H-GlcN, ³H-Man, and ³H-Fuc metabolic-radiolabeling of HL-60 cells was performed

-60-

essentially as described by Moore, et al. (1992) *J. Cell. Biol.* **118**, 445-456; Wilkins, et al. (1995) *J. Biol. Chem.* **270**, 22677-22680. 3 H-PSGL-1 was purified from cell extracts using affinity chromatography on a column of immobilized, recombinant soluble P-selectin (Ushiyama, et al. 5 (1993) *J. Biol. Chem.* **268**, 15229-15237) coupled to Emphaze™ at a density of 5 mg/ml (1.0 ml bed volume) (Moore, et al. (1994) *J. Biol. Chem.* **269**, 23318-23327). The enriched EDTA-eluted samples were re-chromatographed on the P-selectin column after dialysis into a Ca^{2+} -containing buffer. Purified 3 H-PSGL-1 was treated with reducing or 10 nonreducing SDS sample buffer [Laemmli, U. K., (1970) *Nature* **227**, 680-685] and electrophoresed in a 7.5% polyacrylamide gel. Gels were dried and then autoradiographed by exposure to Fuji RX X-ray film at -80°C. The approximately 250 kDa 3 H-PSGL-1 in gel slices from nonreducing lanes was analyzed.

15 *Immunoprecipitation of CD43 -* CD43 was immunoprecipitated from 3 H-sugar-labeled HC-60 cells using a CD43-specific mAb, H5H5 (IgG1). The H5H5 hybridoma cell line was produced by Dr. T. August and obtained from the Developmental hybridoma Bank (The Johns Hopkins University School of Medicine). CD43 was immunoprecipitated 20 by first preloading 50 μl Protein A-Sepharose beads with 100 μg of a rabbit anti-mouse IgG1 antibody at 37°C for 1 h. After washing, the CD43 specific antibody was absorbed to the beads by adding 500 μl of undiluted H5H5 culture media at 37°C for 1 h. After washing, the preloaded beads were added to the 3 H-sugar-labeled HC-60 cell extract 25 and incubated at 4°C overnight. The beads were washed 5 x with Hanks balanced salt solution, boiled 5 min in nonreducing SDS sample buffer and analyzed in a 10% SDS-polyacrylamide gel. The purified CD43 of 120 kDa in gel slices identified after autoradiography was excised and analyzed.

30 *β -Elimination of Radiolabeled O-Glycans from 3 H-PSGL-1 and 3 H-CD43 and Chromatography of Glycans on Bio-Gel and QAE-Sephadex* - The gel slices containing radiolabeled PSGL-1 and CD43 were directly

treated with mild base/borohydride as described [Iyer, et al. (1971) *Arch. Biochem. Biophys.* **142**, 101-105; Cummings, et al. (1983) *J. Biol. Chem.* **258**, 15261-15273]. Released radiolabeled glycans were desalted by chromatography on a column of Dowex 50 (H⁺ form) and the glycans 5 were further analyzed. Chromatography was performed on Bio-Gel P-4 (medium mesh) in a 1 x 90 cm column and on Bio-Gel P-10 (medium mesh) in a 1 x 48 cm column in 0.1 M pyridyl/acetate buffer, pH 5.5 and 1 ml fractions were collected. The anionic character and sialylation of 10 glycans was assessed partly by chromatography on column of QAE-Sephadex, with step elution in 2 mM Tris-base containing 20, 70, 140, 200, and 250 mM NaCl.

Preparation of Radiolabeled O-Glycan Standards - The simple core-1 O-glycan Gal β 1 \rightarrow 3GalNAc was radiolabeled by reduction in the presence of NaB[³H]₄ to form Gal β 1 \rightarrow 3GalNAcOH[³H] [Li, et al. (1978) 15 *J. Biol. Chem.* **253**, 7762-777]. A set of ³H-GlcN-labeled standard glycans was prepared from ³H-GlcN-labeled HL-60 cell total glycoproteins following previously described procedures [Carlsson, et al. (1986) *J. Biol. Chem.* **261**, 12787-12795]. The standards included the disialylated core-2 hexasaccharide 20 NeuAc α 2 \rightarrow 3Gal β 1 \rightarrow 4GlcNAc β 1 \rightarrow 6(NeuAc α 2 \rightarrow 3Gal β 1 \rightarrow 3)GalNAcOH, the neutral core-2 tetrasaccharide Gal β 1 \rightarrow 4GlcNAc β 1 \rightarrow 6(3Gal β 1 \rightarrow 3)GalNAcOH, and the trisaccharide GlcNAc β 1 \rightarrow 6(Gal β 1 \rightarrow 3)GalNAcOH. To prepare these standards, the ³H-GlcN-labeled cell extracts were dialyzed against 0.01 M NaHCO₃, pH 7.5 25 and treated with 100 μ g/ml TPCK-treated trypsin overnight at 37°C. The dried tryptic digest was treated with mild base/borohydride to effect β -elimination and the released O-glycans were size-fractionated on a column of Bio-Gel P-4. Pooled fractions corresponding to the peaks of radioactivity were analyzed for charge distribution by QAE-Sephadex 30 column chromatography. The structure of each glycan was then confirmed by sensitivity to *Arthrobacter* neuraminidase, β -galactosidase

-62-

and β -N-acetylhexosaminidase, as determined by descending paper chromatography as described below.

The core-2 fucosylated pentasaccharide standard $\text{Gal}\beta 1\rightarrow 4(\text{Fucal}\rightarrow 3)\text{GlcNAc}\beta 1\rightarrow 6(\text{Gal}\beta 1\rightarrow 3)\text{GalNAcOH}$ was prepared by
5 incubating the ^3H -GlcN-tetrasaccharide $\text{Gal}\beta 1\rightarrow 4\text{GlcNAc}\beta 1\rightarrow 6(\text{Gal}\beta 1\rightarrow 3)\text{GalNAcOH}$ with unlabeled GDP-Fuc (100 μM) in the presence of 50 μg of extract from COS7 cells stably transfected with the human FTIV gene. This $\alpha 1,3$ FT adds Fuc in a $\alpha 1,3$ linkage to GlcNAc residues in acceptors containing terminal N-
10 acetyllactosamine sequences [Goelz, et al. (1990) *Cell* **63**, 1349-1356; Lowe, et al. (1991) *J. Biol. Chem.* **266**, 21777-21783; Kumar, et al. (1991) *J. Biol. Chem.* **266**, 21777-21783]. The product of the reaction migrated as a pentasaccharide, as expected, and was converted to the corresponding tetrasaccharide by treatment with the *Streptomyces* $\alpha 1,3/4$ -fucosidase.
15

To define the Man:Fuc ratio and provide standard high mannose-type N-glycans, radiolabeled glycoproteins were prepared from [^3H]Man-labeled HL-60 cell extracts. For the preparation of glycopeptides, the cell extracts were precipitated with TCA and treated with Pronase as described
20 by Cummings, et al. (1983) *J. Biol. Chem.* **258**, 15261-15273. From these glycopeptides, high mannose-type N-glycan standards ($\text{Man}_9\text{GlcNAc}_1$, $\text{Man}_8\text{GlcNAc}_1$, $\text{Man}_7\text{GlcNAc}_1$, $\text{Man}_6\text{GlcNAc}_1$, $\text{Man}_5\text{GlcNAc}_1$) were prepared following treatment with endo- β -N-acetylglucosaminidase H [Cummings, R. D. (1993) *Glycobiology: A Practical Approach* (M. Fukuda and A. Kobata, eds), pp. 243-289, IRL Press at Oxford University Press, Oxford].

Relative Molar Ratios of Radiolabeled Monosaccharides in ^3H -GlcN-labeled Glycans - In the ^3H -GlcN-labeled, disialylated hexasaccharide
30 NeuAc $\alpha 2\rightarrow 3$ Gal $\beta 1\rightarrow 4$ GlcNAc $\beta 1\rightarrow 6$ (NeuAc $\alpha 2\rightarrow 3$ Gal $\beta 1\rightarrow 3$)GalNAcOH, prepared from HL-60 cells, the NeuAc, GlcNAc and GalNAcOH residues are radiolabeled [Varki, A. (1994) *Meth. Enzymol.* **230**, 16-32]. During

-63-

equilibrium radiolabeling the relative molar ratios for these residues should be close to unity. To determine the relative molar ratio for GlcNAc and GalNAc residues, this ³H-GlcN-hexasaccharide was desialylated to generate a ³H-GlcN-tetrasaccharide; this tetrasaccharide 5 was hydrolyzed in strong acid and the released radioactivity was identified by HPAEC as described below. All radioactivity was recovered in ³H-GlcN and ³H-GalNOH in the ratio of ³H-GlcN:³H-GalNOH of 1.0:0.8. This ratio was used as a correction factor for calculating relative molar 10 ratios of isolated glycans, i.e. the radioactivity recovered in ³H-GalN(OH) in a sample after hydrolysis was divided by 0.8. The relative molar ratio is consistent with other studies on HL-60 cells metabolically-radiolabeled with the ³H-GlcN precursor [Maemura, K. and Fukuda, M. (1992) *J. Biol. Chem.* **267**, 24379-24386]. To determine the relative molar ratio for NeuAc and GlcNAc residues, the ³H-GlcN-hexasaccharide was 15 desialylated with neuraminidase and the released radioactivity in NeuAc was determined. The radioactivity recovered in GlcN and NeuAc was in the ratio of 1.0:1.4, respectively. Since NeuAc radioactivity was derived from two residues of the disialylated hexasaccharide, this gave a final value for the relative molar ratio for GlcN:NeuAc of 1.0:0.7.

20 *Determination of Monosaccharide Composition* - The ratios of GlcN:GalN and Man:Fuc in PSGL-1 and CD43 were determined following strong acid hydrolysis of excised gel slices containing purified PSGL-1 and CD43. The gel slices were treated with 250 μ l 2N trifluoroacetic acid (TFA) at 121°C for 2 h. The released, radiolabeled 25 monosaccharides in the hydrolysate were identified by a pH anion exchange chromatography (HPAEC) on a CarboPac PA-1 column (4 x 250 mm) in a Dionex system and elution with 16 mM NaOH for 30 min. Fractions (0.3 min) were collected and radioactivity was determined by scintillation counting. The radiolabeled monosaccharides were identified 30 by their retention times compared to those of standard monosaccharides. The relative molar ratio for GlcN:GalN in glycans was calculated by dividing the radioactivity in the GlcN peak by the radioactivity in the

GalN peak and correcting for differences in specific activity of ^3H -GlcN versus ^3H -GalN. The Man:Fuc ratio in ^3H -Man-glycans was also determined following acid hydrolysis as described above, using a Man:Fuc ratio of 1.0:1.0, which is typically observed after equilibrium 5 radiolabelling of cells with [2- ^3H]Man.

Miscellaneous procedures - Enzymatic treatments of glycans with β -N-acetylhexosaminidase, β -galactosidase, *Arthrobacter* neuraminidase, *Streptomyces* α 1,3/4-fucosidase and *E. freundii* endo- β -galactosidase were performed as described by Cummings, et al. (1989) *Methods in Cell 10 Biol.* **32**, 142-180; Srivatsan, et al. (1992) *J. Biol. Chem.* **267**, 20196-20203. Digestion with NDV-neuraminidase was performed in 20 μl 10 mM phosphate, pH 7.0 with 20 mU of enzyme for 24 h at 37°C, followed by addition of another 20 mU of enzyme and further incubation for 24 h. O-glycans were analyzed and purified by descending paper 15 chromatography on Whatman filter paper for the times noted using the pyridine/ethyl acetate/water/acetic acid (5:5:1:3) solvent system, as described by Cummings, R. D. (1993) *Glycobiology*. Glycans were chemically defucosylated by treatment with 0.1 N TFA at 100°C for 1 h as described by Spooncer, et al. (1984) *J. Biol. Chem.* **259**, 4792-4801.

20 II. Results

Purification of PSGL-1 and CD43 - PSGL-1 and CD43 were purified from HL-60 cells metabolically labeled with ^3H -GlcN. PSGL-1 from ^3H GlcN-labeled HL-60 cell extracts was purified by affinity chromatography on immobilized P-selectin, following by re- 25 chromatography of the bound, EDTA-eluted fraction. Virtually all (90-99%) of the radiolabel in the 2X-purified material bound P-selectin. This two-step procedure was necessary to remove a contaminating glycoprotein of approximately 120 kDa under nonreducing conditions that remained after the first step. The purified PSGL-1 behaved as a dimer of 30 approximately 250 kDa in nonreducing conditions and approximately 120 kDa in reducing conditions. The purified PSGL-1 was resolved by SDS-

-65-

PAGE under reducing or nonreducing conditions in 7.5% polyacrylamide, and then analyzed by autoradiography.

CD43 3 H-GlcN-labeled HL-60 cell extracts were immunoprecipitated with a CD43-specific mAb plus a secondary rabbit

5 anti-mouse IgG1 antibody or secondary antibody alone. The immunoprecipitates were electrophoresed under nonreducing conditions in a 10% acrylamide gel and then analyzed by autoradiography. A single band of 120 kDa for CD43 was observed in both reducing and in nonreducing conditions.

10 *Composition of Radiolabeled Sugars in PSGL-1 and CD43* - In the initial assessment of the glycosylation of PSGL-1 and CD43, the ratio of GlcN:GalN in the purified glycoproteins was determined. 3 H-GlcN is metabolized by animal cells into radiolabeled GlcNAc, GalNAc and sialic acids. Gel slices containing the 3 H-GlcN-glycoproteins were treated with 15 strong acid (which results in destruction of sialic acids) and the radiolabeled GlcN and GalN were identified by Dionex HPAEC. The GlcN:GalN ratio was determined to be approximately 2:1 for PSGL-1 and approximately 1:1 for CD43. The GlcN:GalN ratio of approximately 1:1 for CD43 is consistent with published evidence that a majority of the 20 glycans in CD43 have the simple core-2 motif Gal β 1 \rightarrow 4GlcNAc β 1 \rightarrow 6 (Gal β 1 \rightarrow 3)GalNAcOH. These results demonstrate that the glycans of PSGL-1 contain higher amounts of GlcNAc relative to GalNAc than the glycans of CD43.

25 The presence of the sLe α determinant on PSGL-1 has led to expectations that PSGL-1 might be heavily fucosylated. The amount of Fuc present on PSGL-1 and CD43 isolated from HL-60 cells metabolically radiolabeled with [2- 3 H]Man was assessed. This labeled precursor is metabolized by cells to 2- 3 H-Fuc and the relative specific activity of Man and Fuc after equilibrium labeling is equivalent. 3 H-Man-30 PSGL-1 and -CD43 were isolated by SDS-PAGE and autoradiography. The corresponding bands were subjected to strong acid hydrolysis and the released monosaccharides were separated by HPAEC on a Dionex system.

-66-

The Man:Fuc ratio was determined to be 3:5 for PSGL-1 and 3:2 for CD43. As a control, the Man:Fuc ratio was also determined for the total unpurified glycoproteins from HL-60 cells and found to be 3:1. The relative amounts of ³H-Man and ³H-Fuc in PSGL-1, CD43, and total HL-5 60 glycoproteins was determined after acid hydrolysis of ³H-Man-PSGL-1, ³H-Man-CD43, and total ³H-Man-labeled HL-60 cell glycoproteins. Acid hydrolysates were analyzed by HPAEC using a Carbo-Pac PA-1 column eluted with 16 mM NaOH. The total radioactivity in each shaded peak was determined. PSGL-1 contains more Fuc residues than CD43 10 and more Fuc residues than average glycoproteins in HL-60 cells.

Using this information it is possible to estimate the number of Fuc residues on PSGL-1. The cDNA sequence of PSGL-1 predicts that PSGL-1 has three potential N-glycosylation sites. PSGL-1 contains only complex-type N-glycans, each of which should have 3 Man residues. 15 Thus, 3 complex-type N-glycans on PSGL-1 represent 9 Man residues per mol and, correspondingly, there are approximately 15 Fuc residues per mol of PSGL-1. In contrast, CD43 contains only a single N-linked glycan and has much less Fuc in comparison to PSGL-1. Taken together, the compositional analyses from ³H-Man-glycoproteins demonstrate that 20 PSGL-1 is glycosylated differently than CD43.

β-Elimination of O-Glycans from PSGL-1 and CD43 and Chromatography on Bio-Gel P-4 and P-10 — The O-glycans of PSGL-1 and CD43 were directly released by treating gel slices containing purified glycoproteins with mild base-borohydride to effect β-elimination and were 25 fractionated on a column of Bio-Gel P-4. The released ³H-GlcN-glycans from both PSGL-1 and CD43 were recovered in three fractions represented 47%, 41% and 11% of the total radioactivity, respectively. Compared to CD43, PSGL-1 contains a high proportion of relatively large glycans.

30 The P-4-I samples from both PSGL-1 and CD43 were further purified by chromatography on a column of Bio-Gel P-10. Radioactivity in the P-4-I fraction from PSGL-1 separated into two major peaks on Bio-

Gel P-10 designated P-10-1 and P-10-2. This population of glycans is larger in size than the disialylated core-2 hexasaccharide standard NeuAc α 2 \rightarrow 3 Gal β 1 \rightarrow 4GlcNAc β 1 \rightarrow 6(NeuAc α 2 \rightarrow 3Gal β 1 \rightarrow 3)GalNAcOH. The major material in the P-4-II fraction from PSGL-1 eluted on Bio-Gel 5 P-10 in a position slightly smaller (designated P-10-3) than the disialylated core-2 hexasaccharide standard. In contrast, most of the material in both the P-4-I and P-4-II fractions of CD43 eluted identically on Bio-Gel P-10 as the disialylated core-2 hexasaccharide standard.

The glycans recovered in P-4-I from CD43 were analyzed using 10 exoglycosidase treatments and anion exchange chromatography. The glycans were shown to be the expected disialylated core-2 hexasaccharide. A small fraction of the P-4-I sample from CD43 was recovered in larger-sized glycans on Bio-Gel P-10, consistent with previous studies showing that a small fraction of O-glycans from CD43 have an extended 15 polylactosamine structure on the core-2 motif. Since the structures of glycans in CD43 have been described, they were not further analyzed.

Compositional Analyses of P-10-1, P-10-2 and P-10-3 Glycans from PSGL-1 — O-glycans released from Ser/Thr residues by β -elimination should contain 3 H-GalNAcOH at the reducing terminus, which 20 is recoverable as 3 H-GalNOH following strong acid hydrolysis. To identify which glycans in the mixture from PSGL-1 represent the O-glycans, the GalNOH, GalN and GlcN content of each glycan from Bio-Gel P-10 was determined by HPAEC on a Dionex system following strong acid hydrolysis of the 3 H-GlcN-PSGL-1 P-10-1, P-10-2, and P-10-25 3 fractions. Acid hydrolysates were analyzed by HPAEC using a Carbo-Pac PA-1 column eluted with 16 mM NaOH. The total radioactivity in each peak was determined. More than 95% of the total GalN(OH) in PSGL-1 P-10-2 and P-10-3 fractions was recovered as GalNOH, demonstrating that β -elimination was efficient and that the O-glycans of 30 PSGL-1 are represented in these fractions. The glycans in P-10-1 lack GalNOH and do not represent O-glycans released by β -elimination. Instead, the P-10-1 fraction contains N-glycans still attached to the

peptide. This was confirmed by the presence of ^3H -Man recovered in these glycans from ^3H -Man-PSGL-1. The small amount of GalN recovered in P-10-1 samples might arise from GalNAc residues present in N-glycans, or it could result from a small amount of residual O-glycans still linked to peptide and not released during β -elimination procedures.

Sialylation of P-10-1, P-10-2 and P-10-3 Glycans from PSGL-1 —
The sialylation patterns of the O-glycans in P-10-2 and P-10-3 and of the N-glycans in P-10-1 were determined by an anion-exchange column chromatography on QAE-Sephadex, before and after neuraminidase treatment. In this system, glycans with 1 negative charge (1 sialic acid) elute with 20 mM NaCl, those with 2 negative charges (2 sialic acids) elute with 70 mM NaCl, and those with 3 negative charges (3 sialic acids) elute with 140 mM NaCl [24,25]. ^3H -GlcN-PSGL-1 P-10-1, P-10-2, and P-10-3 fractions were applied to columns of QAE-Sephadex and the columns were eluted with the indicated concentrations of NaCl (mM) either before or after treatment with Newcastle disease virus (NDV) neuraminidase. The glycans in the P-10-1 fraction were heterogeneously charged, consistent with the occurrence of N-glycans in glycopeptides in this fraction. Following treatment with NDV neuraminidase, the radioactivity eluted with 20 mM NaCl was analyzed by descending paper chromatography for 20 h. Authentic NeuAc migrated 25 cm. The P-10-2 glycans were mono-and disialylated species and the P-10-3 glycans were a mixture of neutral and monosialylated species.

To determine whether the anionic character of the glycans was due to sialic acid and to define the linkage of sialic acid, portions of the ^3H -GlcN-labeled O-glycans were treated with neuraminidase from Newcastle Disease virus (NDV). This enzyme displays high specificity for α 2,3-linked sialic acid residues and will not efficiently cleave sialic acid in other linkages [Paulson, et al. (1982) *J. Biol. Chem.* **257**, 12734-12738]. After NDV-neuraminidase treatment, the glycans in P-10-1 were less charged, consistent with a loss of sialic acid. However, the presence of residually charged glycans is indicative of the profile expected for N-

glycans in glycopeptides. In contrast, the glycans in P-10-2 and P-10-3 became neutral following NDV-neuraminidase treatment, demonstrating that all sialic acids in these glycans are α 2,3-linked. The peak of material eluting with 20 mM NaCl following NDV-neuraminidase treatment was 5 quantitatively recovered as free 3 H-NeuAc, as shown by its co-elution with standard NeuAc on descending paper chromatography. It was previously established that the sialic acid on PSGL-1 from 3 H-GlcN-labeled HL-60 cells is Neu5Ac. These results demonstrate that the O-glycans in P-10-2 are mono-and disialylated species and that the O-glycans in P-10-3 are a combination of neutral and sialylated species, with 10 sialic acid in α 2,3-linkage to glycans. Furthermore, these results demonstrate that the O-glycans of PSGL-1 are not sulfated, since neutral species result following treatment with NDV-neuraminidase.

Descending Paper Chromatography of P-10-2 Glycans from PSGL-1 — The glycans in P-10-2 were further purified using preparative descending paper chromatography and two major species were recovered. One peak contained larger-sized glycans that migrated slowly from the origin (designated P-10-2a). A small peak of faster migrating material (approximately 24 cm) was also observed, which represented some 20 residual glycans derived from the P-4-II peak. Upon anion-exchange chromatography on QAE-Sephadex, the P-10-2a material eluted as monosialylated species, whereas the P-10-2b material eluted as disialylated species.

Exoglycosidase Treatments of P-10-2b Glycans — The smaller-sized, disialylated glycans in P-10-2b were desialylated by treatment with NDV-neuraminidase, and released sialic acid was removed by chromatography on QAE-Sephadex. The desialylated glycans were analyzed by descending paper chromatography before and after sequential exoglycosidase treatments (Figures 4A-E). Following treatment with 25 neuraminidase, the desialylated P-10-2b glycans fractionated as two species (Figure 4A). A minor peak (approximately 10%) co-migrated with the fucosylated pentasaccharide standard

-70-

Gal β 1 \rightarrow 4(Fuc β 1 \rightarrow 3)GlcNAc β 1 \rightarrow 6 (Gal β 1 \rightarrow 3)GalNAcOH and was designated P-10-2b₁. A major peak (approximately 90%) of material co-migrated with the standard core-2 tetrasaccharide

Gal β 1 \rightarrow 4GlcNAc β 1 \rightarrow 6(Gal β 1 \rightarrow 3)GalNAcOH and was designated P-10-

5 2b₂.

The desialylated P-10-2b glycans were treated with β -N-acetylhexosaminidase. This treatment did not alter the migration of either species in the sample, indicating that the glycans lack terminal GlcNAc residues. However, when the desialylated P-10-2b mixture was treated

10 with β -galactosidase, P-10-2b₁ glycans were unaffected, whereas P-10-2b₂ glycans were degraded and co-migrated with the standard trisaccharide

GlcNAc β 1 \rightarrow 6(Gal β 1 \rightarrow 3)GalNAcOH (Figure 4B). The

Gal β 1 \rightarrow 3GalNAcOH structure is resistant to jack bean β -galactosidase, since the enzyme does not efficiently cleave terminal galactosyl residues

15 in β 1,3 linkage [Li, et al. (1975) *J. Biol. Chem.* 250, 6786-6791]. This was confirmed in control studies, in which no cleavage of the standard disaccharide Gal β 1 \rightarrow 3GalNAcOH occurred with the concentrations of jack bean β -galactosidase used in these analyses. Since the P-10-2b₂ glycans are derived from disialylated species, these results demonstrate that the P-

20 10-2b₂ glycans are derived from the core-2 hexasaccharide

NeuAc α 2 \rightarrow 3Gal β 1 \rightarrow 4GlcNAc β 1 \rightarrow 6 (NeuAc α 2 \rightarrow 3Gal β 1 \rightarrow 3)GalNAcOH.

When the disialylated ³H-GlcN-P-10-2b glycans were treated with the *Streptomyces* α 1,3/4-fucosidase, the P-10-2b₁ glycans were lost and all recovered glycans co-migrated with the core-2 tetrasaccharide standard

25 Gal β 1 \rightarrow 4GlcNAc β 1 \rightarrow 6(Gal β 1 \rightarrow 3)GalNAcOH (Figure 4C). Combined


treatment of the desialylated P-10-2b glycans with *Streptomyces* α 1,3/4-fucosidase, β -N-acetylhexosaminidase and β -galactosidase resulted in complete degradation of both P-10-2b₁ and P-10-2b₂ glycans to free ³H-GlcNAc and the ³H-disaccharide Gal β 1 \rightarrow 3GalNAcOH (Figure 4D). These

30 results demonstrate that the P-10-2b₁ glycans have the fucosylated

pentasaccharide structure

Gal β 1 \rightarrow 4(Fuc α 1 \rightarrow 3)GlcNAc β 1 \rightarrow 6(Gal β 1 \rightarrow 3)GalNAcOH. Since the

original P-10-2b glycans are desialylated species, the P-10-2b₁ glycans contain the sLe^x antigen NeuAc α 2 \rightarrow 3Gal β 1 \rightarrow 4(Fuc α 1 \rightarrow 3)GlcNAc β 1 \rightarrow R and have the heptasaccharide structure



5 (NeuAc α 2 \rightarrow 3Gal β 1 \rightarrow 3)GalNAcOH (Figure 7).

To further confirm the presence of fucose in these glycans and to facilitate the identification of Fuc in other glycans, HL-60 cells were metabolically radiolabeled with [³H-Fuc]. The ³H-Fuc-labeled O-glycans were recovered by β -elimination as described for the ³H-GlcN-glycans, following the same procedures described above. The ³H-Fuc recovered in the desialylated P-10-2b fraction co-migrated with the desialylated ³H-GlcN-P-10-2b₁ glycans (Figure 4E). Taken together, these results demonstrate that the P-10-2b₁ glycans from PSGL-1 are fucosylated and contain the sLe^x structure.

15 *Endo- and Exoglycosidase Treatments of P-10-2a Glycans —*

Because of their relatively large size, the possibility that the P-10-2a glycans contain polylactosamine [3Gal β 1 \rightarrow 4GlcNAc β 1-]_n was considered.

To assess this possibility, the P-10-2a glycans were desialylated and the sialic acid was removed by QAE-Sephadex chromatography. The

20 resulting neutral glycans were treated with endo- β -galactosidase, an enzyme that cleaves internal β 1 \rightarrow 4 galactosyl residues within a type 2 polylactosamine [Kobata and Takasaki (1993) *Glycobiology: A Practical Approach* (M. Fukuda and A. Kobata, eds), pp. 165-185, IRL Press at Oxford University Press, Oxford]. The glycans were resistant to this


25 treatment (Figure 5A). The desialylated P-10-2a glycans were also resistant to combined treatment with β -galactosidase and β -N-acetylhexosaminidase (Figure 5A). The possibility that the P-10-2a glycans might contain a polylactosamine backbone in which all internal GlcNAc residues are fucosylated was then considered. Such

30 polyfucosylated, polylactosamine structures are resistant to endo- β -galactosidase [Fukuda, et al. (1984) *J. Biol. Chem.* **259**, 10925-10935].

-72-

Complete defucosylation is necessary before endo- β -galactosidase can digest the chains.

The desialylated P-10-2a glycans were chemically defucosylated and then treated with endo- β -galactosidase. After defucosylation, the 5 enzyme quantitatively digested the P-10-2a glycans to release three major compounds in an approximate equimolar ratio identified as the trisaccharide $\text{Gal}\beta 1\rightarrow 4\text{GlcNAc}\beta 1\rightarrow 3\text{Gal}$, the residual core-2 trisaccharide $\text{GlcNAc}\beta 1\rightarrow 6(\text{Gal}\beta 1\rightarrow 3)\text{GalNAcOH}$, and the disaccharide $\text{GlcNAc}\beta 1\rightarrow 3\text{Gal}$ (Figure 5B). The generation of such products from the 10 specific action of endo- β -galactosidase is predicted for a glycan with the backbone structure:



where the arrows indicate specific cleavage sites for endo- β -galactosidase of a defucosylated polylactosamine-containing glycan [Kobata and Takasaki (1993) *Glycobiology*]. The length of the polylactosamine chain 15 can be deduced from the radioactivity recovered in the disaccharide $\text{GlcNAc}\beta 1\rightarrow 3\text{Gal}$ relative to the other fragments. The recovery of the trisaccharide $\text{GlcNAc}\beta 1\rightarrow 6(\text{Gal}\beta 1\rightarrow 3)\text{GalNAcOH}$ following endo- β -galactosidase treatment demonstrates that the polylactosamine is extended from the GlcNAc in $\beta 1\text{-}6$ linkage to the GalNAcOH residue. The 20 recovery of the $\text{Gal}\beta 1\rightarrow 4\text{GlcNAc}\beta 1\rightarrow 3\text{Gal}$ demonstrates that Gal residues are present in the nonreducing terminus of the polylactosamine. As expected, treatment of the desialylated and defucosylated P-10-2a glycans with a combination of β -galactosidase and β -N-acetylhexosaminidase resulted in complete digestion to free $^3\text{H}\text{-GlcNAc}$ and the ^3H -disaccharide 25 $\text{Gal}\beta 1\rightarrow 3\text{GalNAcOH}$ (Figure 4B).

The resistance of the desialylated P-10-2a glycans to endo- β -galactosidase prior to defucosylation is consistent with the possibility that each GlcNAc residue within the glycan contains an $\alpha 1,3$ -linked fucosyl residue in the structure



30 $\text{Gal}\beta 1\rightarrow 4(\text{Fuc}\alpha 1\rightarrow 3)\text{GlcNAc}\beta 1\rightarrow 3\text{Gal}\beta 1\rightarrow 4(\text{Fuc}\alpha 1\rightarrow 3)\text{GlcNAc}\beta 1\rightarrow 3\text{Gal}\beta 1\rightarrow$

-73-

4(Fuc α 1 \rightarrow 3)GlcNAc β 1 \rightarrow 6(Gal β 1 \rightarrow 3)GalNAcOH. O-glycans containing incompletely fucosylated polylactosamines are sensitive to endo- β -galactosidase.

To confirm the fucosylation pattern predicted for the P-10-2a glycans, the ^3H -Fuc-P-10-2a glycans were prepared from ^3H -Fuc-PSGL-1. Following treatment with neuraminidase, the desialylated ^3H -Fuc-glycans co-migrated with the desialylated ^3H -GlcN-P-10-2a glycans. When the desialylated ^3H -Fuc-P-10-2a glycans were treated with the *Streptomyces* α 1,3/4 fucosidase, approximately one-third of the radioactive Fuc was released. This result is predicted based on the postulated glycan structure and on the specificity of the *Streptomyces* α 1,3/4 fucosidase, which can remove Fuc only from penultimate GlcNAc residues within a polyfucosylated glycan [Maemura and Fukuda (1992) *J. Biol. Chem.* **267**, 24379-24386, Sano, et al. (1992) *J. Biol. Chem.* **267**, 1522-1527]. Taken together, these data demonstrate that the P-10-2a glycans contain three Fuc residues, one of which is in the terminal lactosaminyl unit.

The P-10-2a glycans are monosialylated and could have one of two possible structures (a or b) shown below:



35 To distinguish between these possibilities, the ^3H -GlcN-P-10-2a sialylated glycans were treated with a mixture of β -galactosidase, β -N-acetylhexosaminidase, and *Streptomyces* $\alpha 1,3/4$ fucosidase with or without

-74-

neuraminidase. It was predicted that structure (a) would require neuraminidase in addition to the other enzymes for complete digestion, whereas structure (b) would be degraded by the exoglycosidases in the absence of neuraminidase. Treatment of the ^3H -GlcN-P-10-2a with 5 neuraminidase alone, released radiolabeled NeuAc. When the glycans were treated with a mixture of β -galactosidase, β -N-acetylhexosaminidase, and *Streptomyces* α 1,3/4 fucosidase, in the absence of neuraminidase, no radioactivity was released. Inclusion of neuraminidase with other exoglycosidases resulted in complete degradation of the glycans to free 10 ^3H -NeuAc, free ^3H -GlcNAc and the ^3H -disaccharide $\text{Gal}\beta 1\rightarrow 3\text{GalNAcOH}$. Taken together, these results demonstrate that the single sialic acid residue on the P-10-2a glycans is present in a terminal position on the polylactosamine chain, shown in structure (a), and that these glycans have the sLe x structure (Figure 7).

15 *Endo- and Exoglycosidase Treatments of P-10-3 Glycans* — The P-10-3 glycans were primarily neutral or monosialylated species. the predominant species (approximately 80% of the radioactivity) was non-sialylated. The P-10-3 fraction was treated with neuraminidase, and released sialic acid was removed by chromatography on the QAE- 20 Sephadex. Approximately 90% of the desialylated species co-migrated with the standard core-2 tetrasaccharide $\text{Gal}\beta 1\rightarrow 4\text{GlcNAc}\beta 1\rightarrow 6(\text{Gal}\beta 1\rightarrow 3)\text{GalNAcOH}$ and the remainder co-migrated with the core-1 disaccharide $\text{Gal}\beta 1\rightarrow 3\text{GalNAcOH}$ (Figure 6). Treatment of the desialylated P-10-3 glycans with β -galactosidase caused 25 a shift in the migration of the major peak to that of the expected trisaccharide standard (Figure 6). Treatment with a combination of β -galactosidase and β -N-acetylhexosaminidase generated free ^3H -GlcNAc and ^3H - $\text{Gal}\beta 1\rightarrow 3\text{GalNAcOH}$ (Figure 6). Approximately 20% of the radiolabel in the P-10-3 glycans was found in predominately 30 monosialylated species that were converted to neutral species by neuraminidase treatment. These results demonstrate that the P-10-3 glycans are primarily neutral core-2 tetrasaccharides and some

-75-

monosialylated core-2 pentasaccharides (Figure 7). In addition, some of the neutral glycans have the core-2 disaccharide structure (Figure 7).

Relative Percentages of O-Glycans In PSGL-1 from HL-60 Cells

— The relative percentages of O-Glycans in PSGL-1 can be calculated from the amount of radioactivity recovered in each O-glycan and from the relative molar ratio for $^3\text{H-GlcN} : ^3\text{H-GalNOH} : ^3\text{H-NeuAc}$ of 1.0:0.8:0.7 (Figure 7). The majority of O-glycans in PSGL-1 contain the core-2 structure. A minority of the glycans, recovered in P-10-2b₁ and P-10-2a, contain the sLe^x determinant.

10 III. Conclusions.

This study demonstrates that PSGL-1 from human HL-60 cells contains O-glycans with a core-2 motif. A majority of the glycans are not fucosylated and are mixtures of neutral and sialylated species. A minority of the O-glycans (approximately 14%) are fucosylated and contain the terminal sLe^x structure. Two types of fucosylated O-glycans are present; one type is a disialylated heptasaccharide lacking polylactosamine and the other is a unique monosialylated, tri-fucosylated glycan containing polylactosamine. The presence of core-2 O-glycans on PSGL-1 from HL-60 cells is consistent with results of studies on the glycosylation of PSGL-1 from human neutrophils. Desialylated PSGL-1 from both human neutrophils and HL-60 cells is resistant to treatment with endo- α -N-acetylgalactosaminidase (O-glycanase), which cleaves only desialylated core-1 glycans. The direct demonstration of sLe^x determinants and polylactosamine on O-glycans of HL-60-derived PSGL-1 reinforces indirect evidence that these structures are present on O-glycans from neutrophil-derived PSGL-1.

Significant differences were observed in the O-glycans on PSGL-1 and CD43 from HL-60 cells. Although both proteins have primarily core-2 O-glycans, PSGL-1 has many neutral core-2 tetrasaccharides, whereas CD43 has mostly disialylated, core-2 hexasaccharides. The core-2 structure is a precursor for polylactosamine synthesis in O-glycans, but only PSGL-1 has significant amounts of polylactosamine. This indicates

-76-

that the core-2 structure is necessary but not sufficient for polygalactosamine addition. Furthermore, CD43 lacks the two species of fucosylated O-glycans found in PSGL-1. Although a monofucosylated O-glycan was identified in CD43, this species represents only 0.5% of the O-glycans in

5 CD43. The tri-fucosylated monosialylated O-glycan identified on PSGL-1 is not found on CD43.

As discussed above, P-selectin can bind weakly to a variety of sulfated glycans and these glycans inhibit binding of P-selectin to human myeloid cells. However, the O-glycans of PSGL-1 are not sulfated.

10 Instead, PSGL-1 contains tyrosine sulfate that is required for interactions with P-selectin but not with E-selectin. Three consensus sites for tyrosine sulfation occur at the amino terminus of PSGL-1 at residues 46, 48 and 51. PL1, a mAb to PSGL-1, blocks binding to P-selectin and recognizes an epitope spanning residues 49-62 that overlaps the tyrosine sulfation

15 sites. Near the tyrosine sulfation sites are two Thr residues that represent potential O-glycosylation sites at residues 44 and 57. Mutations in these residues reduce binding of PSGL-1 to P-selectin when PSGL-1 is co-expressed in COS cells with either FTIII or FTVII. These results suggest that only one or two O-glycans in conjunction with tyrosine sulfate

20 residues may be sufficient to promote high affinity binding of PSGL-1 to P-selectin. However, O-glycans in other regions of the molecule may also contribute to interactions with P- and E-selectin.

It was originally suggested that mucin-like glycoproteins act as convenient scaffolds upon which many O-glycans can be clustered for

25 recognition by selectins. However, this data, in conjunction with other studies, indicate that mucin-like glycoproteins are differentially glycosylated.

-77-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Board of Regents of the University of Oklahoma

(ii) TITLE OF INVENTION: Peptide and O-Glycan Inhibitors of Selectin Mediated Inflammation

(iii) NUMBER OF SEQUENCES: 5

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Patrea L. Pabst

(B) STREET: 2800 One Atlantic Center, 1201 West Peachtree Street

(C) CITY: Atlanta

(D) STATE: Georgia

(E) COUNTRY: US

(F) ZIP: 30306-3450

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

(D) SOFTWARE: PatentIn Release #1.0, Version #1.25

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Pabst, Patrea L.

(B) REGISTRATION NUMBER: 31,284

(C) REFERENCE/DOCKET NUMBER: OMRF110CIP7

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (404) 873-8794

(B) TELEFAX: (404) 873-8795

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 402 amino acids

(B) TYPE: amino acid

-78-

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Met Pro Leu Gln Leu Leu Leu Leu Leu Ile Leu Leu Gly Pro Gly Asn
1 5 10 15

Ser Leu Gln Leu Trp Asp Thr Trp Ala Asp Glu Ala Glu Lys Ala Leu
20 25 30

Gly Pro Leu Leu Ala Arg Asp Arg Arg Gln Ala Thr Glu Tyr Glu Tyr
35 40 45

Leu Asp Tyr Asp Phe Leu Pro Glu Thr Glu Pro Pro Glu Met Leu Arg
50 55 60

Asn Ser Thr Asp Thr Thr Pro Leu Thr Gly Pro Gly Thr Pro Glu Ser
65 70 75 80

Thr Thr Val Glu Pro Ala Ala Arg Arg Ser Thr Gly Leu Asp Ala Gly
85 90 95

Gly Ala Val Thr Glu Leu Thr Thr Glu Leu Ala Asn Met Gly Asn Leu
100 105 110

Ser Thr Asp Ser Ala Ala Met Glu Ile Gln Thr Thr Gln Pro Ala Ala
115 120 125

Thr Glu Ala Gln Thr Thr Pro Leu Ala Ala Thr Glu Ala Gln Thr Thr
130 135 140

Arg Leu Thr Ala Thr Glu Ala Gln Thr Thr Pro Leu Ala Ala Thr Glu
145 150 155 160

Ala Gln Thr Thr Pro Pro Ala Ala Thr Glu Ala Gln Thr Thr Gln Pro
165 170 175

Thr Gly Leu Glu Ala Gln Thr Thr Ala Pro Ala Ala Met Glu Ala Gln
180 185 190

-79-

Thr Thr Ala Pro Ala Ala Met Glu Ala Gln Thr Thr Pro Pro Ala Ala
195 200 205

Met Glu Ala Gln Thr Thr Gln Thr Thr Ala Met Glu Ala Gln Thr Thr
210 215 220

Ala Pro Glu Ala Thr Glu Ala Gln Thr Thr Gln Pro Thr Ala Thr Glu
225 230 235 240

Ala Gln Thr Thr Pro Leu Ala Ala Met Glu Ala Leu Ser Thr Glu Pro
245 250 255

Ser Ala Thr Glu Ala Leu Ser Met Glu Pro Thr Thr Lys Arg Gly Leu
260 265 270

Phe Ile Pro Phe Ser Val Ser Ser Val Thr His Lys Gly Ile Pro Met
275 280 285

Ala Ala Ser Asn Leu Ser Val Asn Tyr Pro Val Gly Ala Pro Asp His
290 295 300

Ile Ser Val Lys Gln Cys Leu Leu Ala Ile Leu Ile Leu Ala Leu Val
305 310 315 320

Ala Thr Ile Phe Phe Val Cys Thr Val Val Leu Ala Val Arg Leu Ser
325 330 335

Arg Lys Gly His Met Tyr Pro Val Arg Asn Tyr Ser Pro Thr Glu Met
340 345 350

Val Cys Ile Ser Ser Leu Leu Pro Asp Gly Gly Glu Gly Pro Ser Ala
355 360 365

Thr Ala Asn Gly Gly Leu Ser Lys Ala Lys Ser Pro Gly Leu Thr Pro
370 375 380

Glu Pro Arg Glu Asp Arg Glu Gly Asp Asp Leu Thr Leu His Ser Phe
385 390 395 400

Leu Pro

(2) INFORMATION FOR SEQ ID NO:2:

-80-

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2042 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

AGGAAACCTG CCATGGCCTC CTGGTGAGCT GTCCTCATCC ACTGCTCGCT GCCTCTCCAG	60
ATACTCTGAC CCATGGATCC CCTGGGTGCA GCCAAGGCCAC AATGGCCATG GCGCCGCTGT	120
CTGGCCGCAC TGCTATTCA GCTGCTGGTG GCTGTGTGTT TCTTCTCCTA CCTGCGTGTG	180
TCCCGAGACG ATGCCACTGG ATCCCCTAGG GCTCCCAGTG GGTCCCTCCCG ACAGGACACC	240
ACTCCCACCC GCCCCACCCCT CCTGATCCTG CTATGGACAT GGCCTTTCCA CATCCCTGTG	300
GCTCTGTCCC GCTGTTCAGA GATGGTGCCTC GGCACAGCCG ACTGCCACAT CACTGCCGAC	360
CGCAAGGTGT ACCCACAGGC AGACACGGTC ATCGTGCACC ACTGGGATAT CATGTCCAAC	420
CCTAAGTCAC GCCTCCCACC TTCCCCGAGG CCGCAGGGGC AGCGCTGGAT CTGGTTCAAC	480
TTGGAGCCAC CCCCTAACTG CCAGCACCTG GAAGCCCTGG ACAGATACTT CAATCTCACC	540
ATGTCTTACCC GCAGCGACTC CGACATCTTC ACGCCCTACG GCTGGCTGGA GCCGTGGTCC	600
GGCCAGCCTG CCCACCCACC GCTCAACCTC TCGGCCAAGA CCGAGCTGGT GGCCCTGGCG	660
GTGTCCAAGT GGAAGCCGGA CTCAGCCAGG GTGCGCTACT ACCAGAGCCT GCAGGCTCAT	720
CTCAAGGTGG ACGTGTACGG ACGCTCCCAC AAGCCCTGC CCAAGGGGAC CATGATGGAG	780
ACGCTGTCCC GGTACAAGTT CTACCTGGCC TTGAGAACT CCTTGCACCC CGACTACATC	840
ACCGAGAACG TGTGGAGGAA CGCCCTGGAG GCCTGGCCG TGCCCGTGGT GCTGGGCC	900
AGCAGAACG ACTACGAGAG GTTCCCTGCCA CCCGACGCCT TCATCCACGT GGACGACTTC	960
CAGAGCCCCA AGGACCTGGC CCGGTACCTG CAGGAGCTGG ACAAGGACCA CGCCCGCTAC	1020

-81-

CTGAGCTACT TTGCTGGCG GGAGACGCTG CGGCCTCGCT CCTTCAGCTG GGACTGGATT	1080
TCTGCAAGGC CTGCTGGAAA CTGCAGCAGG AATCCAGGTA CCAGACGGTG CGCAGCATAG	1140
CGGCTTGGTT CACCTGAGAG GCCGGCATGG TGCCTGGCT GCCGGGAACC TCATCTGCCT	1200
GGGGCCTCAC CTGCTGGAGT CCTTTGTGGC CAACCCTCTC TCTTACCTGG GACCTCACAC	1260
GCTGGGCTTC ACGGCTGCCA GGAGCCTCTC CCCTCCAGAA GACTTGCCTG CTAGGGACCT	1320
CGCCTGCTGG GGACCTCGCC TGTTGGGAC CTCACCTGCT GGGGACCTCA CCTGCTGGGG	1380
ACCTTGGCTG CTGGAGGCTG CACCTACTGA GGATGTCGGC GGTCGGGAC TTTACCTGCT	1440
GGGACCTGCT CCCAGAGACC TTGCCACACT GAATCTCACC TGCTGGGAC CTCACCCCTGG	1500
AGGGCCCTGG GCCCTGGGA ACTGGCTTAC TTGGGGCCCC ACCCGGGAGT GATGGTTCTG	1560
GCTGATTGT TTGTGATGTT GTTAGCCGCC TGTGAGGGGT GCAGAGAGAT CATCACGGCA	1620
CGGTTCCAG ATGTAATACT GCAAGGAAAA ATGATGACGT GTCTCCTCAC TCTAGAGGGG	1680
TTGGTCCCAT GGGTTAAGAG CTCACCCAG GTTCTCACCT CAGGGTTAA GAGCTCAGAG	1740
TTCAGACAGG TCCAAGTTCA AGCCCAGGAC CACCACTTAT AGGGTACAGG TGGGATCGAC	1800
TGTAAATGAG GACTTCTGGA ACATTCCAAA TATTCTGGGG TTGAGGGAAA TTGCTGCTGT	1860
CTACAAAATG CCAAGGGTGG ACAGGCCTG TGGCTCACGC CTGTAATTCC AGCACTTTGG	1920
GAGGCTGAGG TAGGAGGATT GATTGAGGCC AAGAGTTAAA GACCAGCCTG GTCAATATAG	1980
CAAGACCACG TCTCTAAATA AAAAATAATA GGCCGGCCAG GAAAAAAA AAAAAAAA	2040
AA	2042

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2102 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GTGAAGTGCT CAGAATGGGG CAGGATGTCA CCTGGAATCA GCACTAAGTG ATTCA	60
GACTTTACTTT TAAATGTGCG CTCTTCATTT CAAGATGCCG TTGCAGCTCT GATAAATGCA	120
AACTGACAAC CTTCAAGGCC ACGACGGAGG GAAAATCATT GGTGCTTGGAA GCATAGAAGA	180
CTGCCCTTCA CAAAGGAAAT CCCTGATTAT TGTTGAAAT GCTGAGGACG TTGCTGCGAA	240
GGAGACTTTT TTCTTATCCC ACCAAATACT ACTTTATGGT TCTTGTTTA TCCCTAATCA	300
CCTTCTCCGT TTTAAGGATT CATCAAAAGC CTGAATTGT AAGTGTAGA CACTTGGAGC	360
TTGCTGGGA GAATCCTAGT AGTGATATTA ATTGCACCAA AGTTTACAG GGTGATGTAA	420
ATGAAATCCA AAAGGTAAAG CTTGAGATCC TAACAGTGAA ATTTAAAAG CGCCCTCGGT	480
GGACACCTGA CGACTATATA AACATGACCA GTGACTGTTA TTCTTCATC AAGAGACGCA	540
AATATATTGT AGAACCCCTT AGTAAAGAAG AGGCGGAGTT TCCAATAGCA TATTCTATAG	600
TGGTTCATCA CAAGATTGAA ATGCTTGACA GGCTGCTGAG GGCCATCTAT ATGCCTCAGA	660
ATTTCTATTG CGTTCATGTG GACACAAAAT CCGAGGATTG CTATTTAGCT GCAGTGATGG	720
GCATCGCTTC CTGTTTTAGT AATGCTTTG TGGCCAGCCG ATTGGAGAGT GTGGTTTATG	780
CATCGTGGAG CCGGGTTCAAG GCTGACCTCA ACTGCATGAA GGATCTCTAT GCAATGAGTG	840
CAAACGGAA GTACTTGATA AATCTTGAGT GTATGGATTT TCCCATTAAA ACCAACCTAG	900
AAATTGTCAG CAAGCTCAAG TTGTTAATGG GAGAAAACAA CCTGGAAACG GAGAGGATGC	960
CATCCCATAA AGAACAAAGG TGGAAGAACG CCTATGAGGT CGTTAATCGA AAGCTGACAA	1020
ACACAGGGAC TGTCAAAATG CTTCCCTCAC TCGAAACACC TCTCTTTCT GGCAGTGCCT	1080
ACTTCGTGCT CAGTAGGGAC TATGTGGGGT ATGTACTACA GAATGAAAAA ATCCAAAAGT	1140
TGATGGAGTG GGCACAAAGAC ACATACAGCC CTGATGAGTA TCTCTGCGCC ACCATCCAA	1200
GGATTCCCTGA AGTCCCGGCC TCACTCCCTG CCAGCCATAA GTATGATCTA TGTGACATGC	1260

-83-

AAGCAGTTGC CAGGTTGTC AAGTGGCACT ACTTTCAGGG TGATGTTCC AAGGGTGCTC	1320
CCTACCCGCC CTGCGATGGA GTCCATGTGC GCTCAGTGTG CATTTCGGA GCTGGTGACT	1380
TGAACCTGGAT GCTGCGCAAA CACCACTTGT TTGCCAATAA CTTTGACGTG CATGTTGACC	1440
TCTTGCCAT CCAGTGTGTTG CATGAGCATT TGAGACACAA AGCTTGCGAG ACATTAAC	1500
ACTGACCATT ACGGGCAATT TTATGAACAA GAAGAAGGAT ACACAAAACG TACCTTATCT	1560
GTTTCCCCTT CCTTGTCAAGC GTCGGGAAGA TGGTATGAAG TCCTCTTGG GGCAGGGACT	1620
CTAGTAGATC TTCTTGTAGA GAAGCTGCAT GGTTCTGCA GAGCACAGTT AGCTAGAAAG	1680
GTGATAGCAT TAAATGTTCA TCTAGAGTTA ATAGTGGGAG GAGTAAAGGT AGCCTTGAGG	1740
CCAGAGCAGG TAGCAAGGCA TTGTGCAAAG AGGGGACCAG GGTGGCTGGG CAAGAGGCCG	1800
ATGCATAAAAG TCAGCCTGTT CCAAGTGCTC AGGGACTTAG CAAAATGAGA AGATGTGACC	1860
TGTGCCAAAA CTATTTGAC AATTTAAAT GTGACCATT TTCTGGTATG AATAAACTTA	1920
CAGCAACAAA TAATCAAAGA TACAATTAAT CTGATATTAT ATTTGTTGAA ATAGAAATT	1980
GATTGTACTA TAAATGATT TTGTAAATAA TTTATATTCT GCTCTAATAC TGTACTGTGT	2040
AGTCTGTCTC CGTATGTCAT CTCAGGGAGC TTAAAATGGG CTTGATTAA CATTGAAAAAA	2100
AA	2102

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GTCCCTCTAG ACCACCATGA TTGCTTCACA GTTT

-84-

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 29 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA


(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TCCCGGTCGA CTTAGGGAGC TTCACAGGT


29

We claim:

1. An O-glycan selected from the group consisting of a disialylated, monofucosylated glycan:



and the other is a monosialylated, trifucosylated glycan having a polylactosamine backbone:




wherein R = H, OH, another sugar or an aglycone such as an amino acid.


2. The O-glycan of claim 1 further comprising a pharmaceutically acceptable carrier for administration to a patient in need thereof.
3. A method for inhibiting inflammation mediated by P-selectin binding comprising administering to an individual in need thereof an effective amount to inhibit inflammation of an O-glycan selected from the group consisting of disialylated, monofucosylated glycans and monosialylated, trifucosylated glycans having a polylactosamine backbone.

-86-

4. The method of claim 3 wherein the O-glycan is selected from the group consisting of
a disialylated, monofucosylated glycan:




and a monosialylated, trifucosylated glycan having a polylactosamine backbone:




wherein R = H, OH, another sugar or an aglycone such as an amino acid.

5. The method of claim 4 wherein the glycans are selected from the group consisting of:



and:



6. An expression system comprising eukaryotic cells expressing a fucosyltransferase and a core 2 β 1-6-N-acetylglucosaminyltransferase, wherein the cells do not naturally express both enzymes.
7. The system of claim 6 wherein the eukaryotic cells are mammalian cells.
8. The system of claim 6 wherein the enzymes are human glycosyltransferases.
9. The system of claim 6 wherein both of the glycosyltransferases are expressed from recombinant DNA or RNA molecules.
10. The system of claim 7 wherein the cells are transfected with a recombinant DNA molecule encoding PSGL-1 or a fragment thereof.
11. The system of claim 10 wherein the fucosyltransferase is an α 1/3-1-4 fucosyltransferase (FTIII).
12. A recombinant PSGL-1 or fragment thereof binding to P-selectin expressed in eukaryotic cells expressing a fucosyltransferase and a core 2 β 1-6-N-acetylglucosaminyltransferase.
13. The PSGL-1 of claim 12 wherein the eukaryotic cells are mammalian cells.
14. The PSGL-1 of claim 12 wherein the glycosyltransferases are human transferases.
15. The PSGL-1 of claim 12 wherein the fucosyltransferase is an α 1/3-1-4 fucosyltransferase (FTIII).
16. Sulfated, glycosylated peptides which bind to P-selectin comprising tyrosine residues 46, 48, and 51 of Sequence Listing ID No. 1 wherein at least one of the tyrosine residues is sulfated (-SO₄) and there is at least one site for O-linked glycosylation.
17. The peptides of claim 16 wherein all three tyrosine residues are sulfated.
18. The peptides of claim 16 comprising threonines or serines which are suitable for O-linked glycosylation.

19. The peptides of claim 18 wherein the threonines correspond to the threonine residues at 44, 57, 69, and 70 of Sequence Listing ID No. 1.

20. The peptides of claim 16 comprising amino acid sequence corresponding to the amino acid sequence of Sequence ID No. 1 from tyrosine residue 46 through the next site for N or O-linked glycosylation.

21. The peptides of claim 16 selected from the group consisting of peptides corresponding to Sequence ID No. 1 from 42 to 57, 42 to 67, 42 to 69, 42 to 70, 42 to 73, 42 to 77, 42 to 80, 42 to 81, 42 to 82, 42 to 90, 42 to 91, 42 to 100, 42 to 103, 42 to 104, and these peptides having conservative substitutions in these peptides which do not alter the sulfated tyrosines or attachment of carbohydrate in a way decreasing binding of the peptide to P-selectin.

22. The peptides of claim 16 further comprising a pharmaceutically acceptable carrier.

23. The peptides of claim 16 further comprising a detectable label.

24. A method for inhibiting inflammation mediated by P-selectin binding comprising administering to an individual in need thereof an effective amount to inhibit inflammation of PSGL-1 or a fragment thereof, selected from the group consisting of recombinant PSGL-1 or fragments thereof binding to P-selectin expressed in eukaryotic cells expressing an α 1-3 fucosyltransferase, an α 1-4 fucosyltransferase, or an α 1/3-1-4 fucosyltransferase (FTIII) and a core 2 β 1-6-N-acetylglucosaminyltransferase and sulfated, glycosylated peptides which bind to P-selectin comprising tyrosine residues 46, 48, and 51 of Sequence Listing ID No. 1 wherein at least one of the tyrosine residues is sulfated (-SO₄) and there is at least one site for O-linked glycosylation.

-89-


25. The method of claim 24 wherein the sulfated, glycosylated peptides are expressed in the eukaryotic cells expressing an α 1-3 fucosyltransferase, an α 1-4 fucosyltransferase, or an α 1/3-1-4 fucosyltransferase (FTIII) and a core 2 β 1-6-N-acetylglucosaminyltransferase.

26. The method of claim 24 wherein the cells are mammalian and the glycosyltransferases are human.

27. The method of claim 24 wherein the peptides are selected from the group consisting of peptides corresponding to Sequence ID No. 1 from 42 to 57, 42 to 67, 42 to 69, 42 to 70, 42 to 73, 42 to 77, 42 to 80, 42 to 81, 42 to 82, 42 to 90, 42 to 91, 42 to 100, 42 to 103, 42 to 104, and these peptides having conservative substitutions in these peptides which do not alter the sulfated tyrosines or attachment of carbohydrate in a way decreasing binding of the peptide to P-selectin.


1/6

Tyr Tyr- SO_4 SO_4




SHAM

FIG. 1b



ARYLSULFATASE

FIG. 1c



SUBSTITUTE SHEET (RULE 26)

2/6




FIG. 2




FIG. 3a







FIG. 3b



4/6






FIG. 6

FIG. 7

O-Glycan Structures	Fraction %
Gal $\xrightarrow{\beta 1,3}$ GalNAcOH	P-10-3 14
Gal $\xrightarrow{\beta 1,4}$ GlcNAc ↓ Gal $\xrightarrow{\beta 1,3}$ GalNAcOH	P-10-3 52
NeuAc $\xrightarrow{\alpha 2,3}$ { Gal $\xrightarrow{\beta 1,4}$ GlcNAc ↓ Gal $\xrightarrow{\beta 1,3}$ GalNAcOH}	P-10-3 6
NeuAc $\xrightarrow{\alpha 2,3}$ Gal $\xrightarrow{\beta 1,4}$ GlcNAc ↓ NeuAc $\xrightarrow{\alpha 2,3}$ Gal $\xrightarrow{\beta 1,3}$ GalNAcOH	P-10-2b ₂ 14
NeuAc $\xrightarrow{\alpha 2,3}$ Gal $\xrightarrow{\beta 1,4}$ GlcNAc ↓ NeuAc $\xrightarrow{\alpha 2,3}$ Gal $\xrightarrow{\beta 1,3}$ GalNAcOH	P-10-2b ₁ 2
NeuAc $\xrightarrow{\alpha 2,3}$ Gal $\xrightarrow{\beta 1,4}$ ↓ Fuc Gal $\xrightarrow{\beta 1,3}$ GlcNAc $\xrightarrow{\beta 1,4}$ GlcNAc ↓ Fuc Gal $\xrightarrow{\beta 1,3}$ GlcNAc $\xrightarrow{\beta 1,4}$ GlcNAc ↓ Fuc Gal $\xrightarrow{\beta 1,3}$ GalNAcOH	P-10-2a 12