

(19)



Europäisches
Patentamt
European
Patent Office
Office européen
des brevets



(11)

EP 3 238 696 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
01.11.2017 Bulletin 2017/44

(21) Application number: 17174953.4

(22) Date of filing: 10.10.2008

(51) Int Cl.:

A61K 8/30 (2006.01) A61P 17/00 (2006.01)
A61P 17/02 (2006.01) A61P 17/06 (2006.01)
A61P 17/10 (2006.01) A61P 17/14 (2006.01)
A61K 9/70 (2006.01) A61K 31/085 (2006.01)

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT
RO SE SI SK TR

(30) Priority: 10.10.2007 US 998345 P
05.05.2008 US 126478

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
08837020.0 / 2 244 690

(71) Applicant: Therametrics, Llc
Oklahoma City, OK 73104 (US)

(72) Inventor: FULLER, Bryan B
Edmond, OK 73034 (US)

(74) Representative: Stuttard, Garry Philip
Urquhart-Dykes & Lord LLP
Arena Point
Merrion Way
Leeds LS2 8PA (GB)

Remarks:

This application was filed on 08-06-2017 as a
divisional application to the application mentioned
under INID code 62.

(54) METHODS AND COMPOSITIONS FOR TREATING DERMATOLOGICAL DISEASES AND CONDITIONS

(57) The present invention contemplates compositions comprising compounds, in particular dihydroeugenol (DHE) and/or isoeugenol (IE) and/or ethyl vanillin (EV) or salts, esters, ethers, or derivatives thereof, and methods for topically or systemically delivering them for treatment against inflammation-related and other dermatological conditions such as described herein. These DHE and/or IE and/or EV or salts, esters, ethers or derivatives thereof also are delivered to a measurable extent transepidermally or transdermally and thus may have systemic and/or localized anti-inflammatory effects within the body. In one of its method aspects, this invention is directed to a method for treating a patient with a dermatological disease or condition by topically admin-

istering to said patient a pharmaceutical or cosmetic composition comprising a pharmaceutically or cosmetically acceptable topical carrier and an effective dermatological disease, disorder or condition-treating amount of a formulation of DHE and/or IE and/or EV, or salts, esters, ethers, or derivatives thereof. In particular, the present invention is directed to a method of using a composition comprising one or more of DHE and/or IE and/or EV or salts, esters, ethers, or derivatives for treating dermatological diseases, disorders or conditions, aging effects related to decreasing production of collagen, elastin, or hyaluronic acid, or to hyperpigmentation. The formulations may further contain cinnamaldehyde.

Description**CROSS-REFERENCE TO RELATED APPLICATIONS**

[0001] The present application claims the benefit under 35 U.S.C. 119(e) of U.S. Provisional Application Serial No. 60/998,345, filed October 10, 2007, and U.S. Provisional Application Serial No. 61/126,478, filed May 5, 2008, the entireties of which are hereby expressly incorporated herein by reference.

5

10

BACKGROUND OF THE INVENTION

[0002] Human skin comprises an epidermis layer, which is predominantly composed of keratinocytes and a small number of melanocytes and Langerhans cells (antigen presenting cells), and a dermis layer, which is primarily composed of fibroblasts. The majority of skin disorders involve inflammation triggered by some insult to the skin. Keratinocytes respond quickly to environmental stimuli (e.g., UV radiation (UVR), allergens, irritants or physical damage) by producing a variety of inflammatory mediators, including cytokines (e.g., IL-1, TNF-alpha, and IL-6) and chemokines (e.g., IL-8). One of the most active inflammatory mediators is PGE-2 (Prostaglandin E2) and, of course, many topical dermatology drugs have been designed to lower levels of PGE-2. The fibroblasts in the dermis also produce PGE-2 along with a variety of chemokines, cytokines and matrix destroying enzymes such as collagenase (MMP-1).

15

[0003] The identification of compounds that can suppress the production of inflammatory mediators in the skin would allow effective topical products to be developed to treat a variety of inflammatory skin diseases or disorders including eczema, radiation dermatitis, atopic dermatitis, actinic keratosis, seborrheic dermatitis, other dermatic diseases and acne. Compositions able to treat other dermatological conditions such as aging effects and hyperpigmentation would also be desirable.

20

25

30

35

40

BRIEF DESCRIPTION OF THE DRAWINGS**[0004]**

Figure 1 is a graph showing the effects of dihydroeugenol (DHE) on UV-induced PGE-2 in fibroblasts.

45

Figure 2 is a graph showing the effects of DHE on UV-induced PGE-2 in keratinocytes.

Figure 3 is a graph showing the effects of DHE on TPA-induced TNF- α in keratinocytes.

50

Figure 4 is a graph showing the effects of DHE on LPS-induced inflammatory cytokines in monocytes.

Figure 5 is a photograph of an arm of a human subject which was exposed to erythema-inducing UVB radiation and treated with a lotion of the present invention.

55

Figure 6 shows photographs of a human subject's

face which has symptoms of rosacea. Facial areas of rosacea are shown in (A) while these areas, 12 weeks after initiation of treatment with DHE, IE, and cinnamaldehyde, are shown in (B).

Figure 7 shows the effects of topical DHE application on patients undergoing radiation treatment. (A) Radiation dermatitis in a typical radiation patient untreated for radiation dermatitis. (B) A patient was treated with a topical 2% DHE lotion daily during 35 radiation treatments. The photograph was taken after the 35th treatment and the skin shows no evidence of radiation dermatitis. (C) A close-up photograph of the patient of (B), one month after the end of radiation treatments. There is no evidence of radiation damage to the skin.

Figure 8 is a graph showing stimulation of collagen 1 mRNA in fibroblasts by DHE, isoeugenol (IE), and Ethyl vanillin (EV).

Figure 9 is a graph showing stimulation of elastin mRNA in fibroblasts by DHE, IE and EV.

Figure 10 is a graph showing stimulation of hyaluronic acid synthase-2 in RNA in fibroblasts by DHE, IE and EV.

Figure 11 is a graph showing stimulation of tissue inhibitor of metalloproteinase-1 mRNA (TIMP-1) in fibroblasts by DHE, IE and EV.

Figure 12 is a graph showing inhibition of matrix metalloproteinases (MMPs) in fibroblasts by ethyl vanillin.

Figure 13 is a graph showing inhibition of melanin synthesis *in vitro* by isoeugenol (referred to in the figure as TH-212).

Figure 14 shows photographs which demonstrate the loss of pigment in human melanocyte cultures treated with isoeugenol (TH-212) and isoeugenol acetate (referred to in the figure as TH-213) for 9 days. Figure 15 shows photographs which demonstrate how a topically-applied formulation of isoeugenol acetate (referred to in the figure as TH-212 ester) plus salicylic acid causes a clearing of acne and hyperpigmentation after a 4 week treatment.

Figure 16 shows photographs which demonstrate how a topically-applied formulation of isoeugenol acetate (TH-212 ester) plus 2% salicylic acid removes hyperpigmentation after a 30-day treatment period. Figure 17 shows photographs which demonstrate how a topically-applied formulation of isoeugenol acetate (TH-212 ester) plus 2% salicylic acid reduces sun-induced hyperpigmentation after a 30-day treatment period.

Figure 18 shows a photograph of cell extracts treated with isoeugenol (TH-212) and isoeugenol acetate (TH-213) then provided with L-DOPA, a tyrosinase substrate.

Figure 19 shows photographs which demonstrate psoriatic skin before (A) and after (B) treatment with a formulation comprising DHE and IE.

DETAILED DESCRIPTION OF THE INVENTION

[0005] The present invention is directed to methods and compositions for treating a variety of skin disorders, diseases, and conditions including for example, rosacea, radiation dermatitis, erythemas (sunburns), psoriasis, atopic dermatitis, allergic and irritant contact dermatitis, actinic keratitis, acne, scarring, hyperpigmentation, and seborrheic dermatitis or eczema, or other eczemas, and alopecia areata, wherein the compounds act, for example, on skin keratinocytes and fibroblasts and on immune cells such as monocytes and on melanocytes. The invention further comprises methods and compositions for mitigating the effects of aging, e.g., by enhancing production of collagen, elastin and hyaluronic acid synthases in skin fibroblast cells.

[0006] Inflammatory skin diseases are the most common problem in dermatology. They come in many forms, from occasional rashes accompanied by itching and redness to chronic conditions such as dermatitis (eczema), rosacea, seborrheic dermatitis, and psoriasis. However, they are all linked by one common factor, inflammation. It has been found that the inflammatory markers (cytokines) produced by skin and immune cells that are required for the development of an inflammatory response, such as atopic dermatitis and radiation dermatitis. The present invention comprises agents that suppress the production of a variety of inflammatory responses in cultured skin cells (keratinocytes and fibroblasts), and immune cells (monocytes and T-lymphocytes) and in intact living skin. As a result of blocking these inflammatory processes in the skin, the present compounds are able to effectively reduce or eliminate a variety of inflammatory symptoms that occur with common skin problems.

[0007] Rosacea is a vascular, inflammatory skin disorder that affects approximately 5% of the population and is characterized by frequent periods of facial redness or flushing caused by over-active capillaries. Over time, this chronic state of skin inflammation gives rise to a variety of rosacea symptoms. Rosacea is sometimes characterized mistakenly as adult-acne because patients present with a reddened face and acne-like symptoms. However, individuals affected with this skin disease also may have persistent redness with accompanying pain and itching in areas such as the forehead, chin, nose, ears, chest and back. As the disease progresses, small blood vessels and tiny pimples (called papules or pustules) begin to appear on and around the reddened area. In severe cases rosacea can affect the eyes (ocular rosacea) and cause disfigurement of the nose (rhynophyma). In addition to the physical symptoms associated with rosacea, patients also suffer significant psychological and social problems if left untreated.

[0008] Regarding skin aging, research shows that there are, in fact, two distinct types of skin aging. Aging caused by the genes we inherit is called *intrinsic* (internal) aging. The other type of aging is known as *extrinsic* (external) aging and is caused by environmental factors,

such as exposure to the sun's rays. Intrinsic aging, also known as the natural aging process, is a continuous process that normally begins in our mid-20s. Within the skin the dermis of normal (wrinkle-free) skin is composed of abundant amounts of type I collagen and type VII collagen, as well as elastin, which provides tissue strength, resiliency and recoil. However, as the dermal fibroblasts which produce collagen and elastin begin to age, they produce decreasing amounts of these proteins. Further, aging fibroblasts produce increased amounts of enzymes called matrix metalloproteinases (MMP's), which degrade collagen and elastin. This results in a drastic loss of collagen and elastin over time and results in skin laxity and fragility, visible in the form of fine lines and wrinkles. The signs of intrinsic aging are: fine wrinkles, thin and transparent skin, loss of underlying fat, leading to hollowed cheeks and eye sockets as well as noticeable loss of firmness on the hands and neck, dry skin that may itch, graying hair that eventually turns white, and hair loss. Genes control how quickly the normal aging process unfolds. Some persons notice their first gray hairs in their 20s; others do not see graying until their 40s.

[0009] A number of extrinsic aging factors often act together with the normal aging process to prematurely age our skin. Most premature aging is caused by sun exposure. Other external factors that prematurely age our skin are repetitive facial expressions, gravity and smoking. Without protection from the sun's rays, just a few minutes of exposure each day over the years can cause noticeable changes to the skin. Freckles, age spots, spider veins on the face, rough and leathery skin, fine wrinkles that disappear when stretched, loose skin, a blotchy complexion and skin cancer can all be traced to sun exposure. "Photoaging" is the term dermatologists use to describe this type of aging caused by exposure to the sun's rays. The amount of photoaging that develops depends on: (1) a person's skin color and (2) his or her history of long-term or intense sun exposure. Persons with fair skin who have a history of sun exposure develop more signs of photoaging than those with dark skin. In the darkest skin, the signs of photoaging are usually limited to fine wrinkles and a mottled complexion.

[0010] Photoaging occurs over a period of years. With repeated exposure to the sun, the skin loses the ability to repair itself, and the damage accumulates. Scientific studies have shown that repeated ultraviolet (UV) exposure impairs the synthesis of new collagen and increases the expression of MMP enzymes, which break down collagen. The sun also causes excessive elastin production but the elastin made is abnormal and aggregates into clumps, leading to a condition referred to by dermatologists as elastosis. Due to the loss of collagen and the production of abnormal elastin, sun-weakened skin ceases to spring back compared to skin protected from UV rays. Skin also becomes loose, wrinkled, and leathery much earlier with unprotected exposure to sunlight.

[0011] While there is nothing currently available to stop the aging process, it is possible to slow the rate at which

the skin ages. As described above, both intrinsic and photoaging are due to the breakdown and loss of collagen and elastin in the skin. The present compositions contain ingredients that have been scientifically proven herein to increase the production of collagen and elastin, as well as reduce the expression of MMP enzymes. These two effects slow the aging process and can even aid in rebuilding the dermal matrix, which reduces the appearance of new and existing wrinkles and fine lines.

[0012] Another skin condition, acne, is the most common skin disorder seen by doctors and affects almost everyone at some time. Teenagers are affected most often. Acne can cause a great deal of embarrassment and anxiety and can even cause people to become depressed which can lead to withdrawing from friends, and performing poorly at school or work. The exact cause of acne is unknown, but the following factors are considered important. (1) Acne is the visible end result of hormonal, bacterial and inflammatory disturbances that take place at the level of the oil pore (pilosebaceous follicle). (2) As the process advances, greater amounts of oil may be produced within the sebaceous glands, though the change in composition and quality of the oil may be more important than the quantity, the scale produced on the inside walls of the hair follicle becomes stickier and it builds up and blocks the pore which shows up as whiteheads and blackheads (comedones). (3) The acne bacteria (*Propionobacterium acnes*) grow and multiply in the retained oil. The sebum acts as a nutrition source for the bacterial, which in turn release chemicals within the pore. These alert and attract white cells from the blood leading to inflammation. (4) As these inflamed hair follicles (pores) and glands enlarge, the surrounding skin also becomes inflamed and may lead to even larger lumps and cysts (also called nodules). (5) Inflammation may damage the cells that make collagen. Less collagen production causes thinning of the skin, which is seen as depressed scars. Occasionally, collagen production will increase, which then causes the scars to become thickened. The present formulations may also be used to heal scars from acne and stretch marks by improving their structure and coloration.

[0013] Another skin disorder, psoriasis, is a chronic (long-lasting) skin disease characterized by scaling and inflammation. Scaling occurs when cells in the outer layer of skin reproduce faster than normal and pile up on the skin's surface. Psoriasis affects 2 to 2.6 percent of the United States population, or almost 5.8 to 7 million people. It occurs in all age groups and about equally in men and women. People with psoriasis may suffer discomfort, restricted motion of joints, and emotional distress. When psoriasis develops, patches of skin thicken, redden, and become covered with silvery scales. These patches are sometimes referred to as plaques. They may itch or burn. The skin at joints may crack. Psoriasis most often occurs on the elbows, knees, scalp, lower back, face, palms, and soles of the feet. The disease also may affect the fingernails, toenails, and the soft tissues inside the mouth

and genitalia. About 10 percent of people with psoriasis have joint inflammation that produces symptoms of arthritis. This condition is called psoriatic arthritis.

[0014] Research indicates that psoriasis may be a disorder of the immune system. The immune system includes a type of white blood cell, called a T cell, that normally helps protect the body against infection and disease. In psoriasis, the immune system produces too many T cells, in the skin. These T cells trigger the inflammation and excessive skin cell reproduction seen in people with psoriasis. This leads to inflammation and flaking of skin. The present invention inhibits the production of the same inflammatory mediators that other psoriasis therapies target, but since it is in topical form, it does not require frequent injections nor does it lower body's overall immune function.

[0015] Eczema is a general term for many types of skin inflammation (dermatitis). Atopic dermatitis is the most common of the many types of eczema. Several other forms have very similar symptoms. The diverse types of eczema are listed and briefly described below.

(1) Atopic dermatitis is a chronic skin disease characterized by itchy, inflamed skin. The word "dermatitis" means inflammation of the skin. "Atopic" refers to diseases that are hereditary, tend to run in families, and often occur together. These diseases include asthma, hay fever, and atopic dermatitis. In atopic dermatitis, the skin becomes extremely itchy and inflamed, causing redness, swelling, cracking, weeping, crusting, and scaling. Atopic dermatitis most often affects infants and young children, but it can continue into adulthood or first show up later in life. In most cases, there are periods of time when the disease is worse, called exacerbations or flares, which are followed by periods when the skin improves or clears up entirely, called remissions. Many children with atopic dermatitis enter into a permanent remission of the disease when they get older, although their skin often remains dry and easily irritated. Environmental factors can activate symptoms of atopic dermatitis at any time in the lives of individuals who have inherited the atopic disease trait. The cause of atopic dermatitis is unknown, but the disease seems to result from a combination of genetic and environmental factors. Evidence suggests that the disease is associated with other so-called atopic disorders such as hay fever and asthma, which many people with atopic dermatitis also have. In addition, many children who outgrow the symptoms of atopic dermatitis go on to develop hay fever or asthma. Although one disorder does not cause another, they may be related, thereby giving researchers clues to understanding atopic dermatitis. Atopic dermatitis is very common and affects males and females equally and accounts for 10 to 20 % of all referrals to dermatologists. Atopic dermatitis occurs most often in infants and children and its onset decreases sub-

stantially with age. Scientists estimate that 65 percent of patients develop symptoms in the first year of life, and 90 percent develop symptoms before the age of 5. Onset after age 30 is less common and often occurs after exposure of the skin to harsh conditions. People who live in urban areas and in climates with low humidity seem to be at an increased risk for developing atopic dermatitis. About 10% of all infants and young children experience symptoms of the disease. Roughly 60 percent of these infants continue to have one or more symptoms of atopic dermatitis even after they reach adulthood. This means that more than 15 million people in the United States have symptoms of the disease.

(2) Contact eczema is a localized reaction that includes redness, itching, and burning where the skin has come into contact with an allergen (an allergy-causing substance) or with an irritant such as an acid, a detergent (soap, bodywash), or other chemical.

(3) Allergic contact eczema is a red, itchy, weepy reaction where the skin has come into contact with a substance that the immune system recognizes as foreign, such as poison ivy or certain preservatives in creams and lotions.

(4) Seborrheic eczema is a form of skin inflammation of unknown cause but which is associated with a certain type of yeast that lives on the skin. Seborrheic eczema presents as yellowish, oily, scaly patches of skin on the scalp, face, and occasionally other parts of the body (called cradle cap in infants).

(5) Nummular eczema is coin-shaped patches of irritated skin-most commonly on the arms, back, buttocks, and lower legs-that may be crusted, scaling, and extremely itchy.

(6) Neurodermatitis is scaly patches of skin on the head, lower legs, wrists, or forearms caused by a localized itch (such as an insect bite) that becomes intensely irritated when scratched.

(7) Stasis dermatitis is a skin irritation on the lower legs, generally related to circulatory problems.

(8) Dyshidrotic eczema is irritation of the skin on the palms of hands and soles of the feet characterized by clear, deep blisters that itch and burn.

[0016] Radiation therapy, another skin disorder, can have some unpleasant side effects in the skin. The following are the most common side effects, both acute and chronic, resulting from radiation. Unforeseen side effects may occur because of the unique and varied tolerance of individual persons. Late effects of treatment may not always be predictable and may be influenced by concurrent and/or subsequent treatment for this and other diseases.

[0017] Specific side effects of radiotherapy depend on the part of the body being treated as well as the dose given. In general, the first change is a reddening of the skin, resembling a sunburn. In many patients this is all that is experienced. However, in most patients the burn

can be severe and in many cases equivalent to second degree burns. Like a sunburn, the involved area is often sensitive and even painful to the touch. In addition, the overlying skin may break down and the area may remain open until several days to weeks after the course of radiation is completed. Once the course of radiotherapy is completed, the redness will gradually go away and any open areas normally will heal. However, the skin in this area will most likely develop features of aged skin including pronounced wrinkling, skin thinning, stiffness and/or dryness, as well as possible pigmentation changes.

[0018] Most of the current treatment options for radiation dermatitis involve the use of emollients or aloe gels in an attempt to keep the skin moisturized. However, as most know who have had the experience of a sunburn, moisturization helps the skin from drying out but does not reduce the pain or redness, which are caused by inflammation. The present invention comprises a moisturizing lotion that contains an active agent, a bioactive

that is able to reduce skin redness and pain associated with radiation therapy.

[0019] Another skin disorder, hyperpigmentation, is a common and distressing condition afflicting a large subset of the population. Hyperpigmentation is the result of an increased amount of melanin in the epidermis, the dermis, or both. This pigmentary change can be divided into 2 pathophysiologic processes: melanocytosis (increased number of melanocytes) and melanosis (increased amount of melanin).

[0020] In a preferred embodiment, the invention comprises a formulation comprising dihydroeugenol. The DHE formulation may be administered topically, systemically, or orally administered to a subject having rosacea for example for treating and reducing the rosacea on the subject's skin. The DHE formulation may also be used for example to treat and/or inhibit psoriasis and radiation dermatitis in a subject. The formulation may comprise, with or instead of the DHE, a salt, ester, or ether, thereof as described elsewhere herein. The formulation may further comprise one or more of isoeugenol (IE) or of a salt, ester, or ether thereof, for example isoeugenyl acetate or methyl isoeugenol. Any of these formulations may also be used to treat inflammatory skin conditions such as actinic keratitis, acne, scarring, allergic and irritant contact dermatitis, atopic dermatitis, erythema (sunburn), hand eczema, seborrheic dermatitis, alopecia areata, and other inflammatory irritations of the skin. Formulations comprising isoeugenol and salts, esters, and ethers thereof can be used in particular to reduce the effects of hyperpigmentation of the skin.

[0021] In preferred embodiments, the compositions of the present invention comprise synthetic and/or natural versions of the compounds dihydroeugenol (DHE) and/or isoeugenol (IE) and/or ethyl vanillin (EV), and/or salts, esters, ethers or derivatives of DHE, IE and EV, or combinations thereof and particularly salts, esters, ethers, and derivatives of DHE, IE and EV including isoeugenol acetate, for example.

[0022] In one of its compositional aspects, this invention is directed to pharmaceutical compositions for topical, transdermal or other systemic administration containing a pharmaceutically-acceptable carrier and DHE and/or IE and/or EV and/or salts, esters, ethers or derivatives thereof or combinations thereof as described herein. In particular, the present invention is directed to a composition comprising DHE and/or IE and/or EV and/or salts, esters, ethers or derivatives thereof or combinations thereof for use in treating dermatologic diseases, disorders or conditions.

[0023] In one of its method aspects, this invention is directed to a method for treating a patient with a dermatological disease, disorder, or condition by topically administering to said patient a pharmaceutical composition comprising a pharmaceutically acceptable topical carrier and an effective dermatological disease-treating or condition-treating amount of a formulation of DHE and/or IE and/or EV and/or salts, esters, ethers, or derivatives thereof or combinations thereof.

[0024] In another one of its method aspects, this invention is directed to a method for treating a dermatological condition, in particular radiation dermatitis, by topically applying to a human a cosmetic composition comprising a pharmaceutically acceptable topical carrier and an effective amount of a formulation of DHE and/or IE and/or EV and/or salts, esters, ethers or derivatives thereof or combinations thereof. In still another of its method aspects, this invention is directed to a method for treating a patient with an inflammatory disease, disorder, or condition by systemically administering to said patient a pharmaceutical composition comprising a pharmaceutically acceptable carrier and an effective inflammatory disease-treating amount of a formulation of DHE and/or IE and/or EV and/or salts, esters, ethers, or derivatives thereof or combinations thereof.

[0025] In yet another of its method aspects, this invention is directed to a method for treating a human with an inflammatory disease, disorder or condition by topically applying to said human a pharmaceutical composition comprising a pharmaceutically acceptable carrier and an effective amount of a formulation of DHE and/or IE and/or EV and/or salts, esters, ethers or derivatives thereof or combinations thereof.

[0026] In yet another of its method aspects, this invention is directed to a method for improving the skin appearance of a person such as reducing skin aging by topically administering to said person a pharmaceutical or cosmetic composition comprising a pharmaceutically acceptable carrier and a pharmaceutically or cosmetically effective amount of a formulation of DHE and/or IE and/or EV and/or salts, esters, ethers, or derivatives thereof or combinations thereof.

[0027] In another one of its method aspects, this invention is directed to a method for reducing areas of hyperpigmentation on the skin by topically applying to said person's skin a pharmaceutical or cosmetic composition comprising a pharmaceutically or cosmetically accepta-

ble carrier and an effective amount of a formulation of IE and/or salts, esters, ethers, or derivatives thereof (e.g., isoeugenol acetate) or combinations thereof.

[0028] An optional component of any of the formulations of the present invention is cinnamaldehyde at a concentration which preferably is in the range of 0.01% to 5%. Cinnamaldehyde is preferably included in the present formulations for use in treatment of rosacea, acne, and seborrheic dermatitis. The cinnamaldehyde provides both anti-inflammatory and anti-bacterial/anti-microbial effects against, for example, bacteria which contribute to acne and yeasts which contribute to seborrheic dermatitis.

15 Definitions

[0029] When describing the cosmetic and pharmaceutical compositions and methods of this invention as well as the compositions and methods themselves, the following terms have the following meanings:

[0030] "Ionizing radiation" refers to any radiation that ionizes the atoms or molecules of matter. It may consist of particles (such as electrons) or it may be electromagnetic (ultraviolet radiation; X-rays; gamma radiation). Ionizing radiation occurs naturally, for example as a component of sunlight, and is emitted by radioactive substances. It is also produced artificially in X-ray machines, particle accelerators, and nuclear reactors, for example.

[0031] "Isolated", when used to define the state of purity of the synthetic or natural compounds used in the practice of this invention, means that the compounds described herein and/or salts, esters, ethers, or derivatives thereof or combinations thereof has been substantially freed of (i.e. at least about 90% and especially at least about 95%) or separated from related feedstocks, raw materials, co-products, or in the case of naturally-occurring mixtures, related materials with which the compound appears in nature.

[0032] "Pharmaceutically-acceptable topical carrier" and equivalent terms refer to an inactive liquid or cream vehicle capable of suspending or dissolving the compounds described herein and/or salts, esters, ethers, or derivatives thereof or combinations thereof and having the properties of being generally nontoxic and noninflammatory when applied to the skin. This term is specifically intended to encompass carrier materials approved for use in topical cosmetics and topical and systemic pharmaceuticals. Representative carriers include water, silicone fluids, oils, both vegetable and mineral, cream bases, lotion bases, ointment bases and the like. These bases include suspending agents, thickeners, penetration enhancers, and the like. Their formulation is well known to those in the art of cosmetics and topical pharmaceuticals. Additional information concerning carriers can be found in Remington's Pharmaceutical Sciences, 20th edition, 2000, Lippincott, Williams and Wilkins, which is hereby expressly incorporated herein by reference in its entirety.

[0033] "Therapeutically effective dose" means a dose of a composition of this invention which, when applied topically to the skin of a patient afflicted with a dermatologic or other cosmetic or medical disease, disorder, or condition, or when administered by another route such as systemically results in an observable improvement in the patient's condition.

[0034] "Topical", when used to define a mode of administration, means that a material is administered by being applied to the skin or to an internal epithelial layer such as within the rectum, or colon, or nasal or respiratory passage.

[0035] "Topically effective" means that a material, when applied to the skin or epithelial layer described above, produces a desired pharmacological result either locally at the place of application or systemically as a result of transdermal passage of an active ingredient in the material.

[0036] Where used herein, the term "oil-in-water formulation" refers to a formulation wherein a continuous water phase surrounds droplets of oil or lipid in an emulsion.

[0037] Where used herein, the term "water-in-oil formulation" refers to a formulation wherein a continuous lipid or oil phase surrounds droplets of water in an emulsion.

[0038] Where used herein the term "non-aqueous formulation" refers to a formulation having less than 1%, and preferably less than 0.1%, by weight of water in the formulation. The term "non-aqueous" is also intended to include formulations having a negligible amount of water due to absorption of atmospheric moisture.

[0039] Where used herein the term "emulsifier" refers to a compound which is used to promote and maintain a stable mixture or dispersion (emulsion) of oil droplets in a water phase, or water droplets in an oil phase.

[0040] Emulsifiers are, essentially, surfactants. These surfactants can be ionic (cationic or anionic) or non-ionic, and they can be used alone or in combination. Emulsifiers contemplated for use herein include, but are not limited to, cetearyl alcohol and sodium cetearyl sulfate, PEG-1000 monocetyl ether, glycol stearate, glyceryl stearate, cetyl alcohol, PEG-100 stearate, ceteareth-20, or quaternary ammonium salts such as alkyl trimethyl ammonium bromide, the polyol ester glycerol monostearate, potassium stearate, sodium lauryl sulfate, and ethoxylated fatty alcohols. Fatty acids like stearic acids may be included to regulate the consistency of the emulsion. Finally, polymers such as carbomers can be included in small amounts to stabilize the emulsion.

[0041] Penetration enhancers are substances which enhance passage of topically-applied compounds into the stratum, corneum of the skin and therefrom into the epidermis and dermis. Examples include, but are not limited to: dimethylisosorbide, ethoxydiglycol, 1-dodecylazacycloheptan-2-one, propylene glycol, oleyl alcohol, polyoxyethylene ester, sorbitan mono-9-octadecenoate, poly(oxy-1,2-ethanediyl) and derivatives there-

of, ethanol, glyceryl monoethyl ether, monoglycerides, isopropylmyristate, lauryl alcohol, lauric acid, lauryl lactate, terpinol, menthol, D-limonene, beta-cyclodextrin, DMSO (dimethyl sulfoxide), polysorbates, fatty acids (e.g., oleic), bile salts, N-methylpyrrolidone, polyglycosylated glycerides, 1-dodecylazacycloheptan-2-one (Azone®), Cyclopentadecalactone (CPE-215®), Alkyl-2-(N,N-disubstituted amino)-alkanoate ester (NexAct®), 2-(n-nonyl)-1,3-dioxolane (DEPA®), and penetration enhancers shown for example in U.S. Patents 3,909,816; 4,405,616; 4,801,586; 4,861,764; 4,886,783; 4,983,396; 5,118,845; and 5,196,410, each of which is hereby expressly incorporated herein by reference in its entirety.

[0042] Where the compositions and methods of the present invention comprise isoeugenol or dihydroeugenol, the present invention in particular contemplates methods of inhibiting or treating dermatological conditions, disorders, or diseases by epithelial application of compositions comprising esters and ethers of isoeugenol and dihydroeugenol, including, but not limited to, isoeugenyl formate, isoeugenyl acetate, isoeugenyl propionate, isoeugenyl butyrate, isoeugenyl isobutyrate, isoeugenyl oleate (and other unsaturated fatty acid esters), isoeugenyl benzoate, isoeugenyl phthalate, isoeugenyl hexanoate, isoeugenyl heptanoate, isoeugenyl octanoate, isoeugenyl pentanoate, isoeugenyl decanoate, isoeugenyl lactate, isoeugenyl cinnamate, isoeugenyl valerate, isoeugenyl isovalerate, isoeugenyl nonanoate, isoeugenyl caprylate, isoeugenyl phenylacetate, isoeugenyl anthranilate, isoeugenyl salicylate, isoeugenyl methyl ether (methyl isoeugenol), benzyl isoeugenyl ether, isoeugenyl ethyl ether (ethyl isoeugenol), and dihydroeugenyl formate, dihydroeugenyl acetate, dihydroeugenyl propionate, dihydroeugenyl butyrate, dihydroeugenyl dihydrobutyrate, dihydroeugenyl oleate (and other unsaturated fatty acid esters), dihydroeugenyl benzoate, dihydroeugenyl phthalate, dihydroeugenyl hexanoate, dihydroeugenyl heptanoate, dihydroeugenyl octanoate, dihydroeugenyl pentanoate, dihydroeugenyl decanoate, dihydroeugenyl lactate, dihydroeugenyl cinnamate, dihydroeugenyl valerate, dihydroeugenyl isovalerate, dihydroeugenyl nonanoate, dihydroeugenyl caprylate, dihydroeugenyl phenylacetate, dihydroeugenyl anthranilate, dihydroeugenyl salicylate, dihydroeugenyl methyl ether (methyl dihydroeugenol), benzyl dihydroeugenyl ether and dihydroeugenyl ethyl ether (ethyl dihydroeugenol). These esters and ethers of isoeugenol and dihydroeugenol can be combined with various carriers, vehicles, diluents, and excipients to form topical formulations as described elsewhere herein.

[0043] While the invention will now be described herein in connection with certain examples and embodiments so that aspects thereof may be more fully understood and appreciated, it is not intended that the invention be limited to these particular embodiments or examples. On the contrary, it is intended that all alternatives, modifications and equivalents are included within the scope of the invention as defined by the appended claims. Thus

the examples described herein, which include preferred embodiments, will serve to illustrate the practice of this invention, it being understood that the particulars shown are by way of example and for purposes of illustrative discussion of preferred embodiments of the present invention only and are presented in the cause of providing what is believed to be the most useful and readily understood description of procedures as well as of the principles and conceptual aspects of the invention.

EXPERIMENTAL EXAMPLES

Example 1.

[0044] In one study, various concentrations of DHE were added to culture medium and these media were placed on human fibroblast cell cultures (results shown in Fig. 1). Cells were either not induced or induced with UVB radiation to stimulate cytokine/PGE-2 production. At 24 hours after treatment, the cell culture medium was removed and assayed by ELISA methods for the expression of PGE-2, various cytokines and chemokines.

[0045] As is shown in Fig. 1, UV radiation (UVR) treatment of fibroblasts results in a 6 fold increase in PGE-2. When DHE at a concentration as low as 10 micromolar was put into the culture media after irradiation, it completely blocked the UVR induction of PGE-2.

[0046] When similar experiments were carried out with human keratinocytes, DHE was again found to markedly suppress the UVR induction of PGE-2, with a concentration of 50 uM inhibiting the production of PGE-2 by over 60% (results shown in Fig. 2). Since PGE-2 is the major inflammatory mediator responsible for sunburn these data indicate the efficacy of DHE in the treatment of sunburn. Further, since PGE-2 is known to be a principal factor in the development of actinic keratosis and skin cancer, DHE can also be used in treating these conditions.

[0047] Inflammatory mediators produced in the skin contribute to the development and propagation of such diseases as rosacea, psoriasis and atopic dermatitis. While many inflammatory mediators are involved in these diseases, TNF- α is known to be a major cytokine involved in psoriasis. For atopic dermatitis, TNF- α , IL-8, and MCP-1 are important mediators of inflammation in these diseases. DHE is effective in suppressing the TPA-induced (TPA is tetradecanoyl phorbol ester) production of TNF- α by approximately 50% at a 100 micromolar concentration (results shown in Fig. 3).

[0048] In monocytes (as shown in Fig. 4), dihydroeugenol can inhibit the production of the cytokine TNF- α as well as the chemokines, IL-8, and MCP-1 (monocyte chemotactic protein 1). Since these inflammatory mediators are critically important for the development of immune driven inflammatory diseases such as atopic dermatitis and psoriasis, the results indicate that DHE can be used to treat these diseases.

[0049] A summary of some of the inflammatory medi-

ators blocked in keratinocytes, fibroblasts and monocytes is shown below in Tables 1, 2 and 3.

[0050] Table 1 shows the inhibitory effects of DHE and IE on production of several inflammatory mediators (PGE-2, IL-6, IL-8, and TNF- α) induced in keratinocytes by exposure to TPA and UV light. DHE and IE both have inhibitory effects on production of the inflammatory mediators in keratinocytes.

10 TABLE 1. Percent Inhibition Of Inflammatory
Mediators of Keratinocytes by DHE and IE

Active Compounds	DHE	IE
Concentration	100 uM	100 uM
Keratinocytes (TPA)		
PGE2	77%	82%
IL-6	30%	94%
IL-8	45%	75%
TNF-α	12%	71%
Keratinocytes (UV)		
PGE2	68%	76%
IL-6	30%	93%
IL-8	40%	76%
TNF-α	31%	60%

[0051] Table 2 shows the inhibitory effects of DHE and IE on production of several inflammatory mediators (PGE-2, IL-6, and IL-8) induced in fibroblasts by exposure to IL-1 and UV light. DHE and IE both have inhibitory effects on the production of the inflammatory mediators in fibroblasts.

45 TABLE 2. Percent Inhibition Of Inflammatory
Mediators of Fibroblasts By DHE and IE

Active Compounds	DHE	IE
Concentration	100 uM	100 uM
Fibroblasts (IL-1)		
PGE-2	86%	81%
IL-6	47%	53%
IL-8	46%	98%

(continued)

Fibroblasts (UV)		
PGE-2	67%	77%
IL-6	49%	39%
IL-8	11%	2%

[0052] Table 3 shows the inhibitory effects of DHE and IE on production of several inflammatory mediators (MCP-1 (monocyte chemotactic protein-1), IL-12, TNF- α , and IL-8) induced in monocytes by LPS (lipopolysaccharide) and TNF- α . DHE and IE both have inhibitory effects on the production of the inflammatory mediators in monocytes.

TABLE 3. Percent Inhibition Of Inflammatory Mediators of Monocytes By DHE and IE

Active Compounds	DHE	IE
Concentration	100 μ M	100 μ M
Monocytes (LPS)		
MCP-1	60%	87%
IL-12	66%	80%
TNF- α	78%	80%
IL-8	70%	75%
Monocytes (TNF- α)		
MCP-1	46%	82%
IL-8	72%	78%
IL-12	98%	100%

Example 2.

[0053] In one human clinical efficacy study the forearm of a clinical subject was irradiated at two sites with a dose of UVB radiation sufficient to induce erythema (Fig. 5). Immediately after the irradiation dose, but not before, one irradiated site was treated with a topical lotion containing 2% by weight of DHE while the other irradiated site was treated with the same topical formulation without DHE (vehicle control). After 4 hours, pronounced erythema was visible at the site treated with the vehicle control while the site treated with 2% DHE lotion displayed only minimal erythema. This result was unexpected because stronger anti-inflammatory compounds, such as steroids, when used typically are unable to prevent UVB irradiation-induced erythema even though they are more effective than DHE in blocking inflammatory mediators, including PGE-2, in vitro. Therefore, the actual mechanism of action of DHE in reducing UVB radiation induced ery-

thema likely involves more than just inhibiting inflammatory cytokines and hormones such as PGE-2. The formulations of the present invention not only are able to prevent induction of erythema but are also able to reverse erythema that is pre-existing.

Example 3.

[0054] In addition to its ability to block radiation induced erythema (as shown in Example 2) another unexpected finding is that topically-applied DHE is effective in reducing the symptoms of rosacea, a disease whose etiology is for the most part not understood. Typical treatments for rosacea include topical metronidazole which is an antibiotic effective against bacteria and some parasites, and oral antibiotics. When topical DHE and/or a combination of DHE, IE, and cinnamaldehyde was topically applied, over 8-12 weeks, to patients suffering from rosacea, the results revealed an overall improvement in their condition with use of a DHE formulation versus the same lotion base without the DHE or DHE/IE/cinnamaldehyde. The results of this study are shown in Fig. 6A-B.

Example 4.

[0055] A clinical study carried out with cancer patients undergoing radiation therapy showed that topically-applied 2% DHE lotion (applied twice daily) can almost completely prevent the onset of radiation dermatitis. For example in a patient who underwent 35 radiation treatments for breast cancer and who was treated with a 2% DHE lotion (e.g., Formulation 5 shown below) daily during the 35 radiation treatments, there was no radiation damage after the 35 days of treatment or one month after the end of radiation treatment (see Fig. 7A-C).

Example 5.

[0056] The present invention further contemplates use of various DHE, IE and EV compositions and salts, esters, ethers, and derivatives thereof to mitigate the effects of aging on the skin. For all aging studies, normal human fibroblast cell cultures were used. These cells normally produce collagen, elastin and hyaluronic acid. As the skin ages, the fibroblasts in the dermis lose their ability to produce these three key components of skin, and the skin consequently loses elasticity, thickness and smooth texture. In addition, as fibroblasts age, they increase the expression of certain enzymes that destroy collagen and elastin. There are about 13 of these enzymes, collectively referred to as MMPs (matrix metalloproteinases). One of the principal MMPs, MMP-1, is responsible for collagen degradation. For experiments herein to determine the effects of DHE, IE and EV on collagen, elastin and hyaluronic acid, fibroblasts were treated with the compounds for 72 hours, mRNA was then isolated, and analyzed for abundance by RT-PCR (Reverse transcriptase- Polymerase Chain Reaction). IE, DHE and EV

were shown to stimulate collagen production in fibroblasts.

[0057] Fibroblast cell cultures were treated with either IE, DHE or EV for 72 hours at which time the amount of collagen mRNA present in the cells was determined. DHE, IE and EV each caused stimulation of collagen production in fibroblasts, as indicated by increased collagen mRNA production (Fig. 8).

[0058] In addition to stimulating collagen mRNA synthesis and collagen protein synthesis, DHE, IE and EV and salts, esters, ethers, and derivatives thereof can increase the level of elastin mRNA in human dermal fibroblasts as shown in Fig. 9. The image is from an electrophoresis gel of DNA amplified by PCR from fibroblast mRNA. This gel was analyzed and the density of each band quantified by image analysis software. The bar graph showing the densitometric analysis is shown above the image. DHE, IE and EV increase elastin levels in human dermal fibroblasts.

[0059] DHE, IE and EV, are also shown herein to stimulate mRNA for HAS-2. HAS-2 (Hyaluronic acid synthase-2) is the enzyme that manufactures hyaluronic acid (HA) in the skin. HA is a glycosaminoglycan important for maintaining moisture and suppleness in the skin. HA has a half-life of 24 hours and thus, must be continually replaced. As the skin ages, the production of HA decreases and this causes the skin to sag and become thin. We have discovered that DHE, either alone or in combination with EV, can also stimulate the HAS-2 gene as shown in Fig. 10. In addition EV alone can stimulate HAS-2, but the combination of DHE and salts and esters thereof and EV synergistically works better than either one alone.

[0060] We have also shown that TIMP production can be upregulated by DHE, IE and EV. Fibroblasts in the skin not only produce enzymes (MMPs) that destroy the skin matrix but they also produce proteins (Tissue Inhibitors of Metalloproteinases, i.e., TIMPs) that inhibit these enzymes. There are several TIMPs, but one that is most important for inhibiting MMPs is TIMP-1. We have discovered that DHE, IE and EV (and salts, esters, ethers, and derivatives thereof), either alone or in combination, can significantly stimulate the production of TIMP-1, thereby providing the skin with protection against the matrix destroying activity of collagenase (MMP-1), the enzyme that TIMP-1 inhibits. Shown in Fig. 11 is PCR data showing the up-regulation of TIMP-1 mRNA by these compounds.

[0061] We have also demonstrated inhibition of MMPs by EV. As mentioned above, as fibroblasts age they increase their production of MMP enzymes that destroy collagen and elastin, particularly collagenase, MMP-1. We utilized a protein array blotting technique to determine the effect of DHE, IE and EV on the protein abundance of these enzymes. As shown below, EV is able to reduce the level of several MMPs in human dermal fibroblasts, including MMP-1, MMP-3, MMP-8, MMP-9 MMP-10 and MMP-13 (Fig. 12). This remarkable effect makes this compound an ideal addition to an anti-aging

product since MMP levels in aging skin are quite high.

Example 6.

5 IE Inhibits Melanin Synthesis.

[0062] IE has been found to have a surprising and unexpected effect on blocking the synthesis of melanin catalyzed by tyrosinase. Two ml of a sodium phosphate buffer, pH 6.8 containing 0.2% L-DOPA (dihydroxyphenylalanine- a tyrosinase substrate) were placed in each of 10 two test tubes. To one test tube was added 20 microliters of ethanol and to the other tube was added 20 microliters of a 1M stock of IE made up in ethanol (labeled TH-212). To start melanin synthesis from the L-DOPA substrate, 15 20 microliters of a tyrosinase preparation (0.5 mg/ml) was added to both tubes, the tubes were mixed and left at room temperature. The photograph in Fig. 13 was taken 2 hours after the start of the reaction. As can be seen in 20 Fig. 13, melanin synthesis was almost completely blocked in the tube containing IE.

[0063] Without wishing to be bound by theory, it does not appear that IE is directly inhibiting tyrosinase, but rather is interfering with the post-tyrosinase steps required for melanin synthesis. IE does not appear to interfere with the tyrosinase mediated conversion of DOPA to dopaquinone or dopaquinone to dopachrome since the assay tube containing tyrosinase, DOPA and IE temporarily turns reddish. Since dopachrome is red, this indicates that IE is allowing tyrosinase to convert DOPA to dopaquinone. The conversion of dopaquinone to dopachrome is spontaneous and does not require tyrosinase. Isoeugenol may be converting dopachrome to some chemical entity that cannot proceed down the melanin synthesis pathway but remains as a colorless intermediate. Alternatively, IE may allow dopachrome to be converted to 5,6 dihydroxyindole, the next intermediate in the melanin pathway. It may then prevent the polymerization of 5,6 dihydroxyindole (a colorless intermediate) to melanin.

[0064] Figs. 14-17 show the skin-lightening effects of topically-applied formulations of isoeugenol and its ester isoeugenol acetate (a.k.a. isoeugenyl acetate) in the treatment of hyperpigmented areas in the skin, wherein 45 a reduction of melanin pigment in the skin is caused by the topical application of the formulations. In addition to blocking a post-tyrosinase step in melanin synthesis, IE and IE acetate can inhibit the expression of the tyrosinase protein in human melanocytes. This inhibitory effect can be demonstrated by preparing cell extracts from melanocytes treated for 9 days with either IE or IE-acetate or left untreated. If the cell extracts contain tyrosinase when an aliquot of extract is placed in a tube containing L-DOPA (a tyrosinase substrate) in buffer, the enzyme will convert 50 the DOPA to melanin. The melanin will appear brownish-black in the assay tube. If IE or IE-acetate has inhibited tyrosinase synthesis in melanocytes, then the extracts from these cells will contain no enzyme and will not be 55

able to produce melanin from DOPA in the assay tube. As shown in Fig. 18 very little melanin was made in assay tubes that contained DOPA plus cell extracts from either IE or IE-acetate treated cells. An inhibitory effect of IE or IE-acetate on tyrosinase synthesis can also be demonstrated by the use of western immunoblots. In this assay, cell extracts from human melanocytes either left untreated or treated with 200 micromolar of either IE or isoeugenol acetate for 9 days are run on SDS polyacrylamide electrophoresis gels to separate tyrosinase from other proteins. The tyrosinase in the gel is then blotted to a membrane and the tyrosinase detected by staining with a specific anti-tyrosinase antibody. The antibody-bound tyrosinase is then visualized by chemiluminescence. The results show that while untreated melanocytes have a high abundance of tyrosinase, the melanocytes treated for 9 days with either IE or isoeugenol acetate had almost no detectable tyrosinase present. This indicates that these compounds suppress the synthesis of the enzyme in human melanocytes.

[0065] Western immunoblots of tyrosinase abundance in human melanocyte extracts treated for 9 days with isoeugenol or isoeugenol acetate showed a considerable reduction in tyrosinase (as measured by RDU-relative densitometric units). The control showed approximately 7.25 RDU, while 200 μ M concentrations of isoeugenol and isoeugenol acetate resulted in about 3 RDU and 3.75 RDU, respectively (data not shown).

[0066] Regardless of the mechanism by which IE or IE acetate or other compounds described herein have their effect, the inhibition of melanin production is essentially permanently blocked since test tubes containing DOPA, IE and tyrosinase fail to darken even after 2 weeks. The invention thus further contemplates topically applying an IE composition which has a skin-penetrating vehicle to reduce or prevent the formation of melanin in vivo in keratinocytes, thereby acting as a skin lightening agent.

[0067] The invention in particular contemplates methods of inhibiting skin pigmentation (hyperpigmentation) and causing skin lightening by topical application of compositions comprising isoeugenol and salts, esters, and ethers of isoeugenol, including, but not limited to, isoeugenyl formate, isoeugenyl acetate, isoeugenyl propionate, isoeugenyl butyrate, isoeugenyl isobutyrate, isoeugenyl oleate (and other unsaturated fatty acid esters), isoeugenyl benzoate, isoeugenyl phthalate, isoeugenyl hexanoate, isoeugenyl heptanoate, isoeugenyl octanoate, isoeugenyl pentanoate, isoeugenyl decanoate, isoeugenyl lactate, isoeugenyl cinnamate, isoeugenyl valerate, isoeugenyl isovalerate, isoeugenyl nonanoate, isoeugenyl caprylate, isoeugenyl phenylacetate, isoeugenyl anthranilate, isoeugenyl salicylate, isoeugenyl methyl ether (methyl isoeugenol), isoeugenyl ethyl ether (ethyl isoeugenol) and benzyl isoeugenyl ether. These esters and ethers of isoeugenol can be combined with various carriers, vehicles, diluents, and excipients to form topical formulations as described elsewhere herein.

Example 7

Treatment of Psoriasis

5 **[0068]** The IE and DHE compositions (and salts, esters and ethers thereof) contemplated herein can be used to treat the scaly patches which occur in the skin of sufferers of psoriasis. Clinical treatments have shown significant reduction of scaling and inflammation after a 30-day 10 course of treatment using the compositions of Examples 4-7. For example, Fig. 19 shows before (A) and after (B) pictures of a single psoriatic skin lesion after a 30 day treatment with the formulation of the present invention comprising DHE and IE.

Example 8.

Effect of Water on Stability of EV.

20 **[0069]** In a preferred embodiment, the present invention comprises a non-aqueous formulation of ethyl vanillin, preferably comprising a silicone as the primary component (e.g., for example in the range of 20%-90% silicone). The non-aqueous formulation of ethyl vanillin can 25 be topically applied to the skin to stimulate fibroblast production of collagen, elastin, and TIMPs in the dermis and to reduce production of MMPs in the dermis, thereby inhibiting and counteracting the effects of aging of the skin and causing improved appearance thereof. Preferably 30 the formulation comprises .1 to 5% of ethyl vanillin. The formulation may further be used to "re-model" scars in the skin wherein scars in the skin (including stretch marks and scarring due to acne) are treated so as to cause them to "disappear" i.e., to regain normal skin coloration and 35 texture. The formulation may further comprise DHE, IE, and/or salts, esters, and/or ethers thereof and cinnamaldehyde.

40 **[0070]** In a novel discovery of the present invention, it has been found that EV cannot be placed in a water-oil emulsion without the EV significantly decomposing within 24-48 hours. This surprising and unexpected finding 45 meant that it was necessary to identify a carrier/vehicle system that would accommodate EV in a soluble form and at the same time maintain its chemical stability (defined herein as at least 95% of the ethyl vanillin remaining chemically intact for at least 3 months). In one formulation contemplated herein, EV is dissolved in caprylic/capric triglyceride or other fatty acid triglyceride and then mixed into a silicone fluid wherein the silicone is the primary 50 (greatest percentage) component of the formulation. The cosmetic silicone fluid used in this embodiment accepts the caprylic/capric triglyceride and allows the EV to remain in solution. Because the formulation contains no water (or a negligible amount, such as less than .5%, due to absorption of atmospheric moisture) and no emulsifiers, the EV remains stable. Therefore, the present invention 55 contemplates formulations of EV in non-aqueous solutions, particularly comprising at least 20% to 25% to

30% to 35% to 40% to 45% to 50%, or up to 60%, 70%, 80% or 90% silicone thereby explicitly excluding water-in-oil, or oil-in-water, emulsions. Additionally, besides caprylic capric triglyceride, EV can be solubilized in jojoba oil, sunflower oil or squalane or any other solubilizer which is accepted by the silicones.

[0071] For example, a water-in-oil formulation in which EV decomposes within 24 hours (and thus is not a formulation of the present invention) is:

water (62.8%), butanediol (5%), glycerin (4%), ethoxydiglycol (3%), glycereth-7 (2%), polysorbate 20 (0.2%), glyceryl stearate (and) PEG-100 stearate (4%), isocetyl stearate (3.5%), jojoba oil (3.5%), mineral oil (3%), isostearyl palmitate (3%), PEG-7 glyceryl cocoate (2%), isocetyl alcohol (2%), cetyl ricinoleate (1%), ethyl vanillin (1%).

Pharmaceutical and Cosmetic Compositions

[0072] The DHE and/or IE and/or EV compositions, and/or salts, esters, ethers, or derivatives thereof described herein are preferably administered in the form of pharmaceutical or cosmetic compositions. Such compositions can be prepared in a manner well known to those of ordinary skill in the pharmaceutical and cosmetic arts. As noted above, the compositions may further comprise cinnamaldehyde.

[0073] Generally, the compositions of this invention are administered in a cosmetic amount or a therapeutically or cosmetically effective dose. The amount of the compound actually administered in a therapeutic setting may, typically be determined by a physician, such as a dermatologist in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like. In cosmetic settings, the amount to be applied is selected to achieve a desired cosmetic effect.

[0074] The cosmetic compositions of this invention are to be administered topically (or via other epithelial administration, where desired). The pharmaceutical compositions of this invention are to be administered topically, transdermally or systemically such as orally or by injection or other suitable methods known by those of ordinary skill in the art.

[0075] In such compositions, the DHE and/or IE and/or EV and/or salts, esters, ethers, or derivatives thereof is usually a minor component (or components) (from about 0.001 to about 20% by weight, or preferably from about 0.01 to about 10% or .1% to 5%, or 1.0% to 3%, by weight), with the remainder being various vehicles or carriers and processing aids helpful for forming the desired dosing form and for carrying the active agent into the epidermis. Cinnamaldehyde may be added in amounts of .001% to %5, or more preferably .01% to 1%, or more preferably .05% to .5%. One preferred formulation

comprises .4% DHE, .16% IE acetate, and .04% EV, and optimally .1% cinnamaldehyde.

[0076] Topical cosmetic forms and topical pharmaceutical dosing forms can include lotions, shampoos, soaks, gels, creams, ointments and pastes. Lotions commonly employ a water or alcohol base. Gels are semi-solid emulsions or suspensions. Creams generally contain a significant proportion of water in their base while ointments are commonly more oily.

[0077] Liquid forms, such as lotions suitable for topical administration or for cosmetic application, may include a suitable aqueous or non-aqueous vehicle with buffers, suspending and dispensing agents, thickeners, penetration enhancers, and the like. More solid forms such as creams or pastes or the like may include, for example, any of the following ingredients: water, oil, alcohol or grease as a substrate with surfactant, polymers such as polyethylene glycol, thickeners, solids and the like. Liquid or solid formulations may include enhanced delivery technologies such as liposomes, microsomes, microsponges and the like.

[0078] The above-described components for liquid, semisolid and solid topical compositions are merely representative. Other materials as well as processing techniques and the like are set forth in Remington's Pharmaceutical Sciences, cited above, which is hereby expressly incorporated herein by reference in its entirety.

[0079] When pharmaceutical compositions are to be administered transdermally they typically are employed as liquid solutions or as gels. In these settings the concentration of DHE and/or IE and or EV and/or salts, esters, ethers, or derivatives thereof ranges, individually or in combination, from about 0.1% to about 20%, and preferably from about 0.1% to about 10%, and more particularly from 1% to 5%, of the composition with the remainder being aqueous mixed or nonaqueous vehicle, such as alcohols and the like, suspending agents, gelling agents, surfactant, and the like. Examples of suitable such materials are described below.

[0080] The compositions comprising DHE and/or IE and/or EV or salts, esters, ethers, or derivatives thereof as defined herein can also be administered in sustained release transdermal forms or from transdermal sustained release drug delivery systems. A description of representative sustained release materials can be found in the incorporated materials in Remington's Pharmaceutical Sciences, cited above.

[0081] The compositions for systemic administration include compositions for oral administration, that is liquids and solids, and liquid compositions or suspensions for injection and formulations for rectal, colonic, vaginal, nasal, and parenteral administration.

[0082] Compositions for oral administration can take the form of bulk liquid solutions or suspensions, or bulk powders. More commonly, however, the compositions are presented in unit dosage forms to facilitate accurate dosing. The term "unit dosage forms" refers to physically discrete units suitable as unitary dosages for human sub-

jects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical occupant. Typical unit dosage forms include profiled, premeasured ampules or syringes of the liquid compositions or pills, tablets, capsules or the like in the case of solid compositions. In such compositions, the DHE and/or IE and/or EV and/or salts, esters, ethers, or derivatives thereof alone or in combination is usually a minor component (from about 0.01 to about 20% by weight or preferably from about 0.1 to about 10% by weight or 1% to 5% by weight) with the remainder being various vehicles or carriers and processing aids helpful for forming the desired dosing form.

[0083] Liquid forms suitable for oral administration may include, but are not limited to, a suitable aqueous or non-aqueous vehicle with buffers, suspending and dispensing agents, colorants, flavors and the like. Solid forms may include, but are not limited to, any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an occupant such as starch or lactose; a disintegrating agent such as alginic acid, Primogel™, or corn starch; a lubricant such as magnesium stearate; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

[0084] Injectable compositions are typically based upon injectable sterile saline or phosphate-buffered saline or other injectable carriers known in the art. As before, the DHE and/or IE and/or EV and/or salts, esters, ethers, or derivatives thereof in such compositions are typically minor components, often being from about 0.005 to 10% by weight, or .1 to 2% by weight, for example, with the remainder being the injectable carrier and the like.

[0085] The above-described compositions for orally or epidermally administrable or injectable compositions are merely representative. Other materials as well as processing techniques and the like are set forth in the part of Remington's Pharmaceutical Sciences, cited above.

[0086] The following formulation examples illustrate representative cosmetic and pharmaceutical compositions of this invention. The present invention, however, is not limited to the following pharmaceutical compositions.

EXEMPLARY FORMULATIONS

Formulation 1-Lotion

[0087] In a preferred embodiment, DHE was formulated into a topical lotion a 2% final concentration and tested for its ability to block a UVB-irradiation induced sunburn. Volunteers were irradiated with a UVB radiation source sufficient to produce a sunburn (approximately a 3 MED dose) and after irradiation 1 ml of a 2% topical DHE lotion was applied to one of two sites. Another irradiated site

was left untreated. By 2-6 hours erythema was noted in the untreated UVB site while the side treated with the topical DHE lotion showed no erythema.

5 Formulation 2-Liquid

[0088] A formulation of DHE (125 mg total), and xanthan gum (4 mg) are blended, passed through a No. 10 mesh U.S. sieve, and then mixed with a previously made 10 solution of microcrystalline cellulose and sodium carboxymethyl cellulose (11:89, 50 mg) in a water/isopropanol (75:25) mixture. Sufficient water/isopropanol and salicylic acid to 2% by weight or a sufficient amount to maintain a pH of 3-6.5 are then added to produce a total 15 volume of 5 mL.

Formulation 3-Cream

[0089] A commercial mineral oil-water cold cream 20 base is obtained. To 100 grams of this base, 1 gm of DHE and/or IE and/or EV and/or salts, esters or derivatives thereof as a fine powder or liquid, is added with continuous mixing and stirring to suspend the powder in the base yielding a cosmetic or pharmaceutical composition.

[0090] In one embodiment, this composition includes the following: deionized water (55.6% by weight); niacinamide (2.0%); glycerin (4.0%); phenonip (1.0%); propylene glycol (5.0%); transcutol (3.2%); jojoba Oil (3.5%); isocetyl alcohol (2.0%); isocetyl stearate (3.5%); mineral oil (3.0%); dihydroeugenol (1.0 %); salicylic acid (2%); isostearyl palmitate (3.0%); PEG-7 glyceryl cocoate (2.0%); Glycereth-7 (2.0%); POLYSORBATE-20® (0.2%); cetyl ricinoleate (1.0%); glyceryl stearate/PEG-35 100 stearate (4.0%); and SEPIGEL® (2.0%).

Formulation 4-Cream

[0091] Deionized water (56.4 % by weight); caffeine 40 (1.0 %); butanediol (4.0 %); glycerin (1.0 %); phenonip (1.0 %); POLYSORBATE-20® (0.2 %); niacinamide (2.0 %); arlacet (6.0 %); isocetyl stearate (3.5 %); cetyl ricinoleate (1.0 %); protaderm B (10.0 %); jojoba oil (3.5 %); stearyl alcohol (3.0 %); cetearyl 20 (0.4%); PEG-12 (3.0 %); dihydroeugenol (2.0 %); SEPIGEL™ (2.0%).

Formulation 5-Cream

[0092] Water (44.4% by weight); niacinamide (2.0%); 50 propylene glycol (3%); PEG-100 stearate (1%); ajidew (1%); glycerin (1%); EDTA (0.1%); carbopol (20%); squalene (2%); jojoba oil (2%); stearic acid (2%); glyceryl stearate (1%); cetyl alcohol (1.5%); vitamin E (1%); dimethicone (1%); caprylic/capric triglyceride (2%); dihydroeugenol (2%); petrolatum (1%); promulgen D (2%); PP2 (2%); glycol stearate (1%); dimethyl isosorbide-DMI (3%); and added after emulsion: germaben (1%) < 55°; and triethanolamine (1%).

Formulation 6-Tablets

[0093] A formulation of DHE and/or IE and/or EV and/or salts, esters or derivatives thereof and salicylic acid to maintain a pH of 3-6.5 is mixed with dry gelatin binder and starch diluent in a 0.1: 1:1 weight ratio. A lubricating amount of magnesium stearate is added and the mixture is tabletted into 210 mg tablets containing 10 mg of DHE, IE or EV or other compounds described herein.

Formulation 7-Injection

[0094] A formulation of DHE and/or IE and/or salts, esters or derivatives thereof, and salicylic acid to maintain a pH of 3-6.5, is dissolved in injectable aqueous saline medium at a concentration of 1 mg/ml.

Ethyl Vanillin (EV) Formulations - Topical

[0095] The following are five non-aqueous formulations comprising EV which maintain EV in a chemically-stable condition.

1. Cyclomethicone (and) dimethiconol (20-90%), caprylic/capric triglyceride (1-20%), dimethicone (1-10%), ethyl vanillin (0.1-5%).
2. Cyclomethicone (and) dimethiconal (5-60%), cyclpentasiloxane (and) dimethicone crosspolymer (5-60%), caprylic/capric triglyceride (1-20%), dimethicone (1-10%), jojoba oil (1-10%), squalane (1-10%), ethyl vanillin (0.1-5%).
3. SD alcohol (5-50%), cetearyl octanoate (1-10%), vitamin E (0.5%), cyclomethicone (10-50%), PPG-26 oleate (1-10%), caprylic/capric triglyceride (1-10%), ethyl vanillin (0.1-5%).
4. PPG-15 Stearylether cyclomethicone (10-50%), Sunflower oil (10-50%), isopropyl alcohol (1-10%), isostearyl palmitate (1-10%), ethyl vanillin (0.1-5%).
5. Benzyl laurate (5-25%), PPG-10 butanediol (1-10%), mineral oil (10-70%), squalane (1-10%), caprylic/capric triglyceride (1-10%), ethyl vanillin (0.1-5%).

Utility and Dosing

[0096] The composition and methods of this invention can be used topically to treat dermatological conditions such as actinic keratitis, acne, scarring, allergic contact dermatitis, atopic dermatitis, contact dermatitis, erythema (sunburn), hand eczema, itch, irritant contact dermatitis, psoriasis, seborrheic eczema (dermatitis), other eczemas, rosacea, hyperpigmentation, alopecia areata, damage from radiation (radiation dermatitis) including UV radiation, IR radiation and any other ionizing radiation and the like, and other dermatological conditions described elsewhere herein.

[0097] The compositions, both cosmetic and pharma-

ceutical, can also be used to treat and inhibit sunburn and to treat and prevent other forms of UV-induced inflammation and damage as well as damage from other forms of ionizing radiation.

[0098] In these applications the cosmetic and pharmaceutical compositions are administered topically to achieve a desired cosmetic effect or a topical therapeutic effect.

[0099] In these uses the dose levels or application levels can be expressed in terms of the amount of DHE and/or IE and/or EV and/or salts, esters, ethers, or derivatives thereof delivered to the skin. For example, 1 to about 5 doses or applications per day, each containing from about 0.001 g to about 1 gram of each of DHE and/or IE and/or EV or salts, esters, ethers, or derivatives thereof or combinations thereof can be used.

[0100] Alternatively, dose levels can be expressed in terms of the volume of formulated composition administered. For example, 1 to about 5 doses or applications per day, each containing from about 1 to about 30 grams of composition containing alone or in combination from about 0.01% to about 10% by weight of each of DHE and/or IE and/or EV and/or salts, esters, ethers, or derivatives thereof and especially from 0.02% to about 8% by weight or .1% to 5% by weight, or more preferably 1.0-4%.

[0101] When used in sun care products, such as sun-care lotion, the concentration of DHE and/or IE and/or EV or salts, esters, ethers, or derivatives thereof can be as set forth above and the product can be applied as needed based on the intensity and duration of sun exposure before, during, or after sun exposure.

[0102] Additionally, since the DHE and/or IE and/or EV or salts, esters, ethers, or derivatives thereof have been discovered to effectively inhibit the release of cytokines, such as IL-1 α or others cited herein, such compounds are useful for treating diseases characterized by an overproduction or a dysregulated production of cytokines, particularly IL-1 α wherein treatment causes reduction of said cytokines. Elevated levels of IL-1 α and other cytokines, as noted above, are associated with a wide variety of inflammatory conditions, including rheumatoid arthritis, septic shock, erythema nodosum leprosy, septicemia, adult respiratory distress syndrome (ARDS), inflammatory bowel disease (IBD), uveitis, damage from ionizing radiation and the like.

[0103] In the case of transdermal administration to treat such inflammatory conditions, one can administer a quantity of composition of the invention to a surface area of skin suitable to achieve an effective systemic bloodstream concentration of DHE and/or IE and/or EV, or salts, esters, ethers, or derivatives thereof, e.g., of from about 0.5 μ M to about 1000 μ M and more preferably from about 1 μ M to about 500 μ M, or other concentrations noted herein. In formulations to be applied topically or systemically it is preferred that the skin layer (epidermis and/or dermis) to be affected by the active agent maintain a concentration of active agent therein in a range of from

1 μM to 1000 μM , more preferably from 10 μM to 500 μM , more preferably from 50 μM to 300 μM , and still more preferably between 100 μM to 200 μM .

[0104] Injection dose levels for treating inflammatory conditions can range from (but are not limited to) about 0.01 mg/kg/hour to at least 1 mg/kg/hour, all for from about 1 to about 120 hours and especially 24 to 96 hours. A preloading bolus of from, for example, about 0.01 mg/kg to about 1 mg/kg or more may also be administered to achieve adequate steady state levels.

[0105] With oral dosing, one to five, two to four, or typically three oral doses per day are representative regimens. Using these dosing patterns, each dose preferably provides from about 0.01 to about 10 mg/kg of the DHE and/or IE and/or EV or salts, esters, ethers, or derivatives thereof, with preferred doses each providing from about 0.01 to about 5 mg/kg, or other dosages described herein.

[0106] The DHE and/or IE and/or EV or salts, esters, ethers, or derivatives thereof can be administered as the sole active agent or they can be administered alone or in combination, or in combination with other active agents.

[0107] Changes may be made in the formulations and methods of use of the various compositions and formulations described herein or in the steps or the sequence of steps of the methods described herein without departing from the spirit and scope of the claims as described below.

[0108] All patents, published patent applications, articles or references cited herein are expressly incorporated herein by reference in their entireties.

[0109] Viewed from a first aspect the present invention provides a composition for topical treatment of a dermatological disorder, disease or condition in a subject, comprising:

an active agent comprising at least one of isoeugenol, dihydroeugenol, or ethyl vanillin, or a salt, ester, or ether thereof; and

a pharmaceutically-acceptable carrier or vehicle effective in carrying or enabling passage of the active agent into the epidermis or dermis of the skin of the subject where the active agent has its effect.

[0110] Preferably the dermatological disorder, disease or condition of the subject is at least one of rosacea, allergic contact dermatitis, irritant contact dermatitis, seborrheic dermatitis, radiation dermatitis, erythema, psoriasis, atopic dermatitis, eczemas, actinic keratitis, acne, hyperpigmentation, scarring, aging and alopecia areata.

[0111] Preferably the pharmaceutically-acceptable carrier or vehicle comprises a penetration enhancer. Particularly preferably the carrier primarily comprises at least 25% by weight of a silicone.

[0112] Preferably the active agent primarily comprises soluble ethyl vanillin and the pharmaceutically-acceptable carrier or vehicle is non-aqueous and is absent emulsifiers and wherein the soluble ethyl vanillin remains sta-

ble for at least 3 months.

[0113] Preferably the composition comprises cinnamaldehyde.

[0114] Preferably the composition is effective in percutaneous transmission such that the active agent can be maintained in the epidermis or dermis at a concentration in a range of from 10 μM to 500 μM .

[0115] Viewed from a further aspect the present invention provides a method of treating a dermatological disorder, disease or condition in a subject, comprising:

providing a composition comprising:

an active agent comprising at least one of isoeugenol, dihydroeugenol, or ethyl vanillin, or salts, esters or ethers thereof; and a pharmaceutically-acceptable carrier or vehicle effective in carrying or enabling passage of the active agent into the epidermis or dermis of the skin of the subject; and

topically applying a quantity of the composition to a portion of the skin of the subject affected by the disorder, disease, or condition.

[0116] Preferably the dermatological disorder, disease, or condition is at least one of rosacea, allergic contact dermatitis, irritant contact dermatitis, radiation dermatitis, seborrheic dermatitis, erythema, psoriasis, atopic dermatitis, eczemas, actinic keratitis, acne, hyperpigmentation, aging, scarring, and alopecia areata.

[0117] Preferably the composition comprises cinnamaldehyde.

[0118] Preferably the composition comprises a penetration enhancer.

[0119] Preferably the active agent of the composition is ethyl vanillin or a salt, ester, or ether thereof, and wherein the dermatological condition is aging and/or scarring. Particularly preferably the composition stimulates dermal production of collagen, elastin and TIMPs and inhibits production of MMPs.

[0120] Preferably the active agent comprises dihydroeugenol or a salt, ester, or ether thereof and the dermatological condition is rosacea or atopic dermatitis. Particularly preferably the composition further comprises cinnamaldehyde. Alternatively particularly preferably the active agent further comprises isoeugenol or a salt, ester, or ether thereof. Alternatively particularly preferably the active agent comprises dihydroeugenol.

[0121] Preferably the active agent comprises dihydroeugenol or salt, ester, or ether thereof and the dermatological condition is radiation dermatitis and/or erythema. Particularly preferably the active agent comprises dihydroeugenol. More preferably the dihydroeugenol comprises .5% to 5% by weight of the composition. Alternatively more preferably the dihydroeugenol comprises 1% to 4% by weight of the composition. Alternatively more preferably the active agent further comprises iso-

eugenol and/or salt, ester, or ether thereof.

[0122] Preferably the active agent is isoeugenol or a salt, ester, or ether thereof and the dermatological condition is hyperpigmentation.

[0123] Preferably the active agent is dihydroeugenol and isoeugenol or salts, esters, and methods thereof and cinnamaldehyde and the dermatological condition is acne.

[0124] Preferably the active agent comprises dihydroeugenol, or a salt, ester or ether thereof, and isoeugenol, or a salt, ester or ether thereof, and wherein the dermatological condition is psoriasis.

[0125] Preferably the active agent is maintained in the epidermis or dermis of the skin at a concentration in a range of from 10 μ M to 500 μ M.

Claims

1. Transdermal composition for use in treating a dermatological disorder, disease or condition in a subject, the transdermal composition comprising:

an active agent comprising at least one of isoeugenol, dihydro-isoeugenol, or ethyl vanillin, or salts, esters or ethers thereof; and a pharmaceutically-acceptable carrier or vehicle effective in carrying or enabling passage of the active agent into the epidermis or dermis of the skin of the subject.

2. Transdermal composition for use as claimed in claim 1 wherein the dermatological disorder, disease, or condition is at least one of rosacea, allergic contact dermatitis, irritant contact dermatitis, radiation dermatitis, seborrheic dermatitis, erythema, psoriasis, atopic dermatitis, eczemas, actinic keratitis, acne, hyperpigmentation, aging, scarring, and alopecia areata.

3. Transdermal composition for use as claimed in claim 1 wherein the active agent is isoeugenol or a salt, ester or ether thereof and the dermatological condition is hyperpigmentation.

4. Transdermal composition for use as claimed in claim 1 or 2 wherein the dermatological disorder, disease, or condition is hyperpigmentation.

5. Transdermal composition for use as claimed in any of claims 1 to 4 wherein the active agent comprises at least one of an isoeugenol ester and a salt thereof.

6. Transdermal composition for use as claimed in claim 5 wherein the isoeugenol ester comprises isoeugenyl acetate.

7. Transdermal composition for use as claimed in claim

5 or 6, wherein the active agent further comprises at least one member chosen from dihydroeugenol, a salt thereof, and an ester thereof.

5 8. Transdermal composition for use as claimed in any of claims 5, 6 or 7, wherein the active agent further comprises at least one member chosen from isoeugenol and a salt thereof.

10 9. Transdermal composition for use as claimed in any of claims 5 to 8, wherein the active agent further comprises at least one member chosen from soluble ethyl vanillin, a salt thereof, and an ester thereof.

15 10. Transdermal composition for use as claimed in any of claims 5 to 9, wherein the pharmaceutically-acceptable carrier or vehicle comprises a penetration enhancer.

20 11. Transdermal composition for use as claimed in any of claims 5 to 10, wherein the pharmaceutically-acceptable carrier or vehicle comprises a silicone.

25 12. Transdermal composition for use as claimed in any of claims 5 to 11 further comprising cinnamaldehyde.

13. Transdermal composition for use as claimed in any of claims 5 to 12 further comprising salicylic acid.

30 14. Transdermal composition for use as claimed in any of claims 5 to 13 further defined as being effective in percutaneous transmission such that the active agent can be maintained in the epidermis or dermis at a concentration in a range of from 1 μ M to 1000 μ M.

35 15. Transdermal composition for use as claimed in any of claims 5 to 14, wherein the active agent is present at about 0.001% to about 20% by weight of the transdermal composition.

45

50

55




Figure 1




Figure 2

**Effects of Dihydroeugenol on TPA induced TNF- α
In Keratinocytes**




Figure 3

**Effects of dihydroeugenol on LPS induced
inflammatory cytokines in Monocytes**




Figure 4





Figure 5




C



B

Figure 7

A



B



A

Figure 6

Stimulation of Collagen I mRNA by Dihydroeugenol, Isoeugenol, and Ethyl Vanillin




Figure 8

Stimulation of Elastin mRNA in Fibroblasts by Dihydroeugenol, Isoeugenol, and Ethyl Vanillin




Figure 9

**Stimulation of HAS-2 mRNA in fibroblasts by
Dihydroeugenol Isoeugenol, and Ethyl Vanillin**




Figure 10

**Stimulation of TIMP-1 mRNA in Fibroblasts by
Dihydroeugenol, Isoeugenol and Ethyl Vanillin**




Figure 11




Figure 12





Figure 13

Depigmentation of Human Melanocytes
Treated with TH-212 and TH-213



Note the pronounced loss of melanin in cells treated with
either TH-212 or TH-213 for only 9 days

Figure 14

Topical Formulation of TH-212-ester plus Salicylic Acid
Clears Acne and Hyperpigmentation in 4 Weeks





Figure 15

**TH 212-ester/ 2% Salicylic Acid
Eliminates Hyperpigmentation In 30 Days**



**Hyperpigmentation
completely cleared in 30 days**

**Note hyperpigmentation
of upper mid-forehead**

Figure 16

Reduction of Sun-Induced Hyperpigmentation
with TH-212-ester/ 2% salicylic acid for 30 days







Figure 17

Close up After Treatment
Close up Before Treatment


30 days Treatment



Before Treatment

TH-212/TH-213 Reduces the Amount of Tyrosinase in Human Melanocytes

In this assay there is NO TH-212 or TH-213 in the assay tube. The absence of tyrosinase activity (dopa conversion to melanin) is therefore likely due to a lack of tyrosinase in the cell lysates used in the assay.



Note the lack of tyrosinase activity in cell lysates from TH-212 or TH-213 treated melanocytes.

Figure 18




Figure 19

B

A



EUROPEAN SEARCH REPORT

Application Number

EP 17 17 4953

5

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
10	X WO 2004/000305 A1 (GARDOVIC MILENKA [SE]) 31 December 2003 (2003-12-31) * claims; examples *	1-3	INV. A61K8/30 A61P17/00 A61P17/02
15	X WO 03/043621 A1 (CUTANIX CORP [US]; ENGLES CHARLES R [US]; FULLER BRYAN [US]; PILCHER B) 30 May 2003 (2003-05-30) * paragraph [0145] *	1,2,9	A61P17/06 A61P17/10 A61P17/14 A61K9/70 A61K31/085
20	A MURAKAMI Y ET AL: "Dehydrodiisoeugenol, an isoeugenol dimer, inhibits lipopolysaccharide-stimulated nuclear factor kappa B activation and cyclooxygenase-2 expression in macrophages", ARCHIVES OF BIOCHEMISTRY AND BIOPHYSICS, ACADEMIC PRESS, US, vol. 434, no. 2, 15 February 2005 (2005-02-15), pages 326-332, XP004705012, ISSN: 0003-9861, DOI: DOI:10.1016/J.ABB.2004.11.013 * abstract *	1-15	TECHNICAL FIELDS SEARCHED (IPC)
25			A61K A61P A61Q
30	A S. TANAKA ET AL: "Contact allergy to isoeugenol and its derivatives: problems with allergen substitution", CONTACT DERMATITIS: ENVIRONMENTAL AND OCCUPATIONAL DERMATITIS, vol. 51, no. 5-6, 1 November 2004 (2004-11-01), pages 288-291, XP055407587, US ISSN: 0105-1873, DOI: 10.1111/j.0105-1873.2004.00446.x * abstract *	1-15	
35			
40			
45			
4	The present search report has been drawn up for all claims		
50	Place of search Munich	Date of completion of the search 20 September 2017	Examiner Zimmer, Barbara
	CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document	T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
55			

EPO FORM 1503 03.82 (P04C01)



EUROPEAN SEARCH REPORT

Application Number

EP 17 17 4953

5

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (IPC)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
10	A DE 202 21 578 U1 (HOENZELAER NORBERT [DE]) 8 June 2006 (2006-06-08) * page 8; claims *	1-15	
15	A US 2007/053849 A1 (DOYLE MATTHEW J [US] ET AL) 8 March 2007 (2007-03-08) * claims *	1-15	
20			
25			
30			
35			
40			
45			
46	The present search report has been drawn up for all claims		
47	Place of search	Date of completion of the search	Examiner
48	Munich	20 September 2017	Zimmer, Barbara
49	CATEGORY OF CITED DOCUMENTS		
50	X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document

EPO FORM 1503 03.82 (P04C01)

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 17 17 4953

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

20-09-2017

10	Patent document cited in search report	Publication date		Patent family member(s)	Publication date
15	WO 2004000305 A1	31-12-2003	AU	2003237742 A1	06-01-2004
			WO	2004000305 A1	31-12-2003
20	WO 03043621 A1	30-05-2003	AT	330591 T	15-07-2006
			AU	2002326363 A1	10-06-2003
			BR	0214208 A	26-10-2004
			CA	2464867 A1	30-05-2003
			CN	1589136 A	02-03-2005
			CN	102973417 A	20-03-2013
			DE	60212683 T2	28-06-2007
			DK	1443915 T3	23-10-2006
			EA	200400689 A1	24-02-2005
			EP	1443915 A1	11-08-2004
			ES	2266554 T3	01-03-2007
			IL	161545 A	30-11-2010
			JP	4587669 B2	24-11-2010
			JP	2005509662 A	14-04-2005
			JP	2010059203 A	18-03-2010
			KR	20050018633 A	23-02-2005
			MX	PA04004673 A	17-05-2005
			NZ	552216 A	31-07-2008
			PT	1443915 E	30-11-2006
			TW	I297604 B	11-06-2008
			WO	03043621 A1	30-05-2003
			ZA	200403282 B	03-05-2005
35	DE 20221578 U1	08-06-2006		NONE	
40	US 2007053849 A1	08-03-2007	AU	2007318251 A1	15-05-2008
			BR	PI0718699 A2	15-01-2013
			CA	2669019 A1	15-05-2008
			CN	101534823 A	16-09-2009
			CN	104958187 A	07-10-2015
			EP	2079468 A1	22-07-2009
			EP	2415473 A1	08-02-2012
			ES	2395854 T3	15-02-2013
			JP	5280366 B2	04-09-2013
			JP	2010508346 A	18-03-2010
			MY	157928 A	15-08-2016
			RU	2009114589 A	20-12-2010
			US	2007053849 A1	08-03-2007
			WO	2008057136 A1	15-05-2008
55	EPO FORM P0459				

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 99834507 P [0001]
- US 61126478 A [0001]
- US 3909816 A [0041]
- US 4405616 A [0041]
- US 4801586 A [0041]
- US 4861764 A [0041]
- US 4886783 A [0041]
- US 4983396 A [0041]
- US 5118845 A [0041]
- US 5196410 A [0041]

摘要

本發明提供了包括化合物，特別是二氫丁子香酚 (DHE) 和/或異丁子香酚 (IE) 和/或乙基香蘭素 (EV) 或其鹽、酯、醚或衍生物的組合物，以及用於將它們局部或全身傳遞以治療炎症相關的病症和其他如本文所述的皮膚病症的方法。這些 DHE 和/或 IE 和/或 EV 或其鹽、酯、醚或衍生物也是以可測量程度經表皮或經皮膚傳遞，因此可在體內具有全身和/或局部的消炎作用。在該方法的其中一個方面，本發明是針對治療患有皮膚疾病或病症的病人的方法，該方法對所述患者局部施加藥物或美容組合物，該組合物包括藥物上或美容上可接受的局部載體和 DHE 和/或 IE 和/或 EV 或其鹽、酯、醚或其衍生物的製劑的對皮膚疾病、失調或病症有效的治療量。特別是，本發明針對使用包括一種或多種 DHE 和/或 IE 和/或 EV 或其鹽、酯、醚或衍生物的組合物來治療皮膚疾病、失調或病症，或與減少產生膠原蛋白、彈性蛋白或透明質酸相關的老化效應或色素沉著過度的方法。該製劑可進一步包括肉桂醛。