

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
5 February 2009 (05.02.2009)

PCT

(10) International Publication Number
WO 2009/018186 A2

(51) International Patent Classification:

A01N 43/56 (2006.01) *A61K 31/4155* (2006.01)
A01P 7/04 (2006.01)

(21) International Application Number:

PCT/US2008/071288

(22) International Filing Date:

28 July 2008 (28.07.2008)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/962,519 30 July 2007 (30.07.2007) US

(71) Applicant (for all designated States except US): **E. I. DU PONT DE NEMOURS AND COMPANY** [US/US]; 1007 Market Street, Wilmington, Delaware 19898 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **TAYLOR, Wendy, Sue** [US/US]; 29 Mallard Drive, Mount Laurel, New Jersey 08054 (US).

(74) Agent: **REHBERG, Edward, F.**; E. I. du Pont de Nemours and Company, Legal Patent Records Center, 4417 Lancaster Pike, Wilmington, Delaware 19805 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BI, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

(54) Title: FLY CONTROL METHOD



1

(57) **Abstract:** This invention relates to a method of treating myiasis of an animal by applying to the animal a composition comprising an parasitically effective amount of a compound of Formula 1, an N-oxide or a pharmaceutically or veterinarily acceptable salts salt thereof wherein R¹ is Me, Cl, Br or F; R² is F, Cl, Br, C₁-C₄ haloalkyl or C₁-C₄ haloalkoxy; R³ is F, Cl or Br, R⁴ is H; C₁-C₄alkyl, C₃-C₄ alkynyl, C₃-C₅cycloalkyl, or C₄-C₆ cycloalkylalkyl, each optionally substituted with one substituent selected from the group consisting of halogen, CN, SMe, S(O)Me, S(O)₂Me, and OMe; R⁵ is H or Me; R⁶ is H, F or Cl; and R⁷ is H, F or Cl.

WO 2009/018186 A2

TITLE
FLY CONTROL METHOD

FIELD OF THE INVENTION

This invention relates to certain methods for combatting biting flies and blowflies on
5 animals.

BACKGROUND OF THE INVENTION

Flies are not just a nuisance; they carry diseases which pose a serious health hazard to people and animals. Globally, they cause livestock and poultry production losses estimated in the billions of dollars. The growth and performance of nearly all farmed animals
10 are adversely affected by flies, especially when they are present in high numbers. Infested animals become harassed and feed intake is drastically reduced. The result: significant reductions of meat, milk and egg production and serious economic losses.

Non-biting flies often feed on secretions from the eyes, nose and any small wounds of livestock. This distracts animals from grazing, causing a reduction in growth and
15 productivity. Non-biting flies are not key vectors of any specific disease organisms, but because of their feeding and reproduction habits, and the structure of their feet and mouthparts, they can act as mechanical vectors for a whole range of pathogens, from viruses to helminthes.

Biting flies can cause even greater irritation to domestic animals, and they too
20 are vectors for disease transmission. However, because they feed on blood, they can also cause anemia and hypersensitivity. Biting flies therefore are considered by some to be a more serious problem in livestock production than non-biting flies.

However some non-biting flies often designated "blowflies" cause significant damage in their own right by virtue of their propensity to cause "myiisis" in susceptible
25 animals. Myiasis is an animal or human disease caused by parasitic dipterous fly larvae feeding on the host's necrotic or living tissue. Blowflies are the single most important parasite of the sheep industry in Australia. Losses are estimated at more than \$50 million yearly. These losses are caused by reduced growth of the sheep, reduced and inferior wool production, and extremely high labour costs expended in attempts to control the parasite.


30 Under normal conditions, blowflies do not attack live healthy sheep. If animals suffer open wounds, for example from branding or castration, some species of blowflies, deposit eggs in the wounds. These will hatch into maggots which eat the flesh of the animal.

The principal control method of adult populations of biting flies and blowflies involves topical insecticide applications to the livestock. Organophosphorus or organochlorine compounds are often used, usually in a spraying formulation.

There is a compelling need for improved insecticide formulations useful in the treatment and prevention of myiasis and the control of biting flies which the present invention addresses.

SUMMARY OF THE INVENTION

This invention relates to a method of controlling or preventing infestations of flies on an animal by applying to the animal a composition comprising a parasitically effective amount of a compound of Formula 1, or an *N*-oxide, or a salt thereof,



1

wherein

R¹ is Me, Cl, Br or F;

R² is F, Cl, Br, C₁–C₄ haloalkyl or C₁–C₄ haloalkoxy;

R³ is F, Cl or Br;

R⁴ is H; C₁–C₄ alkyl, C₃–C₄ alkenyl, C₃–C₄ alkynyl, C₃–C₅ cycloalkyl, or C₄–C₆ cycloalkylalkyl, each optionally substituted with one substituent selected from the group consisting of halogen, CN, SMe, S(O)Me, S(O)₂Me, and OMe;

R⁵ is H or Me;

R⁶ is H, F or Cl; and

R⁷ is H, F or Cl.

Of note are compounds of Formula I wherein:

R⁴ is H or C₁–C₄ alkyl optionally substituted with one substituent selected from the group consisting of CN, SMe and OMe;

R⁵ is H or Me;

R⁶ is H; and

R⁷ is H.

Of further note are:

a) Compounds of Formula 1 wherein

R¹ is Me or Cl;

5 R² is Cl, Br, CF₃, OCF₂H, OCF₃ or OCH₂CF₃; and

R⁴ is H, Me, Et, *i*-Pr, *t*-Bu, CH₂CN, CH(Me)CH₂SMe or C(Me)₂CH₂SMe.

b) Compounds of a) above wherein

R² is Cl, Br, CF₃ or OCH₂CF₃;

R⁴ is H, Me, Et or *i*-Pr; and

10 R⁵ is H.


c) Compounds of b) wherein:

R¹ is Me; R² is Br; R³ is Cl; R⁴ is Me.

Of further note are compounds of a, b, c above wherein R⁶ is H; and R⁷ is H.

A compound of special interest is:

15



or 3-bromo-1-(3-chloro-2-pyridinyl)-N-[4-cyano-2-methyl-6-[(methylamino)carbonyl]-phenyl]-1H-pyrazole-5-carboxamide.

The preferred compositions of the present invention are those, which comprise the 20 above preferred compounds. The preferred methods of use are those involving the above-preferred compounds.

This invention also relates to a method of treating myiasis of an animal by applying to the animal a composition comprising a parasitically effective amount of the compounds noted above.

25 The invention also comprises a compound of Formula 1 for use as a medicament.

The invention also relates to the use of a compound of Formula 1 in the manufacture of a medicament for the treatment of myiasis or the infestation of flies on an animal.

DETAILS OF THE INVENTION

As used herein, the terms “comprises,” “comprising,” “includes,” “including,” “has,” “having,” “contains” or “containing,” or any other variation thereof, are intended to cover a non-exclusive inclusion. For example, a composition, a mixture, process, method, article, or 5 apparatus that comprises a list of elements is not necessarily limited to only those elements but may include other elements not expressly listed or inherent to such composition, mixture, process, method, article, or apparatus. Further, unless expressly stated to the contrary, “or” refers to an inclusive or and not to an exclusive or. For example, a condition A or B is satisfied by any one of the following: A is true (or present) and B is false (or not present), A 10 is false (or not present) and B is true (or present), and both A and B are true (or present).

Also, the indefinite articles “a” and “an” preceding an element or component of the invention are intended to be nonrestrictive regarding the number of instances (i.e. occurrences) of the element or component. Therefore “a” or “an” should be read to include one or at least one, and the singular word form of the element or component also includes the 15 plural unless the number is obviously meant to be singular.

In the above recitations, the term “alkyl”, used either alone or in compound words such as “alkylthio” or “haloalkyl” includes straight-chain or branched alkyl, such as, methyl, ethyl, *n*-propyl, *i*-propyl, or the different butyl isomers. The term “halogen”, either alone or in compound words such as “haloalkoxy”, includes fluorine, chlorine, bromine or iodine. 20 Further, when used in compound words such as “haloalkyl”, or “haloalkoxy”, said alkyl or alkoxy may be partially or fully substituted with halogen atoms which may be the same or different. Examples of “haloalkyl” include F_3C , ClCH_2 , CF_3CH_2 and CF_3CCl_2 . Examples of “haloalkoxy” include CF_3O , HCF_2O , $\text{CCl}_3\text{CH}_2\text{O}$, $\text{HCF}_2\text{CH}_2\text{CH}_2\text{O}$ and $\text{CF}_3\text{CH}_2\text{O}$.

One skilled in the art will appreciate that not all nitrogen-containing heterocycles can 25 form *N*-oxides since the nitrogen requires an available lone pair for oxidation to the oxide; one skilled in the art will recognize those nitrogen-containing heterocycles which can form *N*-oxides. One skilled in the art will also recognize that tertiary amines can form *N*-oxides. Synthetic methods for the preparation of *N*-oxides of heterocycles and tertiary amines are very well known by one skilled in the art including the oxidation of heterocycles and tertiary 30 amines with peroxy acids such as peracetic and *m*-chloroperbenzoic acid (MCPBA), hydrogen peroxide, alkyl hydroperoxides such as *t*-butyl hydroperoxide, sodium perborate, and dioxiranes such as dimethyldioxirane. These methods for the preparation of *N*-oxides have been extensively described and reviewed in the literature, see for example:

T. L. Gilchrist in *Comprehensive Organic Synthesis*, vol. 7, pp 748–750, S. V. Ley, Ed., Pergamon Press; M. Tisler and B. Stanovnik in *Comprehensive Heterocyclic Chemistry*, vol. 3, pp 18–20, A. J. Boulton and A. McKillop, Eds., Pergamon Press; M. R. Grimmett and B. R. T. Keene in *Advances in Heterocyclic Chemistry*, vol. 43, pp 149–161, A. R. Katritzky, 5 Ed., Academic Press; M. Tisler and B. Stanovnik in *Advances in Heterocyclic Chemistry*, vol. 9, pp 285–291, A. R. Katritzky and A. J. Boulton, Eds., Academic Press; and G. W. H. Cheeseman and E. S. G. Werstiuk in *Advances in Heterocyclic Chemistry*, vol. 22, pp 390–392, A. R. Katritzky and A. J. Boulton, Eds., Academic Press.

Compounds of this invention can exist as one or more stereoisomers. The various 10 stereoisomers include enantiomers, diastereomers, atropisomers and geometric isomers. One skilled in the art will appreciate that one stereoisomer may be more active and/or may exhibit beneficial effects when enriched relative to the other stereoisomer(s) or when separated from the other stereoisomer(s). Additionally, the skilled artisan knows how to separate, enrich, and/or to selectively prepare said stereoisomers. Accordingly, the present 15 invention comprises compounds selected from Formula 1, *N*-oxides and salts thereof. Pharmaceutically or veterinarianily acceptable salts, suitable to the mode of administration, are contemplated. The compounds of the invention may be present as a mixture of stereoisomers, individual stereoisomers, or as an optically active form..

The salts of the compounds of the invention include acid-addition salts with inorganic 20 or organic acids such as hydrobromic, hydrochloric, nitric, phosphoric, sulfuric, acetic, butyric, fumaric, lactic, maleic, malonic, oxalic, propionic, salicylic, tartaric, 4-toluenesulfonic or valeric acids. In the compositions and methods of this invention, the salts of the compounds of the invention are preferably acceptable for the veterinary/pharmaceutical uses described herein.

25 The preferred compositions of the present invention are those, which comprise the above preferred compounds. The preferred methods of use are those involving the above-preferred compounds.

“Flies” are insects in the order Diptera, meaning “two-winged”. True flies have one 30 pair of wings used for flying. Posterior to the wings is a pair of stalked knob-like structures (called halteres), which are organs of balance. Flies undergo complete metamorphosis, i.e. the life cycle consists of the following stages: egg, larva (called a maggot), pupa, and adult. Each stage of the life cycle may be a target for control and intervention.

Flies may be categorized into two functional categories “biting” and “non-biting”.

“Biting flies” have specially adapted mouthparts well suited for piercing the host animal integument. The stable fly *Stomoxys calcitrans* is a good example of a biting fly. The stable fly has a proboscis which is used to pierce the skin and imbibe blood. Both the males and the females are bloodfeeders. The stable fly is often the only biting, blood-sucking fly 5 breeding in any appreciable numbers in and around confined-animal production facilities. Another example of a biting fly is the horn fly, *Haematobia irritans irritans*, which like the stable fly is a bloodsucker and has great economic impact. Like the stable fly the horn fly has piercing/sucking mouthparts.

“Blowflies” are defined as flies which are the etiologic agent of myiasis. By way of 10 example the Calliphoridae family, together with the Sarcophagidae and the Oestridae families, contain the species largely responsible for many of the important myiases of domestic animals and man. Major species of blowflies include *Lucilia sericata* (greenbottles), *Phormia terraenovae* (blackbottles), *Calliphora erythrocephala* and *Calliphora vomitoria* (bluebottles) in Europe. These flies are characterized by the color of 15 the metallic sheen on their body sections. *Lucilia cuprina*, *L. caeser*, *L. illustris*, *Phormia regina*, *Calliphora stygia*, *C. australis*, *C. fallax*, *Chrysomyia albiceps*, *C. chlorophyga*, *C. micropogon*, and *C. rufifacies* are examples of major species of blowflies in the tropics and subtropics. Blowflies are a particularly important problem in sheep farming.

The blowflies that attack sheep fall into two main categories:

- 20 (1) Primary flies, which are capable of initiating a strike on living sheep. These include *Lucilia* and *Phormia* spp. and some *Calliphora* spp.
- (2) Secondary flies, which cannot initiate a strike, but attack an area already struck or otherwise damaged. They frequently extend the injury, rendering the strike one of great severity. Examples include many *Calliphora* spp. and, in warmer climates, *Chrysomyia* spp.

25 A “parasitically effective amount” is the amount of active ingredient needed to achieve an observable effect diminishing the occurrence or activity of the target invertebrate parasite pest. One skilled in the art will appreciate that the parasitically effective dose can vary for the various compounds and compositions of the present invention, the desired parasitical effect and duration, the target invertebrate pest species, the animal to be protected, 30 the mode of application and the like, and the amount needed to achieve a particular result can be determined through simple experimentation

“Myiasis” is an animal disease caused by parasitic dipterous fly larvae feeding on the animal host's necrotic or living tissue. Colloquialisms for myiasis include “fly-strike” and

"fly-blown". Blowfly myiasis is often associated with sheep; however, many other animals may be affected.

"Treating" or "Treatment" as it applies to myiasis or infestation refers to both the prevention and control of myiasis or infestation respectively

5 Embodiments of the present invention include:

Embodiment 1. The method or uses described in the Summary of the Invention wherein the fly is a biting fly.

Embodiment 2. The method or use described in Embodiment 1 wherein the fly is a stable fly.

10 **Embodiment 3.** The method or use described in Embodiment 1 wherein the fly is horn fly.

Embodiment 4. The method or use described in any of Embodiments 1 through 3 wherein the animal is a herd animal.

Embodiment 5. The method or use described in Embodiment 4 wherein the animal is a cattle or sheep.

15 **Embodiment 6.** The method or use of any of Embodiments 1-5 wherein the composition comprises at least one additional component selected from the group consisting of solvents and/or carriers, emulsifiers and/or dispersing agents.

Embodiment 7. The method or use of Embodiment 6 and wherein the composition comprises at least one additional biologically active compound or agent.

20 **Embodiment 8.** The method or use of Embodiment 7 wherein the additional biologically active compound or agent is selected from the group consisting of macrocyclic lactones, acetyl cholinesterase inhibitors, arthropod growth regulators, GABA-gated chloride channel antagonists, mitochondrial electron transport inhibitors, nicotinic acetylcholine agonists/antagonists/activator, oxidative phosphorylation inhibitors, anthelmintics, sodium channel modulators or other antiparasitic compounds.

Embodiment 9. The method or use of Embodiment 8 wherein said biologically active compound is a macrocyclic lactone.

30 **Embodiment 10.** The method or use of Embodiment 8 wherein said biologically active compound is an acetyl cholinesterase inhibitor selected from the group of organophosphates and carbamates.

Embodiment 11. The method or use of Embodiment 8 wherein said biologically active compound is an arthropod growth regulator selected from the group of chitin synthesis

inhibitors, ecdysone agonists/disruptors, lipid biosynthesis inhibitor and juvenile hormone mimics.

Embodiment 12. The method or use of Embodiment 8 wherein said biologically active compound is a GABA-gated chloride channel antagonist.

5 **Embodiment 13.** The method or use of Embodiment 8 wherein said biologically active compound is a mitochondrial electron transport inhibitor.

Embodiment 14. The method or use of Embodiment 8 wherein said biologically active compound is a nicotinic acetylcholine agonist/antagonist/activator.

10 **Embodiment 15.** The method or use of Embodiment 8 wherein said biologically active compound is an oxidative phosphorylation inhibitor.

Embodiment 16. The method or use of Claim 8 wherein said biologically active compound is an anthelmintic.

Embodiment 17. The method or use of Claim 8 wherein said biologically active compound is a sodium channel modulator.

15 **Embodiment 18.** The method of treatment of myiasis or use described in the Summary of the Invention wherein the myiasis is caused at least in part by larvae selected from the taxonomic families Calliphoridae, Sarcophagidae or Oestridae.

20 **Embodiment 19.** The method or use of Embodiment 18 wherein the myiasis is caused at least in part by larvae selected from the taxonomic families Calliphoridae, Sarcophagidae or Oestridae.

Embodiment 20. The method or use of Embodiment 18 wherein the myiasis is caused at least in part by larvae of the family Calliphoridae.

25 **Embodiment 21.** The method or use of Embodiment 18 wherein the myiasis is caused at least in part by larvae which are selected from the group consisting of *Lucilia cuprina* and *Lucilia sericata*.

Embodiment 22. The method or use Embodiment18 wherein the myiasis is caused at least in part by larvae of *Lucilia cuprina*.

Embodiment 23. The method or use of Embodiment 18 wherein the myiasis is caused at least in part by larvae of *Lucilia sericata*.

30 **Embodiment 24.** The method or use Embodiment 18 wherein the animal is a cattle or sheep.

Embodiment 25. The method or use of any of Embodiments 18-24 wherein the composition comprises at least one additional component selected from the group consisting of solvents and/or carriers, emulsifiers and/or dispersing agents.

Embodiment 26. The method or use of Embodiment 25 and wherein the composition comprises at least one additional biologically active compound or agent.

Embodiment 27. The method or use of Embodiment 26 wherein the additional biologically active compound or agent is selected from the group consisting of macrocyclic lactones, acetyl cholinesterase inhibitors, arthropodgrowth regulators, GABA-gated chloride channel antagonists, mitochondrial electron transport inhibitors, nicotinic acetylcholine agonists/antagonists/activator, oxidative phosphorylation inhibitors, anthelmintics, sodium channel modulators or other antiparasitic compounds.

Embodiment 28. The method or use of Embodiment 27 wherein said biologically active compound is a macrocyclic lactone.

Embodiment 29. The method or use of Embodiment 27 wherein said biologically active compound is an acetyl cholinesterase inhibitor selected from the group of organophosphates and carbamates.

Embodiment 30. The method or use of Embodiment 27 wherein said biologically active compound is an arthropodgrowth regulator selected from the group of chitin synthesis inhibitors, ecdysone agonists/disruptors, lipid biosynthesis inhibitor and juvenile hormone mimics.

Embodiment 31. The method or use of Embodiment 27 wherein said biologically active compound is a GABA-gated chloride channel antagonist.

Embodiment 32. The method or use of Claim 27 wherein said biologically active compound is a mitochondrial electron transport inhibitor.

Embodiment 33. The method or use of Claim 27 wherein said biologically active compound is a nicotinic acetylcholine agonist/antagonist/activator.

Embodiment 34. The method or use of Claim 27 wherein said biologically active compound is an oxidative phosphorylation inhibitor.

Embodiment 35. The method or use of Claim 27 wherein said biologically active compound is an anthelmintic.

The embodiments above are intended to be illustrative and not limiting. Further aspects of the invention are discussed throughout the specification.

This invention also relates to a method of treating myiasis of an animal by applying to the animal a composition comprising a parasitically effective amount of a compound of **Formula 1**, or an *N*-oxide, or a pharmaceutically or veterinarianily acceptable salts salt thereof,

Therefore, the invention is understood to include the compounds described in the 5 Summary of the Invention (and compositions containing them) for use as an animal medicament, or more particularly an anti-myiasis animal medicament. The animals to be protected include those delineated in Embodiments 4 and 5. The flies include those delineated in Embodiments 2 and 3. The medicament may be presented in topical forms.

The invention is also understood to include the compounds described in the Summary 10 of the Invention in the manufacture of medicaments for the protection of an animal myiasis. The animals to be protected include those delineated in Embodiments 4 and 5. The flies include those delineated in Embodiments 2 and 3. The medicament may be presented in topical forms.

The invention is also understood to include the compounds described in the Summary 15 of the Invention for use in the manufacture of medicaments for the protection of an animal from myiasis. The animals to be protected include those delineated in Embodiments 4 and 5. The flies include those delineated in Embodiments 2 and 3. The medicament may be presented in topical forms.

The invention is also understood to include the compounds described in the Summary 20 of the Invention packaged and presented for the protection of an animal myiasis. The animals to be protected include those delineated in Embodiments 4 and 5. The flies include those delineated in Embodiments 2 and 3. The compounds of the invention may be packaged and presented as topical dosage forms.

The invention is also understood to include a process for manufacturing a composition 25 for protecting an animal from an invertebrate parasitic pest characterized in that a compound of **Formula 1** is admixed with at least one pharmaceutically or veterinarianily acceptable carrier. The animals to be protected include those delineated in Embodiments 4 and 5. The flies include those delineated in Embodiments 2 and 3. The compositions of the invention may be packaged and presented in topical dosage forms.

30 The compounds of **Formula 1** which can be used according to the invention, have an excellent action against biting flies and blowflies, whilst being very well tolerated by animals. The invention thus represents a genuine enrichment of the art.

The compounds according to the invention possess a good ectoparasiticidal activity, whilst being of low toxicity to animals.


Since the compounds of Formula 1 are both effective adulticides, and larvacides i.e. since they are effective in both the adult stage of the target parasites, and the juvenile stages 5 of the parasites they are particularly advantageous in the treatment of myiasis.

The compounds of Formula 1 can be prepared by the methods as described in US Patent Publication 2006/0111403A1 (herein incorporated by reference to the extent not inconsistent with the disclosure herein) and variations readily apparent to the skilled artisan. Synthetic methods for the preparation of *N*-oxides of heterocycles and tertiary amines are 10 very well known by one skilled in the art including the oxidation of heterocycles and tertiary amines with peroxy acids such as peracetic and *m*-chloroperbenzoic acid (MCPBA), hydrogen peroxide, alkyl hydroperoxides such as *t*-butyl hydroperoxide, sodium perborate, and dioxiranes such as dimethyldioxirane. These methods for the preparation of *N*-oxides have been extensively described and reviewed in the literature, see for example:

15 T. L. Gilchrist in *Comprehensive Organic Synthesis*, vol. 7, pp 748-750, S. V. Ley, Ed., Pergamon Press; M. Tisler and B. Stanovnik in *Comprehensive Heterocyclic Chemistry*, vol. 3, pp 18-20, A. J. Boulton and A. McKillop, Eds., Pergamon Press; M. R. Grimmett and B. R. T. Keene in *Advances in Heterocyclic Chemistry*, vol. 43, pp 149-161, A. R. Katritzky, Ed., Academic Press; M. Tisler and B. Stanovnik in *Advances in Heterocyclic Chemistry*, 20 vol. 9, pp 285-291, A. R. Katritzky and A. J. Boulton, Eds., Academic Press; and G. W. H. Cheeseman and E. S. G. Werstiuk in *Advances in Heterocyclic Chemistry*, vol. 22, pp 390-392, A. R. Katritzky and A. J. Boulton, Eds., Academic Press.

The following compounds, by way of example and not by limitation, are expected to be advantageous in the practice of the invention.

Table 1



1									
<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁵</u>
Me	Cl	F	H	H	Cl	Cl	F	H	H
Me	Cl	F	Me	H	Cl	Cl	F	Me	H
Me	Cl	F	Et	H	Cl	Cl	F	Et	H
Me	Cl	F	<i>i</i> -Pr	H	Cl	Cl	F	<i>i</i> -Pr	H
Me	Cl	F	<i>t</i> -Bu	H	Cl	Cl	F	<i>t</i> -Bu	H
Me	Cl	F	CH ₂ CN	H	Cl	Cl	F	CH ₂ CN	H
Me	Cl	F	CH(Me)CH ₂ SMe	H	Cl	Cl	F	CH(Me)CH ₂ SMe	H
Me	Cl	F	C(Me) ₂ CH ₂ SMe	H	Cl	Cl	F	C(Me) ₂ CH ₂ SMe	H
Me	Cl	F	Me	Me	Cl	Cl	F	Me	Me
Me	Cl	Cl	H	H	Cl	Cl	Cl	H	H
Me	Cl	Cl	Me	H	Cl	Cl	Cl	Me	H
Me	Cl	Cl	Et	H	Cl	Cl	Cl	Et	H
Me	Cl	Cl	<i>i</i> -Pr	H	Cl	Cl	Cl	<i>i</i> -Pr	H
Me	Cl	Cl	<i>t</i> -Bu	H	Cl	Cl	Cl	<i>t</i> -Bu	H
Me	Cl	Cl	CH ₂ CN	H	Cl	Cl	Cl	CH ₂ CN	H
Me	Cl	Cl	CH(Me)CH ₂ SMe	H	Cl	Cl	Cl	CH(Me)CH ₂ SMe	H
Me	Cl	Cl	C(Me) ₂ CH ₂ SMe	H	Cl	Cl	Cl	C(Me) ₂ CH ₂ SMe	H
Me	Cl	Cl	Me	Me	Cl	Cl	Cl	Me	Me
Me	Cl	Br	H	H	Cl	Cl	Br	H	H
Me	Cl	Br	Me	H	Cl	Cl	Br	Me	H
Me	Cl	Br	Et	H	Cl	Cl	Br	Et	H
Me	Cl	Br	<i>i</i> -Pr	H	Cl	Cl	Br	<i>i</i> -Pr	H
Me	Cl	Br	<i>t</i> -Bu	H	Cl	Cl	Br	<i>t</i> -Bu	H
Me	Cl	Br	CH ₂ CN	H	Cl	Cl	Br	CH ₂ CN	H
Me	Cl	Br	CH(Me)CH ₂ SMe	H	Cl	Cl	Br	CH(Me)CH ₂ SMe	H


<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁵</u>		<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁵</u>
Me	Cl	Br	C(Me) ₂ CH ₂ SMe	H		Cl	Cl	Br	C(Me) ₂ CH ₂ SMe	H
Me	Cl	Br	Me	Me		Cl	Cl	Br	Me	Me
Me	Br	F	H	H		Cl	Br	F	H	H
Me	Br	F	Me	H		Cl	Br	F	Me	H
Me	Br	F	Et	H		Cl	Br	F	Et	H
Me	Br	F	<i>i</i> -Pr	H		Cl	Br	F	<i>i</i> -Pr	H
Me	Br	F	<i>t</i> -Bu	H		Cl	Br	F	<i>t</i> -Bu	H
Me	Br	F	CH ₂ CN	H		Cl	Br	F	CH ₂ CN	H
Me	Br	F	CH(Me)CH ₂ SMe	H		Cl	Br	F	CH(Me)CH ₂ SMe	H
Me	Br	F	C(Me) ₂ CH ₂ SMe	H		Cl	Br	F	C(Me) ₂ CH ₂ SMe	H
Me	Br	F	Me	Me		Cl	Br	F	Me	Me
Me	Br	Cl	H	H		Cl	Br	Cl	H	H
Me	Br	Cl	Me	H		Cl	Br	Cl	Me	H
Me	Br	Cl	Et	H		Cl	Br	Cl	Et	H
Me	Br	Cl	<i>i</i> -Pr	H		Cl	Br	Cl	<i>i</i> -Pr	H
Me	Br	Cl	<i>t</i> -Bu	H		Cl	Br	Cl	<i>t</i> -Bu	H
Me	Br	Cl	CH ₂ CN	H		Cl	Br	Cl	CH ₂ CN	H
Me	Br	Cl	CH(Me)CH ₂ SMe	H		Cl	Br	Cl	CH(Me)CH ₂ SMe	H
Me	Br	Cl	C(Me) ₂ CH ₂ SMe	H		Cl	Br	Cl	C(Me) ₂ CH ₂ SMe	H
Me	Br	Cl	Me	Me		Cl	Br	Cl	Me	Me
Me	Br	Br	H	H		Cl	Br	Br	H	H
Me	Br	Br	Me	H		Cl	Br	Br	Me	H
Me	Br	Br	Et	H		Cl	Br	Br	Et	H
Me	Br	Br	<i>i</i> -Pr	H		Cl	Br	Br	<i>i</i> -Pr	H
Me	Br	Br	<i>t</i> -Bu	H		Cl	Br	Br	<i>t</i> -Bu	H
Me	Br	Br	CH ₂ CN	H		Cl	Br	Br	CH ₂ CN	H
Me	Br	Br	CH(Me)CH ₂ SMe	H		Cl	Br	Br	CH(Me)CH ₂ SMe	H
Me	Br	Br	C(Me) ₂ CH ₂ SMe	H		Cl	Br	Br	C(Me) ₂ CH ₂ SMe	H
Me	Br	Br	Me	Me		Cl	Br	Br	Me	Me
Me	CF ₃	F	H	H		Cl	CF ₃	F	H	H
Me	CF ₃	F	Me	H		Cl	CF ₃	F	Me	H
Me	CF ₃	F	Et	H		Cl	CF ₃	F	Et	H
Me	CF ₃	F	<i>i</i> -Pr	H		Cl	CF ₃	F	<i>i</i> -Pr	H
Me	CF ₃	F	<i>t</i> -Bu	H		Cl	CF ₃	F	<i>t</i> -Bu	H
Me	CF ₃	F	CH ₂ CN	H		Cl	CF ₃	F	CH ₂ CN	H
Me	CF ₃	F	CH(Me)CH ₂ SMe	H		Cl	CF ₃	F	CH(Me)CH ₂ SMe	H
Me	CF ₃	F	C(Me) ₂ CH ₂ SMe	H		Cl	CF ₃	F	C(Me) ₂ CH ₂ SMe	H

<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁵</u>
Me	CF ₃	F	Me	Me	Cl	CF ₃	F	Me	Me
Me	CF ₃	Cl	H	H	Cl	CF ₃	Cl	H	H
Me	CF ₃	Cl	Me	H	Cl	CF ₃	Cl	Me	H
Me	CF ₃	Cl	Et	H	Cl	CF ₃	Cl	Et	H
Me	CF ₃	Cl	<i>i</i> -Pr	H	Cl	CF ₃	Cl	<i>i</i> -Pr	H
Me	CF ₃	Cl	<i>t</i> -Bu	H	Cl	CF ₃	Cl	<i>t</i> -Bu	H
Me	CF ₃	Cl	CH ₂ CN	H	Cl	CF ₃	Cl	CH ₂ CN	H
Me	CF ₃	Cl	CH(Me)CH ₂ SMe	H	Cl	CF ₃	Cl	CH(Me)CH ₂ SMe	H
Me	CF ₃	Cl	C(Me) ₂ CH ₂ SMe	H	Cl	CF ₃	Cl	C(Me) ₂ CH ₂ SMe	H
Me	CF ₃	Cl	Me	Me	Cl	CF ₃	Cl	Me	Me
Me	CF ₃	Br	H	H	Cl	CF ₃	Br	H	H
Me	CF ₃	Br	Me	H	Cl	CF ₃	Br	Me	H
Me	CF ₃	Br	Et	H	Cl	CF ₃	Br	Et	H
Me	CF ₃	Br	<i>i</i> -Pr	H	Cl	CF ₃	Br	<i>i</i> -Pr	H
Me	CF ₃	Br	<i>t</i> -Bu	H	Cl	CF ₃	Br	<i>t</i> -Bu	H
Me	CF ₃	Br	CH ₂ CN	H	Cl	CF ₃	Br	CH ₂ CN	H
Me	CF ₃	Br	CH(Me)CH ₂ SMe	H	Cl	CF ₃	Br	CH(Me)CH ₂ SMe	H
Me	CF ₃	Br	C(Me) ₂ CH ₂ SMe	H	Cl	CF ₃	Br	C(Me) ₂ CH ₂ SMe	H
Me	CF ₃	Br	Me	Me	Cl	CF ₃	Br	Me	Me
Me	OCF ₂ H	F	H	H	Cl	OCF ₂ H	F	H	H
Me	OCF ₂ H	F	Me	H	Cl	OCF ₂ H	F	Me	H
Me	OCF ₂ H	F	Et	H	Cl	OCF ₂ H	F	Et	H
Me	OCF ₂ H	F	<i>i</i> -Pr	H	Cl	OCF ₂ H	F	<i>i</i> -Pr	H
Me	OCF ₂ H	F	<i>t</i> -Bu	H	Cl	OCF ₂ H	F	<i>t</i> -Bu	H
Me	OCF ₂ H	F	CH ₂ CN	H	Cl	OCF ₂ H	F	CH ₂ CN	H
Me	OCF ₂ H	F	CH(Me)CH ₂ SMe	H	Cl	OCF ₂ H	F	CH(Me)CH ₂ SMe	H
Me	OCF ₂ H	F	C(Me) ₂ CH ₂ SMe	H	Cl	OCF ₂ H	F	C(Me) ₂ CH ₂ SMe	H
Me	OCF ₂ H	F	Me	Me	Cl	OCF ₂ H	F	Me	Me
Me	OCF ₂ H	Cl	H	H	Cl	OCF ₂ H	Cl	H	H
Me	OCF ₂ H	Cl	Me	H	Cl	OCF ₂ H	Cl	Me	H
Me	OCF ₂ H	Cl	Et	H	Cl	OCF ₂ H	Cl	Et	H
Me	OCF ₂ H	Cl	<i>i</i> -Pr	H	Cl	OCF ₂ H	Cl	<i>i</i> -Pr	H
Me	OCF ₂ H	Cl	<i>t</i> -Bu	H	Cl	OCF ₂ H	Cl	<i>t</i> -Bu	H
Me	OCF ₂ H	Cl	CH ₂ CN	H	Cl	OCF ₂ H	Cl	CH ₂ CN	H
Me	OCF ₂ H	Cl	CH(Me)CH ₂ SMe	H	Cl	OCF ₂ H	Cl	CH(Me)CH ₂ SMe	H
Me	OCF ₂ H	Cl	C(Me) ₂ CH ₂ SMe	H	Cl	OCF ₂ H	Cl	C(Me) ₂ CH ₂ SMe	H
Me	OCF ₂ H	Cl	Me	Me	Cl	OCF ₂ H	Cl	Me	Me

<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁵</u>	<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁵</u>
Me	OCF ₂ H	Br	H	H	Cl	OCF ₂ H	Br	H	H
Me	OCF ₂ H	Br	Me	H	Cl	OCF ₂ H	Br	Me	H
Me	OCF ₂ H	Br	Et	H	Cl	OCF ₂ H	Br	Et	H
Me	OCF ₂ H	Br	<i>i</i> -Pr	H	Cl	OCF ₂ H	Br	<i>i</i> -Pr	H
Me	OCF ₂ H	Br	<i>t</i> -Bu	H	Cl	OCF ₂ H	Br	<i>t</i> -Bu	H
Me	OCF ₂ H	Br	CH ₂ CN	H	Cl	OCF ₂ H	Br	CH ₂ CN	H
Me	OCF ₂ H	Br	CH(Me)CH ₂ SMe	H	Cl	OCF ₂ H	Br	CH(Me)CH ₂ SMe	H
Me	OCF ₂ H	Br	C(Me) ₂ CH ₂ SMe	H	Cl	OCF ₂ H	Br	C(Me) ₂ CH ₂ SMe	H
Me	OCF ₂ H	Br	Me	Me	Cl	OCF ₂ H	Br	Me	Me
Me	OCH ₂ CF ₃	F	H	H	Cl	OCH ₂ CF ₃	F	H	H
Me	OCH ₂ CF ₃	F	Me	H	Cl	OCH ₂ CF ₃	F	Me	H
Me	OCH ₂ CF ₃	F	Et	H	Cl	OCH ₂ CF ₃	F	Et	H
Me	OCH ₂ CF ₃	F	<i>i</i> -Pr	H	Cl	OCH ₂ CF ₃	F	<i>i</i> -Pr	H
Me	OCH ₂ CF ₃	F	<i>t</i> -Bu	H	Cl	OCH ₂ CF ₃	F	<i>t</i> -Bu	H
Me	OCH ₂ CF ₃	F	CH ₂ CN	H	Cl	OCH ₂ CF ₃	F	CH ₂ CN	H
Me	OCH ₂ CF ₃	F	CH(Me)CH ₂ SMe	H	Cl	OCH ₂ CF ₃	F	CH(Me)CH ₂ SMe	H
Me	OCH ₂ CF ₃	F	C(Me) ₂ CH ₂ SMe	H	Cl	OCH ₂ CF ₃	F	C(Me) ₂ CH ₂ SMe	H
Me	OCH ₂ CF ₃	F	Me	Me	Cl	OCH ₂ CF ₃	F	Me	Me
Me	OCH ₂ CF ₃	Cl	H	H	Cl	OCH ₂ CF ₃	Cl	H	H
Me	OCH ₂ CF ₃	Cl	Me	H	Cl	OCH ₂ CF ₃	Cl	Me	H
Me	OCH ₂ CF ₃	Cl	Et	H	Cl	OCH ₂ CF ₃	Cl	Et	H
Me	OCH ₂ CF ₃	Cl	<i>i</i> -Pr	H	Cl	OCH ₂ CF ₃	Cl	<i>i</i> -Pr	H
Me	OCH ₂ CF ₃	Cl	<i>t</i> -Bu	H	Cl	OCH ₂ CF ₃	Cl	<i>t</i> -Bu	H
Me	OCH ₂ CF ₃	Cl	CH ₂ CN	H	Cl	OCH ₂ CF ₃	Cl	CH ₂ CN	H
Me	OCH ₂ CF ₃	Cl	CH(Me)CH ₂ SMe	H	Cl	OCH ₂ CF ₃	Cl	CH(Me)CH ₂ SMe	H
Me	OCH ₂ CF ₃	Cl	C(Me) ₂ CH ₂ SMe	H	Cl	OCH ₂ CF ₃	Cl	C(Me) ₂ CH ₂ SMe	H
Me	OCH ₂ CF ₃	Cl	Me	Me	Cl	OCH ₂ CF ₃	Cl	Me	Me
Me	OCH ₂ CF ₃	Br	H	H	Cl	OCH ₂ CF ₃	Br	H	H
Me	OCH ₂ CF ₃	Br	Me	H	Cl	OCH ₂ CF ₃	Br	Me	H
Me	OCH ₂ CF ₃	Br	Et	H	Cl	OCH ₂ CF ₃	Br	Et	H
Me	OCH ₂ CF ₃	Br	<i>i</i> -Pr	H	Cl	OCH ₂ CF ₃	Br	<i>i</i> -Pr	H
Me	OCH ₂ CF ₃	Br	<i>t</i> -Bu	H	Cl	OCH ₂ CF ₃	Br	<i>t</i> -Bu	H
Me	OCH ₂ CF ₃	Br	CH ₂ CN	H	Cl	OCH ₂ CF ₃	Br	CH ₂ CN	H
Me	OCH ₂ CF ₃	Br	CH(Me)CH ₂ SMe	H	Cl	OCH ₂ CF ₃	Br	CH(Me)CH ₂ SMe	H
Me	OCH ₂ CF ₃	Br	C(Me) ₂ CH ₂ SMe	H	Cl	OCH ₂ CF ₃	Br	C(Me) ₂ CH ₂ SMe	H
Me	OCH ₂ CF ₃	Br	Me	Me	Cl	OCH ₂ CF ₃	Br	Me	Me
Me	OCF ₃	F	H	H	Cl	OCF ₃	F	H	H


<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁵</u>		<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁵</u>
Me	OCF ₃	F	Me	H		Cl	OCF ₃	F	Me	H
Me	OCF ₃	F	Et	H		Cl	OCF ₃	F	Et	H
Me	OCF ₃	F	<i>i</i> -Pr	H		Cl	OCF ₃	F	<i>i</i> -Pr	H
Me	OCF ₃	F	<i>t</i> -Bu	H		Cl	OCF ₃	F	<i>t</i> -Bu	H
Me	OCF ₃	F	CH ₂ CN	H		Cl	OCF ₃	F	CH ₂ CN	H
Me	OCF ₃	F	CH(Me)CH ₂ SMe	H		Cl	OCF ₃	F	CH(Me)CH ₂ SMe	H
Me	OCF ₃	F	C(Me) ₂ CH ₂ SMe	H		Cl	OCF ₃	F	C(Me) ₂ CH ₂ SMe	H
Me	OCF ₃	F	Me	Me		Cl	OCF ₃	F	Me	Me
Me	OCF ₃	Cl	H	H		Cl	OCF ₃	Cl	H	H
Me	OCF ₃	Cl	Me	H		Cl	OCF ₃	Cl	Me	H
Me	OCF ₃	Cl	Et	H		Cl	OCF ₃	Cl	Et	H
Me	OCF ₃	Cl	<i>i</i> -Pr	H		Cl	OCF ₃	Cl	<i>i</i> -Pr	H
Me	OCF ₃	Cl	<i>t</i> -Bu	H		Cl	OCF ₃	Cl	<i>t</i> -Bu	H
Me	OCF ₃	Cl	CH ₂ CN	H		Cl	OCF ₃	Cl	CH ₂ CN	H
Me	OCF ₃	Cl	CH(Me)CH ₂ SMe	H		Cl	OCF ₃	Cl	CH(Me)CH ₂ SMe	H
Me	OCF ₃	Cl	C(Me) ₂ CH ₂ SMe	H		Cl	OCF ₃	Cl	C(Me) ₂ CH ₂ SMe	H
Me	OCF ₃	Cl	Me	Me		Cl	OCF ₃	Cl	Me	Me
Me	OCF ₃	Br	H	H		Cl	OCF ₃	Br	H	H
Me	OCF ₃	Br	Me	H		Cl	OCF ₃	Br	Me	H
Me	OCF ₃	Br	Et	H		Cl	OCF ₃	Br	Et	H
Me	OCF ₃	Br	<i>i</i> -Pr	H		Cl	OCF ₃	Br	<i>i</i> -Pr	H
Me	OCF ₃	Br	<i>t</i> -Bu	H		Cl	OCF ₃	Br	<i>t</i> -Bu	H
Me	OCF ₃	Br	CH ₂ CN	H		Cl	OCF ₃	Br	CH ₂ CN	H
Me	OCF ₃	Br	CH(Me)CH ₂ SMe	H		Cl	OCF ₃	Br	CH(Me)CH ₂ SMe	H
Me	OCF ₃	Br	C(Me) ₂ CH ₂ SMe	H		Cl	OCF ₃	Br	C(Me) ₂ CH ₂ SMe	H
Me	OCF ₃	Br	Me	Me		Cl	OCF ₃	Br	Me	Me

Table 2



<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁶</u>	<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁶</u>
Me	CF ₃	Cl	Me	F	Me	CF ₃	Cl	Me	Cl
Cl	CF ₃	Cl	Me	F	Cl	CF ₃	Cl	Me	Cl
Br	CF ₃	Cl	Me	F	Br	CF ₃	Cl	Me	Cl
Me	Cl	Cl	Me	F	Me	Cl	Cl	Me	Cl
Cl	Cl	Cl	Me	F	Cl	Cl	Cl	Me	Cl
Br	Cl	Cl	Me	F	Br	Cl	Cl	Me	Cl
Me	Br	Cl	Me	F	Me	Br	Cl	Me	Cl
Cl	Br	Cl	Me	F	Cl	Br	Cl	Me	Cl
Br	Br	Cl	Me	F	Br	Br	Cl	Me	Cl
Me	CF ₃	Cl	<i>i</i> -Pr	F	Me	CF ₃	Cl	<i>i</i> -Pr	Cl
Cl	CF ₃	Cl	<i>i</i> -Pr	F	Cl	CF ₃	Cl	<i>i</i> -Pr	Cl
Br	CF ₃	Cl	<i>i</i> -Pr	F	Br	CF ₃	Cl	<i>i</i> -Pr	Cl
Me	Cl	Cl	<i>i</i> -Pr	F	Me	Cl	Cl	<i>i</i> -Pr	Cl
Cl	Cl	Cl	<i>i</i> -Pr	F	Cl	Cl	Cl	<i>i</i> -Pr	Cl
Br	Cl	Cl	<i>i</i> -Pr	F	Br	Cl	Cl	<i>i</i> -Pr	Cl
Me	Br	Cl	<i>i</i> -Pr	F	Me	Br	Cl	<i>i</i> -Pr	Cl
Cl	Br	Cl	<i>i</i> -Pr	F	Cl	Br	Cl	<i>i</i> -Pr	Cl
Br	Br	Cl	<i>i</i> -Pr	F	Br	Br	Cl	<i>i</i> -Pr	Cl

Table 3



<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁷</u>	<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁷</u>
Me	CF ₃	F	Me	F	Me	CF ₃	Cl	Me	Cl
Cl	CF ₃	F	Me	F	Cl	CF ₃	Cl	Me	Cl
Br	CF ₃	F	Me	F	Br	CF ₃	Cl	Me	Cl
Me	Cl	F	Me	F	Me	Cl	Cl	Me	Cl
Cl	Cl	F	Me	F	Cl	Cl	Cl	Me	Cl
Br	Cl	F	Me	F	Br	Cl	Cl	Me	Cl
Me	Br	F	Me	F	Me	Br	Cl	Me	Cl

<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁷</u>		<u>R¹</u>	<u>R²</u>	<u>R³</u>	<u>R⁴</u>	<u>R⁷</u>
Cl	Br	F	Me	F		Cl	Br	Cl	Me	Cl
Br	Br	F	Me	F		Br	Br	Cl	Me	Cl
Me	CF ₃	F	<i>i</i> -Pr	F		Me	CF ₃	Cl	<i>i</i> -Pr	Cl
Cl	CF ₃	F	<i>i</i> -Pr	F		Cl	CF ₃	Cl	<i>i</i> -Pr	Cl
Br	CF ₃	F	<i>i</i> -Pr	F		Br	CF ₃	Cl	<i>i</i> -Pr	Cl
Me	Cl	F	<i>i</i> -Pr	F		Me	Cl	Cl	<i>i</i> -Pr	Cl
Cl	Cl	F	<i>i</i> -Pr	F		Cl	Cl	Cl	<i>i</i> -Pr	Cl
Br	Cl	F	<i>i</i> -Pr	F		Br	Cl	Cl	<i>i</i> -Pr	Cl
Me	Br	F	<i>i</i> -Pr	F		Me	Br	Cl	<i>i</i> -Pr	Cl
Cl	Br	F	<i>i</i> -Pr	F		Cl	Br	Cl	<i>i</i> -Pr	Cl
Br	Br	F	<i>i</i> -Pr	F		Br	Br	Cl	<i>i</i> -Pr	Cl

The invention described herein relates to a method of controlling or prevention of infestations of flies on an animal by applying to the animal a parasitically effective amount of a compound of Formula 1.

5

Application of Compounds of the Invention

The compound of Formula 1 of this invention can be applied to any animal, including herd animals, that can be bothered by flies or afflicted by myiasis. The compositions can be applied, for example, to cattle, sheep, goats, horses, donkeys, camels, pigs, reindeer, caribou and buffalo. Humans may also be treated.

10

The “applying” can be accomplished by way of non limiting example, whole-animal sprays, self-applicating devices, pour on treatments and controlled-release devices, such as ear tags and tapes, neck collars, ear tags, tail bands, limb bands or halters which comprise compounds or compositions comprising compounds of Formula 1. In addition to sprays and pour on treatments, application may be by other forms of topical administration, for

15

example, in the form of immersion or dipping, washing, coating with powder, or application to a small area of the animal.

Application of the compositions according to the invention to the animals to be treated is done topically via solutions, emulsions, suspensions, (drenches), powders, and pour-on formulations.

20

The pour-on or spot-on method consists in applying the compound of Formula 1 to a specific location of the skin or coat, advantageously to the neck or backbone of the animal. This takes place e.g. by applying a swab or spray of the pour-on or spot-on formulation to a relatively small area of the coat, from where the active substance is dispersed almost

automatically over wide areas of the fur owing to the spreading nature of the components in the formulation and assisted by the animal's movements.

The compounds of Formula 1 may be indirectly applied to an animal by applying it to the local environment in which the animal dwells (such as bedding, enclosures, or the like).

5 Whole-animal sprays provide rapid relief from fly pressure. Animal sprays are applied either as a dilute coarse spray, often applied under high pressure to soak the skin, or as a fine low-volume, more concentrated mist.

10 Self-applicating devices include back rubber covered with an absorbent material treated with an insecticide-oil solution, or dust bags filled with an insecticidal dust. Back rubbers and dustbags should be placed in gateways, near water and feed source, and in other areas where animals will make frequent contact with them.

Controlled-release ear tags and tapes are generally very effective for fly control in certain farm areas.

15 Pour-on treatments involves the application of an insecticide along the backline of the animal at a prescribed dosage of topical products. The pour-on or spot-on method is especially advantageous for use on herd animals such as cattle, horses, sheep or pigs, in which it is difficult or time-consuming to treat all the animals by more labor intensive methods of administration.

20 The compounds according to the invention are especially effective against fly larvae. The active compounds are employed in known manner, preferably by dermal or topical use, for example in the form of dipping, spraying, pour-on and spot-on, and powdering.

25 Blowfly strikes are almost always fatal unless the sheep is caught, the wool clipped from the infected area, the maggots scraped out, disinfectant or antibiotic and insecticide applied to prevent further strikes. Treatment of blowfly strike should aim to kill any maggots present, prevent the likelihood of further fly strike and assist the wound heal. The wool should be carefully clipped away from around the wound and surrounding area. A cream containing the compound of Formula 1 can be applied to the infected areas. Mild cases should heal quickly with correct treatment.

30 It is also effective to treat myiasis afflicted sheep or those at risk of being afflicted by dipping. It is particularly important to immerse sheep for at least a full minute so as to ensure the dip saturates the whole fleece and regular replenishment of dip baths is important to maintain the strength of dip concentrate. Pour-ons would also be an effective treatment

Compositions of the Invention

The compounds of the invention may be applied alone but are typically formulated into a veterinary or pharmaceutical composition. The compounds are prepared or formulated into compositions in a known manner, for example by extending the active compounds with solvents and/or carriers, if appropriate using emulsifiers and/or dispersing agents; if, for 5 example, water is used as the diluent, organic solvents can, if appropriate, be used as auxiliary solvents.

Typically a composition used in the present invention comprises a mixture of a compound of Formula 1, an *N*-oxide or a salt thereof, with one or more pharmaceutically or 10 veterinarily acceptable carriers comprising excipients and auxiliaries selected with regard to their suitability for topical administration and in accordance with standard practice. In addition, a suitable carrier is selected on the basis of compatibility with the one or more active ingredients in the composition, including such considerations as stability relative to 15 pH and moisture content. The typical application medium will be a composition for protecting an animal from an invertebrate parasitic pest comprising a parasitically effective amount of a compound of Formula 1 and at least one carrier.

Formulations for topical administration are typically in the form of a powder, cream, suspension, spray, emulsion, foam, paste, aerosol, ointment, salve or gel. More typically a topical formulation is a water-soluble solution, which can be in the form of a concentrate that is diluted before use. Parasiticidal compositions suitable for topical administration typically 20 comprise a compound of the present invention and one or more topically suitable carriers. In applications of a parasiticidal composition topically to the exterior of an animal as a line or spot (i.e. "spot-on" treatment), the active ingredient migrates over the surface of the animal to cover most or all of its external surface area. As a result, the treated animal is particularly 25 protected from invertebrate pests that feed off the epidermis of the animal such as ticks, fleas and lice. Therefore formulations for topical localized administration often comprise at least one organic solvent to facilitate transport of the active ingredient over the skin and/or penetration into the epidermis of the animal. Carriers in such formulations include propylene glycol, paraffins, aromatics, esters such as isopropyl myristate, glycol ethers, alcohols such as ethanol, *n*-propanol, 2-octyl dodecanol or oleyl alcohol; solutions in esters 30 of monocarboxylic acids, such as isopropyl myristate, isopropyl palmitate, lauric acid oxalic ester, oleic acid oleyl ester, oleic acid decyl ester, hexyl laurate, oleyl oleate, decyl oleate, caproic acid esters of saturated fatty alcohols of chain length C₁₂-C₁₈; solutions of esters of dicarboxylic acids, such as dibutyl phthalate, diisopropyl isophthalate, adipic acid

diisopropyl ester, di-*n*-butyl adipate or solutions of esters of aliphatic acids, e.g., glycols. It may be advantageous for a crystallization inhibitor or a dispersant known from the pharmaceutical or cosmetic industry also to be present.

A pour-on formulation may also be prepared for control of parasites in an animal of 5 agricultural worth. The pour-on formulations of this invention can be in the form of a liquid, powder, emulsion, foam, paste, aerosol, ointment, salve or gel. Typically, the pour-on formulation is liquid. These pour-on formulations can be effectively applied to sheep, cattle, goats, other ruminants, camelids, pigs and horses. The pour-on formulation is typically applied by pouring in one or several lines or in a spot-on the dorsal midline (back) or 10 shoulder of an animal. More typically, the formulation is applied by pouring it along the back of the animal, following the spine. The formulation can also be applied to the animal by other conventional methods, including wiping an impregnated material over at least a small area of the animal, or applying it using a commercially available applicator, by means of a 15 syringe, by spraying or by using a spray race. The pour-on formulations include a carrier and can also include one or more additional ingredients. Examples of suitable additional ingredients are stabilizers such as antioxidants, spreading agents, preservatives, adhesion promoters, active solubilisers such as oleic acid, viscosity modifiers, UV blockers or absorbers, and colourants. Surface active agents, including anionic, cationic, non-ionic and 20 ampholytic surface active agents, can also be included in these formulations.

The formulations of this invention often include an antioxidant, such as BHT (butylated hydroxytoluene). The antioxidant is generally present in amounts of at 0.1-5% (wt/vol). Some of the formulations require a solubilizer, such as oleic acid, to dissolve the active agent. Common spreading agents used in these pour-on formulations are: isopropyl myristate (IPM), isopropyl palmitate (IPP), caprylic/capric acid esters of saturated C₁₂-C₁₈ 25 fatty alcohols, oleic acid, oleyl ester, ethyl oleate, triglycerides, silicone oils and dipropylene glycol monomethyl ether (DPM). The pour-on formulations of this invention are prepared according to known techniques. Where the pour-on is a solution, the parasiticide/insecticide is mixed with the carrier or vehicle, using heat and stirring where required. Auxiliary or additional ingredients can be added to the mixture of active agent and carrier, or they can be 30 mixed with the active agent prior to the addition of the carrier. If the pour-on is an emulsion or suspension, these formulations are similarly prepared using known techniques.

Other delivery systems for relatively hydrophobic pharmaceutical compounds may be employed. Liposomes and emulsions are well-known examples of delivery vehicles or

carriers for hydrophobic drugs. In addition, organic solvents such as dimethylsulfoxide may be used.

The compounds of Formula 1 are generally present in the compositions in concentrations of 0.1 to 95 percent by weight, preferably 0.5 to 90 percent by weight.

5 Preparations which are intended for direct application contain the active compound according to the invention in concentrations of between 0.001 and 5 percent by weight, preferably 0.005 to 3 percent by weight.

Dosages may range from 0.0001 mg/kg of animal body weight to about 1000 mg/kg. of the compound of Formula 1. Sometimes dosages may be from 0.1 mg/kg of animal body 10 weight to about 200 mg/kg. Often times it would be advantageous to administer amounts of about 0.01 to about 100 mg or between 0.02 to about 50mg/kg. and frequently between 0.1 and 75 mg/kg. Preferably, the treatment is carried out so as to administer to the animal a dose of from 0.1 to 40 mg/kg and in particular from 1 to 30 mg/kg. Administration may be given as a single dose or intermittent in time and may be administered daily, weekly, 15 monthly, bimonthly or quarterly in order to achieve effective results in order to achieve effective results.

Nevertheless it can at times be necessary to deviate from the amounts mentioned, and in particular to do so in accordance with the body weight of the test animal and/or the method of application, but also because of the species of animal and its individual behavior 20 towards the medicament, or the nature of the formulation of the latter and the time or interval at which it is administered. Thus it can suffice in some cases to manage with less than the above mentioned minimum amount while in other cases the upper limit mentioned must be exceeded. Where substantial amounts are applied, it can be advisable to divide these into several individual administrations over the course of the day. The general sense of the other 25 statements made above also applies.

Pour-on or spot-on formulations suitably contain carriers, which promote rapid dispersement over the skin surface or in the coat of the host animal, and are generally regarded as spreading oils. Suitable carriers are e.g. oily solutions; alcoholic and isopropanolic solutions such as solutions of 2-octyldodecanol or oleyl alcohol; solutions in 30 esters of monocarboxylic acids, such as isopropyl myristate, isopropyl palmitate, lauric acid oxalate, oleic acid oleyl ester, oleic acid decyl ester, hexyl laurate, oleyl oleate, decyl oleate, capric acid esters of saturated fat alcohols of chain length C_{sub}.12-C_{sub}.18; solutions of esters of dicarboxylic acids, such as dibutyl phthalate, diisopropyl isophthalate, adipic acid

diisopropyl ester, di-n-butyl adipate or also solutions of esters of aliphatic acids, e.g. glycols. It may be advantageous for a dispersing agent to be additionally present, such as one known from the pharmaceutical or cosmetic industry. Examples are 2-pyrrolidone, 2-(N-alkyl)pyrrolidone, acetone, polyethylene glycol and the ethers and esters thereof, propylene 5 glycol or synthetic triglycerides.

The oily solutions include e.g. vegetable oils such as olive oil, groundnut oil, sesame oil, pine oil, linseed oil or castor oil. The vegetable oils may also be present in epoxidised form. Paraffins and silicone oils may also be used.

A pour-on or spot-on formulation generally contains 1 to 20% by weight of a 10 compound of Formula 1, 0.1 to 50% by weight of dispersing agent and 45 to 98.9% by weight of solvent.

Importantly the compounds of Formula 1 may be indirectly applied to an animal by applying it to the local environment in which the animal dwells (such as bedding, enclosures, or the like). Effective use rates will range from about 1.0 to 50 mg/square meter but as little 15 as 0.1 mg/square meter may be sufficient or as much as 150 mg/square meter may be required. One skilled in the art can easily determine the biologically effective amount necessary for the desired level of pest control.

The Compositions of the Invention may comprise Additional Active Compounds:

It is contemplated that additional biologically active compounds may be be 20 administered at the same time or separately over time to obtain broader spectrum of pest control or to attack adult fleas. Such additional biologically active compounds may be packaged together with the compound of Formula 1 as a kit. For convenience sake such additional biologically active compounds may be formulated into the same composition containing the compound of Formula 1. Therefore the present invention contemplates the 25 use of compositions characterised in that they contain, in addition to a compound of Formula 1, further auxiliaries and/or active compounds, such as additional biologically active compounds, disinfectants or antibiotics may be admixed to the formulations, or the ready-to-use solutions, in addition to the customary solid or liquid extenders, diluents and/or surface-active agents.

30 Of note are additional biologically active compounds or agents selected from art-known anthelmintics, such as, for example, avermectins (e.g. ivermectin, moxidectin, milbemycin), benzimidazoles (e.g. albendazole, triclabendazole), salicylanilides (e.g.

closantel, oxyclozanide), substituted phenols (e.g. nitroxynil), pyrimidines (e.g. pyrantel), imidazothiazoles (e.g. levamisole) and praziquantel.

Other biologically active compounds or agents useful in the compositions of the present invention can be selected from Insect Growth Regulators (IGRs) and Juvenile 5 Hormone Analogues (JHAs) such as diflubenzuron, trilumuron, fluazuron, cyromazine, methoprene, etc., thereby providing both initial and sustained control of parasites (at all stages of insect development, including eggs) on the animal subject, as well as within the environment of the animal subject.

The compounds of Formula I according to the invention may be used alone or in 10 combination with other biocides. They may be combined with pesticides having the same sphere of activity e.g. to increase activity, or with substances having another sphere of activity e.g. to broaden the range of activity. It can also be sensible to add so-called repellents. If the range of activity is to be extended to endoparasites, e.g. wormers, the compounds of Formula 1 are suitably combined with substances having endoparasitic 15 properties. Of course, they can also be used in combination with antibacterial compositions.

Preferred groups of combination partners and especially preferred combination partners are named in the following, whereby combinations may contain one or more of these partners in addition to a compound of Formula 1.

Suitable partners in the mixture may be biocides, e.g. the insecticides and acaricides 20 with a varying mechanism of activity, which are named in the following and have been known to the person skilled in the art for a long time, e.g. chitin synthesis inhibitors, growth regulators; active ingredients which act as juvenile hormones; active ingredients which act as adulticides; broad-band insecticides, broad-band acaricides and nematicides; and also the well known anthelmintics and insect- and/or acarid-deterring substances, and also 25 repellents or detachers.

Examples of such biologically active compounds include but are not restricted to the following: Organophosphates, a class which are generally known to be inhibitors of acetyl cholinesterase: acephate, azamethiphos, azinphos-ethyl, azinphos-methyi, bromophos, bromophos-ethyl, cadusafos, chlorethoxyphos, chlorpyrifos, chlorgenvinphos, chlormephos, 30 demeton, demeton-S-methyl, demeton-S-methyl sulphone, dialifos, diazinon, dichlorvos, dicrotophos, dimethoate, disulfoton, ethion, ethoprophos, etrimfos, famphur, fenamiphos, fenitrothion, fensulfothion, fenthion, flupyrazofos, fonofos, formothion, fosthiazate, heptenophos, isazophos, isothioate, isoxathion, malathion, methacriphos, methamidophos,

methidathion, methyl-parathion, mevinphos, monocrotophos, naled, omethoate, oxydemeton-methyl, paraoxon, parathion, parathion-methyl, phentoate, phosalone, phosfolan, phosphocarb, phosmet, phosphamidon, phorate, phoxim, pirimiphos, pirimiphos-methyl, profenofos, propaphos, proetamphos, prothiofos, pyraclofos, pyridapenthion, 5 quinalphos, sulprophos, temephos, terbufos, tebupirimfos, tetrachlorvinphos, thimeton, triazophos, trichlorfon, vamidothion.

Carbamates, a class which are generally known to be inhibitors of acetyl cholinesterase: alanycarb, aldicarb, 2-sec-butylphenyl methylcarbamate, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenoxy carb, fenthiocarb, furathiocarb, HCN-801,

10 isoprocarb, indoxacarb, methiocarb, methomyl, 5 methyl-m-cumenylbutyryl(methyl) carbamate, oxamyi, pirimicarb, propoxur, thiodicarb, thifanox, triazamate, UC-51717

Pyrethroids, a class which are generally known to be modulators of sodium channels: acrinathrin, allethrin, alphametrin, 5-benzyl-3-furylmethyl (E)-(1 R)-cis-2,2-dimethyl-3-(2-oxothiolan-3-ylidene)methyl cyclopropanecarboxylate, bifenthrin, ,8 cyfluthrin, cyfluthrin,

15 ,oc-cypermethrin, ,8-cypermethrin, bioallethrin, bioallethrin((S)-I cyclopentylisomer), bioresmethrin, bifenthrin, NCI-85193, cycloprothrin, cyhalothrin, cythithrin, cyphenothrin, deltamethrin, empenthrin, esfenvalerate, ethofenprox, fenfluthrin, fenpropathrin, fenvalerate, flucythrinate, flumethrin, fluvalinate (D isomer), imiprothrin, cyhalothrin, λ -cyhalothrin, permethrin, phenothrin, prallethrin, pyrethrins (natural products), resmethrin, tetramethrin, 20 transfluthrin, theta-cypermethrin, silafluofen, T-fluvalinate, tefluthrin, tralomethrin, Zeta-cypermethrin.

Arthropod growth regulators including: a) chitin synthesis inhibitors: benzoylureas: chlorfluazuron, diflubenzuron, fluazuron, flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron, teflubenzuron, triflumuron, buprofezin, diofenolan, hexythiazox,

25 etoxazole, chlorfentazine; b) ecdysone agonists/disruptors: halofenozide, methoxyfenozide, tebufenozide; c) juvenoid hormone mimes: pyriproxyfen, methoprene, fenoxy carb; d) lipid biosynthesis inhibitors: spirodiclofen. Other antiparasitics: acequinocyl, amitraz, AKD-1022, ANS-118, azadirachtin, Bacillus thuringiensis, bensultap, bifenazate, binapacryl, bromopropylate, BTG-504, I BTG-505, camphechlor, cartap, chlorobenzilate,

30 chlordimeform, chlorfenapyr, chromafenozide, clothianidine, cyromazine, diacloden, diafenthiuron, DBI-3204, dinactin, dihydroxymethyidihydroxypyrrolidine, dinobuton, dinocap, endosulfan, ethiprole, ethofenprox, fenazaquin, flumite, MTI-800, fenpyroximate,

fluacrypyrim, flubenzimine, flubrocythrinate, flufenzine, flufenprox, fluproxyfen, halofenprox, hydramethyinon, IKI-220, kanemite, NC-196, neem guard, nidinorterfuran, nitenpyram, SD-35651, WL-108477, pirydaryl, propargite, protrifenbute, pymethrozine, pyridaben, pyrimidifen, NC-1111, R-195, RH-0345, RH-2485, RYI-210, S-1283, S-1833,

5 SI-8601, silafluofen, silomadine, spinosad, tebufenpyrad, tetradifon, tetranactin, thiacloprid, thiocyclam, thiamethoxam, tolfenpyrad, triazamate, triethoxyspinosyn, trinactin, verbutin, vertalec, YI-5301 Fungicides: acibenzolar, aldimorph, ampropylfos, andoprim, azaconazole, azoxystrobin, benalaxyl, benomyl, bialaphos, blasticidin-S, Bordeaux mixture, bromuconazole, bupirimate, carpropamid, captafol, captan, carbendazim, chlорfenazole, 10 chloroneb, chloropicrin, chlorothalonil, chlozolinate, copper oxychloride, copper salts, cyflufenamid, cymoxanil, cyproconazole, cyprodinil, cyprofuram, RH-7281, diclocymet, diclobutrazole, diclomezine, dicloran, difenoconazole, RP-407213, dimethomorph, domoxystrobin, diniconazole, diniconazole-M, dodine, edifenphos, epoxiconazole, famoxadone, fenamidone, fenarimol, fenbuconazole, fencaramid, fenpiclonil, fenpropidin, 15 fenpropimorph, fentin acetate, fluazinam, fludioxonil, flumetover, flumorf/flumorlin, fentin hydroxide, fluoxastrobin, fluquinconazole, flusilazole, flutolanil, flutriafol, folpet, fosetyl-aluminium, furalaxyl, furametapyr, hexaconazole, ipconazole, iprobenfos, iprodione, isoprothiolane, kasugamycin, krsoxim-methyl, mancozeb, maneb, mefenoxam, mepronil, metalaxyl, metconazole, metominostrobin/fenominostrobin, metrafenone, myclobutanil, neo- 20 asozin, nicobifen, orysastrobin, oxadixyl, penconazole, pencycuron, probenazole, prochloraz, propamocarb, propiconazole, proquinazid, prothioconazole, pyrifenox, pyraclostrobin, pyrimethanil, pyroquilon, quinoxyfen, spiroxamine, sulfur, tebuconazole, tetriconazole, thiabendazole, thifluzamide, thiophanate-methyl, thiram, tiadinil, triadimefon, triadimenol, tricyclazole, trifioxystrobin, triticonazole, validamycin, vinclozin Biological 25 agents: *Bacillus thuringiensis* ssp alzawai, kurstaki, *Bacillus thuringiensis* delta endotoxin, baculovirus, entomopathogenic bacteria, virus and fungi Bactericides: chlortetracycline, oxytetracycline, streptomycin,

Additional more specific examples of partner insecticides and acaricides are listed below:

30	Compound	Class
	Compound	Class
	Abamectin	macrocyclic lactones
	AC 303 630	energy production modulator
	Acephate	acetyl cholinesterase inhibitor

Acrinathrin	sodium channel modulator
Alanycarb	acetyl cholinesterase inhibitor
Aldicarb	acetyl cholinesterase inhibitor
alpha.-Cypermethrin	sodium channel modulator
Alphamethrin	sodium channel modulator
Amitraz	octopamine receptor ligand
Avermectin	macrocyclic lactones
Azinphos A	acetyl cholinesterase inhibitor
Azinphos M	acetyl cholinesterase inhibitor
Azinphos-methyl	acetyl cholinesterase inhibitor
Azocyclotin	oxidative phosphorylation inhibitor
Bacillus subtil. toxin	
Bendiocarb	acetyl cholinesterase inhibitor
Benfuracarb	acetyl cholinesterase inhibitor
Bensultap	nicotinic acetylcholine agonist/antagonist
beta.-Cyfluthrin	sodium channel modulator
Bifenthrin	sodium channel modulator
Brofenprox	sodium channel modulator
Bromophos A	acetyl cholinesterase inhibitor
Bufencarb	acetyl cholinesterase inhibitor
Buprofezin	chitin synthesis inhibitor
Butocarboxin	acetyl cholinesterase inhibitor
Cadusafos	acetyl cholinesterase inhibitor
Carbaryl	acetyl cholinesterase inhibitor
Carbofuran	acetyl cholinesterase inhibitor
Carbophenthion	acetyl cholinesterase inhibitor
Cartap	nicotinic acetylcholine agonist/antagonist
Chloethocarb	acetyl cholinesterase inhibitor
Chlorethoxyfos	acetyl cholinesterase inhibitor
Chlorfenapyr	oxidative phosphorylation inhibitor
Chlorfluazuron	chitin synthesis inhibitor
Chlormephos	acetyl cholinesterase inhibitor
Chlorpyrifos	acetyl cholinesterase inhibitor
Cis-Resmethrin	sodium channel modulator
Clofentezine	
Cyanophos	acetyl cholinesterase inhibitor
Cycloprothrin	sodium channel modulator
Cyfluthrin	sodium channel modulator
Cyhexatin	oxidative phosphorylation inhibitor
D 2341 (bifenazate)	
Deltamethrin	sodium channel modulator
Demeton M	acetyl cholinesterase inhibitor
Demeton S	acetyl cholinesterase inhibitor
Demeton-S-methyl	acetyl cholinesterase inhibitor
Dichlofenthion	acetyl cholinesterase inhibitor
Diaphorin	acetyl cholinesterase inhibitor
Diethion	acetyl cholinesterase inhibitor
Diflubenzuron	chitin synthesis inhibitor
Dimethoate	acetyl cholinesterase inhibitor
Dimethylvinphos	acetyl cholinesterase inhibitor

Dioxathion	acetyl cholinesterase inhibitor
Doramectin	macrocyclic lactones
DPX-MP062 (indoxacarb)	sodium channel modulator
Edifenphos	acetyl cholinesterase inhibitor
Emamectin	macrocyclic lactones
Endosulfan	gaba-gated chloride channel antagonist
Eprinomectin	macrocyclic lactones
Esfenvalerate	sodium channel modulator
Ethiofencarb	acetyl cholinesterase inhibitor
Ethion	acetyl cholinesterase inhibitor
Ethofenprox	sodium channel modulator
Ethoprophos	acetyl cholinesterase inhibitor
Etrimphos	acetyl cholinesterase inhibitor
Fenamiphos	acetyl cholinesterase inhibitor
Fenazaquin	mitochondrial electron transport inhibitor
Fenbutatin oxide	oxidative phosphorylation inhibitor
Fenitrothion	acetyl cholinesterase inhibitor
Fenobucarb (BPMC)	acetyl cholinesterase inhibitor
Fenothiocarb	acetyl cholinesterase inhibitor
Fenoxy carb	juvenile hormone mimic
Fenpropathrin	sodium channel modulator
Fenpyrad	mitochondrial electron transport inhibitor
Fenpyroximate	mitochondrial electron transport inhibitor
Fenthion	acetyl cholinesterase inhibitor
Fenvalerate	sodium channel modulator
Fipronil	gaba-gated chloride channel antagonist
Fluazinam	oxidative phosphorylation uncoupler
Fluazuron	chitin synthesis inhibitor
Flucycloxuron	chitin synthesis inhibitor
Flucythrinate	sodium channel modulator
Flufenoxuron	chitin synthesis inhibitor
Flufenprox	sodium channel modulator
Fonophos	acetyl cholinesterase inhibitor
Formothion	acetyl cholinesterase inhibitor
Fosthiazate	acetyl cholinesterase inhibitor
HCH	gaba-gated chloride channel antagonist
Heptenophos	acetyl cholinesterase inhibitor
Hexaflumuron	chitin synthesis inhibitor
Hexythiazox	
Hydroprene	juvenile hormone mimic
Imidacloprid	nicotinic acetylcholine agonist/antagonist
insect-active fungi	
insect-active nematodes	
insect-active viruses	
Iprobenfos	acetyl cholinesterase inhibitor
Isofenphos	acetyl cholinesterase inhibitor
Isoprocarb	acetyl cholinesterase inhibitor
Isoxathion	acetyl cholinesterase inhibitor
Ivermectin	chloride channel activator
lambda.-Cyhalothrin	sodium channel modulator

Lufenuron	chitin synthesis inhibitor
Malathion	acetyl cholinesterase inhibitor
Mecarbam	acetyl cholinesterase inhibitor
Mesulfenphos	acetyl cholinesterase inhibitor
Metaldehyd	
Methamidophos	acetyl cholinesterase inhibitor
Methiocarb	acetyl cholinesterase inhibitor
Methomyl	acetyl cholinesterase inhibitor
Methoprene	juvenile hormone mimic
Metolcarb	acetyl cholinesterase inhibitor
Mevinphos	acetyl cholinesterase inhibitor
Milbemectin	macrocyclic lactones
Moxidectin	macrocyclic lactones
Naled	acetyl cholinesterase inhibitor
NI-25, Acetamiprid	nicotinic acetylcholine agonist/antagonist
Nitenpyram	nicotinic acetylcholine agonist/antagonist
Nodulisporic acid/derivatives	macrocyclic lactones
Omethoat	acetyl cholinesterase inhibitor
Oxamyl	acetyl cholinesterase inhibitor
Oxydemethon M	acetyl cholinesterase inhibitor
Oxydeprofos	acetyl cholinesterase inhibitor
Parathion	acetyl cholinesterase inhibitor
Parathion-methyl	acetyl cholinesterase inhibitor
Permethrin	sodium channel modulator
Phenthroate	acetyl cholinesterase inhibitor
Phorat	acetyl cholinesterase inhibitor
Phosalone	acetyl cholinesterase inhibitor
Phosmet	acetyl cholinesterase inhibitor
Phoxim	acetyl cholinesterase inhibitor
Pirimicarb	acetyl cholinesterase inhibitor
Pirimiphos A	acetyl cholinesterase inhibitor
Pirimiphos M	acetyl cholinesterase inhibitor
Promecarb	acetyl cholinesterase inhibitor
Propaphos	acetyl cholinesterase inhibitor
Propoxur	acetyl cholinesterase inhibitor
Prothiofos	acetyl cholinesterase inhibitor
Prothoat	acetyl cholinesterase inhibitor
Pyrachlophos	acetyl cholinesterase inhibitor
Pyradaphenthion	acetyl cholinesterase inhibitor
Pyresmethrin	sodium channel modulator
Pyrethrim	sodium channel modulator
Pyridaben	mitochondrial electron transport inhibitor
Pyrimidifen	mitochondrial electron transport inhibitor
Pyriproxyfen	juvenile hormone mimic
RH 5992	ecdysone agonist
RH-2485	ecdysone agonist
Salithion	acetyl cholinesterase inhibitor
selamectin	macrocyclic lactones
Silafluofen	sodium channel modulator
Spinosad	nicotinic acetylcholine activator

Sulfotep	acetyl cholinesterase inhibitor
Sulprofos	acetyl cholinesterase inhibitor
Tebufenozide	ecdysone agonist
Tebufenpyrad	mitochondrial electron transport inhibitor
Tebupirimphos	acetyl cholinesterase inhibitor
Teflubenzuron	chitin synthesis inhibitor
Tefluthrin	sodium channel modulator
Temephos	acetyl cholinesterase inhibitor
Terbufos	acetyl cholinesterase inhibitor
Tetrachlorvinphos	acetyl cholinesterase inhibitor
Thiafenox	
Thiodicarb	acetyl cholinesterase inhibitor
Thifanox	acetyl cholinesterase inhibitor
Thionazin	acetyl cholinesterase inhibitor
Thuringiensin	
Tralomethrin	sodium channel modulator
Triarathen	
Triazamate	acetyl cholinesterase inhibitor
Triazophos	acetyl cholinesterase inhibitor
Trichlorfon	acetyl cholinesterase inhibitor
Triflumuron	chitin synthesis inhibitor
Trimethacarb	acetyl cholinesterase inhibitor
Vamidothion	acetyl cholinesterase inhibitor
XMC (3,5,-Xylyl-methylcarbamate)	acetyl cholinesterase inhibitor
Xylylcarb	acetyl cholinesterase inhibitor
YI 5301/5302	
zeta.-Cypermethrin	sodium channel modulator
Zetamethrin	sodium channel modulator

Non-limitative examples of suitable anthelmintics are named in the following, a few representatives have insecticidal and acaricidal activity in addition to the anthelmintic activity, and are partly already in the above list.

5 (A1) Praziquantel=2-cyclohexylcarbonyl-4-oxo-1,2,3,6,7,11b-hexahydr- o-4H-pyrazino[2,1-.alpha.]isoquinoline

(A2) Closantel=3,5-diido-N-[5-chloro-2-methyl-4-(a-cyano-4-chlorob- enzyl)phenyl]-salicylamide

(A3) Triclabendazole=5-chloro-6-(2,3-dichlorophenoxy)-2-methylthio-- 1H-benzimidazole

10 (A4) Levamisol=L-(--)-2,3,5,6-tetrahydro-6-phenylimidazo[2,1b]thiazo- le

(A5) Mebendazole=(5-benzoyl-1H-benzimidazol-2-yl)carbaminic acid methylester

(A6) Omphalotin=a macrocyclic fermentation product of the fungus Omphalotus olearius described In WO 97/20857

(A7) Abamectin=avermectin B1

15 (A8) Ivermectin=22,23-dihydroavermectin B1

(A9) Moxidectin=5-O-demethyl-28-deoxy-25-(1,3-dimethyl-1-butenyl)-6- ,28-epoxy-23-

(methoxyimino)-milbemycin B

(A10) Doramectin=25-cyclohexyl-5-O-demethyl-25-de(1-methylpropyl)-a- vermectin A1a

(A11) Milbemectin=mixture of milbemycin A3 and milbemycin A4

(A12) Milbemycinoxim=5-oxime of milbemectin

5 Non-limitative examples of suitable repellents and detachers are:

(R1) DEET (N,N-diethyl-m-toluamide)

(R2) KBR 3023 N-butyl-2-oxycarbonyl-(2-hydroxy)-piperidine

(R3) Cymiazole=N,-2,3-dihydro-3-methyl-1,3-thiazol-2-ylidene-2,4-xy- lidene

The aforementioned partners in the mixture are best known to specialists in this field.

10 Most are described in various editions of the Pesticide Manual, The British Crop Protection Council, London, and others in the various editions of The Merck Index, Merck & Co., Inc., Rahway, N.J., USA or in patent literature. Therefore, the following listing is restricted to a few places where they may be found by way of example.

(I) 2-Methyl-2-(methylthio)propionaldehyde-O-methylcarbamoyloxime (Aldicarb), from

15 The Pesticide Manual, 11.sup.th Ed. (1997), The British Crop Protection Council, London, page 26;

(II) S-(3,4-dihydro-4-oxobenzo[d]-[1,2,3]-triazin-3-ylmethyl)O,O-di- methyl-phosphorodithioate (Azinphos-methyl), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 67;

20 (III) Ethyl-N-[2,3-dihydro-2,2-dimethylbenzofuran-7-yl]oxycarbonyl-(methyl)aminothio]-N-isopropyl-.beta.-alaninate (Benfuracarb), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 96;

(IV) 2-Methylbiphenyl-3-ylmethyl-(Z)-(1RS)-cis-3-(2-chloro-3,3,3-tr- ifluoroprop-1-enyl)-2,2-dimethylcyclopropanecarboxylate (Bifenthrin), from The Pesticide Manual, 11.sup.thEd. (1997),

25 The British Crop Protection Council, London, page 118;

(V) 2-tert-butylimino-3-isopropyl-5-phenyl-1,3,5-thiadiazian-4-one (Buprofezin), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 157;

(VI) 2,3-Dihydro-2,2-dimethylbenzofuran-7-yl-methylcarbamate (Carbofuran), from The

30 Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 186;

(VII) 2,3-Dihydro-2,2-dimethylbenzofuran-7-yl-(dibutylaminothio)met- hylcarbamate (Carbosulfan), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 188;

35 (VIII) S,S'-(2-dimethylaminotrimethylene)-bis(thiocarbamate) (Cartap), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 193;

(IX) 1-[3,5-Dichloro-4-(3-chloro-5-trifluoromethyl-2-pyridyloxy)phe- nyl]-3-(2,6-difluorobenzoyl)-urea (Chlorfluazuron), from The Pesticide Manual, 11.sup.thEd. (1997),

The British Crop Protection Council, London, page 213;

(X) O,O-diethyl-O-3,5,6-trichloro-2-pyridyl-phosphorothioate (Chlorpyrifos), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 235;

5 (XI) (RS)-.alpha.-cyano-4-fluoro-3-phenoxybenzyl-(1RS,3RS;1RS,3RS)-- 3-(2,2-dichlorovinyl)-2,2-di-methylcyclopropanecarboxylate (Cyfluthrin), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 293;

(XII) Mixture of (S)-.alpha.-cyano-3-phenoxybenzyl-(Z)-(1R,3R)-3-(2- -chloro-3,3,3-trifluoropropenyl)-2,2-dimethylcyclopropanecarboxylate and (R)-.alpha.-cyano-3-phenoxybenzyl-(Z)-(1R,3)-3-(2-chloro-3,3,3-trifluoropropenyl)-2,2-

10 dimethylcyclopropanecarboxylate (Lambda-Cyhalothrin), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 300;

(XIII) Racemate consisting of (S)-.alpha.-cyano-3-phenoxybenzyl-(2)- -(1R,3R)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate and (R)-.alpha.-cyano-3-phenoxybenzyl-(1S,3S)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate

15 (Alpha-cypermethrin), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 308;

(XIV) a mixture of the stereoisomers of (S)-.alpha.-cyano-3-phenoxy- benzyl (1RS,3RS,1RS,3RS)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (zeta-Cypermethrin),

20 from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 314;

(XV) (S)-.alpha.-cyano-3-phenoxybenzyl-(1R,3R)-3-(2,2-dibromovinyl)- -2,2-dimethylcyclopropanecarboxylate (Deltamethrin), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 344;

25 (XVI) (4-chlorophenyl)-3-(2,6-difluorobenzoyl)urea (Diflubenzuron), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 395;

(XVII) (1,4,5,6,7,7-Hexachloro-8,9,10-trinorborn-5-en-2,3-ylenebism- ethylene)-sulphite (Endosulfan), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 459;

30 (XVIII) .alpha.-ethylthio-o-tolyl-methylcarbamate (Ethiofencarb), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 479;

(XIX) O,O-dimethyl-O-4-nitro-m-tolyl-phosphorothioate (Fenitrothion), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 514;

(XX) 2-sec-butylphenyl-methylcarbamate (Fenobucarb), from The Pesticide Manual,

35 11.sup.thEd. (1997), The British Crop Protection Council, London, page 516;

(XXI) (RS)-.alpha.-cyano-3-phenoxybenzyl-(RS)-2-(4-chlorophenyl)-3-- methylbutyrate (Fenvalerate), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 539;

(XXII) S-[formyl(methyl)carbamoylmethyl]-O,O-dimethyl-phosphorodithioate (Formothion), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 625;

5 (XXIII) 4-Methylthio-3,5-xylyl-methylcarbamate (Methiocarb), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 813;

(XXIV) 7-Chlorobicyclo[3.2.0]hepta-2,6-dien-6-yl-dimethylphosphate (Heptenophos), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 670;

10 (XXV) 1-(6-chloro-3-pyridylmethyl)-N-nitroimidazolidin-2-ylidenamin-e (Imidacloprid), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 706;

(XXVI) 2-isopropylphenyl-methylcarbamate (Isoprocarb), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 729;

15 (XXVII) O,S-dimethyl-phosphoramidothioate (Methamidophos), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 808;

(XXVIII) S-Methyl-N-(methylcarbamoyloxy)thioacetimidate (Methomyl), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 815;

20 (XXIX) Methyl-3-(dimethoxyphosphinoyloxy)but-2-enoate (Mevinphos), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 844;

(XXX) O,O-diethyl-O-4-nitrophenyl-phosphorothioate (Parathion), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 926;

25 (XXXI) O,O-dimethyl-O-4-nitrophenyl-phosphorothioate (Parathion-methyl), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 928;

(XXXII) S-6-chloro-2,3-dihydro-2-oxo-1,3-benzoxazol-3-ylmethyl-O,O-- diethyl-phosphordithioate (Phosalone), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 963;

30 (XXXIII) 2-Dimethylamino-5,6-dimethylpyrimidin-4-yl-dimethylcarbamate (Pirimicarb), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 985;

(XXXIV) 2-isopropoxyphenyl-methylcarbamate (Propoxur), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 1036;

35 (XXXV) 1-(3,5-dichloro-2,4-difluorophenyl)-3-(2,6-difluorobenzoyl)urea (Teflubenzuron), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 1158;

(XXXVI) S-tert-butylthiomethyl-O,O-dimethyl-phosphorodithioate (Terbufos), from The

Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 1165;

(XXXVII) ethyl-(3-tert.-butyl-1-dimethylcarbamoyl-1H-1,2,4-triazol-- 5-yl-thio)-acetate,

5 (Triazamate), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 1224;

(XXXVIII) Abamectin, from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 3;

(XXXIX) 2-sec-butylphenyl-methylcarbamate (Fenobucarb), from The Pesticide Manual,

10 11.sup.thEd. (1997), The British Crop Protection Council, London, page 516;

(XL) N-tert.-butyl-N'-(4-ethylbenzoyl)-3,5-dimethylbenzohydrazide (Tebufenozide), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 1147;

(XLI) (+)-5-amino-1-(2,6-dichloro-.alpha.,.alpha.,.alpha.-triflu- oro-p-tolyl)-4-

15 trifluoromethyl-sulphinylpyrazol-3-carbonitrile (Fipronil), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 545;

(XLII) (RS)-.alpha.-cyano-4-fluoro-3-phenoxybenzyl(1RS,3RS;1 RS,3RS)-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate (beta-Cyfluthrin), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 295;

20 (XLIII) (4-ethoxyphenyl)-[3-(4-fluoro-3-phenoxyphenyl)propyl](dimethyl)silane (Silafluofen), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 1105;

(XLIV) tert.-butyl (E)-.alpha.-(1,3-dimethyl-5-phenoxy)pyrazol-4-yl-- methylenamino-oxy)-p-toluate (Fenpyroximate), from The Pesticide Manual, 11.sup.thEd. (1997), The British

25 Crop Protection Council, London, page 530;

(XLV) 2-tert.-butyl-5-(4-tert.-butylbenzylthio)-4-chloropyridazin-3- (2H)-one (Pyridaben), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 1161;

(XLVI) 4-[[4-(1,1-dimethylphenyl)phenyl]ethoxy]-quinazoline (Fenazaquin), from The

30 Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 507;

(XLVII) 4-phenoxyphenyl-(RS)-2-(pyridyloxy)propyl-ether (Pyriproxyfen), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 1073;

35 (XLVIII) 5-chloro-N-{2-[4-(2-ethoxyethyl)-2,3-dimethylphenoxy]ethyl- }-6-ethylpyrimidine-4-amine (Pyrimidifen), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 1070;

(XLIX) (E)-N-(6-chloro-3-pyridylmethyl)-N-ethyl-N'-methyl-2-nitrovinylidenediamine

(Nitenpyram), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 880;

(L) (E)-N.sup.1-[(6-chloro-3-pyridyl)methyl]-N.sup.2-cyano-N.sup.1-- methylacetamidine (NI-25, Acetamiprid), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 9;

(LI) Avermectin B.sub.1, from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 3;

(LII) an insect-active extract from a plant, especially (2R,6aS,12aS)-1,2,6,6a,12,12a-hexhydro-2-isopropenyl-8,9-dimethoxy-chrome- no[3,4-b]furo[2,3-h]chromen-6-one (Rotenone), from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 1097; and an extract from *Azadirachta indica*, especially azadirachtin, from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 59; and

(LII) a preparation which contains insect-active nematodes, preferably *Heterorhabditis bactecophora* and *Heterorhabditis megidis*, from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 671; *Steinemema feltiae*, from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 1115 and *Steinemema scaptedsci*, from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 1116;

(LIV) a preparation obtainable from *Bacillus subtilis*, from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 72; or from a strain of *Bacillus thuringiensis* with the exception of compounds isolated from GC91 or from NCTC11821; The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 73;

(LV) a preparation which contains insect-active fungi, preferably *Verticillium lecanii*, from The Pesticide Manual, 11.sup.th Ed. (1997), The British Crop Protection Council, London, page 1266; *Beauveda brogniartii*, from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 85 and *Beauveda bassiana*, from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 83;

(LVI) a preparation which contains insect-active viruses, preferably *Neodipridon Sertifer* NPV, from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 1342; *Mamestra brassicae* NPV, from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 759 and *Cydia pomonella* granulosis virus, from The Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 291;

(CLXXXI) 7-chloro-2,3,4a,5-tetrahydro-2-[methoxycarbonyl(4-trifluor- omethoxyphenyl)- carbamoyl]indol[1,2e]oxazoline-4a-carboxylate (DPX-MP062, Indoxy carb), from The

Pesticide Manual, 11.sup.thEd. (1997), The British Crop Protection Council, London, page 453;

(CLXXXII) N'-tert.-butyl-N'-(3,5-dimethylbenzoyl)-3-methoxy-2-methy- lbenzohydrazide (RH-2485, Methoxyfenozide), from The Pesticide Manual, 11.sup.thEd. (1997), The British

5 Crop Protection Council, London, page 1094; and

(CLXXXIII) (N'-[4-methoxy-biphenyl-3-yl]-hydrazinecarboxylic acid isopropylester (D 2341), from Brighton Crop Protection Conference, 1996, 487-493;

(R2) Book of Abstracts, 212th ACS National Meeting Orlando, Fla., August 25-29 (1996), AGRO-020. Publisher: American Chemical Society, Washington, D.C. CONEN: 63BFAF.

10 As a rule, the anthelmintic compositions according to the invention contain 0.1 to 99% by weight, especially 0.1 to 95% by weight of active ingredient of Formula 1 mixtures thereof, 99.9 to 1% by weight, especially 99.8 to 5% by weight of a solid or liquid admixture, including 0 to 25% by weight, especially 0.1 to 25% by weight of a surfactant.

15 As noted above, in another embodiment of the process according to the invention, compounds of Formula 1 and the additional compounds noted hereinbefore may be applied in a distinct and separate manner over time.

The following TESTS demonstrate the control efficacy of compounds of this invention on specific pests. The pest control protection afforded by the compounds is not limited, however, to these species.

20

BIOLOGICAL EXAMPLES OF THE INVENTION

TEST A

25 Adult *L. serricata* (blowflies): Adult flies are placed on beds of blood agar (in individual test wells; 4 flies per well) in which test compound (3-bromo-1-(3-chloro-2-pyridinyl)-N-[4-cyano-2-methyl-6-[(methylamino)carbonyl]phenyl]-1*H*-pyrazole-5-carboxamide) was dissolved/suspended prior to hardening of agar. Flies can take up test compound by both ingestion and contact. Assays are scored by number of dead flies at 2 hours, 4 hours, and 24

30 hours.

35

Rate (ppm)	Number flies in test	Number dead flies at 2 hours		Number dead flies at 4 hours		Number dead flies at 24 hours	
		Chlorop yriphos	Test Compound	Chlorop yriphos	Test Compound	Chlorop yriphos	Test Compound
100	4	4	0	4	0	4	3
50	4	4	0	4	0	4	3
25	4	3	0	4	0	4	2
10	4	1	0	4	0	4	1
1	4	0	0	1	0	4	0
Untreated	16	0	0	0	0	0	0

L. serricata (insect; blowfly) larvae.

Test compound (3-bromo-1-(3-chloro-2-pyridinyl)-N-[4-cyano-2-methyl-6-[(methylamino)-carbonyl]phenyl]-1*H*-pyrazole-5-carboxamide) was mixed with dried blood bovine serum

5 and placed on filter paper discs. Newly emerged larvae of blowfly, *Lucilica sericata* were added to the filter paper, which were ingested by larvae. Activity may occur through both feeding and contact. Four replicates were run per data point. Activity was assessed at 24 hours.


Results: Test compound gave 90-100% mortality at 0.5ppm at 24 hours. Fipronil gave 90-10 100% mortality at 0.5ppm at 24 hours.

15 Although the present invention and its advantages have been described in detail, it should be understood that various changes, substitutions and alterations can be made herein without departing from the spirit and scope of the invention as defined by the appended claims.

CLAIMS

What is claimed is:

1. A method of treating myiasis of an animal by applying to the animal a composition
5 comprising an parasitically effective amount of a compound of Formula 1, an *N*-oxide or a pharmaceutically or veterinarianily acceptable salt thereof



1

wherein:

R¹ is Me, Cl, Br or F;

10 R^2 is F, Cl, Br, C_1 – C_4 haloalkyl or C_1 – C_4 haloalkoxy;

R^3 is F, Cl or Br;

R^4 is H; C_1 – C_4 alkyl, C_3 – C_4 alkenyl, C_3 – C_4 alkynyl, C_3 – C_5 cycloalkyl, or C_4 – C_6 cycloalkylalkyl, each optionally substituted with one substituent selected from the group consisting of halogen, CN, SMe, $S(O)Me$, $S(O)_2Me$, and OMe;

15 R⁵ is H or Me;

R^6 is H, F or Cl; and

R^7 is H, F or Cl.

2. The method of Claim 1 wherein

R^1 is Me or Cl;

20 R^2 is Cl, Br, CF_3 , OCF_2H , OCF_3 or OCH_2CF_3 ; and

R^4 is H, Me, Et, *i*-Pr, *t*-Bu, CH_2CN , $CH(Me)CH_2SMe$ or $C(Me)_2CH_2SMe$.

3. The method of Claim 2 wherein:

R^2 is Cl, Br, CF_3 or OCH_2CF_3 ;

25 R⁴ is H, Me, Et or *i*-Pr; and

\mathbb{R}^5 is H

4. The method of Claim 3 wherein the compound of Formula 1 is 3-bromo-1-(3-chloro-2-pyridinyl)-N-[4-cyano-2-methyl-6-[(methylamino)carbonyl]phenyl]-1*H*-pyrazole-5-carboxamide .

5

5. The method of treatment of myiasis of Claim 1 wherein the myiasis is caused at least in part by larvae selected from the taxonomic families Calliphoridae, Sarcophagidae or Oestridae.

10 6. The method of treatment of Claim 1 wherein the myiasis is caused at least in part by larvae of the family Calliphoridae.

7. The method of treatment of Claim 1 wherein the myiasis is caused at least in part by larvae which are selected from the group consisting of *Lucilia cuprina* and *Lucilia sericata*.

15

8. The method of treatment of Claim 1 wherein the myiasis is caused at least in part by larvae of *Lucilia cuprina*.

9. The method of treatment of Claim 1 wherein the myiasis is caused at least in part by 20 larvae of *Lucilia sericata*.

10. The method of treatment of Claim 1 wherein the animal is a cattle or sheep.

25 11. The method of any of Claims 1-10 wherein the composition comprises at least one additional component selected from the group consisting of solvents and/or carriers, emulsifiers and/or dispersing agents.

12. The method of Claim 11 and wherein the composition comprises at least one additional biologically active compound or agent.

30

13. The method of Claim 12 wherein the additional biologically active compound or agent is selected from the group consisting of macrocyclic lactones, acetyl cholinesterase inhibitors, arthropodgrowth regulators, GABA-gated chloride channel antagonists, mitochondrial electron transport inhibitors, nicotinic acetylcholine agonists/antagonists/activator, oxidative

phosphorylation inhibitors, anthelmintics, sodium channel modulators or other antiparasitic compounds.

14. The method of Claim 13 wherein said biologically active compound is a macrocyclic
5 lactone.

15. The method of Claim 13 wherein said biologically active compound is an acetyl
cholinesterase inhibitor selected from the group of organophosphates and carbamates.

10 16. The method of Claim 13 wherein said biologically active compound is an
arthropodgrowth regulator selected from the group of chitin synthesis inhibitors, ecdysone
agonists/disruptors, lipid biosynthesis inhibitor and juvenile hormone mimics.

15 17. The method of Claim 13 wherein said biologically active compound is a GABA-gated
chloride channel antagonist.

18. The method of Claim 13 wherein said biologically active compound is a mitochondrial
electron transport inhibitor.

20 19. The method of Claim 13 wherein said biologically active compound is a nicotinic
acetylcholine agonist/antagonist/activator.

25 20. The method of Claim 13 wherein said biologically active compound is an oxidative
phosphorylation inhibitor.

21. The method of Claim 13 wherein said biologically active compound is an anthelmintic.

22. The method of Claim 13 wherein said biologically active compound is a sodium channel
modulator.


30

35

CLAIMS

What is claimed is:

1. A method of treating myiasis of an animal by applying to the animal a composition comprising an parasitically effective amount of a compound of Formula 1, an *N*-oxide or a pharmaceutically or veterinarily acceptable salt thereof



1

wherein:

R¹ is Me, Cl, Br or F;

10 R² is F, Cl, Br, C₁–C₄ haloalkyl or C₁–C₄ haloalkoxy;

R³ is F, Cl or Br;

R⁴ is H; C₁–C₄ alkyl, C₃–C₄ alkenyl, C₃–C₄ alkynyl, C₃–C₅ cycloalkyl, or C₄–C₆ cycloalkylalkyl, each optionally substituted with one substituent selected from the group consisting of halogen, CN, SMe, S(O)Me, S(O)₂Me, and OMe;

15 R⁵ is H or Me;

R⁶ is H, F or Cl; and

R⁷ is H, F or Cl.

2. The method of Claim 1 wherein

R¹ is Me or Cl;

20 R² is Cl, Br, CF₃, OCF₂H, OCF₃ or OCH₂CF₃; and

R⁴ is H, Me, Et, *i*-Pr, *t*-Bu, CH₂CN, CH(Me)CH₂SM_e or C(Me)₂CH₂SM_e.

3. The method of Claim 2 wherein:

R² is Cl, Br, CF₃ or OCH₂CF₃;

25 R⁴ is H, Me, Et or *i*-Pr; and

R⁵ is H.