

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 January 2007 (25.01.2007)

PCT

(10) International Publication Number
WO 2007/010263 A2

(51) International Patent Classification:
C12Q 1/68 (2006.01)

(21) International Application Number:
PCT/GB2006/002709

(22) International Filing Date: 20 July 2006 (20.07.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0514935.6 20 July 2005 (20.07.2005) GB

(71) Applicant (for all designated States except US): **SOLEXA LIMITED** [GB/GB]; Chesterford Research Park, Little Chesterford, Saffron Walden Essex CB10 1XL (GB).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **SWERDLOW, Harold, Philip** [US/GB]; SOLEXA LIMITED, Chesterford Research Park, Little Chesterford, Saffron Walden Essex CB10 1XL (GB).

(74) Agents: **WHITE, Nina, Louise** et al; BOULT WADE TENNANT, Verulam Gardens, 70 Gray's Inn Road, London WC1X 8BT (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, **BR**, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, **HR**, HU, **ID**, IL, IN, IS, **JP**, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 2007/010263 A2

(54) Title: METHODS FOR SEQUENCING A POLYNUCLEOTIDE TEMPLATE

(57) Abstract: The invention relates to methods for pairwise sequencing of a polynucleotide template which result in the sequential determination of nucleotide sequence in two distinct and separate regions of the polynucleotide template.

Methods for sequencing a polynucleotide template**Field of the invention**

5 The invention relates to methods for pairwise sequencing of a polynucleotide template, which methods result in the sequential determination of nucleotide sequence in two distinct and separate regions of the polynucleotide template .

10

Background to the invention

 Advances in the study of biological molecules have been led, in part, by improvement in technologies used to
15 characterise the molecules or their biological reactions. In particular, the study of the nucleic acids DNA and RNA has benefited from developing technologies used for sequence analysis .

20 US 5,302,509 describes a method for sequencing a polynucleotide template which involves performing multiple extension reactions using a DNA polymerase or DNA ligase to successively incorporate labelled polynucleotides complementary to a template strand. In such a "sequencing
25 by synthesis" reaction a new polynucleotide strand based-paired to the template strand is built up in the 5' to 3' direction by successive incorporation of individual nucleotides complementary to the template strand. The substrate nucleoside triphosphates used in the sequencing
30 reaction are labelled at the 3' position with different 3' labels, permitting determination of the identity of the incorporated nucleotide as successive nucleotides are added.

- 2 -

In order to carry out accurate sequencing a reversible chain-terminating structural modification or "blocking group" may be added to the substrate nucleosides to ensure that nucleotides are incorporated one at a time in a controlled manner. As each single nucleotide is incorporated, the blocking group prevents any further nucleotide incorporation into the polynucleotide chain. Once the identity of the last-incorporated labelled nucleotide has been determined the label moiety and blocking group are removed, allowing the next blocked, labelled nucleotide to be incorporated -in a subsequent round of sequencing.

In certain circumstances the amount of sequence data that can be reliably obtained with the use of sequencing-by-synthesis techniques, particularly when using blocked, labelled nucleotides, may be limited, typically to around 25-30 cycles of incorporation. Whilst sequencing "runs" of this length are extremely useful, particularly in applications such as, for example, SNP analysis and genotyping, it would be advantageous in many circumstances to be able to reliably obtain further sequence data for the same template molecule .

25

The technique of "paired-end" or "pairwise" sequencing is generally known in the art of molecular biology, particularly in the context of whole-genomic shotgun sequencing (Siegel A.F. et al., *Genomics*. 2000, 68: 237-246; Roach J.C. et al., *Genomics*. 1995, 26: 345-353). Paired-end sequencing allows the determination of two "reads" of sequence from two places on a single polynucleotide

30

- 3 -

template. The advantage of the paired-end approach is that there is significantly more information to be gained from sequencing two stretches each of "n" bases from a single template than from sequencing "n" bases from each of two independent templates in a random fashion. With the use of appropriate software tools for the assembly of sequence information (Millikin *s.c. et al.*, *Genome Res.* 2003, 13: 81-90; Kent, W.J. *et al.*, *Genome Res.* 2001, 11: 1541-8) it is possible to make use of the knowledge that the "paired-end" sequences are not completely random, but are known to occur on a single template, and are therefore very close in the genome. This information has been shown to greatly aid the assembly of whole genome sequence into a consensus sequence.

Paired-end sequencing has typically been performed by making use of specialized circular shotgun cloning vectors known in the art. After cutting the vector at a specific single site, the template DNA to be sequenced (typically genomic DNA) is inserted into the vector and the ends resealed to form a new construct. The vector sequences flanking the insert DNA include binding sites for sequencing primers which permit sequencing of the insert DNA on opposite strands.

A disadvantage of this approach is that it requires time-consuming cloning of the DNA templates it is desired to sequence into an appropriate sequencing vector. Furthermore, because of the need to clone the DNA template into a vector in order to position binding sites for sequencing primers at both ends of the template fragment it is extremely difficult to make use of array-based sequencing techniques. With array-based techniques it is generally

- 4 -

only possible to sequence from one end of a nucleotide template, this often being the end proximal to the point of attachment to the array.

5 With the use of hairpin nucleic acid anchors or double stranded nucleic acid anchors (such as those described in the applicant's co-pending International application published as WO 01/57248) , one end of a template immobilised on an array may be "covalently closed" giving a free 3' end
10 which permits sequencing of the 5' overhanging template strand by successive incorporation of nucleotides. However, given that the distal portions (distal from the point of attachment to the array) of such immobilised templates are generally single-stranded and that the sequence of the
15 template is usually unknown prior to immobilisation on the array, it is not straightforward to devise means for determining the sequence of the distal end of the immobilised template, beyond the first "run" of sequence that can be obtained from the free 3' end provided by the
20 anchor. It is not possible simply to design a sequencing primer complementary to a region of the template whose sequence is unknown.

WO 2004/070005 describes a method for double-ended
25 sequencing of a polynucleotide template which can be carried out on a solid support. The method relies on simultaneous hybridisation of two or more primers to a target polynucleotide in a single primer hybridization step. Following the hybridization step, all of the primers
30 hybridized to the template are blocked except for one, which has a free 3' hydroxyl group which serves as an initiation point for a first sequencing reaction. Sequencing proceeds

- 5 -

until no further chain elongation is possible, or else the sequencing reaction is terminated. Then one of the blocked primers is unblocked to give a free 3' hydroxyl and a second sequencing reaction is performed from this initiation point.

5 An advantage of this approach is that there is no need to perform any denaturation and re-hybridization steps between the first and second sequencing reactions, as the two primers providing the initiation points are annealed in a single hybridisation step. Thus, the template remains
10 intact and attached to the solid support throughout.

A major drawback of this approach based in hybridisation of blocked and unblocked primers is that it is necessary to know the sequence of at least two regions of
15 the polynucleotide template to be sequenced in order to design two or more suitable primers capable of binding to the target. If the method is to be used to sequence polynucleotides of unknown sequence then it is necessary to carry out sample preparation steps in order to add regions
20 of known sequence to the polynucleotide to be sequenced in order to provide the necessary primer-binding sites. This can be achieved, for example, by amplification or by sub-cloning a template of unknown sequence into a vector in order to add known sequences onto the 5' and 3' ends of the
25 template.

The present inventors have sought to develop techniques which generally permit the paired-end or pairwise sequencing approach to be used without any knowledge of the sequence at
30 the distal end of the template and without the need for any intermediate cloning of the template into a vector. Such techniques would permit pairwise sequencing to be used in

conjunction with a wide range of array-based sequencing technologies, including single molecule arrays as well as clustered arrays .

5 Description of the invention

In accordance with a first aspect of the invention there is provided a method for pairwise sequencing of a polynucleotide template, the method comprising:

10

(a) providing a polynucleotide template having a first free 3' hydroxyl group which is positioned to initiate sequencing of a first region for sequence determination on a first template strand,

15

(b) carrying out a first sequencing reaction by sequential addition of nucleotides to the first free 3' hydroxyl group to determine the sequence of the first template strand in the first region,

20

(c) generating a second free 3' hydroxyl group within the polynucleotide template at a nucleotide position which is spaced apart from the last nucleotide added in the sequencing reaction of step (b) , wherein the second free 3' hydroxyl group is positioned to initiate sequencing of a second region for sequence determination on a second template strand, and wherein the second free hydroxyl group is in continuous covalent linkage with the last nucleotide added in the sequencing reaction of step (b) , and

30

(d) carrying out a second sequencing reaction by sequential addition of nucleotides to the second free 3' hydroxyl group

- 7 -

to determine the sequence of the second template strand in the second region.

The invention provides a method for sequencing two
5 regions of a polynucleotide template, referred to herein as the first and second regions for sequence determination. The first and second regions for sequence determination may be on the same strand of a single-stranded polynucleotide template, or they may be on complementary strands of a
10 double-stranded or self-complementary polynucleotide template .

If the first and second regions for sequence
determination occur on a single polynucleotide strand then
15 they must not be immediately adjacent to each other, meaning that the last nucleotide to be sequenced in the first region is not in phosphodiester linkage with the first nucleotide to be sequenced in the second region. It is an essential feature that the two regions for sequence determination must
20 be separated by at least one nucleotide, and are preferably separated by at least 50, more preferably at least 100 or more preferably at least 500 nucleotides.

If the first and second regions for sequence
25 determination occur on complementary strands of a double-stranded polynucleotide template, or a self-complementary template, then the two regions may or may not be complementary to each other.

30 A key feature of the method of the invention is that it does not require primer hybridisation to the template to be sequenced in order to provide initiation points for the

first and second sequencing reactions. This is a particular advantage when the method is carried out on a solid support, such as a chip-based array, since there is no need to carry out any hybridisation reactions on the solid support.

5

The starting point for the method of the invention is the provision of a polynucleotide template including a first free 3' hydroxyl group which can serve as an initiation point for a sequencing reaction.

10

In the context of this invention the term "sequencing reaction" refers to any polynucleotide "sequencing-by-synthesis" reaction which involves sequential addition of nucleotides to a growing polynucleotide chain in the 5' to 3' direction using a polymerase in order to form an extended polynucleotide chain complementary to the template region to be sequenced. The identity of the added nucleotide is preferably determined after each addition step, thus the sequence of the template may be inferred using conventional Watson-Crick base-pairing rules.

20

The starting polynucleotide template is at least partially double-stranded, or self-complementary. The first free 3' hydroxyl group is provided on one strand that is base-paired to a complementary strand (which can be the same polynucleotide strand if the template is self-complementary), which forms the first strand of the template to be sequenced (first template strand), in the region immediately upstream of the first 3' hydroxyl group. The region of base-pairing between the strand bearing the first hydroxyl group and the first template strand need only be long enough to hold the two strands together under the

30

- 9 -

conditions used for the subsequent sequencing reaction, in order that the hydroxyl group is positioned to initiate the sequencing reaction. Therefore, this region of base-pairing may be as short as from 10 to 13 base-pairs. In embodiments
5 wherein the template is linked to a hairpin linker (see below), the region of base-pairing is provided by the "stem" portion of the hairpin linker.

The first template strand overhangs the first 3'
10 hydroxyl group in order to define a first region of the template to be sequenced. The overhanging portion of the first template strand may be single-stranded. Alternatively, the overhanging region may be annealed to a complementary strand downstream of the free 3' hydroxyl.
15 The latter arrangement may be used if the first sequencing reaction is to be performed using a strand-displacement polymerase. The length of the overhanging strand may be from 100 nucleotides (or base-pairs if double-stranded) up to 1kb, or even longer. Preferably the overhanging strand
20 will be at least 500 nucleotide (or base-pairs).

The template may be provided as a single polynucleotide molecule which is partially self-complementary. For example, the template polynucleotide molecule may fold back
25 on itself to form a self-complementary hairpin structure at its 3' end.

In one embodiment the starting polynucleotide may be formed by ligating a self-complementary hairpin linker
30 polynucleotide to the desired polynucleotide template to be sequenced, such that the 5' end of the hairpin linker is joined to the 3' end of the first template strand. The

- 10 -

first free 3' hydroxyl group is then provided by the 3' end of the hairpin linker. An advantage of this approach is that it may be used for sequencing templates of unknown sequence. Suitable self-complementary hairpin polynucleotide linkers include those described in the applicant's published International application WO 01/57258.

A first sequencing reaction is carried out by sequential addition of nucleotides to the first free 3' hydroxyl group. The nature of the added nucleotide is preferably determined after each nucleotide addition in order to determine the sequence of the first template strand in the first region for sequence determination. This first sequencing reaction can proceed for as long as is technically possible within the limitations of the chosen sequencing methodology, for example until the sequence obtained is no longer accurate or reliable. The first sequencing reaction will preferably involve sequencing of from 10 to 200 or more consecutive bases, preferably from 15 to 35 consecutive bases of the first template strand, preferably from 20 to 30 consecutive bases and more preferably about 25 consecutive bases of the first template strand. The precise number of bases sequenced will be dependent upon the limitations of the chosen sequencing methodology, and more specifically the methodology used to determine the nature of the base added in each nucleotide addition step.

Once the first sequencing reaction is deemed to be complete, sequencing of a second region of the polynucleotide template can be carried out. In order to sequence a second region of the template it is necessary to

- 11 -

generate a second free 3' hydroxyl group which serves as an initiation point for a second sequencing reaction. The second free 3' hydroxyl group must be provided on a polynucleotide strand which is base-paired to the second
5 template strand to be sequenced in the region immediately upstream of the second free 3' hydroxyl group. The second template strand overhangs the second free 3' hydroxyl group to define a second region of the template to be sequenced. The first and second template strands may be the same strand
10 of the polynucleotide template if the template is single-stranded, or self-complementary.

It is an essential feature of the invention that the second free 3' hydroxyl group is "spaced apart from" the
15 last nucleotide added in the first sequencing reaction, meaning that the second free 3' hydroxyl group must not be provided by the last nucleotide added in the first sequencing reaction, and that there must be at least one nucleotide separating the last nucleotide added in the first
20 sequencing reaction and the nucleotide which provides the second free 3' hydroxyl group. Preferably the last nucleotide added in the first sequencing reaction and the nucleotide which provides the second free 3' hydroxyl group will be separated by at least 50 nucleotides, more
25 preferably at least 100 nucleotides and more preferably at least 500 nucleotides.

It is also a feature of the invention that the last nucleotide added in the first sequencing reaction and the
30 nucleotide which provides the second free 3' hydroxyl group should be generated in "continuous covalent linkage". This means that the last nucleotide added in the first sequencing

- 12 -

reaction and the nucleotide which provides the second free
3' hydroxyl group should form part of a single
polynucleotide molecule with continuous backbone
phosphodiester linkage, or at least that they should have
5 been so-joined on a single polynucleotide molecule at some
stage in the sequencing method. This polynucleotide
molecule may be a single strand or it may be a partially
self-complementary polynucleotide strand, having a hairpin
or stem-loop structure.

10

The requirement for the last nucleotide added in the
first sequencing reaction and the nucleotide which provides
the second free 3' hydroxyl group to be in continuous
covalent linkage is advantageous for pairwise sequencing of
15 templates of unknown sequence, since it overcomes the
problem of designing a primer to initiate sequencing of
second region of a template of unknown sequence. Thus, the
second free 3' hydroxyl group is NOT provided by
hybridisation of a separate primer. This in turn means that
20 when the method of the invention is carried out on an array
there is no need to carry out any hybridisation steps after
the templates are immobilised on the array.

25 There are several different ways in which the second
free 3' hydroxyl group may be generated, as will be
discussed in further detail below.

30 Non-limiting embodiments of the invention are now
described in detail with reference to the accompanying
drawings, in which:

- 13 -

Figures 1 and 2 illustrate different embodiments of the sequencing method according to the invention which allow sequencing of first and second regions of a single-stranded polynucleotide template;

5

Figures 3 to 6 illustrate different embodiments of the sequencing method according to the invention which allow sequencing of first and second regions on complementary strands of a double-stranded polynucleotide template.

10

Referring to figure IA, a polynucleotide template (1) is provided, having a first free 3' hydroxyl group (2). In this embodiment the template to be sequenced is a single polynucleotide strand (3). In the embodiment shown in Figure IA the first free 3' hydroxyl group is provided by the 3¹ end of a self-complementary hairpin linker (4) which is ligated to the template strand (3) such that the 5' end of the linker (4) is joined to the 3' end of the template strand (3). The linker is also linked to a solid support (5) via any suitable linkage (6). The template strand (3) overhangs the first free 3¹ hydroxyl group to define a first region of the template to be sequenced (7). Although the specific embodiment shown in Figure IA involves use of a hairpin which acts as a primer for sequencing and provides a means for attachment to a solid support the invention is not intended to be limited to the use of hairpins. Any suitable oligonucleotide primer hybridised to the template strand (3) could be used to provide an initiation point for sequencing. Linkage to the solid support could be provided via the primer or via the template strand itself. If linkage to the solid support is provided via the template strand then this may occur at the 5' or the 3' end of the template strand, or

30

- 14 -

even via an internal portion of the template provided that this does not interfere with subsequent sequencing reactions. Template strands generated by solid-phase amplification will generally be linked to a solid support
5 via covalent linkage at their 5' end. Such template strands can be sequenced with the use of "standard" oligonucleotide sequencing primers.

Referring back to the embodiment shown in Figure IA, a
10 first sequencing reaction proceeds by sequential addition of nucleotides to the first free 3' hydroxyl group (2). When this sequencing reaction is determined or deemed to be complete, the strand complementary to the template strand
(3) is further extended by sequential addition of a known or
15 unknown number of nucleotides to the last nucleotide added in the first sequencing reaction in a nucleotide addition step. This nucleotide addition step occurs without sequencing, i.e. nucleotides are added without any determination of the nature of the nucleotide added in each
20 addition. Thus, the nucleotides added in the addition step need not be modified to permit sequencing and are preferably "unmodified", meaning that they do not contain any labels required for sequencing, e.g. fluorescent labels, or any blocking groups. It is most preferred to use naturally
25 occurring nucleotides in this addition step, although nucleotides bearing "modifications" other than labels or blocking groups could be used as long as the modification does not interfere with incorporation into polynucleotide by the chosen polymerase or addition of subsequent nucleotides
30 (e.g. no chain termination effect) to a material extent.

- 15 -

The free 3' hydroxyl group of the last nucleotide added in the nucleotide addition step forms the second free 3' hydroxyl group (9) - shown in Fig. 1B. The template strand (3) now overhangs the second free 3' hydroxyl group to
5 define a second region of the template to be sequenced (8). Thus, in this embodiment the first and second template strands are provided by a single template strand (3). A second sequencing reaction can then proceed from the second free 3' hydroxyl group.

10

The addition of further nucleotides following the first sequencing reaction serves to move the sequencing reaction on to a new position on the template. Constraints in the sequencing methodology may limit the number of sequential
15 base additions which can be made with accurate determination of the added nucleotide, meaning that it would not be possible to reach this point in the template in a single continuous sequencing reaction. By switching to addition of further (unmodified) nucleotides without sequencing and then
20 initiating a second sequencing reaction it is possible to sequence a region farther into the template, by carrying out two sequencing reactions separated by a number of "unsequenced" nucleotides.

25

The number of further nucleotides added between the first and second sequencing reactions is typically at least 50, more preferably at least 100 and more preferably at least 500.

30

Figure 2 illustrates a further embodiment of the invention which is substantially similar to the embodiment shown in Figure 1 and described above, except that the

- 16 -

nucleotides added to further extend the complementary strand after completion of the first sequencing reaction comprise a mixture of unmodified deoxynucleotides (dATP, dCTP, dTTP and dGTP) and a small quantity of deoxyuracil, resulting in
5 incorporation of uracil into the complementary strand at some of the positions opposite an A in the template strand. An enzyme which cleaves DNA immediately adjacent to uracil, for example uracil DNA glycosylase, also known as glycosidase or uracil N-glycosylase or uracil DNA N-
10 glycosylase, may then be used to generate a second free 3' hydroxyl group (9) by cleavage of the complementary strand (10) .

Figure 3 illustrates a still further embodiment of the
15 invention which allows for sequencing of two regions on complementary strands of a double-stranded template. As shown in Figure 3A, formation of the starting polynucleotide template and sequencing of a first region of the template
(7) proceeds as described for the embodiment shown in Figure
20 1. After the first sequencing reaction is deemed to be complete further nucleotides are added to the 3' hydroxyl group of the last nucleotide added in the first sequencing reaction in order to fully extend the complementary strand
(10) to the full length of the first template strand (3) .
25 This extension reaction proceeds without sequencing, i.e. without determination of the nature of the added nucleotides, and results in the production of a blunt-ended fully double-stranded template ligated at one end to a hairpin linker (4) , which itself is linked to a solid
30 support (5) (shown Figure 3B) .

- 17 -

The open end of the double-stranded template (distal from the solid support) is then ligated to a second hairpin linker polynucleotide (11) which has a phosphate group at the 5' end and a free hydroxyl group at the 3' end, such that the 5' end of the second hairpin linker is joined to the free 3' end of the extended complementary strand (10). As illustrated in Figure 3, the 5' end of the first template strand is dephosphorylated (or blocked by some other means) such that it is not capable of forming a phosphodiester linkage with the hydroxyl group at the 3' end of the second hairpin linker. The resulting product has a nicked circular structure (3C). A substantially equivalent structure may also be formed by ligating the second hairpin linker to both the first and second template strands to form a closed circle and then creating a nick in the first template strand (3) or the hairpin linker. The free 3' hydroxyl group (9) of the second hairpin linker provides the second free 3' hydroxyl group for initiation of the second sequencing reaction. The second region of the template to be sequenced (8) is on the complementary strand (10), thus this embodiment allows sequencing of two regions (7, 8) on complementary strands of a double-stranded template. The second free 3' hydroxyl group is formed in continuous covalent linkage with the last nucleotide added in the first sequencing reaction, thus there is no need for hybridisation of a separate sequencing primer in order to initiate sequencing of the second region of the template and no need for any fore-knowledge of the sequence of the template.

The second sequencing reaction can be carried out starting from the nicked circle structure shown in Fig. 3C with the use of a strand displacing polymerase enzyme. In

- 18 -

an alternative embodiment, illustrated in Figs. 3D and 3E, a portion of what was the first template strand (3) may be removed prior to the second sequencing reaction. This may be achieved, for example, by creating a second nick (12) towards the proximal end of the first template strand near the solid support (5). The nick may be created in the first template strand or within the first hairpin linker (4), provided that this does not affect the linkage to the solid support. In a preferred embodiment the nick may be created with the use of an appropriate nicking (single side cutting) restriction enzyme (endonuclease). In order to direct nicking at an appropriate position the first hairpin linker may contain a recognition sequence for a nicking enzyme which directs cleavage of a single strand at a site before, at or beyond the 3' end of the first linker. By including this recognition site in the hairpin linker, generation of the "nick" is independent of the sequence of the template and dependent only on ligation of the linker to the template. This means that the method may be universally applied to templates of unknown, and different, sequence. It will be appreciated that nicking can also be accomplished by blocking one side of a standard restriction enzyme cleavage site (in the hairpin linker sequence or sequences) using methods familiar to those skilled in the art, e.g., by using thiophosphate linkages in one side of the restriction enzyme recognition site, to prevent cutting in that side, but not in the other, or by the use of DNA methylation.

Following the nicking reaction the resulting product may be denatured in order to remove the "free" portion of the first template strand located upstream (5') of the cleavage site, as shown in Fig. 3E.

- 19 -

Figure 4 illustrates a further embodiment of the invention which allows sequencing of two regions on complementary strands of a double-stranded template. This method starts with provision of a fully double-stranded (i.e. blunt ended) polynucleotide template in solution comprising a first template strand (3) annealed to a complementary strand (10). The first template strand has a 5¹ phosphate group, whereas the complementary strand lacks a 5¹ phosphate group (or 5' phosphate may be blocked) such that it cannot be joined in phosphodiester linkage to the 3' hydroxyl group of a further nucleotide.

The double-stranded template can be any double-stranded polynucleotide that it is desired to sequence. In a preferred embodiment it will be a fragment of genomic DNA. Double-stranded templates bearing a 5¹ phosphate on one strand but lacking a 5' phosphate on the complementary strand can be generated in a number of different ways. For example, in one embodiment templates can be generated by limited or controlled digestion of double-stranded DNA fragments bearing 5' phosphates on both strands with a phosphatase enzyme, in order to generate a maximal proportion of strands in which 5' phosphate is removed from one strand only.

In a further embodiment double-stranded DNA fragments may be treated with phosphatase in order to remove 5' phosphate from both strands. The resulting dephosphorylated product may then be subject to limited or controlled treatment with a kinase in order to restore 5' phosphate, limited treatment ensures that a maximal number

- 20 -

of fragments will have 5' phosphate restored on one strand only.

In a still further embodiment an adapter, consisting of
5 paired oligonucleotides having hydroxyl groups at both 3'
and 5' ends of both strands may be ligated to one end only of
a double-stranded DNA fragment which has 5' phosphate on
both strands. A suitable site for attachment of the adapter
to one end only may be generated by cutting the template
10 with a suitable restriction enzyme at a site proximal to the
chosen end.

In a still further embodiment an adapter
oligonucleotide having a 5' phosphate group may be ligated
15 to one end only of a double-stranded DNA template which has
been treated with phosphatase to remove all 5' phosphates.
Again, suitable site for attachment of the adapter to one
end of the template may be generated by cutting the template
with a suitable restriction enzyme at a site proximal to the
20 chosen end.

A second hairpin linker polynucleotide (11) is ligated
to one end of the double-stranded template. This second
linker has hydroxyl groups at both its 3' and 5' ends, thus in
25 the ligation reaction the 3' end of the second linker (OH)
is joined to the 5' end of the first template strand
(phosphorylated), but the 5' end of the second linker (also
OH) remains free. The second hairpin linker may include a
marker moiety, for example biotin, which facilitates
30 selection of template molecules that have been successfully
ligated to the second linker.

- 21 -

A first hairpin linker (4) is ligated to the second end of the double-stranded template. This first linker has a hydroxyl group at the 3' end and a phosphate group at the 5' end. Thus, in the ligation reaction the 5' end of the first linker (phosphate) is joined to the 3' end of the first template strand (OH) but the 3' end of the first linker (OH) remains free. The first hairpin linker may be attached to a solid support (5) via a suitable linkage (6). Attachment to the solid support is preferably carried out after the first linker (4) has been ligated to the double-stranded template. This provides a means of selection, since only templates that have been correctly ligated to the first linker will be linked to the solid support.

The terms "first" and "second" hairpin linker are used for convenience solely for the purposes of identifying the different structure/function of the two linkers. The designations "first" and "second" do not require the linkers to be joined to the template strands in any particular order. In the embodiment shown in Figure 4 it may be convenient for the second linker to be ligated to the template before the first linker, since the presence of biotin, or other suitable selectable marker moiety, will enable selection of templates bound to this linker prior to linkage to the solid support.

The product resulting from ligation of the first and second hairpin linkers is a closed circle which is "nicked" in two places, as illustrated in Fig. 4A. The "free" strand between the two nicks (10) may be removed by denaturation to give the structure illustrated in Fig. 4B. Following

- 22 -

denaturation the 5' end (17) of the second linker (11) may be treated with a kinase to restore the 5' phosphate group.

The 3' end of the first linker provides the first free 3' hydroxyl group (2) for initiation of the first sequencing reaction (illustrated Fig. 4B). When the first sequencing reaction is deemed to be complete, further nucleotides are added to the 3' hydroxyl group of the last nucleotide added in the first sequencing reaction in order to fully extend the complementary strand (10) to the full length of the first template strand. This extension reaction proceeds without sequencing, i.e. without determination of the nature of the added nucleotides. The 3' hydroxyl end of this extended strand (which will subsequently form the second template strand) is ligated to the phosphate group newly added to the 5' end of the second hairpin linker (11). Addition of phosphate to the 5' end of second linker could be carried out after extension of the complementary strand, immediately prior to the ligation reaction, if convenient. The product of this ligation is a fully closed circle structure, as illustrated in Fig. 4C.

In order to generate a second free 3' hydroxyl group for initiation of the second sequencing reaction the closed circle structure shown in Fig. 4C may be nicked at a suitable position in the first template strand (3). The position of this first "nick" (13) will determine the position of the second region to be sequenced on the complementary strand (10, now referred to as the second template strand). Again, the nick may be generated with the use of a suitable nicking or side-cutter enzyme. The second hairpin linker (11) may include a recognition site for a

- 23 -

nicking enzyme which directs nicking/cleavage at a site before, at or beyond the 3' end of the second linker. Thus, generation of the nick (13) is dependent only on the presence of the hairpin linker and is independent of the
5 sequence of the template.

Following generation of a first nick (13), the second sequencing reaction can proceed from the second free 3' hydroxyl group generated as a result of the nicking reaction
10 with the use of a strand displacing polymerase enzyme. Alternatively, a portion of the first template strand downstream from (3' of) this first nick can be removed altogether prior to the second sequencing reaction. This can be achieved by generating a second nick (14) spaced
15 apart from (downstream of) the first nick (13). Again a suitable nicking enzyme can be used for this purpose. In order to form the second nick the first hairpin linker (4) may include a recognition site for a nicking enzyme which directs cleavage at a site before, at or beyond the 5' end
20 of the first hairpin linker. Following generation of the second nick, the portion of the first template strand located between the two nicks may be removed by denaturation, leaving the structure illustrated in Fig. 4E. The second sequencing reaction may then proceed from the
25 second free 3' hydroxyl group (9).

Figure 5 illustrates a further embodiment of the invention which allows sequencing of two regions on complementary strands of a double-stranded template. The
30 starting template for this method is a partially self-complementary covalently closed circle structure as illustrated in Figure 5A. This structure can easily be

- 24 -

formed by ligation of first (4) and second (11) hairpin linkers to the ends of a fully double-stranded, blunt ended template comprising first (3) and second (10) template strands .

5

A first free 3' hydroxyl group is generated by cleavage of one template strand (10) of the covalently closed circle to generate a first "nick" (13) with the use of an appropriate nicking or side-cutter enzyme. Formation of this nick can be directed by including a suitable recognition site in the first hairpin linker (4) which directs cleavage (nicking) at a site before, at or beyond the 3¹ end of the linker. This first "nick" generates the first free 3' hydroxyl group (2) for initiation of the first sequencing reaction. Sequencing may then proceed from this initiation point using a strand displacement polymerase enzyme .

Once this first sequencing reaction is deemed to be complete a second nick (14) may be generated within the opposite (complementary) template strand (3) . Formation of this nick can be achieved by the inclusion of a suitable recognition site in the second hairpin linker (11) which directs cleavage (nicking) at a site before, at or beyond the 3' end of this linker. This second nick generates the second free 3¹ hydroxyl group (9) for initiation of the second sequencing reaction. Sequencing may then proceed from this initiation point using a strand displacement polymerase enzyme. Thus, this method allows sequencing of two target regions (7,8) on opposite strands of a double-stranded template.

30

- 25 -

In this method the nucleotides which will ultimately provide the first and second free 3' hydroxyl groups are in continuous covalent linkage before initiation of the first sequencing reaction through the formation of the closed
5 circle structure illustrated in Figure 5A. Thus, this embodiment provides the same technical advantages as the other embodiments of the invention, since it permits sequencing of two distinct regions of a template of unknown sequence, without the need for hybridisation of two (or
10 more) separate sequencing primers to provide initiation points for sequencing.

Figure 6 illustrates a further embodiment of the invention which allows sequencing of two regions on
15 complementary strands of a double-stranded template. The starting template for this method is a partially self-complementary covalently closed circle structure (6A) identical to that illustrated in Figure 5A. The covalently closed circle may be formed by ligating first (4) and second
20 (11) hairpin linkers to a double-stranded polynucleotide comprising first (3) and second (10) template strands.

The second template strand (10) is then "nicked" at two sites (13,14) spaced apart from each other. These nicks can be formed by the inclusion of suitable recognition sequences
25 in the first (4) and second (11) hairpin linkers. The portion of the template strand (10) located between the two nicks is then removed by denaturation to generate the structure shown in Figure 6C. A first sequencing reaction may then proceed from the first free 3' hydroxyl group (2).

30 When the first sequencing reaction is complete, the method continues substantially as described above with reference to Figure 4, from completion of the first

- 26 -

sequencing reaction. In brief summary, the covalently closed structure is re-generated by extending the strand formed in the first sequencing reaction to produce a fully double-stranded template and then ligating the 3¹ end of this extended strand to the free 5' end of the second linker (11). A further nick may then be formed in the first template strand (3) at a third nicking site (15). Again formation of this nick can be directed by including a suitable recognition sequence in the second linker (11) which directs cleavage before, at or beyond its 3' end. The free 3' hydroxyl group generated as a result of this nicking then provides the second free 3¹ hydroxyl group (9) for initiation of the second sequencing reaction.

The second sequencing reaction may be carried out with the use of a strand displacement polymerase. Alternatively, a portion of the strand downstream of the third nicking site (15) may be removed by nicking this strand at a still further (fourth) nicking site (16). Again formation of this nick can be directed by including a suitable recognition sequence in the first hairpin linker (4) which directs cleavage before, at or beyond its 5' end. The "free" portion of the template strand (3) located between the two nicking sites (15,16) may then be removed by denaturation and sequencing can proceed from the second free 3' hydroxyl group (9). Thus, this embodiment results in sequencing of two regions (7,8) located on opposite strands of a double-stranded template.

The following preferred features apply *mutatis mutandis* to all embodiments of the invention:

- 27 -

The polynucleotide template to be sequenced using the method of the invention may be any polynucleotide that it is desired to sequence. A key advantage of the invention is that the template may be of unknown sequence. However, the
5 method may also be used to sequence templates of known or partially known sequence, for example in re-sequencing applications. With the use of arrays it is possible to sequence multiple templates of the same or different sequence in parallel. A particularly preferred application
10 of the method is in the sequencing of fragments of genomic DNA.

Certain embodiments of the method of the invention make use of hairpin linker polynucleotides, as described above.
15 Self-complementary hairpin polynucleotide linkers generally have a "stem-loop" structure, formed by base-pairing of complementary polynucleotides that are covalently linked at one end. The covalent linkage may be provided by a short single-stranded polynucleotide loop, or may be a non-
20 polynucleotide chemical linkage. Preferred chemical linkages include an arrangement of two hexaethylene glycol (heg) spacers separated by an amino deoxy-thymidine nucleotide. Preferred hairpin linker polynucleotides are those described in the applicant's published International
25 application WO 01/57258. The hairpin linker is effectively "self-priming", including a free 3' hydroxyl group to which further nucleotides may be added. This 3' hydroxyl group provides an initiation point for sequencing of any polynucleotide template strand ligated to the 5' end of the
30 hairpin, independent of the sequence of the template strand. The precise nucleotide sequence of the hairpin linker is not material to the invention.

- 28 -

Hairpin linkers can also provide a means of linking the template strands to be sequenced to a solid support, for example to form an array. For example, hairpins including the above-described linkage consisting of two hexaethylene glycol (heg) spacers separated by an amino deoxy-thymidine nucleotide can be linked to epoxide-modified surfaces (e.g. as described in WO 01/57248). The solid support can be of any suitable material, for example glass, ceramic, silicon, plastics or other polymeric material. Suitable support materials are also described in the applicant's published International applications WO 00/06770. The support may take any convenient form. For example it may be substantially planar, such as a slide or "DNA chip", or may be a three dimensional support, or a support formed from multiple discrete units, for example beads of glass, plastics, magnetic material etc.

Templates may be attached to a solid support via any suitable linkage method known in the art. Preferably linkage will be via covalent attachment. The templates may be attached to the support directly or via a hairpin linker. If hairpin linkers are used then they may be attached to the solid support before they are linked to template (s) to be sequenced. For example, hairpin linkers may be attached to the solid support to form an array prior to attachment of the template (s). Techniques for forming arrays of hairpin linkers are described in WO 01/57248. The templates/ linkers are preferably attached to the solid support via a covalent linkage. Linkage may be made via any part of the template or hairpin, provided that this does not interfere with the ability of the template/ linker to participate in a

- 29 -

sequencing reaction. If hairpins are used, the linkage may-
advantageous ly be made via an internal portion of the
hairpin, leaving the 3' end free to initiate a sequencing
reaction and the 5' end free for attachment to the template .

5 In the case of template strands generated by solid-phase
amplification attachment to the solid support will usually
be via covalent linkage with the 5' end of the template.

If the templates are "arrayed" on a solid support then
10 the array may take any convenient form. An advantage of the
method of the invention is that large numbers of templates
of different sequence can be processed in parallel, since
the method is independent of the sequence of the template
and does not require any sequence-specific hybridisation
15 steps to take place on the array. Thus, the method is
applicable to all types of "high density" arrays, including
single-molecule arrays and clustered arrays.

Single molecule arrays are formed by immobilisation of
20 a single polynucleotide molecule at each discrete site that
is detectable on the array. Single -molecule arrays
comprised of nucleic acid molecules that are individually
resolvable by optical means and the use of such arrays in
sequencing are described, for example, in WO 00/06770.
25 Single molecule arrays comprised of individually resolvable
nucleic acid molecules including a hairpin loop structure
are described in WO 01/57248. The method of the invention
is suitable for sequencing template molecules on single
molecule arrays prepared according to the disclosures of WO
30 00/06770 of WO 01/57248.

- 30 -

However, it is to be understood that the scope of the invention is not intended to be limited to the use of the method in connection with single molecule arrays. The method may be used for sequencing on essentially any type of array formed by immobilisation of nucleic acid molecules on a solid support, and more particularly any type of high-density array. In addition to single molecule arrays suitable arrays may include, for example, multi-polynucleotide or clustered arrays in which distinct regions on the array comprise multiple copies of one individual polynucleotide molecule or even multiple copies of a small number of different polynucleotide molecules (e.g. multiple copies of two complementary nucleic acid strands) .

In particular, the method of the invention may be utilised for sequencing on multi polynucleotide or "clustered" arrays. Multi-polynucleotide or clustered arrays of nucleic acid molecules may be produced using techniques generally known in the art. By way of example, WO 98/44151 and WO 00/18957 both describe methods of nucleic acid amplification which allow amplification products to be immobilised on a solid support in order to form arrays comprised of clusters or "colonies" of immobilised nucleic acid molecules. The nucleic acid molecules present on the clustered arrays prepared according to these methods may be suitable templates for sequencing using the method of the invention. However, the invention is not intended to be limited to use of the method in sequencing reactions carried out on clustered arrays prepared according to these specific methods.

- 31 -

When using arrays formed from nucleic acid "colonies", such as arrays prepared according to the teaching in WO 98/44151 and WO 00/18957, it is not necessary to use a hairpin linker polynucleotide to link the template molecule to the array and provide the first free 3' hydroxyl group for sequencing. The arrays of WO 98/44151 and WO 00/18957 are formed by "on-chip" amplification using primers and templates immobilised on a solid support. Once amplification is complete, the colonies may comprise pluralities of double-stranded nucleic acid molecules. These double-stranded molecules may be processed to form templates suitable for use in the methods of the invention, either by forming a "nick" in one strand only of the double-stranded molecules or by removing a portion of one strand altogether. Either way, a free 3' hydroxyl group will be formed on one strand which serves as an initiation point for the first sequencing reaction. After the first sequencing reaction is complete, formation of the second free 3' hydroxyl group can proceed using some of the alternatives described above, with appropriate modification.

As outlined above, sequencing can be carried out using any suitable "sequencing-by-synthesis" technique, wherein nucleotides are added successively to a free 3' hydroxyl group, resulting in synthesis of a polynucleotide chain in the 5' to 3' direction and the nature of the nucleotide added is preferably determined after each addition.

One preferred sequencing method which can be used in the methods of the invention relies on the use of modified nucleotides that can act as chain terminators. Once the modified nucleotide has been incorporated into the growing

- 32 -

polynucleotide chain complementary to the region of the template being sequenced there is no free 3'-OH group available to direct further sequence extension and therefore the polymerase can not add further nucleotides. Once the nature of the base incorporated into the growing chain has been determined, the 3' block may be removed to allow addition of the next successive nucleotide. By ordering the products derived using these modified nucleotides it is possible to deduce the DNA sequence of the DNA template.

Such reactions can be done in a single experiment if each of the modified nucleotides has attached a different label, known to correspond to the particular base, to facilitate discrimination between the bases added at each incorporation step. Alternatively, a separate reaction may be carried out containing each of the modified nucleotides separately.

The modified nucleotides carry a label to facilitate their detection. Preferably this is a fluorescent label. Each nucleotide type may carry a different fluorescent label. However the detectable label need not be a fluorescent label. Any label can be used which allows the detection of the incorporation of the nucleotide into the DNA sequence .

One method for detecting the fluorescently labelled nucleotides comprises using laser light of a wavelength specific for the labelled nucleotides, or the use of other suitable sources of illumination. The fluorescence from the label on the nucleotide may be detected by a CCD camera or other suitable detection means.

- 33 -

The methods of the invention are not limited to use of the sequencing method outlined above, but can be used in conjunction with essentially any sequencing methodology which relies on successive incorporation of nucleotides into a polynucleotide chain. Suitable techniques include, for example, Pyrosequencing™, FISSEQ (fluorescent in situ sequencing) . . MPSS (massively parallel signature sequencing) and sequencing by ligation-based methods.

The invention will be further understood with reference to the following non-limiting experimental example:

Example

Methods and Materials

(1) Extension of polynucleotide strands with polymerase and all four nucleotides-without sequencing.

A polymerase (e.g., Taq DNA polymerase, Pfu polymerase, Klenow fragment of *E.coli* Pol I, etc.) is added at a suitable concentration in a buffer designed to be used with that enzyme. All four deoxynucleotide triphosphates are added at a concentration in the 1-11 micromolar range. Extension is for a time (10 sec to a few minutes) that will allow approximately 500 base pairs to be incorporated.

(2) Extension of polynucleotide strands with polymerase, deoxyuracil and all four nucleotides.

As in previous description (3) , but with a small amount (1 part in 100, e.g. compared with standard deoxythymidine triphosphate) of deoxyuracil triphosphate mixed in. The

- 34 -

amount of U nucleotide is adjusted to provide incorporation of U at a rate of the order of IU every 500 nucleotides added .

5 (3) Cleavage of U-containing DNA with Uracil DNA Glycosylase (UDG) .

After an appropriate wash, chips containing DNA can be incubated in 14 parts UDG Buffer (70 mM Hepes-KOH pH 8.0, 1 mM dithiothreitol, 1 mM EDTA) , 1 part uracil DNA glycosylase
10 (BRL) . After e.g. two hours of incubation at 37⁰C the chips are optionally washed with a high pH wash solution and heated to 94⁰C for up to 10 min to kill the enzyme. This step may not be necessary for a chip. They are then subjected to appropriate washes in an appropriate buffer and
15 made ready for the next step.

(4) On-chip ligation.

Hairpin DNA on the surface at a density to allow appropriate single molecule array analysis (10^7 - 10^9
20 molecules per square cm) and double -stranded genomic DNA are mixed together. The molecules to be ligated must contain the appropriate phosphate or hydroxyl termini as described above. The DNA molecules are incubated in the presence of a 1:2 dilution of "2X Quick Ligation Reaction Buffer" (NEB =
25 New England Biolabs, Beverly, MA) and a 1:20 dilution of NEB "Quick T4 DNA Ligase", both components of the NEB kit called "Quick Ligation Kit", at room temperature for approximately 1 h. Subsequently, washing is performed in a suitable wash reagent (s) or buffer(s), using suitable times and
30 temperatures.

(5) Nicking reaction.

- 35 -

Nicking of DNA on a surface via specific recognition sequences in the known part of the DNA is described here. The surface equivalent of roughly a solution amount of 1.25 pmoles of DNA is digested at 55°C for 30 minutes with

5 N.BstNBI (50 Units/mL final concentration) in its supplied buffer (NEBuffer N.BstNBI at "IX" final concentration) from the manufacturer (NEB). This experiment can also be performed with little modification in a flow-cell where the substrate comprises DNA ligated to DNA hairpins that are

10 covalently attached to the glass surface of the flow-cell. In this case, the attachment of the DNA to a solid support, the glass, obviates the need to employ a DNA purification step between enzyme steps,- instead, products can be removed and new reagents added by flowing solutions through the

15 cell. Suitable wash buffers as known in the art can be employed.

It will be appreciated that nicking can also be accomplished by blocking one side of a standard restriction

20 enzyme cleavage site (in the hairpin oligonucleotide sequence or sequences) using methods familiar to those skilled in the art, e.g., by using thiophosphate linkages in one side of the restriction enzyme's recognition site, to prevent cutting in that side, but not in the other.

25

(6) Strand-displacement polymerisation.

Strand-displacement polymerisation is carried out using a suitable strand-displacement polymerase (e.g., BST Polymerase). The reaction conditions are, e.g., "IX

30 Thermopol buffer" (final concentrations: 20 mM Tris-HCl pH 8.8 measured @ 25°C, 10mM KCl, 10mM (NH₄)₂SO₄, 2mM MgSO₄, 0.1% Triton X-100), all four dNTPs at a concentration of

- 36 -

between 1 and 200 micromolar, 160 Units/mL (final concentration) of BST large fragment polymerase (NEB). Reactions are performed, for 2 hrs. at 60⁰c, for example, followed by washing in a suitable wash buffer, using
5 suitable times and temperatures .

(7) Denaturation.

Denaturation may be performed by any of the methods known to those skilled in the art. Typically there are
10 three preferred methods:

- Chips are washed in very low ionic strength or distilled water.
- Chips are incubated at high temperature (80-90⁰c, typically) for a few minutes in an appropriate buffer
15 or at 40-70⁰C in 50% deionized formamide dissolved in water .
- Chips are incubated at high pH briefly, i.e. pH greater than 10. This latter method is least preferred as it may be damaging to the surface chemistry of the chip.

20

(8) Phosphatasing and kinasing.

See e.g., Maniatis, T., E.F. Fritsch and J. Sambrook. 1982. Molecular Cloning: A Laboratory Manual. Cold Spring
25 Harbor Laboratory, Cold Spring Harbor, New York.

Claims

1. A method for pairwise sequencing of a polynucleotide template, the method comprising:

5

(a) providing a polynucleotide template having a first free 3' hydroxyl group which is positioned to initiate sequencing of a first region for sequencing determination on a first template strand,

10

(b) carrying out a first sequencing reaction by sequential addition of nucleotides to the first free 3' hydroxyl group to determine the sequence of the first template strand in the first region,

15

(c) generating a second free 3' hydroxyl group within the polynucleotide template at a nucleotide position which is spaced apart from the last nucleotide added in the sequencing reaction of step (b), wherein the second free 3' hydroxyl group is positioned to initiate sequencing of a second region for sequencing determination on a second template strand, and wherein the second free hydroxyl group is in continuous covalent linkage with the last nucleotide added in the sequencing reaction of step (b), and

20

(d) carrying out a second sequencing reaction by sequential addition of nucleotides to the second free 3' hydroxyl group to determine the sequence of the second template strand in the second region.

25

30

- 38 -

2. The method of claim 1 wherein the 3' end of the first template strand is linked to the 5' end of a self-complementary hairpin linker polynucleotide and the 3' end of the linker polynucleotide forms the first free 3' hydroxyl group of part (a).

3. The method of claim 1 or claim 2 wherein the polynucleotide template or linker polynucleotide is linked to a solid support.

4. The method of claim 3 wherein the polynucleotide template forms part of an array.

5. The method according to claim 4 wherein the addressable array is a single molecule array.

6. The method according to claim 4 wherein the addressable array is a clustered array.

7. The method of any one of claims 1 to 6 wherein the first and second regions for sequence determination are in the same strand of the polynucleotide template.

8. The method of claim 7 wherein the second free 3' hydroxyl group is generated by sequential addition of one or more nucleotides to the 3' end of the last nucleotide added in the sequencing reaction of part (b) without determining the identity of the nucleotides to produce an extended complementary strand, wherein the last such nucleotide added provides the second free 3' hydroxyl group which is spaced apart from but in continuous covalent linkage with the last nucleotide added in the sequencing reaction of part (b).

- 39 -

9. The method of claim 8 wherein at least 50, preferably at least 100, and more preferably at least 500 nucleotides are added to the last nucleotide added in the sequencing
5 reaction of part (b) .

10. The method of claim 7 wherein the second free 3' hydroxyl group is generated by sequential addition of nucleotides including dUTP to the 3' end of the last
10 nucleotide added in the sequencing reaction of part (b) to produce an extended complementary strand incorporating one or more uracil bases, followed by treatment with an enzyme which hydrolyses uracil glycosidic bonds, thereby generating a second free 3' hydroxyl group.

15

11. The method of any one of claims 1 to 6 wherein the first and second regions for sequence determination are in complementary strands of the polynucleotide template.

20 12. The method of claim 11 wherein the second free 3' hydroxyl group is generated by (i) removing or blocking any 5' phosphate group present on the first template strand such that it is not capable of being ligated to a free 3' hydroxyl group, (ii) sequentially adding nucleotides to the
25 3' end of the last nucleotide added in the sequencing reaction of part (b) to produce a second template strand complementary to the full length of the first template strand, thereby generating a double-stranded blunt ended polynucleotide, (iii) ligating a self-complementary hairpin
30 polynucleotide having a free 3' hydroxyl group to the free 3' hydroxyl group of the second template strand, whereby the

- 40 -

free 3' hydroxyl group of the linker polynucleotide provides a second free 3' hydroxyl group.

13. The method of claim 12 wherein the second sequencing
5 reaction of part (d) is carried out using a strand-
displacing polymerase enzyme.

14. The method of claim 12 wherein prior to step (d) a
portion of the first template strand is removed.
10

15. The method of claim 14 wherein a portion of the first
template strand is removed by cleavage of the first template
strand with a nicking endonuclease followed by denaturation
of the resulting product to remove the portion of the first
15 template strand upstream of the cleavage site.

16. The method of claim 15 wherein the 3' end of the first
template strand is joined to the 5' end of one strand of a
self-complementary hairpin linker polynucleotide, which
20 linker polynucleotide comprises a recognition site for a
nicking endonuclease which directs cleavage by the nicking
endonuclease at a site before, at or beyond the 5' end of
the linker polynucleotide.

25 17. The method of claim 16 wherein the template is linked
to a solid support via the self-complementary hairpin linker
polynucleotide .

18. The method of claim 11 wherein the polynucleotide
30 template of step (a) is generated by:

(i) providing a fully double-stranded polynucleotide
consisting of a first template strand having a 5' phosphate

- 41 -

group and a complementary strand lacking a 5' phosphate group ;

(ii) ligating a second self-complementary hairpin linker polynucleotide having hydroxyl groups at both the 3' and 5' ends to a first end of the double-stranded polynucleotide such that the 3' end of the second linker is joined to the 5' end of the first template strand but the 5' end of the second linker remains free;

(iii) ligating a first self-complementary hairpin linker polynucleotide having a 5' phosphate group and a 3' hydroxyl group to the second end of the double-stranded polynucleotide such that the 5' end of the first linker is joined to the 3' end of the first template strand but the 3' end of the first linker remains free;

(iv) denaturing the product of step (iii) to remove the complementary strand of the double-stranded polynucleotide, thereby forming a polynucleotide template having a first free 3' hydroxyl group; and

(v) optionally restoring a 5' phosphate group to the 5' end of the second linker polynucleotide.

19. The method of claim 18 wherein following the first sequencing reaction of part (b) a second free 3' hydroxyl group is generated by:

(i) sequentially adding unmodified nucleotides to the 3' end of the last nucleotide added in the first sequencing reaction, thereby forming a second template strand complementary to the entire length of the first template strand;

(ii) ligating the 3' end of the second template strand to the 5' end of the second hairpin polynucleotide linker,

- 42 -

which linker has been treated to restore the 5' phosphate group, thereby forming a self-complementary closed circle, (iii) generating a second free 3' hydroxyl group by cleavage of the first template strand or the second hairpin linker polynucleotide ligated to the 5' end of the first template strand.

20. The method of claim 19 wherein step (iii) comprises cleaving the first template strand at two cleavage sites spaced apart from each other and then denaturing the product to remove the portion of the first template strand located between the two cleavage sites.

21. The method of claim 20 wherein the second linker polynucleotide includes a recognition site for a nicking endonuclease which directs cleavage by the nicking endonuclease at a cleavage site before, at or beyond the 3' end of the second linker.

22. The method of claim 20 or claim 21 wherein the first hairpin polynucleotide linker includes a recognition site for a nicking endonuclease, which directs cleavage by the nicking endonuclease at a site before, at or beyond the 5' end of the first linker.

23. The method of claim 19 wherein step (iii) comprises cleaving the first template strand at a site before, at or beyond the 3' end of the second linker and the second sequencing reaction is carried out using a strand displacement polymerase.

- 43 -

24. The method of any one of claims 18 to 23 wherein the first hairpin polynucleotide linker is linked to a solid support .

5 25. The method of claim 11 wherein the polynucleotide template of step (a) is generated by:

(i) providing a fully double-stranded polynucleotide consisting of a first template strand and a complementary second template strand;

10 (ii) ligating a first self-complementary linker polynucleotide to a first end of the double-stranded polynucleotide, the first linker polynucleotide having a recognition site for a first nicking endonuclease which directs cleavage by the nicking endonuclease at a site
15 before, at or beyond the 3' end of the first linker;

(iii) ligating a second self-complementary linker polynucleotide to a second end of the double-stranded polynucleotide, the second linker polynucleotide having a recognition site for a second nicking endonuclease which
20 directs cleavage by the nicking endonuclease at a site before, at or beyond the 3' end of the second linker, thereby producing a self-complementary closed circle;

(iv) cleaving one strand of the self-complementary closed circle produced in step (iii) with a first nicking
25 endonuclease at a site before, at or beyond the 3¹ end of the first linker, thereby generating a first free 3' hydroxyl group .

26. The method of claim 25 wherein the first sequencing
30 reaction of step (b) is carried out using a strand displacement polymerase enzyme .

- 44 -

27. The method of claim 26 wherein the second free 3' hydroxyl group is generated by nicking one strand of the self-complementary closed circle produced in step (iii) with a second nicking endonuclease at a site before, at or beyond
5 the 3' end of the second linker, thereby generating a second free 3' hydroxyl group .

28. The method of claim 27 wherein the second sequencing reaction of step (d) is carried out using a strand
10 displacement enzyme.

29. The method of claim 11 wherein the polynucleotide template of step (a) is generated by:

- (i) providing a fully double-stranded polynucleotide
15 consisting of a first template strand and a complementary second template strand;
- (ii) ligating a first self-complementary linker polynucleotide to a first end of the double-stranded polynucleotide, the first linker polynucleotide having a
20 recognition site for a first nicking endonuclease which directs cleavage by the nicking endonuclease at a site before, at or beyond the 3' end of the first linker,-
- (iii) ligating a second self-complementary linker polynucleotide to a second end of the double-stranded
25 polynucleotide, the second linker polynucleotide having a recognition site for a second nicking endonuclease which directs cleavage by the nicking endonuclease at a site before, at or beyond the 5' end of the second linker;
- (iv) cleaving one template strand of the ligated product of
30 step (iii) with a first nicking endonuclease at a site before, at or beyond the 3' end of the first linker and cleaving the same strand with a second nicking endonuclease

- 45 -

at a second site before, at or beyond the 5' end of the second linker,-

(v) denaturing the product of step (iv) to remove the portion of the template strand cleaved in step (iv) located
5 between the two cleavage sites, thereby generating a polynucleotide template having a first free 3' hydroxyl group .

30. The method of claim 29 wherein following the first
10 sequencing reaction a second free 3' hydroxyl group is generated by:

(vi) sequentially adding unmodified nucleotides to the 3' end of the last nucleotide added in the first sequencing reaction of step (b) , thereby forming a second template
15 strand complementary to the entire length of the first template strand;

(vii) ligating the 3' end of the second template strand to the 5' end of the second polynucleotide linker;

(viii) generating a second free 3' hydroxyl group by
20 cleavage of the first template strand or the second linker polynucleotide ligated to the 5' end of the first template strand.

31. The method of claim 30 wherein step (viii) comprises
25 cleaving the first template strand at two cleavage sites spaced apart from each other and then denaturing the product to remove the portion of the first template strand located between the two cleavage sites.

30 32. The method of claim 30 or claim 31 wherein the second polynucleotide linker includes a recognition site for a nicking endonuclease which directs cleavage by the nicking

- 46 -

endonuclease at a cleavage site before, at or beyond the 3' end of the second linker.

33. The method of claim 31 or claim 32 wherein the first
5 polynucleotide linker includes a recognition site for a
nicking endonuclease which directs cleavage by the nicking
endonuclease at a site before, at or beyond the 5' end of
the first linker.

10 34. The method of claim 32 wherein the second sequencing
reaction is carried out using a strand displacement
polymerase .




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6