

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 2013221585 B2

(54) Title
Antibodies to integrin alphavbeta6 and use of same to treat cancer

(51) International Patent Classification(s)
C07K 16/00 (2006.01) **A61K 39/395** (2006.01)
A61K 39/00 (2006.01)

(21) Application No: **2013221585** **(22) Date of Filing:** **2013.02.14**

(87) WIPO No: **WO13/123152**

(30) Priority Data

(31) Number	(32) Date	(33) Country
61/602,511	2012.02.23	US
61/600,499	2012.02.17	US

(43) Publication Date: **2013.08.22**

(44) Accepted Journal Date: **2017.03.30**

(71) Applicant(s)
Seattle Genetics, Inc.

(72) Inventor(s)
Ryan, Maureen;Sussman, Django

(74) Agent / Attorney
Spruson & Ferguson, L 35 St Martins Tower 31 Market St, Sydney, NSW, 2000, AU

(56) Related Art
US 7,465,449 B2 (VIOLETTE et al) 16 December 2008
WO 2006/041934 A2 (NEURALAB LIMITED et al) 20 April 2006

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(10) International Publication Number

WO 2013/123152 A2

(43) International Publication Date
22 August 2013 (22.08.2013)

(51) International Patent Classification:

A61K 39/395 (2006.01)

(21) International Application Number:

PCT/US2013/026087

(22) International Filing Date:

14 February 2013 (14.02.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/600,499 17 February 2012 (17.02.2012) US
61/602,511 23 February 2012 (23.02.2012) US

(71) Applicant: SEATTLE GENETICS, INC. [US/US];
21823 30th Drive Southeast, Bothell, Washington 98021
(US).

(72) Inventors: RYAN, Maureen; Seattle Genetics, Inc.,
21823 30th Drive Southeast, Bothell, Washington 98021
(US). SUSSMAN, Django; Seattle Genetics, Inc., 21823
30th Drive Southeast, Bothell, Washington 98021 (US).

(74) Agents: ABERMAN, Leslie et al.; Seattle Genetics, Inc.,
21823 30th Drive Southeast, Bothell, Washington 98021
(US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,

DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: ANTIBODIES TO INTEGRIN α V β 6 AND USE OF SAME TO TREAT CANCER




Figure 12

(57) Abstract: The invention provides antibodies that specifically bind to integrin α V β 6. The antibodies are useful for treatment and diagnoses of various cancers as well as detecting α V β 6.

WO 2013/123152 A2

Antibodies to Integrin $\alpha v\beta 6$ and Use of Same to Treat Cancer

[0001] This application claims the benefit of U.S. Provisional App. No. 61/600,499 filed February 17, 2012 and U.S. Provisional App. No. 61/602,511 filed February 23, 2012, each of which is incorporated by reference in its entirety for all purposes.

BRIEF SUMMARY

[0002] Provided herein are monoclonal antibodies that specifically bind to integrin $\alpha v\beta 6$, and methods of using them to treat cancer. The antibodies described herein include 15H3 antibodies. The 15H3 antibody is a murine monoclonal antibody comprising a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:1 and a light chain variable region comprising the amino acid sequence of SEQ ID NO:2. Chimeric and humanized forms of the 15H3 murine antibody are encompassed by the present invention, as well as human forms thereof. The chimeric, humanized, and human forms of the 15H3 antibody comprise substantially the same complementarity determining regions (CDRs) as the murine 15H3 antibody and specifically bind to integrin $\alpha v\beta 6$. The three heavy chain variable region CDRs of the murine 15H3 antibody have the sequences as provided in SEQ ID NO:3(H-CDR1), SEQ ID NO:4(H-CDR2), and SEQ ID NO:5 (H-CDR3). The three light chain CDRs of the murine 15H3 antibody have the sequences as provided in SEQ ID NO:6(L-CDR1), SEQ ID NO:7(L-CDR2), and SEQ ID NO:8 (L-CDR3). Optionally, in a humanized antibody comprising substantially the same complementarity determining regions (CDRs) as the murine 15H3 antibody that specifically binds to $\alpha v\beta 6$, at least one of positions H28, H30, H72, H73, H75, H78, or H93 of the heavy chain variable region is occupied by the amino acid residue from the corresponding position of the murine 15H3 antibody. Optionally in any such antibody, at least two of positions H28, H30, H72, H73, H75, H78, or H93 are occupied by the amino acid residue from the corresponding position of the murine 15H3 antibody. Optionally in any such antibody, at least four of positions H28, H30, H72, H73, H75, H78, or H93

are occupied by the amino acid residue from the corresponding position of the murine 15H3 antibody. Optionally in any such antibody, at least one of positions H28 and H30 are occupied by S. Optionally, in any such antibody, positions H28 and H30 are occupied by S. Optionally, in any such antibody, positions H28 and H30 are occupied by S; position H72 is occupied by D or Q; position H73 is occupied by T or K, position H75 is occupied by T or S; position H78 is occupied by V or A, and position 93 is occupied by A or V. Optionally in any of these embodiments, not more than 10, not more than 9, not more than 8, not more than 7, not more than 6, or not more than 5 of the amino acid residues in the heavy chain framework region are occupied by the amino acid residue from the corresponding position of the murine 15H3 antibody. Optionally, in any of these embodiments, the CDRs of the heavy chain variable region are those of SEQ ID NO:1 and the CDRs of the light chain variable region are those of SEQ ID NO:2. Optionally, in any of these embodiments, the CDRs of the heavy chain variable region are substantially those of SEQ ID NO:1 and the CDRs of the light chain variable region are substantially those of SEQ ID NO:2. Optionally, in any of these embodiments, the CDRs of the heavy chain variable region are those of SEQ ID NO:1 and the CDRs of the light chain variable region are those of SEQ ID NO:2 having 0, 1, 2 or 3 conservative amino acid substitutions in each CDR. In one aspect, provided herein are chimeric or humanized antibodies that specifically bind integrin $\alpha v\beta 6$ comprising heavy chain complementary determining region (CDR) sequences as set forth in SEQ ID NO:3 (CDR1), SEQ ID NO:4 (CDR2), and SEQ ID NO:5 (CDR3), and light chain CDR sequences as set forth in SEQ ID NO:6 (CDR4), SEQ ID NO:7 (CDR5), and SEQ ID NO:8 (CDR6) and having 0, 1, 2 or 3 conservative amino acid substitutions in each CDR. Here, as elsewhere in this application, Kabat numbering is used to describe positions in the heavy and light chain variable regions.

[0003] The invention also provides antibodies (e.g., humanized antibodies) comprising a heavy chain variable region having an amino acid sequence at least 90% identical to HA (SEQ ID NO:9) and having a light chain variable region at least 90% identical to LA (SEQ ID NO:10). In some aspects, the invention provides antibodies comprising a heavy chain variable region having an amino acid sequence at least 90% identical to HA (SEQ ID NO:9) having at least one murine 15H3 backmutation in the heavy chain

variable framework region and a light chain variable region at least 90% identical to LA (SEQ ID NO:10). By the phrase “having at least one murine 15H3 backmutation in the variable chain framework region,” it is meant that at least one of the amino acid residues in the acceptor variable chain framework region is changed to an amino acid residue that is present at the corresponding position in the donor murine 15H3 antibody. Optionally in any such antibody, at least one of positions H28, H30, H72, H73, H75, H78, or H93 is occupied by the amino acid residue from the corresponding position of the murine 15H3 antibody. Optionally in any such antibody, at least two of positions H28, H30, H72, H73, H75, H78, or H93 are occupied by the amino acid residue from the corresponding position of the murine 15H3 antibody. Optionally in any such antibody, at least four of positions H28, H30, H72, H73, H75, H78, or H93 are occupied by the amino acid residue from the corresponding position of the murine 15H3 antibody. Optionally in any such antibody, all of positions H28, H30, H72, H73, H75, H78, or H93 are occupied by the amino acid residue from the corresponding position of the murine 15H3 antibody.

Optionally in any such antibody, at least one of positions H28 and H30 is occupied by S. Optionally, in any such antibody, positions H28 and H30 are occupied by S. Optionally, in any such antibody, positions H28 and H30 are occupied by S; position H72 is occupied by D or Q; position H73 is occupied by T or K, position H75 is occupied by T or S; position H78 is occupied by V or A, and position 93 is occupied by A or V. Optionally in any of these embodiments, not more than 10, not more than 9, not more than 8, not more than 7, not more than 6, or not more than 5 of the amino acid residues in the heavy chain framework region are occupied by the amino acid residue from the corresponding position of the murine 15H3 antibody. Optionally, in any of these embodiments, the CDRs of the heavy chain variable region are those of SEQ ID NO:1 and the CDRs of the light chain variable region are those of SEQ ID NO:2. Optionally, in any of these embodiments, the CDRs of the heavy chain variable region are substantially those of SEQ ID NO:1 and the CDRs of the light chain variable region are substantially those of SEQ ID NO:2. Optionally, in any of these embodiments, the CDRs of the heavy chain variable region are those of SEQ ID NO:1 and the CDRs of the light chain variable region are those of SEQ ID NO:2 having 0, 1, 2 or 3 conservative amino acid substitutions in each CDR.

[0004] The invention also provides antibodies (e.g., humanized antibodies) comprising a heavy chain variable region having an amino acid sequence at least 90% identical to HB (SEQ ID NO:11), HF (SEQ ID NO:12), HG (SEQ ID NO:13), HK (SEQ ID NO:14), HL (SEQ ID NO:15), HM (SEQ ID NO:16), HN (SEQ ID NO:17), HO (SEQ ID NO:18), HQ (SEQ ID NO:19), HR (SEQ ID NO:20), HS (SEQ ID NO:21), HT (SEQ ID NO:22), or HU (SEQ ID NO:23) and a light chain variable region at least 90% identical to LA (SEQ ID NO:10), LB (SEQ ID NO:24), LC (SEQ ID NO:25), LD (SEQ ID NO:26), LE (SEQ ID NO:27), LF (SEQ ID NO:28) or LG (SEQ ID NO:29), and any combinations thereof (i.e., the antibody can comprise any one of the heavy chain variable regions paired with any one of the light chain variable regions). Optionally, the antibody comprises a heavy chain variable region having an amino acid sequence at least 95% identical to HA, HB, HF, HG, HK, HL, HM, HN, HO, HQ, HR, HS, HT, HU and a light chain variable region at least 95% identical to LA, LB, LC, LD, LE, LF, or LG , and any combinations thereof (i.e., the antibody can comprise any one of the heavy chain variable regions paired with any one of the light chain variable regions). Optionally in any such antibody, at least one of positions H28, H30, H72, H73, H75, H78, or H93 is occupied by the amino acid residue from the corresponding position of the murine 15H3 antibody. Optionally in any such antibody, at least two of positions H28, H30, H72, H73, H75, H78, or H93 is occupied by the amino acid residue from the corresponding position of the murine 15H3 antibody. Optionally, in any such antibody, at least one of positions H28 and H30 are occupied by S. Optionally, in any such antibody, positions H28 and H30 are occupied by S. Optionally, in any such antibody, positions H28 and H30 are occupied by S; position H72 is occupied by D or Q; position H73 is occupied by T or K, position H75 is occupied by T or S; position H78 is occupied by V or A, and position 93 is occupied by A or V. Optionally in any of these embodiments, not more than 10, not more than 9, not more than 8, not more than 7, not more than 6, or not more than 5 of the amino acid residues in the heavy chain framework region are occupied by the amino acid residue from the corresponding position of the murine 15H3 antibody. Optionally, in any of these embodiments, the CDRs of the heavy chain variable region are those of SEQ ID NO:1 and the CDRs of the light chain variable region are those of SEQ ID NO:2. Optionally, in any of these embodiments, the CDRs of the heavy chain variable region

are substantially those of SEQ ID NO:1 and the CDRs of the light chain variable region are substantially those of SEQ ID NO:2. Optionally, in any of these embodiments, the CDRs of the heavy chain variable region are those of SEQ ID NO:1 and the CDRs of the light chain variable region are those of SEQ ID NO:2 having 0, 1, 2 or 3 conservative amino acid substitutions in each CDR. The heavy chain variable region and light chain variable region of the antibodies described herein are optionally fused to a signal peptide. An exemplary heavy chain murine signal peptide is set forth in SEQ ID NO:33; an exemplary heavy chain human signal peptide is set forth in SEQ ID NO:34; an exemplary light chain murine signal peptide is set forth in SEQ ID NO:35; and an exemplary light chain human signal peptide is set forth in SEQ ID NO:36.

[0005] The heavy chain variable region of the antibodies described herein are optionally fused to a heavy chain constant region and the light chain variable region is optionally fused to a light chain constant region. The heavy chain constant region can be a natural human constant region or can be a mutant form of natural human constant region, e.g., one which has reduced binding to an Fc gamma receptor relative to the natural human constant region. Optionally, the heavy chain constant region is of IgG1 isotype. Optionally, the heavy chain constant region has an amino acid sequence comprising SEQ ID NO:30 and the light chain constant region has an amino acid sequence comprising SEQ ID NO:31. Optionally, the heavy chain constant region has an amino acid sequence comprising SEQ ID NO:32 and the light chain constant region has an amino acid sequence comprising SEQ ID NO:31. Optionally, the antibody is conjugated to a cytotoxic agent (*i.e.*, antibody-drug conjugate). Certain antibodies of the present invention have an affinity for human $\alpha v\beta 6$ that is greater than the murine version of the 15H3 antibody. Some other antibodies have an affinity for human $\alpha v\beta 6$ that is no more than 10 times lower, preferably not more than 6 times lower, more preferably no more than 5 times lower, even more preferably no more than 2 times lower than the murine version of the 15H3 antibody as determined by a competition binding assay or saturation binding assay (such as the ones described in the examples). In some aspects, humanized antibodies described herein have an affinity for human $\alpha v\beta 6$ that is within about 10 fold, preferably within about 5 fold, more preferably within about 2 fold, the affinity of the murine 15H3 antibody for human $\alpha v\beta 6$. In some aspect, the humanized antibodies

described herein have an apparent dissociation constant (k_d) for human $\alpha v\beta 6$ within a range of 0.1 nM to 10 nM, preferably 0.1 nM to 5 nM, more preferably 0.1 nM to 2 nM, 0.5 nM to 2 nM or 0.5 nM to 1.5 nM.

[0006] Also provided are bispecific monoclonal antibodies comprising amino acid sequences described herein.

[0007] The invention further provides nucleic acids encoding a heavy chain variable region and/or a light chain variable region of any of the antibodies described above.

[0008] The invention further provides a method of treating a patient having cancer, comprising administering to the patient an effective regime of an antibody as described above. Preferably the antibody is conjugated to a cytotoxic agent. The cancer is one that expresses the $\alpha v\beta 6$ antigen. Optionally, the cancer is bladder cancer, head and neck cancer (including cancers of the lip, oral cavity, nasal cavity, paranasal sinuses, pharynx, and larynx), skin cancer, lung cancer, pancreatic cancer, uterine cancer, breast cancer (including triple negative breast cancer), cervical cancer, colon cancer, prostate cancer, ovarian cancer, gastric cancer, or liver cancer. Optionally the cancer is a squamous cell carcinoma or adenocarcinoma.

[0008a] In a particular aspect the invention relates to an isolated monoclonal antibody that specifically binds integrin $\alpha v\beta 6$ comprising heavy chain complementary determining region (CDR) sequences as set forth in SEQ ID NO:3 (CDR1), SEQ ID NO:4 (CDR2), and SEQ ID NO:5 (CDR3), and light chain CDR sequences as set forth in SEQ ID NO:6 (CDR1), SEQ ID NO:7 (CDR2), and SEQ ID NO:8 (CDR3).

BRIEF DESCRIPTION OF THE FIGURES

[0009] **Figure 1** shows an alignment of the amino acid sequences of the parental murine mAb (referred to as 15H3) with select humanized 15H3 heavy (upper two panels) and light chain variable (lower two panels) regions.

[0010] **Figure 2** shows the results of competition binding studies on 293F cells expressing human $\alpha v\beta 6$ with antibodies having the HB heavy chain and the parental murine antibody (referred to as m15H3).

[0011] **Figures 3A-C** show the results of saturation binding studies on 293F cells and FDC-P1 cells expressing $\alpha v\beta 6$ with antibodies HBLC and HTLC.

[0012] **Figure 4** shows the results of competition binding studies on 293F cells expressing $\alpha\beta\delta$ with antibodies having the HB, HL, and HN heavy chains and the parental murine antibody (referred to as m15H3).

[0013] **Figure 5** shows the results of competition binding studies on 293F cells expressing $\alpha v\beta 6$ with antibodies having the HB, HQ, and HR heavy chains and the parental murine antibody (referred to as m15H3).

[0014] **Figure 6** shows the results of competition binding studies on 293F cells expressing $\alpha v\beta 6$ with antibodies having the HM heavy chains and the parental murine antibody (referred to as m15H3).

[0015] **Figure 7** shows the results of competition binding studies on 293F cells expressing $\alpha v\beta 6$ with antibodies having the HB, HL, HS, HT, and HU heavy chains and the parental murine antibody (referred to as m15H3).

[0016] **Figure 8** shows an analysis of $\alpha v\beta 6$ protein expression by IHC on bladder cancer samples from a Tumor Micro Array.

[0017] **Figures 9A-9F** show that humanized 15H3 anti- $\alpha v\beta 6$ ADCs behaved similarly as the mouse parental in cytotoxicity assays.

[0018] **Figure 10** shows the results of a xenograft study of the Detroit 562 head and neck cancer cell line in nude mice. The dose is indicated on the figure. The first administration was when the tumor reached approximately 100 mm³.

[0019] **Figure 11** shows the results of a xenograft study of the HPAFII pancreatic cancer cell line in nude mice. The dose and time of administration indicated on the figure.

[0020] **Figure 12** shows the results of a xenograft study of the BxPC-3 pancreatic cancer cell line in nude mice. The dose and time of administration is indicated on the figure.

DETAILED DESCRIPTION

[0021] The invention provides, *inter alia*, monoclonal antibodies that specifically bind to $\alpha v\beta 6$ and conjugates thereof. The antibodies are useful for treatment and diagnoses of conditions associated with $\alpha v\beta 6$ expression (including various cancers) as well as for detecting $\alpha v\beta 6$. Antibodies that specifically bind to $\alpha v\beta 6$ specifically bind to the $\beta 6$ integrin subunit alone and/or the $\alpha v\beta 6$ integrin complex but not the αv subunit alone.

[0022] An “isolated” antibody refers to an antibody that has been identified and separated and/or recovered from components of its natural environment and/or an antibody that is recombinantly produced. A “purified antibody” is an antibody that is typically at least 50% w/w pure of interfering proteins and other contaminants arising from its production or purification but does not exclude the possibility that the antibody is combined with an excess of pharmaceutical acceptable carrier(s) or other vehicle intended to facilitate its use. Interfering proteins and other contaminants can include, for example, cellular components of the cells from which an antibody is isolated or recombinantly produced. Sometimes antibodies are at least 60%, 70%, 80%, 90%, 95 or 99% w/w pure of interfering proteins and contaminants from production or purification. The antibodies described herein, including murine, chimeric, humanized, and human antibodies can be provided in isolated and/or purified form.

[0023] The term “monoclonal antibody” as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally occurring mutations that may be present in minor amounts.. The modifier “monoclonal” indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler *et al.* (1975) *Nature* 256:495, or may be made by recombinant DNA methods (see, for example, U.S. Patent No. 4816567). The “monoclonal antibodies” may also be isolated from phage antibody libraries using the techniques described in Clackson *et al.* (1991) *Nature*, 352:624-628 and Marks *et al.* (1991) *J. Mol. Biol.*, 222:581-597, for example or may be made by other methods. The antibodies described herein are monoclonal antibodies.

[0024] Specific binding of a monoclonal antibody to its target antigen means an affinity of at least 10^6 , 10^7 , 10^8 , 10^9 , or 10^{10} M⁻¹ and is detectably higher in magnitude and distinguishable from non-specific binding occurring to at least one unrelated target. Specific binding can be the result of formation of bonds between particular functional groups or particular spatial fit (e.g., lock and key type) whereas nonspecific binding is

usually the result of van der Waals forces. Antibodies can be assayed for specific binding to $\alpha v \beta 6$ by known methods, such as for example, competitive and non-competitive immunoassay systems using techniques such as Western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, or immunoprecipitation assays.

[0025] The basic antibody structural unit of an intact antibody is a tetramer of subunits. Each tetramer includes two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" chain (about 50-70 kDa). The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. This variable region is initially expressed linked to a cleavable signal peptide. The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function.

[0026] Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, and define the antibody's isotype as IgG, IgM, IgA, IgD and IgE, respectively. Within light and heavy chains, the variable and constant regions are joined by a "J" region of about 12 or more amino acids, with the heavy chain also including a "D" region of about 10 or more amino acids. (See generally, *Fundamental Immunology* (Paul, W., ed., 2nd ed. Raven Press, N.Y., 1989, Ch. 7, incorporated by reference in its entirety for all purposes).

[0027] The variable regions of each light/heavy chain pair form the antibody binding site. Thus, an intact antibody has two binding sites. Except in bifunctional or bispecific antibodies, the two binding sites are the same. The chains all exhibit the same general structure of relatively conserved framework regions (FR) joined by three hypervariable regions, also called complementarity determining regions or CDRs. The CDRs from the two chains of each pair are aligned by the framework regions, enabling binding to a specific epitope. From N-terminal to C-terminal, both light and heavy chains comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is in accordance with the definitions of Kabat, *Sequences of Proteins of Immunological Interest* (National Institutes of Health, Bethesda, MD, 1987 and 1991), or Chothia & Lesk, *J. Mol. Biol.* 196:901-917 (1987); Chothia et al., *Nature* 342:878-883 (1989). Kabat also provides a widely used numbering convention (Kabat

numbering) in which corresponding residues between different heavy chains or between different light chains are assigned the same number.

[0028] The term “antibody” includes intact antibodies and antigen binding fragments thereof. Typically, antibody fragments compete with the intact antibody from which they were derived for specific binding to the target including separate heavy chains, light chains Fab, Fab', F(ab')₂, F(ab)c, diabodies, Dabs, nanobodies, and Fv. Fragments can be produced by recombinant DNA techniques, or by enzymatic or chemical separation of intact immunoglobulins. The term “antibody” also includes a diabody (homodimeric Fv fragment) or a minibody (V_L - V_H - C_H 3), a bispecific antibody or the like. A bispecific or bifunctional antibody is an artificial hybrid antibody having two different heavy/light chain pairs and two different binding sites (see, *e.g.*, Songsivilai and Lachmann, *Clin. Exp. Immunol.*, 79:315-321 (1990); Kostelny et al., *J. Immunol.*, 148:1547-53 (1992)). The term “antibody” includes an antibody by itself (naked antibody) or an antibody conjugated to a cytotoxic agent.

[0029] The term “epitope” refers to a site on an antigen to which an antibody binds. An epitope can be formed from contiguous amino acids or noncontiguous amino acids juxtaposed by tertiary folding of one or more proteins. Epitopes formed from contiguous amino acids are typically retained on exposure to denaturing solvents whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents. An epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation. Methods of determining spatial conformation of epitopes include, for example, x-ray crystallography and 2-dimensional nuclear magnetic resonance. See, *e.g.*, *Epitope Mapping Protocols*, in *Methods in Molecular Biology*, Vol. 66, Glenn E. Morris, Ed. (1996).

[0030] Antibodies that recognize the same or overlapping epitopes can be identified in a simple immunoassay showing the ability of one antibody to compete with the binding of another antibody to a target antigen. The epitope of an antibody can also be defined by X-ray crystallography of the antibody bound to its antigen to identify contact residues. Alternatively, two antibodies have the same epitope if all amino acid mutations in the antigen that reduce or eliminate binding of one antibody reduce or eliminate binding of

the other. Two antibodies have overlapping epitopes if some amino acid mutations that reduce or eliminate binding of one antibody reduce or eliminate binding of the other.

[0031] Competition between antibodies is determined by an assay in which an antibody under test inhibits specific binding of a reference antibody to a common antigen (see, e.g., Junghans et al., *Cancer Res.* 50:1495, 1990). A test antibody competes with a reference antibody if an excess of a test antibody (e.g., at least 2x, 5x, 10x, 20x or 100x) inhibits binding of the reference antibody by at least 50% but preferably 75%, 90% or 99% as measured in a competitive binding assay. Antibodies identified by competition assay (competing antibodies) include antibodies binding to the same epitope as the reference antibody and antibodies binding to an adjacent epitope sufficiently proximal to the epitope bound by the reference antibody for steric hindrance to occur.

[0032] The term "patient" includes human and other mammalian subjects that receive either therapeutic treatment.

[0033] The terms "treatment" or "treat" refer to slowing, stopping, or reversing the progression of a $\alpha v \beta 6$ -expressing cancer in a patient, as evidenced by a decrease or elimination of a clinical or diagnostic symptom of the disease, by administration of an anti- $\alpha v \beta 6$ antibody or antibody-drug conjugate to the subject after the onset of the clinical or diagnostic symptom of the $\alpha v \beta 6$ -expressing cancer at any clinical stage. Treatment can include, for example, a decrease in the severity of a symptom, the number of symptoms, or frequency of relapse.

[0034] For purposes of classifying amino acids substitutions as conservative or nonconservative, the following amino acid substitutions are considered conservative substitutions: serine substituted by threonine, alanine, or asparagine; threonine substituted by proline or serine; asparagine substituted by aspartic acid, histidine, or serine; aspartic acid substituted by glutamic acid or asparagine; glutamic acid substituted by glutamine, lysine, or aspartic acid; glutamine substituted by arginine, lysine, or glutamic acid; histidine substituted by tyrosine or asparagine; arginine substituted by lysine or glutamine; methionine substituted by isoleucine, leucine or valine; isoleucine substituted by leucine, valine, or methionine; leucine substituted by valine, isoleucine, or methionine; phenylalanine substituted by tyrosine or tryptophan; tyrosine substituted by tryptophan, histidine, or phenylalanine; proline substituted by threonine; alanine substituted by

serine; lysine substituted by glutamic acid, glutamine, or arginine; valine substituted by methionine, isoleucine, or leucine; and tryptophan substituted by phenylalanine or tyrosine.

[0035] Percentage sequence identities are determined with antibody sequences maximally aligned by the Kabat numbering convention. After alignment, if a subject antibody region (e.g., the entire variable region of a heavy or light chain) is being compared with the same region of a reference antibody, the percentage sequence identity between the subject and reference antibody regions is the number of positions occupied by the same amino acid in both the subject and reference antibody region divided by the total number of aligned positions of the two regions, with gaps not counted, multiplied by 100 to convert to percentage.

[0036] Designation of a range of values includes all integers within or defining the range.

[0037] An antibody effector function refers to a function contributed by an Fc domain(s) of an Ig. Such functions can be, for example, antibody-dependent cellular cytotoxicity, antibody-dependent cellular phagocytosis or complement-dependent cytotoxicity. Such function can be effected by, for example, binding of an Fc effector domain(s) to an Fc receptor on an immune cell with phagocytic or lytic activity or by binding of an Fc effector domain(s) to components of the complement system. Typically, the effect(s) mediated by the Fc-binding cells or complement components result in inhibition and/or depletion of the $\alpha\text{v}\beta\text{6}$ targeted cell. Fc regions of antibodies can recruit Fc receptor (FcR)-expressing cells and juxtapose them with antibody-coated target cells. Cells expressing surface FcR for IgGs including Fc γ RIII (CD16), Fc γ RII (CD32) and Fc γ RI (CD64) can act as effector cells for the destruction of IgG-coated cells. Such effector cells include monocytes, macrophages, natural killer (NK) cells, neutrophils and eosinophils. Engagement of Fc γ R by IgG activates antibody-dependent cellular cytotoxicity (ADCC) or antibody-dependent cellular phagocytosis (ADCP). ADCC is mediated by CD16 $^{+}$ effector cells through the secretion of membrane pore-forming proteins and proteases, while phagocytosis is mediated by CD32 $^{+}$ and CD64 $^{+}$ effector cells (see *Fundamental Immunology*, 4th ed., Paul ed., Lippincott-Raven, N.Y., 1997, Chapters 3, 17 and 30; Uchida *et al.*, 2004, *J. Exp. Med.* 199:1659-69; Akewanlop *et al.*,

2001, *Cancer Res.* 61:4061-65; Watanabe *et al.*, 1999, *Breast Cancer Res. Treat.* 53:199-207). In addition to ADCC and ADCP, Fc regions of cell-bound antibodies can also activate the complement classical pathway to elicit complement-dependent cytotoxicity (CDC). C1q of the complement system binds to the Fc regions of antibodies when they are complexed with antigens. Binding of C1q to cell-bound antibodies can initiate a cascade of events involving the proteolytic activation of C4 and C2 to generate the C3 convertase. Cleavage of C3 to C3b by C3 convertase enables the activation of terminal complement components including C5b, C6, C7, C8 and C9. Collectively, these proteins form membrane-attack complex pores on the antibody-coated cells. These pores disrupt the cell membrane integrity, killing the target cell (see *Immunobiology*, 6th ed., Janeway *et al.*, Garland Science, N. Y., 2005, Chapter 2).

[0038] The term “antibody-dependent cellular cytotoxicity”, or ADCC, is a mechanism for inducing cell death that depends upon the interaction of antibody-coated target cells with immune cells possessing lytic activity (also referred to as effector cells). Such effector cells include natural killer cells, monocytes/macrophages and neutrophils. The effector cells attach to an Fc effector domain(s) of Ig bound to target cells via their antigen-combining sites. Death of the antibody-coated target cell occurs as a result of effector cell activity.

[0039] The term “antibody-dependent cellular phagocytosis”, or ADCP, refers to the process by which antibody-coated cells are internalized, either in whole or in part, by phagocytic immune cells (e.g., macrophages, neutrophils and dendritic cells) that bind to an Fc effector domain(s) of Ig.

[0040] The term “complement-dependent cytotoxicity”, or CDC, refers to a mechanism for inducing cell death in which an Fc effector domain(s) of a target-bound antibody activates a series of enzymatic reactions culminating in the formation of holes in the target cell membrane. Typically, antigen-antibody complexes such as those on antibody-coated target cells bind and activate complement component C1q which in turn activates the complement cascade leading to target cell death. Activation of complement may also result in deposition of complement components on the target cell surface that facilitate ADCC by binding complement receptors (e.g., CR3) on leukocytes.

[0041] A “cytotoxic effect” refers to the depletion, elimination and/or the killing of a target cell. A “cytostatic effect” refers to the inhibition of cell proliferation. A “cytotoxic agent” as used herein refers to an agent that has a cytotoxic and/or cytostatic effect on a cell. Cytotoxic agents can be conjugated to an antibody or administered in combination with an antibody.

[0042] The term “pharmaceutically acceptable” means approved or approvable by a regulatory agency of the Federal or a state government or listed in the U.S.

Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term “pharmaceutically compatible ingredient” refers to a pharmaceutically acceptable diluent, adjuvant, excipient, or vehicle with which an anti- α v β 6 antibody is delivered.

[0043] The phrase “pharmaceutically acceptable salt,” refers to pharmaceutically acceptable organic or inorganic salts of an anti- α v β 6 antibody or conjugate thereof or agent administered with an anti- α v β 6 antibody. Exemplary salts include sulfate, citrate, acetate, oxalate, chloride, bromide, iodide, nitrate, bisulfate, phosphate, acid phosphate, isonicotinate, lactate, salicylate, acid citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucuronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, p toluenesulfonate, and pamoate (i.e., 1,1' methylene bis -(2 hydroxy 3 naphthoate)) salts. A pharmaceutically acceptable salt may involve the inclusion of another molecule such as an acetate ion, a succinate ion or other counterion. The counterion may be any organic or inorganic moiety that stabilizes the charge on the parent compound. Furthermore, a pharmaceutically acceptable salt may have more than one charged atom in its structure. Instances where multiple charged atoms are part of the pharmaceutically acceptable salt can have multiple counter ions. Hence, a pharmaceutically acceptable salt can have one or more charged atoms and/or one or more counterion.

[0044] Unless otherwise apparent from the context, the term “about” encompasses values within a standard deviation of a stated value.

I. Target molecules

[0045] Unless otherwise indicated, $\alpha v\beta 6$ means human $\alpha v\beta 6$. An exemplary $\beta 6$ human sequence is assigned GenBank accession number AAA36122. An exemplary αv human sequence is assigned NCBI NP_002201.1

II. Antibodies of the invention

A. Binding specificity and functional properties

[0046] The invention provides the murine 15H3 antibody and chimeric, humanized, and human 15H3 antibodies.

[0047] The affinity of antibodies of the present invention (e.g., chimeric, humanized and human forms of the mouse 15H3 antibody) for human $\alpha v\beta 6$ is preferably equivalent to the affinity of mouse 15H3 antibody for human $\alpha v\beta 6$, greater than the affinity of mouse 15H3 antibody for human $\alpha v\beta 6$ or within a factor of ten, within a factor of five, or within a factor of two weaker than that of the murine antibody 15H3 for human $\alpha v\beta 6$ e.g., preferably not more than 10 times weaker, more preferably not more than 5 times weaker and even more preferably not more than 2 times weaker than the affinity of mouse 15H3 antibody for human $\alpha v\beta 6$. One method of measuring affinity of an antibody for its target antigen is by determining an antibody's apparent dissociation constant. The present invention encompasses antibodies (e.g., chimeric, humanized and human forms of the mouse 15H3 antibody) having an apparent dissociation constant that is essentially the same as that of murine 15H3 (i.e., within experimental error) as well as antibodies having an dissociation constant lower or higher than that of murine antibody 15H3 for human $\alpha v\beta 6$. In some embodiments, antibodies of the present invention (e.g., chimeric, humanized and human forms of the mouse 15H3 antibody) have an apparent dissociation constant within a range of 0.1 to 10 times, or preferably within a range of 0.1 to 5 times, 0.1 to 2 times, or even 0.5 to 2 times that of the apparent dissociation constant of the murine 15H3 antibody for human $\alpha v\beta 6$. In some aspects, the apparent dissociation constant (k_d) of the antibodies for human $\alpha v\beta 6$ is preferably within a range of 0.1 nM to 5 nM, even more preferably within a range of 0.1 nM to 5 nM, even preferably within a range of 0.1 nM to 2 nM or 0.5 nM to 1.5 nM. Chimeric, humanized and human 15H3

antibodies specifically bind to human $\alpha v\beta 6$ in native form and/or recombinantly expressed from CHO cells as does the mouse 15H3 antibody. Typically, chimeric, humanized and human 15H3 anti- $\alpha v\beta 6$ antibodies compete with murine 15H3 for binding to human $\alpha v\beta 6$.

[0048] Preferred antibodies inhibit cancer (e.g., growth of cells, metastasis and/or lethality to the organisms) alone (*i.e.*, as a naked antibody) or when conjugated to a cytotoxic agent as shown on cancerous cells propagating in culture, in an animal model or clinical trial. Animal models can be formed by implanting $\alpha v\beta 6$ expressing human tumor cell lines into appropriate immunodeficient rodent strains, *e.g.*, athymic nude mice or SCID mice. These tumor cell lines can be established in immunodeficient rodent hosts either as solid tumor by subcutaneous injections or as disseminated tumors by intravenous injections. Once established within a host, these tumor models can be applied to evaluate the therapeutic efficacies of the anti- $\alpha v\beta 6$ antibodies or conjugated forms thereof as described in the Examples.

B. Antibodies

[0049] A humanized antibody is a genetically engineered antibody in which the CDRs from a non-human “donor” antibody are grafted into human “acceptor” antibody sequences (see, *e.g.*, Queen, US 5,530,101 and 5,585,089; Winter, US 5,225,539; Carter, US 6,407,213; Adair, US 5,859,205; and Foote, US 6,881,557). The acceptor antibody sequences can be, for example, a human antibody sequence, a composite of such sequences, a consensus sequence of human antibody sequences, or a germline region sequence. A preferred acceptor sequence for the heavy chain is the germline V_H exon V_H 1-46 and for the J exon (J_H), exon J_H -4. For the light chain, a preferred acceptor sequence is exon VK 2-30 (also referred to in the literature as KV 2-30) and for the J exon J_{κ} -2. Alternative preferred acceptor sequences for the heavy chain include the germline V_H exons V_H 1-8 or VH 1-3 with the J exons (J_H), J_H -1, J_H -2, J_H -3, J_H -4, or J_H -5. Alternative preferred acceptor sequences for the light chain include exon VK 2-29 (also referred to in the literature as KV 2-30) with J exons J_{κ} -1, J_{κ} -2, J_{κ} -3, J_{κ} -4, or J_{κ} -5. Thus, a humanized antibody is an antibody having some or all CDRs entirely or substantially from a non-human donor antibody and variable region framework sequences and constant regions, if present, entirely or substantially from human antibody sequences.

Similarly a humanized heavy chain has at least one, two and usually all three CDRs entirely or substantially from a donor antibody heavy chain, and a heavy chain variable region framework sequence and heavy chain constant region, if present, substantially from human heavy chain variable region framework and constant region sequences. Similarly a humanized light chain has at least one, two and usually all three CDRs entirely or substantially from a donor antibody light chain, and a light chain variable region framework sequence and light chain constant region, if present, substantially from human light chain variable region framework and constant region sequences. In some aspects, a humanized antibody comprises a humanized heavy chain and a humanized light chain. A CDR in a humanized or human antibody is substantially from or substantially identical to a corresponding CDR in a non-human antibody when at least 60%, 85%, 90%, 95% or 100% of corresponding residues (as defined by Kabat) are identical between the respective CDRs. In some embodiments, a CDR in a humanized antibody or human antibody is substantially from or substantially identical to a corresponding CDR in a non-human antibody when there are no more than 3 conservative amino acid substitutions in each CDR. The variable region framework sequences of an antibody chain or the constant region of an antibody chain are substantially from a human variable region framework sequence or human constant region respectively when at least 70%, 80%, 85%, 90%, 95% or 100% of corresponding residues defined by Kabat are identical. In some humanized antibodies of the present invention, there is at least one murine 15H3 backmutation in the heavy chain variable framework region of the antibody.

[0050] Although humanized antibodies often incorporate all six CDRs (preferably as defined by Kabat) from a mouse antibody, they can also be made with less than all CDRs (e.g., at least 3, 4, or 5) CDRs from a mouse antibody (e.g., Pascalis *et al.*, *J. Immunol.* 169:3076, 2002; Vajdos *et al.*, *Journal of Molecular Biology*, 320: 415-428, 2002; Iwahashi *et al.*, *Mol. Immunol.* 36:1079-1091, 1999; Tamura *et al.*, *Journal of Immunology*, 164:1432-1441, 2000).

[0051] Certain amino acids from the human variable region framework residues can be selected for substitution based on their possible influence on CDR conformation and/or binding to antigen. Investigation of such possible influences is by modeling, examination

of the characteristics of the amino acids at particular locations, or empirical observation of the effects of substitution or mutagenesis of particular amino acids.

[0052] For example, when an amino acid differs between a murine variable region framework residue and a selected human variable region framework residue, the human framework amino acid can be substituted by the equivalent framework amino acid from the mouse antibody when it is reasonably expected that the amino acid:

- (1) noncovalently binds antigen directly,
- (2) is adjacent to a CDR region,
- (3) otherwise interacts with a CDR region (e.g. is within about 6 Å of a CDR region); or
- (4) mediates interaction between the heavy and light chains.

[0053] Although the 15H3 antibody was identified as a mouse antibody, the present application also encompasses human 15H3 antibodies. By the term, “human 15H3 antibody” is meant an antibody that is derived from human immunoglobulin gene sequences and that has CDRs that are substantially identical to those of murine 15H3 antibody and displays similar properties, *i.e.*, binding specificity to $\alpha\beta\beta$ 6. In some aspects, a human 15H3 antibody comprises a heavy chain variable region that is substantially identical to a heavy chain variable region described herein and/or a light chain variable region that is substantially identical to a light chain variable region described herein. In some embodiments, a 15H3 antibody of the present invention is not a human antibody, *e.g.*, a 15H3 antibody of the present invention is a murine, chimeric, or humanized antibody.

[0054] The invention provides 15H3 antibodies in which the heavy chain variable region shows at least 90% identity to HA (SEQ ID NO:9) and a light chain variable region at least 90% identical to LA (SEQ ID NO:10). In some aspects, the antibody is a humanized antibody and there is at least one murine 15H3 backmutation in the heavy chain variable framework region. Additionally, the invention provides 15H3 antibodies in which the humanized heavy chain variable region shows at least 90%, 95% or 99% sequence identity to SEQ ID NOS:11-23 and the humanized light chain variable region shows at least 90%, 95% or 99% sequence identity to SEQ ID NOS:10 and 24-29 (and any combinations thereof. The invention provides antibodies in which the heavy chain

variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NOS:11,12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, or 23, and the humanized light chain variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NO:10. In another embodiment, the invention provides antibodies in which the heavy chain variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NOS:11,12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, or 23, and the light chain variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NO:24. In another embodiment, the invention provides antibodies in which the heavy chain variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NOS:11,12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, or 23, and the light chain variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NO:25. In another embodiment, the invention provides antibodies in which the heavy chain variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NOS:11,12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, or 23, and the light chain variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NO:26. In another embodiment, the invention provides antibodies in which the heavy chain variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NOS:11,12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, or 23, and the light chain variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NO:27. In another embodiment, the invention provides antibodies in which the heavy chain variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NOS:11,12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, or 23, and the light chain variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NO:28. In another embodiment, the invention provides antibodies in which the heavy chain variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NOS:11,12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, or 23, and the light chain variable region shows at least 90%, 95% or 99% or 100% sequence identity to SEQ ID NO:29. In some aspects, these antibodies are humanized antibodies and some or all of the backmutations in the respective antibodies are retained. In other words, in such antibodies, preferably, at least one of positions H28, H30, H72, H73, H75, H78, or H93 is occupied by the amino acid from the corresponding position of the murine 15H3

antibody. Optionally, in any such antibody, at least one of positions H28 and H30 are occupied by S. Optionally, in any such antibody, positions H28 and H30 are occupied by S. Optionally, in any such antibody, positions H28 and H30 are occupied by S; position H72 is occupied by D or Q; position H73 is occupied by T or K, position H75 is occupied by T or S; position H78 is occupied by V or A, and position 93 is occupied by A or V. Preferably, in any of the antibodies described about, e.g., 15H3 humanized antibodies in which the heavy chain variable region shows at least 90%, 95% or 99% sequence identity to SEQ ID NOS:11-23 and the light chain variable region shows at least 90%, 95% or 99% sequence identity to SEQ ID NOS:10 and 24-29, the CDR regions are identical or substantially identical to the CDR regions of the mouse donor antibody, *i.e.*, murine 15H3 antibody (SEQ ID NO:3-8). The CDR regions are as defined by Kabat.

Antibodies of the present invention include antibodies HBLA, HBLB, HBLC, HBLD, HBLE, HBLF, HBLG, HLLA, HLLB, HLLC, HLLD, HLLE, HLLF, HLLG, HMLA, HMLB, HMLC, HMLD, HMLE, HMLF, HMLG, HNLA, HNLB, HNLC, HNLD, HNLE, HNLF, HNLG, HRLA, HRLB, HRLC, HRLD, HRLE, HRLF, HRLG, HSLA, HSLB, HSLC, HSLD, HSLE, HSLF, HSLG, HTLA, HTLB, HTLC, HTLD, HTLE, HTLF, HTLG, HULA, HULB, HULC, HULD, HULE, HULF, and HULG.

[0055] Another possible variation is to substitute certain residues in the CDRs with corresponding residues from human CDRs sequences, typically from the CDRs of the human acceptor sequences used in designing the exemplified humanized antibodies. In some antibodies only part of the CDRs, namely the subset of CDR residues required for binding, termed the SDRs, are needed to retain binding in a humanized antibody. CDR residues not contacting antigen and not in the SDRs can be identified based on previous studies (for example residues H60-H65 in CDR H2 are often not required), from regions of Kabat CDRs lying outside Chothia hypervariable loops (Chothia, *J. Mol. Biol.* 196:901, 1987), by molecular modeling and/or empirically, or as described in Gonzales et al., *Mol. Immunol.* 41: 863 (2004). In such humanized antibodies at positions in which one or more donor CDR residues is absent or in which an entire donor CDR is omitted, the amino acid occupying the position can be an amino acid occupying the corresponding position (by Kabat numbering) in the acceptor antibody sequence or a conservative substitution. The number of such substitutions of acceptor for donor amino acids in the

CDRs to include reflects a balance of competing considerations. Such substitutions are potentially advantageous in decreasing the number of mouse amino acids in a humanized antibody and consequently decreasing potential immunogenicity.

[0056] In preferred embodiments herein, the CDR regions of the humanized and human 15H3 antibodies are identical or substantially identical to the CDR regions of the murine 15H3 antibody (SEQ ID NO:3-8). In any of the antibodies described herein, the CDR regions can be those of the murine 15H3 antibody (i.e., SEQ ID NO:3-8) with 0, 1, 2 or 3 conservative amino acid substitutions. In any of the antibodies described herein, the CDR regions can be those of the murine 15H3 antibody (i.e., SEQ ID NO:3-8) with 0, 1, 2 or 3 conservative amino acid substitutions in each CDR. In any of the antibodies described herein, the CDR regions can be those of the murine 15H3 antibody (i.e., SEQ ID NO:3-8) with 0, 1, 2 or 3 conservative amino acid substitutions in CDR1 and CDR2 of the heavy chain and CDR1, CDR2 and optionally CDR3 of the light chain. By molecular modeling and empirical deduction, it has been determined that in any of the antibodies described herein, the conservative amino acid substitution can be, for example, at position 24, 25, 32, 33, and/or 34 in CDR1 of the light chain, positions 53-56 in CDR2 of the light chain, position 31 and/or 32 in CDR1 of the heavy chain, and/or positions 61-65 in CDR2 of the heavy chain. Additionally, in any of the antibodies described herein, the following three positions could be replaced by the corresponding human amino acid: lysine at position 64 in CDR2 of the heavy chain can be replaced by glutamine, lysine at position 24 in CDR1 of the light chain can be replaced by arginine, and leucine at position 54 in CDR2 of the light chain can be replaced by arginine.

C. Selection of Constant Region

[0057] The heavy and light chain variable regions of antibodies described herein can be linked to at least a portion of a human constant region. The choice of constant region depends, in part, whether antibody-dependent cell-mediated cytotoxicity, antibody dependent cellular phagocytosis and/or complement dependent cytotoxicity are desired. For example, human isotopes IgG1 and IgG3 have strong complement-dependent cytotoxicity, human isotype IgG2 weak complement-dependent cytotoxicity and human

IgG4 lacks complement-dependent cytotoxicity. Human IgG1 and IgG3 also induce stronger cell mediated effector functions than human IgG2 and IgG4. Light chain constant regions can be lambda or kappa. Antibodies can be expressed as tetramers containing two light and two heavy chains, as separate heavy chains, light chains, as Fab, Fab', F(ab')2, and Fv, or as single chain antibodies in which heavy and light chain variable domains are linked through a spacer.

[0058] Human constant regions show allotypic variation and isoallotypic variation between different individuals, that is, the constant regions can differ in different individuals at one or more polymorphic positions. Isoallotypes differ from allotypes in that sera recognizing an isoallotype binds to a non-polymorphic region of a one or more other isotypes.

[0059] One or several amino acids at the amino or carboxy terminus of the light and/or heavy chain, such as the C-terminal lysine of the heavy chain, may be missing or derivatized in a proportion or all of the molecules. Substitutions can be made in the constant regions to reduce or increase effector function such as complement-mediated cytotoxicity or ADCC (see, e.g., Winter et al., US Patent No. 5,624,821; Tso et al., US Patent No. 5,834,597; and Lazar et al., Proc. Natl. Acad. Sci. USA 103:4005, 2006), or to prolong half-life in humans (see, e.g., Hinton et al., J. Biol. Chem. 279:6213, 2004).

[0060] Exemplary substitution include the amino acid substitution of the native amino acid to a cysteine residue at amino acid position 234, 235, 237, 239, 267, 298, 299, 326, 330, or 332, preferably an S239C mutation (substitutions of the constant regions are according to the EU index) in a human IgG1 isotype (US 20100158909). The presence of an additional cysteine residue allows interchain disulfide bond formation. Such interchain disulfide bond formation can cause steric hindrance, thereby reducing the affinity of the Fc region-Fc γ R binding interaction. The cysteine residue(s) introduced in or in proximity to the Fc region of an IgG constant region can also serve as sites for conjugation to therapeutic agents (e.g., coupling cytotoxic drugs using thiol specific reagents such as maleimide derivatives of drugs). The presence of a therapeutic agent causes steric hindrance, thereby further reducing the affinity of the Fc region-Fc γ R binding interaction. Other substitutions at any of positions 234, 235, 236 and/or 237

reduce affinity for Fc γ receptors, particularly Fc γ RI receptor (see, e.g., US 6,624,821, US 5,624,821.)

[0061] The in vivo half-life of an antibody can also impact on its effector functions. The half-life of an antibody can be increased or decreased to modify its therapeutic activities. FcRn is a receptor that is structurally similar to MHC Class I antigen that non-covalently associates with β 2-microglobulin. FcRn regulates the catabolism of IgGs and their transcytosis across tissues (Ghetie and Ward, 2000, *Annu. Rev. Immunol.* 18:739-766; Ghetie and Ward, 2002, *Immunol. Res.* 25:97-113). The IgG-FcRn interaction takes place at pH 6.0 (pH of intracellular vesicles) but not at pH 7.4 (pH of blood); this interaction enables IgGs to be recycled back to the circulation (Ghetie and Ward, 2000, *Ann. Rev. Immunol.* 18:739-766; Ghetie and Ward, 2002, *Immunol. Res.* 25:97-113). The region on human IgG₁ involved in FcRn binding has been mapped (Shields *et al.*, 2001, *J. Biol. Chem.* 276:6591-604). Alanine substitutions at positions Pro238, Thr256, Thr307, Gln311, Asp312, Glu380, Glu382, or Asn434 of human IgG₁ enhance FcRn binding (Shields *et al.*, 2001, *J. Biol. Chem.* 276:6591-604). IgG₁ molecules harboring these substitutions have longer serum half-lives. Consequently, these modified IgG₁ molecules may be able to carry out their effector functions, and hence exert their therapeutic efficacies, over a longer period of time compared to unmodified IgG₁. Other exemplary substitutions for increasing binding to FcRn include a Gln at position 250 and/or a Leu at position 428. EU numbering is used for all position in the constant region.

[0062] Oligosaccharides covalently attached to the conserved Asn297 are involved in the ability of the Fc region of an IgG to bind Fc γ R (Lund *et al.*, 1996, *J. Immunol.* 157:4963-69; Wright and Morrison, 1997, *Trends Biotechnol.* 15:26-31). Engineering of this glycoform on IgG can significantly improve IgG-mediated ADCC. Addition of bisecting N-acetylglucosamine modifications (Umana *et al.*, 1999, *Nat. Biotechnol.* 17:176-180; Davies *et al.*, 2001, *Biotech. Bioeng.* 74:288-94) to this glycoform or removal of fucose (Shields *et al.*, 2002, *J. Biol. Chem.* 277:26733-40; Shinkawa *et al.*, 2003, *J. Biol. Chem.* 278:6591-604; Niwa *et al.*, 2004, *Cancer Res.* 64:2127-33) from this glycoform are two examples of IgG Fc engineering that improves the binding between IgG Fc and Fc γ R, thereby enhancing Ig-mediated ADCC activity.

[0063] A systemic substitution of solvent-exposed amino acids of human IgG1 Fc region has generated IgG variants with altered Fc γ R binding affinities (Shields *et al.*, 2001, *J. Biol. Chem.* 276:6591-604). When compared to parental IgG1, a subset of these variants involving substitutions at Thr256/Ser298, Ser298/Glu333, Ser298/Lys334, or Ser298/Glu333/Lys334 to Ala demonstrate increased in both binding affinity toward Fc γ R and ADCC activity (Shields *et al.*, 2001, *J. Biol. Chem.* 276:6591-604; Okazaki *et al.*, 2004, *J. Mol. Biol.* 336:1239-49).

[0064] Complement fixation activity of antibodies (both C1q binding and CDC activity) can be improved by substitutions at Lys326 and Glu333 (Idusogie *et al.*, 2001, *J. Immunol.* 166:2571-2575). The same substitutions on a human IgG2 backbone can convert an antibody isotype that binds poorly to C1q and is severely deficient in complement activation activity to one that can both bind C1q and mediate CDC (Idusogie *et al.*, 2001, *J. Immunol.* 166:2571-75). Several other methods have also been applied to improve complement fixation activity of antibodies. For example, the grafting of an 18-amino acid carboxyl-terminal tail piece of IgM to the carboxyl-termini of IgG greatly enhances their CDC activity. This is observed even with IgG4, which normally has no detectable CDC activity (Smith *et al.*, 1995, *J. Immunol.* 154:2226-36). Also, substituting Ser444 located close to the carboxy-terminal of IgG1 heavy chain with Cys induced tail-to-tail dimerization of IgG1 with a 200-fold increase of CDC activity over monomeric IgG1 (Shopes *et al.*, 1992, *J. Immunol.* 148:2918-22). In addition, a bispecific diabody construct with specificity for C1q also confers CDC activity (Kontermann *et al.*, 1997, *Nat. Biotech.* 15:629-31).

[0065] Complement activity can be reduced by mutating at least one of the amino acid residues 318, 320, and 322 of the heavy chain to a residue having a different side chain, such as Ala. Other alkyl-substituted non-ionic residues, such as Gly, Ile, Leu, or Val, or such aromatic non-polar residues as Phe, Tyr, Trp and Pro in place of any one of the three residues also reduce or abolish C1q binding. Ser, Thr, Cys, and Met can be used at residues 320 and 322, but not 318, to reduce or abolish C1q binding activity. Replacement of the 318 (Glu) residue by a polar residue may modify but not abolish C1q binding activity. Replacing residue 297 (Asn) with Ala results in removal of lytic activity but only slightly reduces (about three fold weaker) affinity for C1q. This alteration

destroys the glycosylation site and the presence of carbohydrate that is required for complement activation. Any other substitution at this site also destroys the glycosylation site. The following mutations and any combination thereof also reduce C1q binding: D270A, K322A, P329A, and P311S (see WO 06/036291).

[0066] Reference to a human constant region includes a constant region with any natural allotype or any permutation of residues occupying polymorphic positions in natural allotypes. Also, up to 1, 2, 5, or 10 mutations may be present relative to a natural human constant region, such as those indicated above to reduce Fcgamma receptor binding or increase binding to FcRN.

D. Expression of Antibodies

[0067] Chimeric or humanized antibodies are typically produced by recombinant expression. Recombinant polynucleotide constructs typically include an expression control sequence operably linked to the coding sequences of antibody chains, including naturally-associated or heterologous promoter regions. Preferably, the expression control sequences are eukaryotic promoter systems in vectors capable of transforming or transfecting eukaryotic host cells. Once the vector has been incorporated into the appropriate host, the host is maintained under conditions suitable for high level expression of the nucleotide sequences, and the collection and purification of the crossreacting antibodies.

[0068] Mammalian cells are a preferred host for expressing nucleotide segments encoding immunoglobulins or fragments thereof. See Winnacker, *From Genes to Clones*, (VCH Publishers, NY, 1987). A number of suitable host cell lines capable of secreting intact heterologous proteins have been developed in the art, and include CHO cell lines (e.g., DG44), various COS cell lines, HeLa cells, HEK293 cells, L cells, and non-antibody-producing myelomas including Sp2/0 and NS0. Preferably, the cells are nonhuman. Expression vectors for these cells can include expression control sequences, such as an origin of replication, a promoter, an enhancer (Queen et al., *Immunol. Rev.* 89:49 (1986)), and necessary processing information sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences. Preferred expression control sequences are promoters derived from endogenous genes,

cytomegalovirus, SV40, adenovirus, bovine papillomavirus, and the like. See Co et al., *J. Immunol.* 148:1149 (1992).

[0069] Human antibodies against $\alpha\beta\beta$ can be provided by a variety of techniques described below. Methods for producing human antibodies include the trioma method of Oestberg et al., *Hybridoma* 2:361-367 (1983); Oestberg, U.S. Patent No. 4,634,664; and Engleman et al., US Patent 4,634,666; use of transgenic mice including human immunoglobulin genes (see, e.g., Lonberg et al., WO93/12227 (1993); US 5,877,397, US 5,874,299, US 5,814,318, US 5,789,650, US 5,770,429, US 5,661,016, US 5,633,425, US 5,625,126, US 5,569,825, US 5,545,806, *Nature* 148, 1547-1553 (1994), *Nature Biotechnology* 14, 826 (1996), Kucherlapati, WO 91/10741 (1991) and phage display methods (see, e.g. Dower et al., WO 91/17271 and McCafferty et al., WO 92/01047, US 5,877,218, US 5,871,907, US 5,858,657, US 5,837,242, US 5,733,743 and US 5,565,332

[0070] Once expressed, antibodies can be purified according to standard procedures of the art, including HPLC purification, column chromatography, gel electrophoresis and the like (see generally, Scopes, *Protein Purification* (Springer-Verlag, NY, 1982)).

III. Nucleic Acids

[0071] The invention further provides nucleic acids encoding any of the heavy and light chains described above. Typically, the nucleic acids also encode a signal peptide fused to the heavy and light chains. Coding sequences on nucleic acids can be in operable linkage with regulatory sequences to ensure expression of the coding sequences, such as a promoter, enhancer, ribosome binding site, transcription termination signal and the like. The nucleic acids encoding heavy and light chains can occur in isolated form or can be cloned into one or more vectors. The nucleic acids can be synthesized by for example, solid state synthesis or PCR of overlapping oligonucleotides. Nucleic acids encoding heavy and light chains can be joined as one contiguous nucleic acid, e.g., within an expression vector, or can be separate, e.g., each cloned into its own expression vector. Exemplary nucleic acids are set forth in SEQ ID NO:37 (murine heavy chain), SEQ ID NO:38 (murine light chain), SEQ ID NO:39 (HA), SEQ ID NO:40 (HB), SEQ ID NO:41 (HL), SEQ ID NO:42 (HT), SEQ ID NO:43 (HN), SEQ ID NO:44 (HU), SEQ ID NO:45 (LA), SEQ ID NO:46 (LC), SEQ ID NO:47 (LF), SEQ ID NO:48 (LG), SEQ ID NO:49

(murine heavy chain signal peptide), SEQ ID NO:50 (human heavy chain signal peptide), SEQ ID NO:51 (murine light chain signal peptide), and SEQ ID NO:52 (human light chain signal peptide). Nucleic acids can be provided in isolated form. An isolated nucleic acid molecule is one that has been identified and separated and/or recovered from components of its natural environment or one that is non-naturally occurring.

IV. Antibody Drug Conjugates

[0072] Anti- α v β 6 antibodies can be conjugated to cytotoxic moieties (including pharmaceutically compatible salts thereof) to form an antibody drug conjugate (ADC). Particularly suitable moieties for conjugation to antibodies are cytotoxic agents (e.g., chemotherapeutic agents), prodrug converting enzymes, radioactive isotopes or compounds, or toxins (these moieties being collectively referred to as a therapeutic agent). For example, an anti- α v β 6 antibody can be conjugated to a cytotoxic agent such as a chemotherapeutic agent or a toxin (e.g., a cytostatic or cytocidal agent such as, e.g., abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin). Examples of useful classes of cytotoxic agents include, for example, DNA minor groove binders, DNA alkylating agents, and tubulin inhibitors. Exemplary cytotoxic agents include, for example, auristatins, camptothecins, duocarmycins, etoposides, maytansinoids (e.g., DM1, DM2, DM3, DM4), taxanes, benzodiazepines including benzodiazepine containing compounds (e.g., pyrrolo[1,4]benzodiazepines, indolinobenzodiazepines, and oxazolidinobenzodiazepines) and vinca alkaloids. Techniques for conjugating therapeutic agents to proteins, and in particular to antibodies, are well-known. (See, e.g., Alley et al., *Current Opinion in Chemical Biology* 2010 14:1-9; Senter, *Cancer J.*, 2008, 14(3):154-169.)

[0073] The therapeutic agent (e.g., cytotoxic agent) can be conjugated in a manner that reduces its activity unless it is cleaved off the antibody (e.g., by hydrolysis, by antibody degradation or by a cleaving agent). Such therapeutic agent can be attached to the antibody with a cleavable linker that is sensitive to cleavage in the intracellular environment of the α v β 6-expressing cancer cell but is not substantially sensitive to the extracellular environment, such that the conjugate is cleaved from the antibody when it is

internalized by the $\alpha v\beta 6$ -expressing cancer cell (e.g., in the endosomal or, for example by virtue of pH sensitivity or protease sensitivity, in the lysosomal environment or in the caveolar environment). Such therapeutic agent can also be attached to the antibody with a non-cleavable linker.


[0074] Typically the ADC comprises a linker region between the therapeutic agent and the anti- $\alpha v\beta 6$ antibody. As noted *supra*, the linker can be cleavable under intracellular conditions, such that cleavage of the linker releases the therapeutic agent from the antibody in the intracellular environment (e.g., within a lysosome or endosome or caveolea). The linker can be, *e.g.*, a peptidyl linker that is cleaved by an intracellular peptidase or protease enzyme, including a lysosomal or endosomal protease. In some aspects, the peptidyl linker is at least two amino acids long or at least three amino acids long. Cleaving agents can include cathepsins B and D and plasmin (see, *e.g.*, Dubowchik and Walker, 1999, *Pharm. Therapeutics* 83:67-123). Most typical are peptidyl linkers that are cleavable by enzymes that are present in $\alpha v\beta 6$ -expressing cells. For example, a peptidyl linker that is cleavable by the thiol-dependent protease cathepsin-B, which is highly expressed in cancerous tissue, can be used (e.g., a linker comprising a Phe-Leu or a Val-Cit peptide). The linker also can be a non-cleavable linker, such as, for example, an maleimido-alkylene- or maleimide-aryl linker that is directly attached to the therapeutic agent (e.g., a drug) and released by degradation of the antibody.


[0075] Typically, the linker is not substantially sensitive to the extracellular environment meaning that no more than about 20%, typically no more than about 15%, more typically no more than about 10%, and even more typically no more than about 5%, no more than about 3%, or no more than about 1% of the linkers in a sample of the ADC is cleaved when the ADC is present in an extracellular environment (e.g., in plasma). Whether a linker is not substantially sensitive to the extracellular environment can be determined, for example, by incubating independently with plasma both (a) the ADC (the “ADC sample”) and (b) an equal molar amount of unconjugated antibody or therapeutic agent (the “control sample”) for a predetermined time period (e.g., 2, 4, 8, 16, or 24 hours) and then comparing the amount of unconjugated antibody or therapeutic agent present in the ADC sample with that present in control sample, as measured, for example, by high performance liquid chromatography.

[0076] The linker can also promote cellular internalization. The linker can promote cellular internalization when conjugated to the therapeutic agent (*i.e.*, in the milieu of the linker-therapeutic agent moiety of the ADC as described herein). Alternatively, the linker can promote cellular internalization when conjugated to both the therapeutic agent and the anti- $\alpha v\beta 6$ antibody (*i.e.*, in the milieu of the ADC as described herein).

[0077] Exemplary antibody-drug conjugates include auristatin based antibody-drug conjugates meaning that the drug component is an auristatin drug. Auristatins bind tubulin, have been shown to interfere with microtubule dynamics and nuclear and cellular division, and have anticancer activity. Typically, the auristatin based antibody-drug conjugate comprises a linker between the auristatin drug and the anti- $\alpha v\beta 6$ antibody. The linker can be, for example, a cleavable linker (*e.g.*, a peptidyl linker) or a non-cleavable linker (*e.g.*, linker released by degradation of the antibody). The auristatin can be auristatin E or a derivative thereof. The auristatin can be, for example, an ester formed between auristatin E and a keto acid. For example, auristatin E can be reacted with paraacetyl benzoic acid or benzoylvaleric acid to produce AEB and AEVB, respectively. Other auristatins include MMAF and MMAE. The synthesis and structure of exemplary auristatins are described in U.S. Publication Nos. 7,659,241, 7,498,298, 2009-0111756, 2009-0018086, and 7,968,687 each of which is incorporated herein by reference in its entirety and for all purposes.

[0078] Exemplary auristatin based antibody drug conjugates include vcMMAE, vcMMAF and mcMMAF antibody drug conjugates as shown below wherein Ab is an antibody as described herein and val-cit represents the valine-citrulline dipeptide:





or a pharmaceutically acceptable salt thereof. The drug loading is represented by p, the number of drug-linker molecules per antibody. Depending on the context, p can represent the average number of drug-linker molecules per antibody, also referred to as the average drug loading. P ranges from 1 to 20 and is preferably from 1 to 8. In some preferred embodiments, when p represents the average drug loading, p ranges from about 2 to about 5. In some embodiments, p is about 2, about 3, about 4, or about 5. The average number of drugs per antibody in a preparation may be characterized by conventional means such as mass spectroscopy, HIC, ELISA assay, and HPLC. In some aspects, the anti- α v β 6 antibody is attached to the drug-linker through a cysteine residue of the antibody. In some aspects, the cysteine residue is one that is engineered into the antibody. In other aspects, the cysteine residue is an interchain disulfide cysteine residue.

V. Other Antibodies to $\alpha v \beta 6$

[0079] As well as the murine 15H3 antibody and chimeric, humanized, or human forms of the 15H3 antibody, other antibodies binding to an extracellular domain of $\alpha v \beta 6$ can be used in some of the methods of the invention, particularly the methods for the treatment of bladder cancer. Preferably, in such embodiments, the antibody is conjugated to a cytotoxic agent. A collection of anti- $\alpha v \beta 6$ antibodies are known, see, for example, U.S. Publication Nos. 20100330103 and 20110294982 and U.S. Patent Nos. 7,465,449;

7,943,742, and 6,692,741, each of which is incorporated by reference herein in its entirety and for all purposes.

[0080] Other antibodies to $\alpha v\beta 6$ can be made de novo by immunizing with $\alpha v\beta 6$ or one or more extracellular domains thereof. The production of other non-human monoclonal antibodies, e.g., murine, guinea pig, primate, rabbit or rat, against an immunogen can be performed by as described by Harlow & Lane, *Antibodies, A Laboratory Manual* (CSHP NY, 1988) (incorporated by reference for all purposes). Such an immunogen can be obtained from a natural source, by peptide synthesis or by recombinant expression.

[0081] Humanized, chimeric or veneered forms of non-human antibodies can be made. General methodology for producing humanized antibodies is described by Queen, US 5,530,101 and 5,585,089; Winter, US 5,225,539; Carter, US 6,407,213; Adair, US 5,859,205; and Foote, US 6,881,557). A chimeric antibody is an antibody in which the variable regions of light and heavy chains of a non-human antibody (e.g., a mouse) are combined with human light and heavy chain constant regions. Such antibodies substantially or entirely retain the binding specificity of the mouse antibody, and are about two-thirds human sequence. A veneered antibody is a type of humanized antibody that retains some and usually all of the CDRs and some of the non-human variable region framework residues of a non-human antibody but replaces other variable region framework residues that may contribute to B- or T-cell epitopes, for example exposed residues (Padlan, Mol. Immunol. 28:489, 1991) with residues from the corresponding positions of a human antibody sequence. The result is an antibody in which the CDRs are entirely or substantially from a non-human antibody and the variable region frameworks of the non-human antibody are made more human-like by the substitutions.

[0082] Human antibodies against $\alpha v\beta 6$ can be provided by a variety of techniques as described above.

[0083] If desired, any of the antibodies can be selected to have the same or overlapping epitope specificity as an exemplary antibody, such as the 15H3 antibody, by a competitive binding assay or otherwise.

VI. Therapeutic Applications

[0084] The antibodies of the invention, alone or as anti- $\alpha v\beta 6$ antibody drug conjugates, can be used to treat cancer. Some such cancers show detectable levels of $\alpha v\beta 6$ measured at either the protein (e.g., by immunoassay using one of the exemplified antibodies) or mRNA level. Some such cancers show elevated levels of $\alpha v\beta 6$ relative to noncancerous tissue of the same type, preferably from the same patient. An exemplary level of $\alpha v\beta 6$ on cancer cells amenable to treatment is 1000-200,000 $\alpha v\beta 6$ molecules per cell, although higher or lower levels can be treated. Generally, higher copy numbers are preferable for treatment of solid tumors, for example, at least about 15,000 $\alpha v\beta 6$ molecules per cell.

Optionally, a level of $\alpha v\beta 6$ in a cancer is measured before performing treatment.

[0085] Examples of cancers associated with $\alpha v\beta 6$ expression and amenable to treatment include, for example, bladder cancer, head and neck cancer (including, for example, cancers of the lip, oral cavity, nasal cavity, paranasal sinuses, pharynx, and larynx), skin cancer, lung cancer, pancreatic cancer, uterine cancer, breast cancer (including triple negative breast cancer), colon cancer, prostate cancer, ovarian cancer, stomach cancer, and liver cancer. In one aspect, squamous cell carcinomas and adenocarcinomas are examples of cancers associated with $\alpha v\beta 6$ expression and amenable to treatment. The treatment can be applied to patients having primary or metastatic tumors of these kinds. The treatment can also be applied to patients who are treatment naïve, who are refractory to conventional treatments (e.g., hormones, tamoxifen, herceptin), or who have relapsed following a response to such treatments. Anti- $\alpha v\beta 6$ antibodies and conjugates thereof can be used to treat cancers that express $\alpha v\beta 6$. In one embodiment, a human, humanized or chimeric anti- $\alpha v\beta 6$ antibody or conjugate thereof is used to treat a subject with a $\alpha v\beta 6$ -expressing bladder cancer, head and neck cancer (including, for example, cancers of the lip, oral cavity, nasal cavity, paranasal sinuses, pharynx, and larynx), skin cancer, lung cancer, pancreatic cancer, uterine cancer, breast cancer (including triple negative breast cancer), colon cancer, prostate cancer, ovarian cancer, stomach cancer, or liver cancer. In one embodiment, a human, humanized or chimeric anti- $\alpha v\beta 6$ antibody or conjugate thereof is used to treat a subject with an $\alpha v\beta 6$ -expressing squamous cell carcinoma or adenocarcinoma.

[0086] This application is believed to be the first credible disclosure that $\alpha v\beta 6$ protein is expressed on the surface of bladder cancer cells. Thus, $\alpha v\beta 6$ -directed therapies can be used to treat patients that are afflicted with $\alpha v\beta 6$ -expressing bladder cancer (e.g., anti- $\alpha v\beta 6$ antibodies, peptides, or conjugates thereof). $\alpha v\beta 6$ -directed therapies for the treatment of bladder cancer includes the antibodies and conjugates disclosed herein, e.g., chimeric, human, and humanized 15H3 antibodies and conjugates thereof, but is not limited to such antibodies or therapies.

[0087] Human, humanized or chimeric antibodies, alone or as conjugates thereof, are administered in an effective regime meaning a dosage, route of administration and frequency of administration that delays the onset, reduces the severity, inhibits further deterioration, and/or ameliorates at least one sign or symptom of cancer. The regime can be referred to as a therapeutically effective regime. In some instances, therapeutic efficacy can be observed in an individual patient relative to historical controls or past experience in the same patient. In other instances, therapeutic efficacy can be demonstrated in a preclinical or clinical trial in a population of treated patients relative to a control population of untreated patients.

[0088] Exemplary dosages for a monoclonal antibody are, for example, 0.1 mg/kg to 50 mg/kg of the patient's body weight, more typically 1 mg/kg to 30 mg/kg, 1 mg/kg to 20 mg/kg, 1 mg/kg to 15 mg/kg, 1 mg/kg to 12 mg/kg, or 1 mg/kg to 10 mg/kg, or 2 mg/kg to 30 mg/kg, 2 mg/kg to 20 mg/kg, 2 mg/kg to 15 mg/kg, 2 mg/kg to 12 mg/kg, or 2 mg/kg to 10 mg/kg, or 3 mg/kg to 30 mg/kg, 3 mg/kg to 20 mg/kg, 3 mg/kg to 15 mg/kg, 3 mg/kg to 12 mg/kg, or 3 mg/kg to 10 mg/kg. Exemplary dosages for antibody drug conjugates are, for example, 0.01 mg/kg to 10 mg/kg, 0.1 mg/kg to 10 mg/kg, 0.3 mg/kg to 3 mg/kg, 0.5 mg/kg to 3 mg/kg, 1 mg/kg to 7.5 mg/kg, or 2 mg/kg to 7.5 mg/kg or 3 mg/kg to 7.5 mg/kg of the subject's body weight, or 0.1-20, or 0.5-5 mg/kg body weight (e.g., 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 mg/kg) or 10-1500 or 200-1500 mg as a fixed dosage. In some methods, the patient is administered a dose of at least 1.5 mg/kg, at least 2 mg/kg or at least 3 mg/kg, administered once every three weeks or greater. The dosage depends on the identity of the cytotoxic drug, frequency of administration, condition of the patient and response to prior treatment, if any, and whether the disorder is acute or chronic, among other factors.

[0089] Administration is typically parenteral, although it need not be. Administration can also be localized directly into a tumor. Administration into the systemic circulation by intravenous or subcutaneous administration is preferred. Intravenous administration can be, for example, by infusion over a period such as 30-90 min or by a single bolus injection.

[0090] The frequency of administration depends on the half-life of the antibody or conjugate in the circulation, the condition of the patient and the route of administration among other factors. The frequency can be, for example, daily, weekly, monthly, quarterly, or at irregular intervals in response to changes in the patient's condition or progression of the cancer being treated. An exemplary frequency for intravenous administration is between twice a week and quarterly over a continuous course of treatment, although more or less frequent dosing is also possible. Other exemplary frequencies for intravenous administration are between weekly or three out of every four weeks, or every three weeks, over a continuous course of treatment, although more or less frequent dosing is also possible. For subcutaneous administration, an exemplary dosing frequency is daily to monthly, although more or less frequent dosing is also possible.

[0091] The number of dosages administered depends on the nature of the cancer (e.g., whether presenting acute or chronic symptoms) and the response of the disorder to the treatment. In some aspects, for acute disorders or acute exacerbations of a chronic disorder between 1 and 10 doses are often sufficient. Sometimes a single bolus dose, optionally in divided form, is sufficient for an acute disorder or acute exacerbation of a chronic disorder. Treatment can be repeated for recurrence of an acute disorder or acute exacerbation. For chronic disorders, an antibody can be administered at regular intervals, e.g., weekly, fortnightly, monthly, quarterly, every six months for at least 1, 5 or 10 years, or the life of the patient.

[0092] Pharmaceutical compositions for parenteral administration are preferably sterile and substantially isotonic and manufactured under GMP conditions. Pharmaceutical compositions can be provided in unit dosage form (i.e., the dosage for a single administration). Pharmaceutical compositions can be formulated using one or more physiologically acceptable carriers, diluents, excipients or auxiliaries. The formulation

depends on the route of administration chosen. For injection, antibodies can be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiological saline or acetate buffer (to reduce discomfort at the site of injection). The solution can contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively antibodies can be in lyophilized form for constitution with a suitable vehicle, e.g., sterile pyrogen-free water, before use. The concentration of antibody in a liquid formulation can be, for example, 1-100 mg/ml.

[0093] Treatment with antibodies and antibody drug conjugates of the invention can be combined with chemotherapy, radiation, stem cell treatment, surgery other treatments effective against the disorder being treated. In some aspects, antibodies and antibody drug conjugates described herein will be administered with other therapies that are standard of care (e.g., front-line standard of care or second or third line treatment or even salvage therapy) for the particular disease to be treated. Useful classes of other agents that can be administered with antibodies and antibody drug conjugates to $\alpha v\beta 6$ include, for example, both other targeted therapies and/or non-targeted therapies. Examples include, antibodies or antibody drug conjugates to other receptors expressed on cancerous cells, $\alpha v\beta 6$ targeted peptides or conjugates thereof, antitubulin agents (e.g., auristatins), DNA minor groove binders, DNA replication inhibitors, alkylating agents (e.g., platinum complexes such as *cis*-platin, mono(platinum), bis(platinum) and tri-nuclear platinum complexes and carboplatin), anthracyclines, antibiotics, antifolates, antimetabolites, chemotherapy sensitizers, duocarmycins, etoposides, fluorinated pyrimidines, ionophores, lexitropsins, nitrosoureas, platinols, pre-forming compounds, purine antimetabolites, puromycins, radiation sensitizers, steroids, taxanes, topoisomerase inhibitors, vinca alkaloids, and the like.

VII. Other Applications

[0094] The anti- $\alpha v\beta 6$ antibodies can be used for detecting $\alpha v\beta 6$ in the context of clinical diagnosis or treatment or in research. Expression of $\alpha v\beta 6$ on a cancer provides an indication that the cancer is amenable to treatment with the antibodies of the present invention. The antibodies can also be sold as research reagents for laboratory research in

detecting cells bearing $\alpha v\beta 6$ and their response to various stimuli. In such uses, monoclonal antibodies can be labeled with fluorescent molecules, spin-labeled molecules, enzymes or radioisotypes, and can be provided in the form of kit with all the necessary reagents to perform the assay for $\alpha v\beta 6$. The antibodies described herein can be used to detect $\alpha v\beta 6$ protein expression and determine whether a cancer is amenable to treatment with anti- $\alpha v\beta 6$ ADCs. The antibodies can also be used to purify $\alpha v\beta 6$, e.g., by affinity chromatography.

[0095] All patent filings, website, other publications, accession numbers and the like cited above or below are incorporated by reference in their entirety for all purposes to the same extent as if each individual item were specifically and individually indicated to be so incorporated by reference. If different versions of a sequence are associated with an accession number at different times, the version associated with the accession number at the effective filing date of this application is meant. The effective filing date means the earlier of the actual filing date or filing date of a priority application referring to the accession number if applicable. Any feature, step, element, embodiment, or aspect of the invention can be used in combination with any other unless specifically indicated otherwise. Although the present invention has been described in some detail by way of illustration and example for purposes of clarity and understanding, it will be apparent that certain changes and modifications may be practiced within the scope of the appended claims.

EXAMPLES

Materials

[0096] Cell lines described in the following examples were maintained in culture according to the conditions specified by the American Type Culture Collection (ATCC), the National Cancer Institute (NCI) or the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany (DMSZ). Detroit 562 cell lines, HPAFII cell lines, and BxPC3 cell lines were obtained from ATCC. FreeStyleTM 293-F (InVitrogen Corp) human epithelial kidney cells and corresponding transfectants were maintained as described by the manufacturer. Cell culture reagents were obtained from

Invitrogen Corp. (Carlsbad, CA.), Molecular Devices (Sunnydale, CA) and other suppliers. Secondary antibody reagents were purchased from Jackson ImmunoResearch Laboratories (West Grove, PA). Recombinant human $\alpha v\beta 6$ was prepared in house. Recombinant $\alpha v\beta 3$ and $\alpha v\beta 8$ were purchased from R&D Systems (Minneapolis, MN). FreeStyleTM 293-F cells express endogenous integrin αv and they were stably transfected with a full length cDNA encoding human, cynomolgus or murine integrin $\beta 6$ to generate HEK293F:hu $\beta 6$, HEK293F:cyno $\beta 6$ and HEK293F:mu $\beta 6$ cell lines, respectively. HEK293F cells transfected with the empty vector (HEK293F:vector) were used as a negative control. Mouse 3T3 and FDC-P1 cells that express endogenous mouse integrin αv were transfected with a full-length cDNA clone for human and mouse integrin $\beta 6$ to generate 3T3:hu $\beta 6$ and FDC-P1:mu $\beta 6$, respectively.

Methodologies:

Saturation binding assays

[0097] Saturation binding studies were done using the following antigen expressing cell lines: 293F:hu $\beta 6$; 293F:cyno $\beta 6$; FDCP1:mu $\beta 6$ and rat NBT-II. $0.25-0.5 \times 10^6$ antigen expressing cells were aliquoted per well into a 96-well v-bottom plate. m15H3, h15H3(HBLC) and h15H3(HTLC) were directly labeled with equivalent levels of biotin and added to cells at concentrations ranging from 0.1pM-150 nM. Cells were incubated on ice for 30 minutes, washed three times in FACS buffer (PBS, 2% fetal bovine serum, 1 mM MgCl₂, 1 mM CaCl₂, 0.02% NaN₃), incubated on ice for 45 minutes with PE conjugated streptavidin (0.5 ug/ml) in FACS buffer, washed three times and resuspended in FACS buffer. Binding was detected using a Becton Dickinson Biosciences FACSCalibur (San Jose, CA). The apparent Kd was calculated using GraphPad Software (LaJolla, CA).

ELISA

[0098] 96-well Maxisorb plates (Nunc) were coated overnight at 4°C with 1 ug/ml of recombinant human $\alpha v\beta 6$, $\alpha v\beta 3$ and $\alpha v\beta 8$ diluted in PBS. Plates were washed in PBS with 0.05% Tween 20 (PBS-T). Wash buffer was removed and plates were blocked for 30 minutes at room temperature in TBS blocking buffer (TBS plus 1 mM MnCl₂, 0.05%

Tween 20, 1% BSA). Plates were washed and then incubated for 1 hour with biotinylated antibody (m15H3 or h15H3) diluted in TBS blocking buffer at concentrations that ranged from 4.6 ng/ml-10 ug/ml. Plates were washed, incubated for 1 hour with 1 ug/ml of HRP-streptavidin, washed, and then incubated with TMB substrate for 10 minutes. The reaction was stopped with 1 M H₂SO₄. Absorbance at 450 nm was read using a Fusion HT plate reader (Perkin Elmer, Waltham, MA).

Competition binding assays

[0099] Competition binding assays were done using the 293F:hu β and HEK293F:cyno β 6 cell lines. 1x10⁵ antigen expressing cells were aliquoted in each well of a 96-well v-bottom plate on ice. The cells were incubated for 1 hour with 2nM AlexaFluor-647 labeled murine 15H3 and increasing concentrations (from 0.024nM – 100nM) of unlabeled humanized 15H3 constructs. Cells were pelleted and washed 3 times with PBS. The cells were pelleted and resuspended in 125uL of PBS/BSA. Fluorescence was analyzed by flow cytometry, using percent of saturated fluorescent signal to determine percent labeled murine 15H3 bound and to subsequently extrapolate the EC50 by fitting the data to a sigmoidal dose-response curve with variable slope.

Quantitative Flow cytometric analysis

[0100] Quantification of α v β 6 copy number on the cell surfaces was determined using murine α v β 6 mAb as primary antibody and the DAKO QiFiKit flow cytometric indirect assay as described by the manufacturer (DAKO A/S, Glostrup, Denmark) and evaluated with a Becton Dickinson FACScan flow cytometer.

Cytotoxicity assay

[0101] Tumor cells were incubated with anti- α v β 6 antibody drug conjugates for 96 hours at 37°C. A non-binding (h00) ADC was used as a negative control. Cell viability was determined using the CellTiter-Glo® luminescent assay (Promega Corporation, Madison, WI) and results were measured on a Fusion HT plate reader (Perkin Elmer, Waltham, MA). Results are reported as IC₅₀, the concentration of compound needed to yield a 50% reduction in viability compared to vehicle-treated cells (control = 100%).

Production of antibody drug conjugates

[0116] Antibody drug conjugates of the anti- α v β 6 antibodies were prepared as described in US20050238649. The drug linkers vcMMAE and mcMMAF are both described in US20050238649 incorporated herein by reference for all purposes.

***In Vivo* Activity Study**

[0102] Nude (*nu/nu*) mice (7-10 animals/group) were implanted with tumor cells grown in culture Detroit 562 (ATCC) were implanted using 5×10^5 cells in 25% matrigel and HPAFII (ATCC) were implanted using 1×10^6 in 25% matrigel. Dosing with humanized α v β 6 ADC or nonbinding control ADC started when tumors reached 100 mm^3 (q4d x 4 intraperitoneal injections). Tumor volumes were monitored using calipers and animals were euthanized when tumor volume reached $\sim 800 \text{ mm}^3$. Median tumor volume plots were continued for each group until one or more animals were euthanized. All animal procedures were performed under a protocol approved by the Institutional Animal Care and Use Committee in a facility accredited by the Association for Assessment and Accreditation of Laboratory Animal Care.

Immunohistochemical (IHC) staining**Method**

[0103] The Multi-Tumor microarrays (TMA) (cat# MC5001) was purchased from US Biomax Inc. All samples were processed on Bond-MaxTM auto-stainer (Leica).

IHC staining of FFPE tissues:

[0104] Slides with paraffin sections from the multi-tumor array block MC5001 were de-paraffinized using BondTM Dewax solution (Leica, cat # AR9222) at 72°C and rehydrated. Antigen retrieval was performed using EDTA based BondTM Epitope Retrieval Solution 2 (Leica, cat # AR9640) for 20 min at 95-100°C. Slides were treated for 30 min with protein block (DAKO cat #X0909) before incubation with the primary antibody murine anti-integrin β 6 clone 437216 (RnD MAB#41551) at 5 μ g/ml for 45 minutes at 25°C. Isotype-matched murine IgG2b (Zymed #02-6300) was used at 5 μ g/ml as negative control for background staining. For automated IHC staining an alkaline

phosphatase – Fast Red based detection kit (BondTM Polymer AP Red Detection kit; Leica, cat # DS9305) was used. After chromogen development, sections were counterstained with hematoxylin and coverslipped. Slides were evaluated and scored by a pathologist and images were taken using a Olympus BX41 microscope.

IHC of frozen tissues:

Slides with frozen sections were incubated in acetone for 10 minutes at -20°C in order to fix the tissue. PeroxAbolish (Biocare cat# PXA96) peroxide blocking reagent was applied for 15 minutes at 25°C to quench endogenous peroxidase. The slides were sequentially incubated in avidin block (Vector cat#SP-2001) followed by biotin block (Vector cat#SP-2001) for 15 minutes each at 25°C. The slides were incubated with the primary antibody humanized anti-integrin β 6 clone 15H3 at 10 μ g/ml for 60 minutes. The primary and secondary antibodies were prepared in BondTM Primary Antibody Diluent (Leica cat#AR9352) containing 50mM MnCl₂. Human IgG (Ancell cat#295-010) was used at 10 μ g/ml as a negative control for background staining. Following primary antibody incubation, a biotinylated goat anti-human secondary antibody (Jackson cat#109-066-098) was used at 5 μ g/ml and incubated for 30 minutes at 25°C. For detection of the biotinylated secondary antibody, VECTASTAIN Elite ABC reagent (Vector cat#PK-7100) was applied for 30 minutes at 25°C. The BondTM Polymer Refine detection kit (Leica cat# DS9800) was used for DAB chromogen development. Sections were counterstained with hematoxylin and coverslipped. Slides were evaluated and scored by a pathologist and images were taken using an Olympus BX41 microscope.

Results

1. Binding of mouse antibody

[0105] The K_D for the murine α v β 6 monoclonal antibody 15H3 antibody and select humanized version thereof was determined for human, cyno, rat, and mouse. Competition and saturation binding studies for integrin β 6 orthologs was done using genetically engineered cell lines (293F:hu β 6, 293F:cyno β 6 or FDC-P1:mu β 6) with the exception of rat. The genetically engineered cell lines express endogenous α v which pairs with the recombinant β 6 chain to produce a heterodimeric receptor composed of

endogenous αv and recombinant $\beta 6$. The affinity for rat $\alpha v\beta 6$ was determined using rat NBT-II bladder carcinoma cells, which express endogenous $\alpha v\beta 6$.

Table 1

Species	h15H3 (HTLC) (nM)	h15H3 (HBLC) (nM)	m15H3 (nM)
Human	1.2	1.4	0.8
Cyno	1.5	1.6	1.0
Rat	0.4	No data	0.2
Mouse	0.3	0.6	0.2

Generation of 15H3 antibody

[0106] BALB/c mice were immunized three times with intraperitoneal injections of $\sim 5 \times 10^6$ 3T3:hu $\beta 6$ transfectants. Three days prior to fusion, mice received a final injection of purified recombinant human $\alpha v\beta 6$ that was given intravenously (6 μ g) and intraperitoneally (30 μ g). Lymphocytes harvested from spleen and lymph nodes were fused to P3X63Ag8.653 myeloma cells using polyethylene glycol. Fused cells were recovered overnight in hybridoma growth media (IMDM containing 4 mM glutamine, 10% Fetal Clone I, 10% Cloning Factor and Penicillin/Streptomycin). Following recovery, cells were spun down and then plated in semi-solid media. Semi-solid media consisted of CloneMatrix media supplemented with hybridoma growth media plus HAT for hybridoma selection and CloneDetect for IgG-production. Hybridomas were incubated for 10 days at 37°C. At day 10, IgG producing hybridoma clones were picked using a ClonePixFL (Molecular Devices) and transferred to 96-well plates containing IgG-depleted hybridoma growth media plus HT. Hybridoma culture supernatants were screened on 293F:hu $\beta 6$ transfectants and positive clones were identified using an Alexifluor-647 labeled secondary antibody for detection. The plates were read in an FMAT 8200 (Applied Biosystems). Hybridomas that bound to 293F:hu $\beta 6$ and 293F:cyno $\beta 6$ but not 293F:vector were expanded for direct conjugation. The directly conjugated antibody panel was tested in binding and cytotoxicity assays. m15H3 was selected as the lead antibody because it showed cytotoxic activity as an ADC on multiple $\alpha v\beta 6$ -positive tumor cell lines and it had comparable affinity to human and cynomolgus

forms of the antigen. The specificity of mouse 15H3 and humanized 15H3 was confirmed in FMAT and FACS binding studies where the antibodies were shown to bind to 293F:hu β 6 transfectants but not to the α v β 5-positive parental line (293F:vector). The binding specificity of m15H3 and h15H3 was also confirmed by ELISA where the antibodies were bound to recombinant human α v β 6 but not α v β 3 or α v β 8.

Design and testing of humanized antibodies

[0107] The starting point or donor antibody for humanization in this example is the murine 15H3 antibody. Genomic sequences provided by VH1-46 and JH4 for the heavy chain and by VK2-30 and Jk2 for the light chain were used.

[0108] In a first humanization round, thirteen positions were identified in the heavy chain (H28, H30, H38, H40, H41, H48, H66, H67, H69, H71, H72, H93, and H108) and 6 position were identified in the light chain (L21, L22, L36, L37, L45 and L63) at which the human acceptor sequence differed from the donor sequence and that may affect antibody binding as a result of contacting antigen directly, affecting conformation of CDRs or affecting packing between heavy and light chains. Eight humanized heavy chains and five humanized light chains were made incorporating back mutations at different permutations of these positions. Backmutations are underlined and in bold. (Tables 2 and 3). The remainder of the positions are occupied by the residues from the human acceptor sequence.

Table 2: Heavy Chain Backmutations

Ab	H28	H30	H38	H40	H41	H48	H66	H67	H69	H71	H72	H93	H108	# back mutations
HA	T	T	R	A	P	M	R	V	M	R	D	A	L	0
HB	<u>S</u>	<u>S</u>	R	A	P	M	R	V	M	R	D	A	L	2
HC	T	T	<u>K</u>	<u>S</u>	<u>H</u>	M	R	V	M	R	D	A	L	3
HD	T	T	R	A	P	<u>I</u>	R	V	M	R	D	A	L	1
HE	T	T	R	A	P	M	<u>K</u>	<u>A</u>	<u>L</u>	<u>V</u>	D	A	L	4

HF	T	T	R	A	P	M	R	V	M	R	D	<u>V</u>	L	1
HG	T	T	R	A	P	M	R	V	M	R	Q	A	L	1
HH	T	T	R	A	P	M	R	V	M	R	D	A	P	1
HI	S	S	K	S	H	I	K	A	L	V	Q	V	P	13

Table 3 Light Chain Backmutations

Antibody	L21	L22	L36	L37	L45	L63	# back mutations
LA	I	S	F	Q	R	S	0
LB	L	F	F	Q	R	S	2
LC	I	S	L	F	R	S	2
LD	I	S	F	Q	K	S	1
LE	I	S	F	Q	R	T	1
LF	L	F	L	F	K	T	6

[0109] Humanized antibodies were then expressed representing permutations of these chains of the humanized heavy and light chains. All of the heavy chains were expressed but only the HB heavy chain demonstrated appreciable binding (Figure 2). The light chains all showed binding in combination with the HB heavy chain.

[0110] Based on the binding data from the first humanization round, a second humanization round was performed. Three positions, H73, H75, H78, in addition to H28, H30, H72, and H93 were identified in the heavy chain at which the human acceptor sequence differed from the donor sequence and that may affect antibody binding as a result of contacting antigen directly, affecting conformation of CDRs or affecting packing between heavy and light chains. One position, L4 in addition to L21, L22, L36, L37, L45, and L63 was identified in the light chain. 10 additional humanized heavy chains and one additional humanized light chain were made incorporating back mutations at

different permutations of these positions. Backmutations are underlined and in bold. (Tables 4 and 5). The remainder of the positions are occupied by the residues from the human acceptor sequence.

Table 4: Heavy Chain Backmutations

Antibody	H28	H30	H72	H73	H75	H78	H93	# back mutations
HA	T	T	D	T	T	V	A	0
HB	<u>S</u>	<u>S</u>	D	T	T	V	A	2
HF	T	T	D	T	T	V	<u>V</u>	1
HG	T	T	Q	T	T	V	A	1
HK	T	T	D	K	<u>S</u>	<u>A</u>	A	3
HL	<u>S</u>	<u>S</u>	D	K	<u>S</u>	<u>A</u>	A	5
HM	<u>S</u>	<u>S</u>	Q	K	<u>S</u>	<u>A</u>	A	6
HN	<u>S</u>	<u>S</u>	Q	T	T	V	A	3
HO	<u>S</u>	<u>S</u>	Q	K	T	V	A	4
HQ	T	T	D	T	<u>S</u>	<u>A</u>	A	2
HR	<u>S</u>	<u>S</u>	D	T	<u>S</u>	<u>A</u>	A	4
HS	<u>S</u>	<u>S</u>	Q	T	<u>S</u>	<u>A</u>	A	5
HT	<u>S</u>	<u>S</u>	Q	T	T	V	<u>V</u>	4
HU	<u>S</u>	<u>S</u>	Q	K	<u>S</u>	<u>A</u>	<u>V</u>	7

Table 5 Light Chain Backmutations

Antibody	L4	L21	L22	L36	L37	L45	L63	# back mutations

LA	M	I	S	F	Q	R	S	0
LB	M	L	F	F	Q	R	S	2
LC	M	I	S	L	F	R	S	2
LD	M	I	S	F	Q	K	S	1
LE	M	I	S	F	Q	R	T	1
LF	M	L	F	L	F	K	T	6
LG	L	I	S	L	F	R	S	3

[0111] All of the heavy chains (except for HO and HK) were expressed. The binding curves are shown in Figures 4-7.

The EC50's are summarized in the Table 6 below.

Table 6: EC₅₀s for select humanized anti- α v β 6 antibodies

Ab	EC50 (nM)
HBLC	8.14
HLLC	0.74
HLLG	0.97
HMLC	0.49
HMLG	1.07
HNLC	11.14
HNLG	3.16
HRLC	26.7
HRLG	23.9
HSLC	32.5
HTLC	3.7
HULC	0.93

Expression Data

[0112] Anti-B6 clones 437216 (R&D systems) was used for immunohistochemical analysis of various tumor types using formalin-fixed paraffin embedded tissues (FFPE).

Summary of the expression data for $\alpha v\beta 6$ in multitumor array

Tumor Origin	$\alpha v\beta 6+$ expression (%)
Bladder	65
Head & Neck	50
Skin	33
Lung	30
Pancreas	30
Uterus	30
Breast	26
Colon	20
Prostate	15
Ovary	14
Stomach	5

[0113] Anti-B6 clones 437216 (R&D systems) was used for immunohistochemical analysis of various tumor types using frozen tissue

Summary of the expression data for $\alpha v\beta 6$ in frozen tissue samples

Tumor Origin	$\alpha v\beta 6+$	# cases	%
Pancreatic	10	9	100%
Head & Neck	10	10	100%

[0114] Anti-B6 clones 437216 (R&D systems) was also used for immunohistochemical analysis of various normal tissue types using formalin-fixed paraffin embedded tissues (FFPE) and frozen tissue. Normal breast, pancreas, esophagus, larynx, lung, skin, uterus, breast, colon, prostate, stomach, and ovary all had no B6 expression.

[0115] The 15H3 antibody was also used to stain frozen tumor and normal tissue samples. The normal tissue samples were negative for B6 expression. 23% (3/13) of triple negative breast cancer tissues showed mild to strong B6 expression. 100% of pancreatic cancer samples (5/5 samples) lung cancer samples (10/10) and gastric cancer samples (10/10) showed strong B6 expression.

***In Vitro* Anti-tumor Activity of murine 15H3 ADC**

[0116] Anti-tumor activity of anti- α v β 6 ADCs in vitro was measured using cytotoxicity assays. A survey of α v β 6 expression in various cell lines by quantitative FACS analysis was performed. Referring to the table below, the murine 15H3 ADCs (the murine 15H3 antibody conjugated with vcMMAE or mcMMAF (both small molecules and/or linkers described in US20050238649)) were effective in killing α v β 6 cells.

Cancer cell line	Copies/cell (x10 ³)	M15H3-vc-MMAE IC ₅₀ (ng/ml)	M15H3-mc-MMAF IC ₅₀ (ng/ml)
HupT4	184	35	16
BxPC3	37	34	7
HPAFII	22	76	>2000
Panc08.13	137	93	1400
Capan-1	19	365	1508
SU8686	33	No effect	No effect
5637	36	>1000	>1000
SW780	80	614	86
RT4	42	>1000	>1000

Figure 9 shows that humanized 15H3 anti- α v β 6 ADCs behaved similarly as the mouse parental in cytotoxicity assays.

***In Vivo* Anti-tumor Activity of humanized anti- α v β 6 ADCs**

[0117] Using pancreatic cancer (HPAFII, BxPC-3) and head and neck cancer (Detroit 562) models, the anti-tumor activity of select humanized α v β 6 ADCs (average of 4 drugs per antibody) *in vivo* (Figs. 10-12) was demonstrated. α v β 6 ADCs conjugated to vcMMAE showed significant tumor delay or tumor regression compared to untreated and

control ADCs. h15H3-vcMMAE(4) refers to antibody drug conjugates of the humanized form of the parental murine antibody having an average of 4 vcMMAE drug linker molecules per antibody. h00- vcMMAE (4) refers to an antibody drug conjugate of a nonbinding control antibody having an average of 4 vcMMAE drug linker molecules per antibody. h15H3-1 is the HBLC construct, h15H3-2 is the HLLC construct and h15H3-3 is the HTLC construct.

Sequences

SEQ ID NO:1 - murine heavy chain variable region - protein

EVQLQQSGPELVKPGASVKISCKASGYSFSGYFMNWVKQSHGQSLEWIGLINPYNGDSFYNQKFKGKATLT
VQKSSSTAHMELQSLTSEDSAFCVRLRRDFDYWGQGTPLTVSS

SEQ ID NO:2 - murine light chain variable region - protein

DVVLTIQIPLTLSVTIGQPASLFCSSQSLSDGKTYLNWLFQRPGQSPKRLIYLVSELDGVPDRFTGSG
SGTDFTLKISRVEAEDLGVYYCWQGTHFPRTFGGGTKLEIKR

SEQ ID NO:3 - murine heavy chain CDR1 - protein

GYFMN

SEQ ID NO:4 - murine heavy chain CDR2 - protein

LINPYNGDSFYNQKFKG

SEQ ID NO:5 - murine heavy chain CDR3 - protein

GLRRDFDY

SEQ ID NO:6 - murine light chain CDR1 - protein

KSSQSLSDGKTYLN

SEQ ID NO:7 - murine light chain CDR2 - protein

LVSELDs

SEQ ID NO:8 - murine light chain CDR3 - protein

WQGTHFPRT

SEQ ID NO:9 - HA heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVSCKASGYTFTGYFMNWVRQAPGQGLEWMGLINPYNGDSFY
NQKFKGRVTMTRDTSTVYMESSLRSEDTAVYYCARGLRRDFDYWGQGTPLTVSS

SEQ ID NO:10 - LA light chain variable region - protein

DVVMTQSPLSLPVTLGQPASISCKSSQSLSDGKTYLNWFQQRPGQSPRRLIYLVSELD
SGVPDRFSGSGTDFTLKISRVEAEDGVYYCWQGTHFPRTFGGGTKLEIKR

SEQ ID NO:11 - HB heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVSCKASGYSFSGYFMNWVRQAPGQGLEWMGLINPYNGDSFY

NQKFKGRVTMTRDTSTSTVYMEPLLSEDTAVYYCARGLRRDFDYWGQGTLTVSS

SEQ ID NO:12 - HF heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVSKASGYTFTGYFMNWVRQAPGQGLEWMGLINPYNGDSFY
NQKFKGRVTMTRDTSTSTVYMEPLLSEDTAVYYCARGLRRDFDYWGQGTLTVSS

SEQ ID NO:13 - HG heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVSKASGYTFTGYFMNWVRQAPGQGLEWMGLINPYNGDSFY
NQKFKGRVTMTRQTSTSTVYMEPLLSEDTAVYYCARGLRRDFDYWGQGTLTVSS

SEQ ID NO:14 - HK heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVSKASGYTFTGYFMNWVRQAPGQGLEWMGLINPYNGDSFY
NQKFKGRVTMTRDKSSSTAYMELSSLRSEDTAVYYCARGLRRDFDYWGQGTLTVSS

SEQ ID NO:15 - HL heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVSKASGYFSGYFMNWVRQAPGQGLEWMGLINPYNGDSFY
NQKFKGRVTMTRDKSSSTAYMELSSLRSEDTAVYYCARGLRRDFDYWGQGTLTVSS

SEQ ID NO:16 - HM heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVSKASGYFSGYFMNWVRQAPGQGLEWMGLINPYNGDSFY
NQKFKGRVTMTRQKSSSTAYMELSSLRSEDTAVYYCARGLRRDFDYWGQGTLTVSS

SEQ ID NO:17 - HN heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVSKASGYFSGYFMNWVRQAPGQGLEWMGLINPYNGDSFY
NQKFKGRVTMTRQTSTSTVYMEPLLSEDTAVYYCARGLRRDFDYWGQGTLTVSS

SEQ ID NO:18 - HO heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVSKASGYFSGYFMNWVRQAPGQGLEWMGLINPYNGDSFY
NQKFKGRVTMTRQKSTSTVYMEPLLSEDTAVYYCARGLRRDFDYWGQGTLTVSS

SEQ ID NO:19 - HQ heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVSKASGYTFTGYFMNWVRQAPGQGLEWMGLINPYNGDSFY
NQKFKGRVTMTRDTSSSTAYMELSSLRSEDTAVYYCARGLRRDFDYWGQGTLTVSS

SEQ ID NO:20 - HR heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVSKASGYFSGYFMNWVRQAPGQGLEWMGLINPYNGDSFY
NQKFKGRVTMTRDTSSSTAYMELSSLRSEDTAVYYCARGLRRDFDYWGQGTLTVSS

SEQ ID NO:21 - HS heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVSKASGYFSGYFMNWVRQAPGQGLEWMGLINPYNGDSFY
NQKFKGRVTMTRQTSSSTAYMELSSLRSEDTAVYYCARGLRRDFDYWGQGTLTVSS

SEQ ID NO:22 - HT heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVKASGYFSGYFMNWRQAPGQGLEWMGLINPYNGDSFY
NQKFKGRVTMTRQTSTVYMEMLSSLRSEDTAVYYCVRGLRRDFDYWGQGTLTVSS

SEQ ID NO:23 - HU heavy chain variable region - protein

QVQLVQSGAEVKKPGASVKVKASGYFSGYFMNWRQAPGQGLEWMGLINPYNGDSFY
NQKFKGRVTMTRQKSSSTAYMELSSLRSEDTAVYYCVRGLRRDFDYWGQGTLTVSS

SEQ ID NO:24 - LB light chain variable region - protein

DVVMQTQSPSLPVTLGQPASLFCRKSSQSLSDGKTYLNWFQQRPGQSPRRLIYLVSELD
SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCWQGTHFPRTFGGGTKLEIKR

SEQ ID NO:25 - LC light chain variable region - protein

DVVMQTQSPSLPVTLGQPASICKSSQSLSDGKTYLNWLFQQRPGQSPRRLIYLVSELD
SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCWQGTHFPRTFGGGTKLEIKR

SEQ ID NO:26 - LD light chain variable region - protein

DVVMQTQSPSLPVTLGQPASICKSSQSLSDGKTYLNWLFQQRPGQSPKRLIYLVSELD
SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCWQGTHFPRTFGGGTKLEIKR

SEQ ID NO:27 - LE light chain variable region - protein

DVVMQTQSPSLPVTLGQPASICKSSQSLSDGKTYLNWLFQQRPGQSPRRLIYLVSELD
SGVPDRFTGSGSGTDFTLKISRVEAEDVGVYYCWQGTHFPRTFGGGTKLEIKR

SEQ ID NO:28 - LF light chain variable region - protein

DVVMQTQSPSLPVTLGQPASICKSSQSLSDGKTYLNWLFQQRPGQSPKRLIYLVSELD
SGVPDRFTGSGSGTDFTLKISRVEAEDVGVYYCWQGTHFPRTFGGGTKLEIKR

SEQ ID NO:29 - LG light chain variable region - protein

DVVLQTQSPSLPVTLGQPASICKSSQSLSDGKTYLNWLFQQRPGQSPRRLIYLVSELD
SGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCWQGTHFPRTFGGGTKLEIKR

SEQ ID NO:30 - human IgG1 heavy chain constant region - protein

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVWSNSGALTSGVHTFPAVLQSS
GLYSLSSVVTVPSSSLGTQTYICNVNWKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGG
PSVFLFPPKPKDTLmisRTPEVTCVVVDVSHEDPEVFKFNWYVDGVEVHNAKTKPREEQYN
STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSRDE
LTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSGSFFLYSKLTVDKSRW
QQGNVFSCSVMHEALHNHYTQKSLSLSPGK

SEQ ID NO:31 - human kappa light chain constant domain - protein

TVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDS
KDSTYSLSSTTLSKADYEKHKVYACEVTHQGLSPVTKSFNRGEC

SEQ ID NO:32 - human IgG1 heavy chain constant region S239C
mutant - protein

ASTKGPSVFPLAPSSKSTSGGTAAALGCLVKDYFPEPVTWSWNSGALTSGVHTFPALQSS
GLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGG
PCVFLFPPKPKDTLMSRTPEVTCVVVDVSHEDPEVFKFNWYVDGVEVHNAKTKPREEQYN
STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDE
LTKNQVSLLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRW
QQGNVFSCSVMHEALHNHYTQKSLSLSPGK

SEQ ID NO:33 - murine heavy chain variable region signal peptide
- protein

MGWSCIIFLFLLSVTVGVFS

SEQ ID NO:34 - human heavy chain variable region signal peptide -
protein

MAWVWTLLFLMAAAQSAQA

SEQ ID NO:35 - murine light chain variable region signal peptide
- protein

MSPAQFLFLLVWLWIRETNG

SEQ ID NO:36 - human light chain variable region signal peptide -
protein

MKLPVRLLVLMFWIPASSS

SEQ ID NO:37 - mouse heavy chain variable region - nucleic acid

gagggttcagctgcagcagtctggacactgagttggtaaaacctggggcttcagtgaagatatcctgcaaggc
ttctggatttattttatggactttatgaactgggtgaaacagagccatggacagagccttgagtgg
ttggacttattatccttacaatggagacttttacaatcagaagttcaaggccacattgact
gtacagaagtcccttagtacagcccacatggaaactccagagcctgacatctgaagactctgcagtctttt
ttgtgttagagggttacgacggactttgactatggggccaaggcacccttcacagtctcctca

SEQ ID NO:38 - mouse light chain variable region - nucleic acid

Gatgttgtttgaccagattccactcaactttgtcggttaccattggacaaccaggcctccctttttaa
gtcaagtccatggacttttagatagtgtatggaaagacatatttgaattgggttatttcagaggccaggccagt
ctccaaagcccttatttacatgggtctgaactggactctggactccctgacaggttcactggcagtgg
tcagggacagatttccacactgaaaatcagcagagttggaggctgaggattgggagttattatgtggca
aggtacacatttccctcgacgttgcggaggcaccagttggaaatcaaacgg

SEQ ID NO:39 - HA heavy chain variable region - nucleic acid

Caggtgcagctgggcagtcggagctgaggtgaagaagcctggagcctctgtgaaggtgtcctgttaaggc
ctctggctacaccctcacaggctacttcatgaactgggtgaggcaggccctggccagggcctggagtgga
tgggcctgatcaacccttacaatggagactccttctacaaccagaagttcaagggcagggtgaccatgacc
agggacacccctccacccatggagactgtacatggagctgtcctccctgaggctgaggacacagctgttacta
ctgtgccagggcctgaggagggactttgactactggggccagggcaccctggtacagtgtcctcc

SEQ ID NO:40 - HB heavy chain variable region - nucleic acid

Caggtgcagctgggcagtcggagctgaggtgaagaagcctggagcctctgtgaaggtgtcctgttaaggc
ctctggctacagctctctggctacttcatgaactgggtgaggcaggccctggccagggcctggagtgga
tgggcctgatcaacccttacaatggagactccttctacaaccagaagttcaagggcagggtgaccatgacc
agggacacccctccacccatggagactgtacatggagctgtcctccctgaggctgaggacacagctgttacta
ctgtgccagggcctgaggagggactttgactactggggccagggcaccctggtacagtgtcctcc

SEQ ID NO:41 - HL heavy chain variable region - nucleic acid

Caggtgcagctgggcagtcggagctgaggtgaagaagcctggagcctctgtgaaggtgtcctgttaaggc
ctctggctacagctctctggctacttcatgaactgggtgaggcaggccctggccagggcctggagtgga
tgggcctgatcaacccttacaatggagactccttctacaaccagaagttcaagggcagggtgaccatgacc
agggacacccctccacccatggagactgtacatggagctgtcctccctgaggctgaggacacagctgttacta
ctgtgccagggcctgaggagggactttgactactggggccagggcaccctggtacagtgtcctcc

SEQ ID NO:42 - HT heavy chain variable region - nucleic acid

Caggtgcagctgggcagtcggagctgaggtgaagaagcctggagcctctgtgaaggtgtcctgttaaggc
ctctggctacagctctctggctacttcatgaactgggtgaggcaggccctggccagggcctggagtgga
tgggcctgatcaacccttacaatggagactccttctacaaccagaagttcaagggcagggtgaccatgacc
agggacacccctccacccatggagactgtacatggagctgtcctccctgaggctgaggacacagctgttacta
ctgtgtcagggcctgaggagggactttgactactggggccagggcaccctggtacagtgtcctcc

SEQ ID NO:43 - HN heavy chain variable region - nucleic acid

Caggtgcagctgggcagtcggagctgaggtgaagaagcctggagcctctgtgaaggtgtcctgttaaggc
ctctggctacagctctctggctacttcatgaactgggtgaggcaggccctggccagggcctggagtgga
tgggcctgatcaacccttacaatggagactccttctacaaccagaagttcaagggcagggtgaccatgacc
agggacacccctccacccatggagactgtacatggagctgtcctccctgaggctgaggacacagctgttacta
ctgtgccagggcctgaggagggactttgactactggggccagggcaccctggtacagtgtcctcc

SEQ ID NO:44 - HU heavy chain variable region - nucleic acid

Caggtgcagctgggcagtcggagctgaggtgaagaagcctggagcctctgtgaaggtgtcctgttaaggc
ctctggctacagctctctggctacttcatgaactgggtgaggcaggccctggccagggcctggagtgga
tgggcctgatcaacccttacaatggagactccttctacaaccagaagttcaagggcagggtgaccatgacc
agggacacccctccacccatggagactgtacatggagctgtcctccctgaggctgaggacacagctgttacta
ctgtgtcagggcctgaggagggactttgactactggggccagggcaccctggtacagtgtcctcc

SEQ ID NO:45 - LA light chain variable region - nucleic acid

gatgtggatgaccaggcccactgtccctgcctgtgaccctggccagggcctgcctccatctcctgtaa
gtcctccaggccctgctggactctgtatggcaagggcacttgcacttgcactgggttccagcagggcctggccagt
ccccttagggctgatctacccctgggtctgagctggactctggagtgccctgacagggttctctggctctggc
tctggcacagacttccaccctgaagatctccagggtggaggctgaggatgtggagtgactactgttggca
gggcaccacttccctaggacccatggaggtggaaaccaagctggagatcaagcg

SEQ ID NO:46 - LC light chain variable region - nucleic acid

gatgtggtgatgaccagtcctctgtccctgcctgtgaccctggccagccctgcctccatctcctgtaa
gtcctccagtcctgctggactctgtatggcaagacacctacctaactggctgtccagaggcctggccagt
cccctaggaggctgatctacctgggtctgagctgactctggagtgccctgacaggttctctggctctggc
tctggcacagacttcaccctgaagatctccagggtgaggctgaggatgtggagtgtaactactgttggca
ggcaccacttccttaggaccttggaggtggaaccaagctggagatcaagcgt

SEQ ID NO:47 - LF light chain variable region - nucleic acid

Gatgtggtgatgaccagtcctctgtccctgcctgtgaccctggccagccctgcctccctgttctgtaa
gtcctccagtcctgctggactctgtatggcaagacacctacctaactggctgtccagaggcctggccagt
cccctaagaggctgatctacctgggtctgagctgactctggagtgccctgacaggttcacagcgtctggc
tctggcacagacttcaccctgaagatctccagggtgaggctgaggatgtggagtgtaactactgttggca
ggcaccacttccttaggaccttggaggtggaaccaagctggagatcaagcgt

SEQ ID NO:48 - LG light chain variable region - nucleic acid

gatgtggtgctgaccagtcctctgtccctgcctgtgaccctggccagccctgcctccatctcctgtaa
gtcctccagtcctgctggactctgtatggcaagacacctacctaactggctgtccagaggcctggccagt
cccctaggaggctgatctacctgggtctgagctgactctggagtgccctgacaggttctctggctctggc
tctggcacagacttcaccctgaagatctccagggtgaggctgaggatgtggagtgtaactactgttggca
ggcaccacttccttaggaccttggaggtggaaccaagctggagatcaagcgt

SEQ ID NO:49 - murine heavy chain variable region signal peptide - nucleic acid

Atggatggagctgtatctttctttcttcctgtcagtaactgttaggtgtgttctct

SEQ ID NO:50 - human heavy chain variable region signal peptide - nucleic acid

Atggctgggtgtggaccttgcattcctgatggcagctgccaaagtgcccaagca

SEQ ID NO:51 - murine light chain variable region signal peptide nucleic acid

atgagtccctgcccagttcctgtttctgttagtgctctggattcggaaaccaacgg

SEQ ID NO:52 - human light chain variable region signal peptide - nucleic acid

Atgaagttgcctgttaggctgttagtgatgttctggattcctgcttcagcagt

SEQ ID NO:53 - human IgG1 heavy chain constant region - nucleic acid

gttagcaccaaggcccateggcttccccctggcaccccttcacagagcacccctggggcacagcggccctggctgcgtcaaggactactcc
ccgaaccggtgacgggtcgatggactcaggcgcctgaccagcggcgtgcacacctcccgctgtctacgtcctcaggactactccctcagcag
cgtggtagccgtgcctccagcagctgggcacccagacactatgcacgtgaatcacaagcccagcaacaccaagggtggacaagaaagttagcc

caaatttgtgacaaaactcacatgcccaccgtgcccagcacctgaactcctgggggaccgtcagtttccttccccccaaaacccaaggacaccc
tcatgatctcccgaccctgaggcacatcgctgggtggacgtgaggcacgaagaccctgaggcacgtcaagtcaactgttacgtggacggcgtggaggt
gcataatgccaagacaaaagecgccccggggaggagcagtcacacagcacgtaccgtgtggcagegtcctcaccgtcctgaccaggactggcgtgaatggca
aggagtacaagtgcaggtctccaacaaagccctccagccccatcgagaaaaccatctccaaagecaaaaggcagccccgagaaccacagggttac
accctgecccccattcccggtgagctgaccagaaccaggcgtcagcgtaccgtcctggtcaaggcttacccagcgtacatcgccgtggagtgagg
agaatggcagccggagaacaactacaagaccgcctcccggtcggactccgacggcgtcccttacagcaagtcaccgtggacaagagc
agggtggcagcagggaacgttcatgtccgtatgcatgaggctgtcacaaccactacacgcagaagagcctccctgtccggtaaa

SEQ ID NO:54 – human kappa light chain constant domain – nucleic acid

Actgtggcgccatctgtttcatcttccgcctatgtatgagcagttgaatctggaaactgectctgtgtgcctgtcaataacttctatcccagagag
gceaaagtacagtggaaagggtggataacgcctccaatcggttaactccaggagagtgtcagcagagcaggacagcacactacgcctcag
cagcacccctgacgctgagcaaagcagactacgagaaacacaaagtctacgcctcgaagtccatcaggcctgagctegccgtcacaagagctt
caacaggggagagtgt

SEQ ID NO:55 – human IgG1 heavy chain constant region S239C mutant – nucleic acid

Gctagcaccaagggcccacatgtttccccctggcacccttcataagagcacctctggggcacagctgcctggcctgtcaaggactacttcc
ctgaacccctgtacagtgttcgttgcactcaggcgccctgaccagcggcgtgcacaccctccggctgttctacagtccctcaggacttactccctcagc
gtgggtaccgtgcctccagcagctggcaccctgaccatctgcaactgtgaatcacaagcccaagacaccatcgagttggacaagaagtgtgagcc
aaatctgtgacaaaactcacatgcctccaccgtgcccagcacctgaactctgggggaccgtgttcttccccccaaaacccaaggacaccctc
atgatctcccgacccttgagggtcacatgcgtgtggacgtgagccacgaagaccctgaggtaagttcaactgttacgtggacggcgtggaggt
cataatgccaagacaaaagecgccccggggaggagcagtcacacagcacgtaccgtgtggcagtcctcaccgtcaccaggactggcgtaaatggca
ggagtacaagtgcaggtctccaacaaagccctcccgccctccatcgagaaaaccatctcaaaagecaaaaggcagccccgagaaccacacagggttaca
ccctgccccatcccggtgagctgaccaagaaccaggcgtaccgtgcctggtaaaggcttacccagcgtacatcgccgtggagtgggaga
gcaatggcagccggagaacaactacaagaccacgcctcccggtcggactccgacggcgttcttacagcaagtcaccgtggacaagagca
ggtggcagcagggaacgttcatgtccgtatgcatgaggctgtcacaaccactacacagaagagcctccctgtccggtaaa

CLAIMS

1. An isolated monoclonal antibody that specifically binds integrin $\alpha v\beta 6$ comprising heavy chain complementary determining region (CDR) sequences as set forth in SEQ ID NO:3 (CDR1), SEQ ID NO:4 (CDR2), and SEQ ID NO:5 (CDR3), and light chain CDR sequences as set forth in SEQ ID NO:6 (CDR1), SEQ ID NO:7 (CDR2), and SEQ ID NO:8 (CDR3).
2. The antibody of claim 1, wherein the antibody is an isolated murine 15H3 monoclonal antibody that specifically binds integrin $\alpha v\beta 6$ and wherein the antibody comprises a heavy chain variable region comprising the sequence set forth in SEQ ID NO:1 and a light chain variable region comprising the sequence set forth in SEQ ID NO:2; and chimeric or humanized forms thereof.
3. The antibody of claim 1 comprising a heavy chain variable region having an amino acid sequence at least 90% identical to HA (SEQ ID NO.9) provided there is at least one murine 15H3 backmutation in the heavy chain variable framework.
4. The antibody of claim 3 wherein there are at least two murine 15H3 backmutations in the heavy chain variable framework region.
5. The antibody of claim 3, wherein at least one of positions H28 and H30 of HA are occupied by S, wherein numbering is according to the Kabat numbering convention.
6. The antibody of claim 5 wherein both of positions H28 and H30 are occupied by S.
7. The antibody of claim 6, wherein position H72 is occupied by D or Q; position H73 is occupied by T or K, position H75 is occupied by T or S; position H78 is occupied by V or A, and position 93 is occupied by A or V.

8. The antibody of any one of the preceding claims, wherein the antibody comprises a light chain variable region comprising the sequence set forth in LA (SEQ ID NO. 10) except for position L4 is occupied by M or L, position L21 is occupied by I or L, position L22 is occupied by S or F, position L36 is occupied by F or L, position L37 is occupied by Q or F, position L45 is occupied by R or K, position L63 is occupied by S or T, wherein numbering is according to the Kabat numbering convention, and/or the heavy chain variable region has the amino acid sequence set forth in HB (SEQ ID NO:11), HT (SEQ ID NO:22), HL (SEQ ID NO:15), HU (SEQ ID NO:23), HM (SEQ ID NO:16) or HN (SEQ ID NO:17), and/or the light chain variable region has the amino acid sequence set forth in LA (SEQ ID NO:10), LB (SEQ ID NO:24), LC (SEQ ID NO:25), LD (SEQ ID NO:26), LE (SEQ ID NO:27), LF (SEQ ID NO:28) or LG (SEQ ID NO:29).
9. The antibody of claim 7 having the heavy chain variable region having the amino acid sequence set forth in HB (SEQ ID NO:11) and a light chain variable region having the amino acid sequence set forth in LA (SEQ ID NO:10), LB (SEQ ID NO:24), LC (SEQ ID NO:25), LD (SEQ ID NO:26), LE (SEQ ID NO:27), LF (SEQ ID NO:28) or LG (SEQ ID NO:29); having the heavy chain variable region having the amino acid sequence set forth in HT (SEQ ID NO:22) and a light chain variable region having the amino acid sequence set forth in LA (SEQ ID NO:10), LB (SEQ ID NO:24), LC (SEQ ID NO:25), LD (SEQ ID NO:26), LE (SEQ ID NO:27), LF (SEQ ID NO:28) or LG (SEQ ID NO:29); having the heavy chain variable region having the amino acid sequence set forth in HL (SEQ ID NO:15) and a light chain variable region having the amino acid sequence set forth in LA (SEQ ID NO:10), LB (SEQ ID NO:24), LC (SEQ ID NO:25), LD (SEQ ID NO:26), LE (SEQ ID NO:27), LF (SEQ ID NO:28) or LG (SEQ ID NO:29); having the heavy chain variable region having the amino acid sequence set forth in HU (SEQ ID NO:23) and a light chain variable region having the amino acid sequence set forth in LA (SEQ ID NO:10), LB (SEQ ID NO:24), LC (SEQ ID NO:25), LD (SEQ ID NO:26), LE (SEQ ID NO:27), LF (SEQ ID NO:28) or LG (SEQ ID NO:29); having the heavy chain variable region having the amino acid

sequence set forth in HN (SEQ ID NO:17) and a light chain variable region having the amino acid sequence set forth in LA (SEQ ID NO:10), LB (SEQ ID NO:24), LC (SEQ ID NO:25), LD (SEQ ID NO:26), LE (SEQ ID NO:27), LF (SEQ ID NO:28) or LG (SEQ ID NO:29); or having the heavy chain variable region having the amino acid sequence set forth in HM (SEQ ID NO:16) and a light chain variable region having the amino acid sequence set forth in LA (SEQ ID NO:10), LB (SEQ ID NO:24), LC (SEQ ID NO:25), LD (SEQ ID NO:26), LE (SEQ ID NO:27), LF (SEQ ID NO:28) or LG (SEQ ID NO:29).

10. The antibody of any of the preceding claims wherein the antibody is an antigen binding fragment.
11. The antibody of any one of the preceding claims, wherein the heavy chain variable region is fused to a human heavy chain constant region and the light chain variable region is fused to a human light chain constant region and optionally wherein the heavy chain constant region is of IgG1 isotype.
12. The antibody of claim 11, wherein the heavy chain constant region is a mutant form of natural human constant region which has reduced binding to an Fc gamma receptor relative to the natural human constant region..
13. The antibody of any one of the preceding claims, wherein the antibody is conjugated to a cytotoxic agent, and/or having within 5 fold affinity for $\alpha v\beta 6$ as compared to the murine 15H3 antibody, and/or having within 2 fold affinity for $\alpha v\beta 6$ as compared to the murine 15H3 antibody, and/or having an apparent dissociation constant (kd) for $\alpha v\beta 6$ within a range of 0.1 nM to 5 nM and/or having an apparent dissociation constant (kd) for $\alpha v\beta 6$ within a range of 0.1 nM to 2 nM, and/or having an apparent dissociation constant (kd) for $\alpha v\beta 6$ within a range of 0.5 nM to 1.5 nM.
14. The antibody of claim 1, wherein the antibody is a chimeric form of the murine 15H3 antibody and comprises the heavy chain variable region comprising the sequence set

	10	20	30	40	50
m15H3 VH				
h15H3 HAQ...P.LV.....I.....S.S.....K.SH..S...I.....				
h15H3 HBS.S.....			
h15H3 HLS.S.....			
h15H3 HNS.S.....			
h15H3 HTS.S.....			
h15H3 HUS.S.....			
	70	80	90	100	110
m15H3 VH				
h15H3 HAKA.L.VQK.S..AH..Q..T..S..F..V.....PL.....				
h15H3 HB				
h15H3 HLK.S..A.....				
h15H3 HNQ.....				
h15H3 HTQ.....	V.....		
h15H3 HUQK.S..A.....	V.....		
	10	20	30	40	50
m15H3 VL				
h15H3 LAL..I..T.S..I..LF.....LF.....K.....		
h15H3 LCLF.....			
h15H3 LFLF.....LF.....K.....	
h15H3 LGL.....LF.....			
	60	70	80	90	100
m15H3 VL				
h15H3 LAT.....L.....				
h15H3 LC				
h15H3 LFT.....				
h15H3 LG				

Figure 1






Figure 2



C

Test Sample	Kd (nM)
m15H3	0.2
h15H3(HBLC)	0.6
h15H3(HTLC)	0.3

Figures 3A-3C




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8





Figure 9A



Figures 9A-9B




Figure 9C




Figure 9D

Figures 9C-9D




Figure 9E
HupT4




Figure 9F
Figures 9E-9F

Activity of h15H3-vcMMAE(4) on Detroit 562 Head and Neck Carcinoma




Figure 10

Activity of anti α V β 6 ADCs on HPAFII Pancreatic Carcinoma Xenografts in Nude Female Mice




Figure 11




Figure 12

AVB6-00112PC_ST25. txt
SEQUENCE LISTING

<110> Seattle Genetics, Inc.

<120> ANTI BODIES TO INTEGRIN ALPHA VBETA6 AND USE OF SAME TO TREAT
CANCER

<130> AVB6-00112PC

<150> US 61/600, 499
<151> 2012-02-17

<150> US 61/602, 511
<151> 2012-02-23

<160> 55

<170> PatentIn version 3.5

<210> 1
<211> 117
<212> PRT
<213> Artificial

<220>
<223> murine heavy chain variable region

<400> 1

Gl u Val Gl n Leu Gl n Gl n Ser Gl y Pro Gl u Leu Val Lys Pro Gl y Al a
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Al a Ser Gl y Tyr Ser Phe Ser Gl y Tyr
20 25 30

Phe Met Asn Trp Val Lys Gl n Ser His Gl y Gl n Ser Leu Gl u Trp Ile
35 40 45

Gl y Leu Ile Asn Pro Tyr Asn Gl y Asp Ser Phe Tyr Asn Gl n Lys Phe
50 55 60

Lys Gl y Lys Al a Thr Leu Thr Val Gl n Lys Ser Ser Ser Thr Al a His
65 70 75 80

Met Gl u Leu Gl n Ser Leu Thr Ser Gl u Asp Ser Al a Val Phe Tyr Cys
85 90 95

Val Arg Gl y Leu Arg Arg Asp Phe Asp Tyr Trp Gl y Gl n Gl y Thr Pro
100 105 110

Leu Thr Val Ser Ser
115

<210> 2
<211> 113
<212> PRT
<213> Artificial

<220>
<223> murine light chain variable region

AVB6-00112PC_ST25. txt

<400> 2

Asp Val Val Leu Thr Glu Ile Pro Leu Thr Leu Ser Val Thr Ile Gly
1 5 10 15

Gl n Pro Ala Ser Leu Phe Cys Lys Ser Ser Glu Ser Leu Leu Asp Ser
20 25 30

Asp Gl y Lys Thr Tyr Leu Asn Trp Leu Phe Glu Arg Pro Gl y Glu Ser
35 40 45

Pro Lys Arg Leu Ile Tyr Leu Val Ser Glu Leu Asp Ser Gl y Val Pro
50 55 60

Asp Arg Phe Thr Gl y Ser Gl y Ser Gl y Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Gl u Ala Gl u Asp Leu Gl y Val Tyr Tyr Cys Trp Gl n Gl y
85 90 95

Thr His Phe Pro Arg Thr Phe Gl y Gl y Thr Lys Leu Gl u Ile Lys
100 105 110

Arg

<210> 3

<211> 5

<212> PRT

<213> Artificial

<220>

<223> murine heavy chain CDR1

<400> 3

Gl y Tyr Phe Met Asn
1 5

<210> 4

<211> 17

<212> PRT

<213> Artificial

<220>

<223> murine heavy chain CDR2

<400> 4

Leu Ile Asn Pro Tyr Asn Gl y Asp Ser Phe Tyr Asn Gl n Lys Phe Lys
1 5 10 15

Gl y

<210> 5

AVB6-00112PC_ST25. txt

<211> 8
<212> PRT
<213> Arti fi ci al

<220>
<223> muri ne heavy chai n CDR3

<400> 5

Gl y Leu Arg Arg Asp Phe Asp Tyr
1 5

<210> 6
<211> 16
<212> PRT
<213> Arti fi ci al

<220>
<223> muri ne light chai n CDR1

<400> 6

Lys Ser Ser Gl n Ser Leu Leu Asp Ser Asp Gl y Lys Thr Tyr Leu Asn
1 5 10 15

<210> 7
<211> 7
<212> PRT
<213> Arti fi ci al

<220>
<223> muri ne light chai n CDR2

<400> 7

Leu Val Ser Gl u Leu Asp Ser
1 5

<210> 8
<211> 9
<212> PRT
<213> Arti fi ci al

<220>
<223> muri ne light chai n CDR3

<400> 8

Trp Gl n Gl y Thr His Phe Pro Arg Thr
1 5

<210> 9
<211> 117
<212> PRT
<213> Arti fi ci al

<220>
<223> HA heavy chai n vari abl e regi on

<400> 9

Gl n Val Gl n Leu Val Gl n Ser Gl y Ala Gl u Val Lys Lys Pro Gl y Ala
1 5 10 15

AVB6-00112PC_ST25. txt

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
20 25 30

Phe Met Asn Trp Val Arg Glu Ala Pro Gly Glu Gly Leu Glu Trp Met
35 40 45

Gly Leu Ile Asn Pro Tyr Asn Gly Asp Ser Phe Tyr Asn Glu Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Gly Leu Arg Arg Asp Phe Asp Tyr Trp Gly Glu Gly Thr Leu
100 105 110

Val Thr Val Ser Ser
115

<210> 10

<211> 113

<212> PRT

<213> Artificial

<220>

<223> Light chain variable region

<400> 10

Asp Val Val Met Thr Glu Ser Pro Leu Ser Leu Pro Val Thr Leu Gly
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Lys Ser Ser Glu Ser Leu Leu Asp Ser
20 25 30

Asp Gly Lys Thr Tyr Leu Asn Trp Phe Glu Glu Arg Pro Gly Glu Ser
35 40 45

Pro Arg Arg Leu Ile Tyr Leu Val Ser Glu Leu Asp Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Glu Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Trp Glu Glu
85 90 95

Thr His Phe Pro Arg Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

Arg

<210> 11
<211> 117
<212> PRT
<213> Artificial

<220>
<223> HB heavy chain variable region

<400> 11

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Ser Gly Tyr
20 25 30

Phe Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Leu Ile Asn Pro Tyr Asn Gly Asp Ser Phe Tyr Asn Gln Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Gly Leu Arg Arg Asp Phe Asp Tyr Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val Ser Ser
115

<210> 12
<211> 117
<212> PRT
<213> Artificial

<220>
<223> HF heavy chain variable region

<400> 12

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
20 25 30

Phe Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Leu Ile Asn Pro Tyr Asn Gly Asp Ser Phe Tyr Asn Gln Lys Phe
Page 5

50

55

60

Lys Gl y Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
 65 70 75 80

Met Gl u Leu Ser Ser Leu Arg Ser Gl u Asp Thr Al a Val Tyr Tyr Cys
 85 90 95

Val Arg Gl y Leu Arg Arg Asp Phe Asp Tyr Trp Gl y Gl n Gl y Thr Leu
 100 105 110

Val Thr Val Ser Ser
 115

<210> 13

<211> 117

<212> PRT

<213> Artifi cial

<220>

<223> HG heavy chain variable region

<400> 13

Gl n Val Gl n Leu Val Gl n Ser Gl y Al a Gl u Val Lys Lys Pro Gl y Al a
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Al a Ser Gl y Tyr Thr Phe Thr Gl y Tyr
 20 25 30

Phe Met Asn Trp Val Arg Gl n Al a Pro Gl y Gl n Gl y Leu Gl u Trp Met
 35 40 45

Gl y Leu Ile Asn Pro Tyr Asn Gl y Asp Ser Phe Tyr Asn Gl n Lys Phe
 50 55 60

Lys Gl y Arg Val Thr Met Thr Arg Gl n Thr Ser Thr Ser Thr Val Tyr
 65 70 75 80

Met Gl u Leu Ser Ser Leu Arg Ser Gl u Asp Thr Al a Val Tyr Tyr Cys
 85 90 95

Al a Arg Gl y Leu Arg Arg Asp Phe Asp Tyr Trp Gl y Gl n Gl y Thr Leu
 100 105 110

Val Thr Val Ser Ser
 115

<210> 14

<211> 117

<212> PRT

<213> Artifi cial

<220>

<223> HK heavy chain variable region

AVB6-00112PC_ST25. txt

<400> 14

Gl n Val Gl n Leu Val Gl n Ser Gl y Al a Gl u Val Lys Lys Pro Gl y Al a
1 5 10 15

Ser Val Lys Val Ser Cys Lys Al a Ser Gl y Tyr Thr Phe Thr Gl y Tyr
20 25 30

Phe Met Asn Trp Val Arg Gl n Al a Pro Gl y Gl n Gl y Leu Gl u Trp Met
35 40 45

Gl y Leu Ile Asn Pro Tyr Asn Gl y Asp Ser Phe Tyr Asn Gl n Lys Phe
50 55 60

Lys Gl y Arg Val Thr Met Thr Arg Asp Lys Ser Ser Ser Thr Al a Tyr
65 70 75 80

Met Gl u Leu Ser Ser Leu Arg Ser Gl u Asp Thr Al a Val Tyr Tyr Cys
85 90 95

Al a Arg Gl y Leu Arg Arg Asp Phe Asp Tyr Trp Gl y Gl n Gl y Thr Leu
100 105 110

Val Thr Val Ser Ser
115

<210> 15

<211> 117

<212> PRT
<213> Artificial

<220>

<223> HL heavy chain variable region

<400> 15

Gl n Val Gl n Leu Val Gl n Ser Gl y Al a Gl u Val Lys Lys Pro Gl y Al a
1 5 10 15

Ser Val Lys Val Ser Cys Lys Al a Ser Gl y Tyr Ser Phe Ser Gl y Tyr
20 25 30

Phe Met Asn Trp Val Arg Gl n Al a Pro Gl y Gl n Gl y Leu Gl u Trp Met
35 40 45

Gl y Leu Ile Asn Pro Tyr Asn Gl y Asp Ser Phe Tyr Asn Gl n Lys Phe
50 55 60

Lys Gl y Arg Val Thr Met Thr Arg Asp Lys Ser Ser Ser Thr Al a Tyr
65 70 75 80

Met Gl u Leu Ser Ser Leu Arg Ser Gl u Asp Thr Al a Val Tyr Tyr Cys
85 90 95

AVB6-00112PC_ST25. txt

Ala Arg Gly Leu Arg Arg Asp Phe Asp Tyr Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val Ser Ser
115

<210> 16
<211> 117
<212> PRT
<213> Artificial

<220>
<223> HM heavy chain variable region

<400> 16

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Ser Gly Tyr
20 25 30

Phe Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Leu Ile Asn Pro Tyr Asn Gly Asp Ser Phe Tyr Asn Gln Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Arg Gln Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Gly Leu Arg Arg Asp Phe Asp Tyr Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val Ser Ser
115

<210> 17
<211> 117
<212> PRT
<213> Artificial

<220>
<223> HN heavy chain variable region

<400> 17

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Ser Gly Tyr
20 25 30

AVB6-00112PC_ST25. txt

Phe Met Asn Trp Val Arg Glu Ala Pro Gly Glu Gly Leu Glu Trp Met
35 40 45

Gly Leu Ile Asn Pro Tyr Asn Gly Asp Ser Phe Tyr Asn Glu Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Arg Glu Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Gly Leu Arg Arg Asp Phe Asp Tyr Trp Gly Glu Gly Thr Leu
100 105 110

Val Thr Val Ser Ser
115

<210> 18

<211> 117

<212> PRT

<213> Artificial

<220>

<223> H0 heavy chain variable region

<400> 18

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Ser Gly Tyr
20 25 30

Phe Met Asn Trp Val Arg Glu Ala Pro Gly Glu Gly Leu Glu Trp Met
35 40 45

Gly Leu Ile Asn Pro Tyr Asn Gly Asp Ser Phe Tyr Asn Glu Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Arg Glu Lys Ser Thr Ser Thr Val Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Gly Leu Arg Arg Asp Phe Asp Tyr Trp Gly Glu Gly Thr Leu
100 105 110

Val Thr Val Ser Ser
115

<210> 19

AVB6-00112PC_ST25. txt

<211> 117
<212> PRT
<213> Artificial

<220>
<223> HQ heavy chain variable region

<400> 19

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
20 25 30

Phe Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Leu Ile Asn Pro Tyr Asn Gly Asp Ser Phe Tyr Asn Gln Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Gly Leu Arg Arg Asp Phe Asp Tyr Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val Ser Ser
115

<210> 20
<211> 117
<212> PRT
<213> Artificial

<220>
<223> HR heavy chain variable region

<400> 20

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Ser Gly Tyr
20 25 30

Phe Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Leu Ile Asn Pro Tyr Asn Gly Asp Ser Phe Tyr Asn Gln Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Ser Thr Ala Tyr
Page 10

65

70

75

80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Gly Leu Arg Arg Asp Phe Asp Tyr Trp Gly Gln Gly Thr Leu
 100 105 110

Val Thr Val Ser Ser
 115

<210> 21
 <211> 117
 <212> PRT
 <213> Artificial

<220>
 <223> HS heavy chain variable region

<400> 21

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Ser Gly Tyr
 20 25 30

Phe Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

Gly Leu Ile Asn Pro Tyr Asn Gly Asp Ser Phe Tyr Asn Gln Lys Phe
 50 55 60

Lys Gly Arg Val Thr Met Thr Arg Gln Thr Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Gly Leu Arg Arg Asp Phe Asp Tyr Trp Gly Gln Gly Thr Leu
 100 105 110

Val Thr Val Ser Ser
 115

<210> 22
 <211> 117
 <212> PRT
 <213> Artificial

<220>
 <223> HT heavy chain variable region

<400> 22

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 Page 11

1

5

10

15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Ser Gly Tyr
 20 25 30

Phe Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

Gly Leu Ile Asn Pro Tyr Asn Gly Asp Ser Phe Tyr Asn Gln Lys Phe
 50 55 60

Lys Gly Arg Val Thr Met Thr Arg Gln Thr Ser Thr Ser Thr Val Tyr
 65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Val Arg Gly Leu Arg Arg Asp Phe Asp Tyr Trp Gly Gln Gly Thr Leu
 100 105 110

Val Thr Val Ser Ser
 115

<210> 23

<211> 117

<212> PRT

<213> Artificial

<220>

<223> HU heavy chain variable region

<400> 23

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Ser Phe Ser Gly Tyr
 20 25 30

Phe Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

Gly Leu Ile Asn Pro Tyr Asn Gly Asp Ser Phe Tyr Asn Gln Lys Phe
 50 55 60

Lys Gly Arg Val Thr Met Thr Arg Gln Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Val Arg Gly Leu Arg Arg Asp Phe Asp Tyr Trp Gly Gln Gly Thr Leu
 100 105 110

AVB6-00112PC_ST25. txt

Val Thr Val Ser Ser
115

<210> 24
<211> 113
<212> PRT
<213> Artificial

<220>
<223> LB Light chain variable region

<400> 24

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly
1 5 10 15

Gln Pro Ala Ser Leu Phe Cys Lys Ser Ser Gln Ser Leu Leu Asp Ser
20 25 30

Asp Gly Lys Thr Tyr Leu Asn Trp Phe Gln Gln Arg Pro Gly Gln Ser
35 40 45

Pro Arg Arg Leu Ile Tyr Leu Val Ser Glu Leu Asp Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Trp Gln Gly
85 90 95

Thr His Phe Pro Arg Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

Arg

<210> 25
<211> 113
<212> PRT
<213> Artifical

<220>
<223> LC Light chain variable region

<400> 25

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly
1 5 10 15

Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Asp Ser
20 25 30

Asp Gly Lys Thr Tyr Leu Asn Trp Leu Phe Gln Arg Pro Gly Gln Ser
35 40 45

AVB6-00112PC_ST25. txt

Pro Arg Arg Leu Ile Tyr Leu Val Ser Glu Leu Asp Ser Glu Val Pro
50 55 60

Asp Arg Phe Ser Glu Ser Glu Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Glu Val Tyr Tyr Cys Trp Glu Glu
85 90 95

Thr His Phe Pro Arg Thr Phe Glu Glu Thr Lys Leu Glu Ile Lys
100 105 110

Arg

<210> 26

<211> 113

<212> PRT

<213> Artificial

<220>

<223> LD Light chain variable region

<400> 26

Asp Val Val Met Thr Glu Ser Pro Leu Ser Leu Pro Val Thr Leu Glu
1 5 10 15

Glut Pro Ala Ser Ile Ser Cys Lys Ser Ser Glu Ser Leu Leu Asp Ser
20 25 30

Asp Glu Lys Thr Tyr Leu Asn Trp Phe Glu Glu Arg Pro Glu Glu Ser
35 40 45

Pro Lys Arg Leu Ile Tyr Leu Val Ser Glu Leu Asp Ser Glu Val Pro
50 55 60

Asp Arg Phe Ser Glu Ser Glu Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Glu Val Tyr Tyr Cys Trp Glu Glu
85 90 95

Thr His Phe Pro Arg Thr Phe Glu Glu Thr Lys Leu Glu Ile Lys
100 105 110

Arg

<210> 27

<211> 113

<212> PRT

<213> Artificial

AVB6-00112PC_ST25. txt

<220>

<223> LE I ght chain variable region

<400> 27

Asp Val Val Met Thr Glu Ser Pro Leu Ser Leu Pro Val Thr Leu Glu
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Lys Ser Ser Glu Ser Leu Leu Asp Ser
20 25 30

Asp Glu Lys Thr Tyr Leu Asn Trp Phe Glu Glu Arg Pro Glu Glu Ser
35 40 45

Pro Arg Arg Leu Ile Tyr Leu Val Ser Glu Leu Asp Ser Glu Val Pro
50 55 60

Asp Arg Phe Thr Glu Ser Glu Ser Glu Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Glu Val Tyr Tyr Cys Trp Glu Glu
85 90 95

Thr His Phe Pro Arg Thr Phe Glu Glu Glu Thr Lys Leu Glu Ile Lys
100 105 110

Arg

<210> 28

<211> 113

<212> PRT

<213> Artificial

<220>

<223> LF I ght chain variable region

<400> 28

Asp Val Val Met Thr Glu Ser Pro Leu Ser Leu Pro Val Thr Leu Glu
1 5 10 15

Glu Pro Ala Ser Leu Phe Cys Lys Ser Ser Glu Ser Leu Leu Asp Ser
20 25 30

Asp Glu Lys Thr Tyr Leu Asn Trp Leu Phe Glu Glu Arg Pro Glu Glu Ser
35 40 45

Pro Lys Arg Leu Ile Tyr Leu Val Ser Glu Leu Asp Ser Glu Val Pro
50 55 60

Asp Arg Phe Thr Glu Ser Glu Ser Glu Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Glu Val Tyr Tyr Cys Trp Glu Glu

Thr His Phe Pro Arg Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110

Arg

<210> 29
 <211> 113
 <212> PRT
 <213> Artificial

<220>
 <223> LG light chain variable region

<400> 29

Asp Val Val Leu Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Leu Gly
 1 5 10 15

Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Asp Ser
 20 25 30

Asp Gly Lys Thr Tyr Leu Asn Trp Leu Phe Gln Arg Pro Gly Gln Ser
 35 40 45

Pro Arg Arg Leu Ile Tyr Leu Val Ser Glu Leu Asp Ser Gly Val Pro
 50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gln Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Trp Gln Gly
 85 90 95

Thr His Phe Pro Arg Thr Phe Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110

Arg

<210> 30
 <211> 330
 <212> PRT
 <213> Artificial

<220>
 <223> human IgG1 heavy chain constant region

<400> 30

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 Page 16

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
 100 105 110

Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
 115 120 125

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
 130 135 140

Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
 145 150 155 160

Tyr Val Asp Glu Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
 165 170 175

Gl u Gl n Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
 180 185 190

His Gl n Asp Trp Leu Asn Gl y Lys Gl u Tyr Lys Cys Lys Val Ser Asn
 195 200 205

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gl y
 210 215 220

Gl n Pro Arg Gl u Pro Gl n Val Tyr Thr Leu Pro Pro Ser Arg Asp Gl u
 225 230 235 240

Leu Thr Lys Asn Gl n Val Ser Leu Thr Cys Leu Val Lys Gl y Phe Tyr
 245 250 255

Pro Ser Asp Ile Ala Val Gl u Trp Gl u Ser Asn Gl y Gl n Pro Gl u Asn
 260 265 270

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gl y Ser Phe Phe
 275 280 285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gl n Gl n Gl y Asn
 Page 17

290

295

300

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
 305 310 315 320

Gl n Lys Ser Leu Ser Leu Ser Pro Gl y Lys
 325 330

<210> 31
 <211> 106
 <212> PRT
 <213> Artificial

<220>

<223> human kappa light chain constant domain

<400> 31

Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Gl u Gl n
 1 5 10 15

Leu Lys Ser Gl y Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
 20 25 30

Pro Arg Gl u Ala Lys Val Gl n Trp Lys Val Asp Asn Ala Leu Gl n Ser
 35 40 45

Gl y Asn Ser Gl n Gl u Ser Val Thr Gl u Gl n Asp Ser Lys Asp Ser Thr
 50 55 60

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Gl u Lys
 65 70 75 80

His Lys Val Tyr Ala Cys Gl u Val Thr His Gl n Gl y Leu Ser Ser Pro
 85 90 95

Val Thr Lys Ser Phe Asn Arg Gl y Gl u Cys
 100 105

<210> 32
 <211> 330
 <212> PRT
 <213> Artificial

<220>

<223> human IgG1 heavy chain constant region S239C mutant

<400> 32

Al a Ser Thr Lys Gl y Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15

Ser Thr Ser Gl y Gl y Thr Ala Ala Leu Gl y Cys Leu Val Lys Asp Tyr
 20 25 30

Phe Pro Gl u Pro Val Thr Val Ser Trp Asn Ser Gl y Ala Leu Thr Ser
 Page 18

AVB6-00112PC_ST25. txt
35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65 70 75 80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
85 90 95

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
100 105 110

Pro Ala Pro Glu Leu Leu Gly Gly Pro Cys Val Phe Leu Phe Pro Pro
115 120 125

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
130 135 140

Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
145 150 155 160

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
165 170 175

Gl u Gl n Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
180 185 190

His Gl n Asp Trp Leu Asn Gly Lys Gl u Tyr Lys Cys Lys Val Ser Asn
195 200 205

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
210 215 220

Gl n Pro Arg Gl u Pro Gl n Val Tyr Thr Leu Pro Pro Ser Arg Asp Gl u
225 230 235 240

Leu Thr Lys Asn Gl n Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
245 250 255

Pro Ser Asp Ile Ala Val Gl u Trp Gl u Ser Asn Gly Gl n Pro Gl u Asn
260 265 270

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
275 280 285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gl n Gl n Gly Asn
290 295 300

Val Phe Ser Cys Ser Val Met His Gl u Ala Leu His Asn His Tyr Thr
Page 19

305

310

315

320

Gl n Lys Ser Leu Ser Leu Ser Pro Gl y Lys
 325 330

<210> 33
 <211> 19
 <212> PRT
 <213> Arti fi ci al

<220>
 <223> muri ne heavy chain vari able region si gnal pepti de

<400> 33

Met Gl y Trp Ser Cys Ile Phe Leu Phe Leu Leu Ser Val Thr Val Gl y
 1 5 10 15

Val Phe Ser

<210> 34
 <211> 19
 <212> PRT
 <213> Arti fi ci al

<220>
 <223> human heavy chain vari abl e region si gnal pepti de

<400> 34

Met Al a Trp Val Trp Thr Leu Leu Phe Leu Met Al a Al a Al a Gl n Ser
 1 5 10 15

Al a Gl n Al a

<210> 35
 <211> 19
 <212> PRT
 <213> Arti fi ci al

<220>
 <223> muri ne l ight chain vari able region si gnal pepti de

<400> 35

Met Ser Pro Al a Gl n Phe Leu Phe Leu Leu Val Leu Trp Ile Arg Gl u
 1 5 10 15

Thr Asn Gl y

<210> 36
 <211> 19
 <212> PRT
 <213> Arti fi ci al

<220>
 <223> human l ight chain vari abl e region si gnal pepti de

AVB6-00112PC_ST25. txt

<400> 36

Met Lys Leu Pro Val Arg Leu Leu Val Leu Met Phe Trp Ile Pro Ala
1 5 10 15

Ser Ser Ser

<210> 37

<211> 351

<212> DNA

<213> Artificial

<220>

<223> mouse heavy chain variable region

<400> 37

gaggttcagc tgcagcagtc tggacctgag ttgggtgaaac ctggggcttc agtgaagata 60
tcctgcaagg cttctggta ttcatttgt ggctacttta tgaactgggt gaaacagagc 120
catggacaga gccttgagtg gatggactt attaatcctt acaatggaga ctcttttac 180
aatcagaagt tcaagggcaa gccacattt actgtacaga agtcctctag tacagccac 240
atggaactcc agagcctgac atctgaagac tctgcagtct ttattgtgt tagagggtta 300
cgacgggact ttgactattt gggcaaggc acccctctca cagtctcctc a 351

<210> 38

<211> 339

<212> DNA

<213> Artificial

<220>

<223> mouse light chain variable region

<400> 38

gatgttgtt tgacccagat tccactca tttgtcggtt ccattggaca accagcctcc 60
ctcttttta agtcaagtca gagccttta gatagtgtatg gaaagacata tttgaattgg 120
ttatttcaga ggccaggcca gtctccaaag cgccttattt atctgggttc tgaactggac 180
tctggagtcc ctgacagggtt cactggcagt ggatcaggga cagatttcac actgaaaatc 240
agcagagtgg aggctgagga tttggagtt tattattgtct ggcaaggtac acatttcct 300
cggacgttcg gtggaggcac caagctggaa atcaaacgg 339

<210> 39

<211> 351

<212> DNA

<213> Artificial

<220>

<223> HA heavy chain variable region

<400> 39

caggtgcagc tggtcagtc tggagctgag gtgaagaagc ctggagcctc tgtgaaggtg 60
tcctgttaagg cctctggcta caccttcaca ggctacttca tgaactgggt gaggcaggcc 120

AVB6-00112PC_ST25.txt

cctggccagg	gcctggagtg	gatgggcctg	atcaaccctt	acaatggaga	ctccttctac	180
aaccagaagt	tcaagggcag	ggtgaccatg	accaggaca	cctccacctc	cacagtgtac	240
atggagctgt	cctccctgag	gtctgaggac	acagctgtgt	actactgtgc	cagggcctg	300
aggagggact	ttgactactg	ggccagggc	accctggta	cagtgtcctc	c	351

<210> 40
 <211> 351
 <212> DNA
 <213> Artificial

<220>
 <223> HB heavy chain variable region

<400> 40	caggtgcagc	tgggcagtc	tggagctgag	gtgaagaagc	ctggagcctc	tgtgaaggtg	60
tcctgttaagg	cctctggcta	cagcttctct	ggctacttca	tgaactgggt	gaggcaggcc	120	
cctggccagg	gcctggagtg	gatgggcctg	atcaaccctt	acaatggaga	ctccttctac	180	
aaccagaagt	tcaagggcag	ggtgaccatg	accaggaca	cctccacctc	cacagtgtac	240	
atggagctgt	cctccctgag	gtctgaggac	acagctgtgt	actactgtgc	cagggcctg	300	
aggagggact	ttgactactg	ggccagggc	accctggta	cagtgtcctc	c	351	

<210> 41
 <211> 351
 <212> DNA
 <213> Artificial

<220>
 <223> HL heavy chain variable region

<400> 41	caggtgcagc	tgggcagtc	tggagctgag	gtgaagaagc	ctggagcctc	tgtgaaggtg	60
tcctgttaagg	cctctggcta	cagcttctct	ggctacttca	tgaactgggt	gaggcaggcc	120	
cctggccagg	gcctggagtg	gatgggcctg	atcaaccctt	acaatggaga	ctccttctac	180	
aaccagaagt	tcaagggcag	ggtgaccatg	accaggaca	agtccctc	cacagttac	240	
atggagctgt	cctccctgag	gtctgaggac	acagctgtgt	actactgtgc	cagggcctg	300	
aggagggact	ttgactactg	ggccagggc	accctggta	cagtgtcctc	c	351	

<210> 42
 <211> 351
 <212> DNA
 <213> Artificial

<220>
 <223> HT heavy chain variable region

<400> 42	caggtgcagc	tgggcagtc	tggagctgag	gtgaagaagc	ctggagcctc	tgtgaaggtg	60
tcctgttaagg	cctctggcta	cagcttctct	ggctacttca	tgaactgggt	gaggcaggcc	120	
cctggccagg	gcctggagtg	gatgggcctg	atcaaccctt	acaatggaga	ctccttctac	180	

AVB6-00112PC_ST25.txt

aaccagaagt	tcaagggcag	ggtgaccatg	accaggcaga	cctccacctc	cacagtgtac	240
atggagctgt	cctccctgag	gtctgaggac	acagctgtgt	actactgtgt	cagggcctg	300
aggagggact	ttgactactg	gggccagggc	accctggta	cagtgtcctc	c	351
<210>	43					
<211>	351					
<212>	DNA					
<213>	Artificial					
<220>						
<223>	HN heavy chain variable region					
<400>	43					
caggtgcagc	tggcagtc	tggagctgag	gtgaagaagc	ctggagcctc	tgtgaaggtg	60
tcctgtaagg	cctctggcta	cagcttctct	ggctacttca	tgaactgggt	gaggcaggcc	120
cctggccagg	gcctggagtg	gatgggcctg	atcaaccctt	acaatggaga	ctccttctac	180
aaccagaagt	tcaagggcag	ggtgaccatg	accaggcaga	cctccacctc	cacagtgtac	240
atggagctgt	cctccctgag	gtctgaggac	acagctgtgt	actactgtgc	cagggcctg	300
aggagggact	ttgactactg	gggccagggc	accctggta	cagtgtcctc	c	351
<210>	44					
<211>	351					
<212>	DNA					
<213>	Artificial					
<220>						
<223>	HU heavy chain variable region					
<400>	44					
caggtgcagc	tggcagtc	tggagctgag	gtgaagaagc	ctggagcctc	tgtgaaggtg	60
tcctgtaagg	cctctggcta	cagcttctct	ggctacttca	tgaactgggt	gaggcaggcc	120
cctggccagg	gcctggagtg	gatgggcctg	atcaaccctt	acaatggaga	ctccttctac	180
aaccagaagt	tcaagggcag	ggtgaccatg	accaggcaga	agtccctcctc	cacagcttac	240
atggagctgt	cctccctgag	gtctgaggac	acagctgtgt	actactgtgt	cagggcctg	300
aggagggact	ttgactactg	gggccagggc	accctggta	cagtgtcctc	c	351
<210>	45					
<211>	339					
<212>	DNA					
<213>	Artificial					
<220>						
<223>	LA light chain variable region					
<400>	45					
gatgtggta	tgaccaggc	ccctctgtcc	ctgcctgtga	ccctgggcca	gcctgcctcc	60
atctccctgt	agtccctcca	gtccctgctg	gactctgatg	gcaagaccta	cctgaactgg	120
ttccacggaga	ggcctggcca	gtcccctagg	aggctgatct	acctgggtgc	tgagctggac	180
tctggagtgc	ctgacagggtt	ctctggctct	ggctctggca	cagacttcac	cctgaagatc	240

AVB6-00112PC_ST25.txt

tccagggtgg	aggctgagga	tgtggagtg	tactactgtt	ggcagggcac	ccactccct	300
aggaccttg	gaggtggaac	caagctggag	atcaagcgt			339
<210> 46						
<211> 339						
<212> DNA						
<213> Artificial						
<220>						
<223> LC Light chain variable region						
<400> 46						
gatgtggta	tgacccagtc	ccctctgtcc	ctgcctgtga	ccctgggcca	gcctgcctcc	60
atctcctgta	agtccccc	gtccctgctg	gactctgatg	gcaagaccta	cctgaactgg	120
ctgttccaga	ggcctggcca	gtcccctagg	aggctgatct	acctgggtgc	tgagctggac	180
tctggagtgc	ctgacagggtt	ctctggctct	ggctctggca	cagacttcac	cctgaagatc	240
tccagggtgg	aggctgagga	tgtggagtg	tactactgtt	ggcagggcac	ccactccct	300
aggaccttg	gaggtggaac	caagctggag	atcaagcgt			339
<210> 47						
<211> 339						
<212> DNA						
<213> Artificial						
<220>						
<223> LF Light chain variable region						
<400> 47						
gatgtggta	tgacccagtc	ccctctgtcc	ctgcctgtga	ccctgggcca	gcctgcctcc	60
ctgttctgta	agtccccc	gtccctgctg	gactctgatg	gcaagaccta	cctgaactgg	120
ctgttccaga	ggcctggcca	gtcccctaag	aggctgatct	acctgggtgc	tgagctggac	180
tctggagtgc	ctgacagggtt	cacaggctct	ggctctggca	cagacttcac	cctgaagatc	240
tccagggtgg	aggctgagga	tgtggagtg	tactactgtt	ggcagggcac	ccactccct	300
aggaccttg	gaggtggaac	caagctggag	atcaagcgt			339
<210> 48						
<211> 339						
<212> DNA						
<213> Artificial						
<220>						
<223> LG Light chain variable region						
<400> 48						
gatgtggtc	tgacccagtc	ccctctgtcc	ctgcctgtga	ccctgggcca	gcctgcctcc	60
atctcctgta	agtccccc	gtccctgctg	gactctgatg	gcaagaccta	cctgaactgg	120
ctgttccaga	ggcctggcca	gtcccctagg	aggctgatct	acctgggtgc	tgagctggac	180
tctggagtgc	ctgacagggtt	ctctggctct	ggctctggca	cagacttcac	cctgaagatc	240
tccagggtgg	aggctgagga	tgtggagtg	tactactgtt	ggcagggcac	ccactccct	300

aggaccttg gaggtggAAC caagctggAG atcaaggcGT

339

<210> 49
 <211> 57
 <212> DNA
 <213> Artificial

<220>
 <223> murine heavy chain variable region signal peptide

<400> 49
 atggatggA gctgtatctt tctttctc ctgtcagtaa ctgttaggtgt gttctct

57

<210> 50
 <211> 57
 <212> DNA
 <213> Artificial

<220>
 <223> human heavy chain variable region signal peptide

<400> 50
 atggcttggg tgtggacctt gctattcctg atggcagctg cccaaagtgc ccaagca

57

<210> 51
 <211> 57
 <212> DNA
 <213> Artificial

<220>
 <223> murine light chain variable region signal peptide

<400> 51
 atgagtcctg cccagttcct gttctgtta gtgctctgga ttcggaaac caacggt

57

<210> 52
 <211> 57
 <212> DNA
 <213> Artificial

<220>
 <223> human light chain variable region signal peptide

<400> 52
 atgaagttgc ctgttaggct gttggtgctg atgttctgga ttccctgcttc cagcagt

57

<210> 53
 <211> 990
 <212> DNA
 <213> Artificial

<220>
 <223> human IgG1 heavy chain constant region

<400> 53
 gctagcacca agggcccatc ggtttcccc ctggcacccct cctccaagag cacctctggg

60

ggcacagcgg ccctgggctg cctggtaag gactacttcc cgcgaaccgggt gacgggtgtcg

120

tggaaactcag ggcgcctgac cagcggcgtg cacaccttcc cggctgtcct acagtcctca

180

ggactctact ccctcagcag cgtggtgacc gtgcctcca gcagcttggg cacccagacc

240

AVB6-00112PC_ST25.txt

tacatctgca acgtgaatca caagcccagc aacaccaagg tggacaagaa agttgagccc	300
aaatcttgt acaaaaactca cacatgccca ccgtgcccag cacctgaact cctgggggga	360
ccgtcagtct tcctttccc cccaaaaccc aaggacaccc tcatgatctc ccggaccct	420
gaggtcacat gcgtggtggt ggacgtgagc cacgaagacc ctgaggtcaa gttcaactgg	480
tacgtggacg gcgtggaggt gcataatgcc aagacaaagc cgccggagga gcagtacaac	540
agcacgtacc gtgtggtcag cgtcctcacc gtcctgcacc aggactggct gaatggcaag	600
gagtacaagt gcaaggtctc caacaaagcc ctcccagccc ccatcgagaa aaccatctcc	660
aaagccaaag ggcagcccg agaaccacag gtgtacaccc tgcccccattc ccggatgag	720
ctgaccaaga accaggtcag cctgacctgc ctggtaaaag gcttctatcc cagcgacatc	780
gccgtggagt gggagagcaa tggcagccg gagaacaact acaagaccac gcctccgtg	840
ctggactccg acggctcctt cttcctctac agcaagctca ccgtggacaa gagcaggtgg	900
cagcagggga acgtttctc atgctccgtg atgcatgagg ctctgcacaa ccactacacg	960
cagaagagcc tctccctgtc tccggtaaa	990

<210> 54

<211> 318

<212> DNA

<213> Artificial

<220>

<223> human kappa light chain constant domain

<400> 54

actgtggcgg cgccatctgt cttcatcttc ccgcctatctg atgagcagtt gaaatctgga	60
actgcctctg ttgtgtgcct gctgaataac ttctatccca gagaggccaa agtacagtgg	120
aagggtggata acgcctcca atcggtaac tcccaggaga gtgtcacaga gcaggacagc	180
aaggacagca cctacagcct cagcagcacc ctgacgctga gcaaaggcaga ctacgagaaa	240
cacaaagtct acgcctgcga agtcacccat cagggcctga gctcgccgt cacaagagc	300
ttcaacaggg gagagtgt	318

<210> 55

<211> 990

<212> DNA

<213> Artificial

<220>

<223> human IgG1 heavy chain constant region S239C mutant

<400> 55

gctagcacca agggcccatc tgtttcccc ctggcaccct cctccaagag cacctctggg	60
ggcacagctg ccctggctg cctggtaag gactacttcc ctgaacctgt gacagtgtcc	120
tggaaacttag ggcgcctgac cagggcgtg cacacccccc cgctgtcct acagtccctca	180
ggactctact ccctcagcag cgtggtgacc gtgcctcca gcagcttggg cacccagacc	240
tacatctgca acgtgaatca caagcccagc aacaccaagg tggacaagaa agttgagccc	300

AVB6-00112PC_ST25.txt

aaatcttgtg acaaaaactca cacatgccca ccgtgcccag cacctgaact cctgggggga	360
ccgtgtgtct tcctttccc cccaaaaccc aaggacaccc tcatgatctc ccggaccct	420
gaggtcacat gcgtgggtggt ggacgtgagc cacgaagacc ctgaggtcaa gttcaactgg	480
tacgtggacg gcgtggaggt gcataatgcc aagacaaagc cgcgggagga gcagtacaac	540
agcacgttacc gtgtggttag cgtcctcacc gtcctgcacc aggactggct gaatggcaag	600
gagtacaagt gcaaggcttc caacaaagcc ctcccaagccc ccatcgagaa aaccatctcc	660
aaagccaaag ggcagcccg agaaccacag gtgtacaccc tgccccatc ccggatgag	720
ctgaccaaga accaggtcag cctgacctgc ctggtaaaag gcttctatcc cagcgacatc	780
gccgtggagt gggagagcaa tggcagccg gagaacaact acaagaccac gcctccgtg	840
ctggactccg acggctcctt cttcctctac agcaagctca ccgtggacaa gagcaggtgg	900
cagcagggga acgtttctc atgctccgtg atgcatgagg ctctgcacaa ccactacaca	960
cagaagagcc tctccctgtc tccggtaaa	990