

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2012201211 B2**

(54) Title
Microbicidal composition

(51) International Patent Classification(s)
A01N 43/78 (2006.01) **A01N 33/08** (2006.01)
A01N 29/04 (2006.01) **A01N 33/12** (2006.01)
A01N 31/02 (2006.01) **A01N 33/20** (2006.01)
A01N 31/04 (2006.01) **A01N 43/50** (2006.01)
A01N 31/08 (2006.01)

(21) Application No: **2012201211** (22) Date of Filing: **2012.02.29**

(43) Publication Date: **2012.03.22**

(43) Publication Journal Date: **2012.03.22**

(44) Accepted Journal Date: **2013.08.15**

(62) Divisional of:
2006222721

(71) Applicant(s)
Rohm and Haas Company

(72) Inventor(s)
Diehl, Megan Anne; Levy, Richard; Shaw, Dolores Ann; Warwick, Eileen Fleck

(74) Agent / Attorney
Phillips Ormonde Fitzpatrick, 367 Collins Street, Melbourne, VIC, 3000

(56) Related Art
JP 10 298012 A

ABSTRACT OF THE DISCLOSURE

Synergistic microbicidal compositions containing N-(n-butyl)-1,2-benzisothiazolin-3-one or N-methyl-1,2-benzisothiazolin-3-one.

2012201211 29 Feb 2012

2012201211 29 Feb 2012

AUSTRALIA

Patents Act

**COMPLETE SPECIFICATION
(ORIGINAL)**

	Class	Int. Class
Application Number: Lodged:		
Complete Specification Lodged: Accepted: Published:		
Priority		
Related Art:		

Name of Applicant:

Rohm and Haas Company

Actual Inventor(s):

Megan Anne Diehl, Richard Levy, Dolores Ann Shaw, Eileen Fleck Warwick

Address for Service and Correspondence:

PHILLIPS ORMONDE FITZPATRICK
Patent and Trade Mark Attorneys
367 Collins Street
Melbourne 3000 AUSTRALIA

Invention Title:

MICROBICIDAL COMPOSITION

Our Ref : 931059
POF Code: 496570/83886

The following statement is a full description of this invention, including the best method of performing it known to applicant(s):

MICROBICIDAL COMPOSITION

29 Feb 2012
2012201211
5 The present invention is a divisional application from Australian Patent Application Number 2006222721, which claims priority from European Patent Application Number 05292073.3, filed 4 October 2005, the entire disclosure of which is incorporated herein by reference.

This invention relates to a synergistic combination of selected microbicides having greater activity than would be observed for the individual microbicides.

0 In some cases, commercial microbicides cannot provide effective control of microorganisms, even at high use concentrations, due to weak activity against certain types of microorganisms, e.g., those resistant to some microbicides, or due to aggressive environmental conditions. Combinations of different microbicides are sometimes used to provide overall control of microorganisms in a particular end use environment. For example, U.S. Pat. App. Pub. No.
5 2004/0014799 discloses a synergistic combination of N-(n-butyl)-1,2-benzisothiazolin-3-one (BBIT) and 2-methyl-4-isothiazolin-3-one (MI) over the limited range of ratios of BBIT to MI of 10:1 to 1.67:1. However, there is a need for additional combinations of microbicides having enhanced activity against various strains of microorganisms to provide effective control of the
0 microorganisms. Moreover, there is a need for combinations containing lower levels of individual microbicides for environmental and economic benefit. The problem addressed by this invention is to provide such additional combinations of microbicides.

25 A reference herein to a patent document or other matter which is given as prior art is not to be taken as an admission that that document or matter was known or that the information it contains was part of the common general knowledge as at the priority date of any of the claims.

30 Throughout the description and claims of the specification, the word "comprise" and variations of the word, such as "comprising" and "comprises", is not intended to exclude other additives, components, integers or steps.

STATEMENT OF THE INVENTION

The present invention is directed to a microbicidal composition comprising: (a) N-(n-butyl)-1,2-benzisothiazolin-3-one; and (b) hexamidine diisethionate.

The present invention is further directed to a microbicidal composition comprising: (a) N-methyl-1,2-benzisothiazolin-3-one; and (b) hexamidine diisethionate.

DETAILED DESCRIPTION OF THE INVENTION

"BBIT" is N-(n-butyl)-1,2-benzisothiazolin-3-one. "EDTA" is ethylenediamine tetraacetic acid. "MBIT" is N-methyl-1,2-benzisothiazolin-3-one.

As used herein, the following terms have the designated definitions, unless the context clearly indicates otherwise. The term "microbicide" refers to a compound capable of killing, inhibiting the growth of or controlling the growth of microorganisms at a locus; microbicides include bactericides, fungicides and algacides. The term "microorganism" includes, for example, fungi (such as yeast and mold), bacteria and algae. The term "locus" refers to an industrial system or product subject to contamination by microorganisms. The following abbreviations are used throughout the specification: ppm = parts per million by weight (weight/weight), mL = milliliter, ATCC = American Type Culture Collection, MBC = minimum biocidal concentration, and MIC = minimum inhibitory concentration. Unless otherwise specified, temperatures are in degrees centigrade (°C), and references to percentages (%) are by weight.

Amounts of organic microbicides are given on an active ingredient basis in ppm (w/w).

The compositions of the present invention unexpectedly have been found to provide enhanced microbicidal efficacy at a combined active ingredient level lower than that of the individual microbicides. In one embodiment of the invention, those antimicrobial compositions which contain halogenated 3-isothiazolones contain relatively low levels thereof, preferably no more than 1000 ppm, more preferably no more than 500 ppm, more preferably no more than 100

2012201211 29 Feb 2012
5 ppm, and most preferably no more than 50 ppm. Concentrations of halogenated 3-isothiazolones in the composition of this invention are based on the total weight of active ingredients in the composition, i.e., the microbicides exclusive of any amounts of solvents, carriers, dispersants, stabilizers or other materials which may be present. In one embodiment of the invention, the antimicrobial composition contains less than 1000 ppm of 5-chloro-2-methyl-4-isothiazolin-3-one, more preferably no more than 500 ppm, more preferably no more than 100 ppm, and most preferably no more than 50 ppm.

0 In one embodiment of the invention, the antimicrobial composition comprises N-(n-butyl)-1,2-benzisothiazolin-3-one and hexamidine diisethionate. Preferably, a weight ratio of N-(n-butyl)-1,2-benzisothiazolin-3-one to hexamidine diisethionate is from 1:0.02 to 1:3.

5 In one embodiment of the invention, the antimicrobial composition comprises N-methyl-1,2-benzisothiazolin-3-one and hexamidine diisethionate. Preferably, a weight ratio of N-methyl-1,2-benzisothiazolin-3-one to hexamidine diisethionate is from 1:0.1 to 1:5, more preferably from 1:0.1 to 1:3.

0 The microbicides in the composition of this invention may be used "as is" or may first be formulated with a solvent or a solid carrier. Suitable solvents include, for example, water; glycols, such as ethylene glycol, propylene glycol, diethylene glycol, dipropylene glycol, polyethylene glycol, and polypropylene glycol; glycol ethers; alcohols, such as methanol, ethanol, propanol, phenethyl alcohol and phenoxypropanol; ketones, such as acetone and methyl ethyl ketone; esters, such as ethyl acetate, butyl acetate, triacetyl citrate, and glycerol triacetate; carbonates, such as propylene carbonate and dimethyl carbonate; and mixtures thereof. It is preferred that the solvent is selected from water, glycols, glycol ethers, esters and mixtures thereof. Suitable solid carriers include, for example, cyclodextrin, silicas, diatomaceous earth, waxes, cellulosic materials, alkali and alkaline earth (e.g., sodium, magnesium, potassium) metal salts (e.g., chloride, nitrate, bromide, sulfate) and charcoal.

25 30 When a microbicide component is formulated in a solvent, the formulation may optionally contain surfactants. When such formulations contain surfactants, they are generally in the form of emulsive concentrates, emulsions,

microemulsive concentrates, or microemulsions. Emulsive concentrates form emulsions upon the addition of a sufficient amount of water. Microemulsive concentrates form microemulsions upon the addition of a sufficient amount of water. Such emulsive and microemulsive concentrates are generally well known in the art; it is preferred that such formulations are free of surfactants. U.S. Patent No. 5,444,078 may be consulted for further general and specific details on the preparation of various microemulsions and microemulsive concentrates.

A microbicide component also can be formulated in the form of a dispersion. The solvent component of the dispersion can be an organic solvent or water, preferably water. Such dispersions can contain adjuvants, for example, co-solvents, thickeners, anti-freeze agents, dispersants, fillers, pigments, surfactants, biodispersants, sulfosuccinates, terpenes, furanones, polycations, stabilizers, scale inhibitors and anti-corrosion additives.

When both microbicides are each first formulated with a solvent, the solvent used for the first microbicide may be the same as or different from the solvent used to formulate the other commercial microbicide, although water is preferred for most industrial biocide applications. It is preferred that the two solvents are miscible.

Those skilled in the art will recognize that the microbicide components of the present invention may be added to a locus sequentially, simultaneously, or may be combined before being added to the locus. It is preferred that the first microbicide and the second microbicide component be added to a locus simultaneously or sequentially. When the microbicides are added simultaneously or sequentially, each may individual components may contain adjuvants, such as, for example, solvent, thickeners, anti-freeze agents, colorants, sequestrants (such as ethylenediamine-tetraacetic acid, ethylenediaminedisuccinic acid, iminodisuccinic acid and salts thereof), dispersants, surfactants, biodispersants, sulfosuccinates, terpenes, furanones, polycations, stabilizers, scale inhibitors and anti-corrosion additives.

The microbicidal compositions of the present invention can be used to inhibit the growth of microorganisms or higher forms of aquatic life (such as protozoans, invertebrates, bryozoans, dinoflagellates, crustaceans, mollusks, etc)

by introducing a microbicidally effective amount of the compositions onto, into, or at a locus subject to microbial attack. Suitable loci include, for example: industrial process water; electrocoat deposition systems; cooling towers; air washers; gas scrubbers; mineral slurries; wastewater treatment; ornamental fountains; reverse osmosis filtration; ultrafiltration; ballast water; evaporative condensers; heat exchangers; pulp and paper processing fluids and additives; starch; plastics; emulsions; dispersions; paints; latices; coatings, such as varnishes; construction products, such as mastics, caulks, and sealants; construction adhesives, such as ceramic adhesives, carpet backing adhesives, and laminating adhesives; industrial or consumer adhesives; photographic chemicals; printing fluids; household products, such as bathroom and kitchen cleaners; cosmetics; toiletries; shampoos; soaps; detergents; industrial cleaners; floor polishes; laundry rinse water; metalworking fluids; conveyor lubricants; hydraulic fluids; leather and leather products; textiles; textile products; wood and wood products, such as plywood, chipboard, wallboard, flakeboard, laminated beams, oriented strandboard, hardboard, and particleboard; petroleum processing fluids; fuel; oilfield fluids, such as injection water, fracture fluids, and drilling muds; agriculture adjuvant preservation; surfactant preservation; medical devices; diagnostic reagent preservation; food preservation, such as plastic or paper food wrap; food, beverage, and industrial process pasteurizers; toilet bowls; recreational water; pools; and spas.

Preferably, the microbicidal compositions of the present invention are used to inhibit the growth of microorganisms at a locus selected from one or more of mineral slurries, pulp and paper processing fluids and additives, starch, emulsions, dispersions, paints, latices, coatings, construction adhesives, such as ceramic adhesives, carpet backing adhesives, photographic chemicals, printing fluids, household products such as bathroom and kitchen cleaners, cosmetics, toiletries, shampoos, soaps, detergents, industrial cleaners, floor polishes, laundry rinse water, metal working fluids, textile products, agriculture adjuvant preservation, surfactant preservation, diagnostic reagent preservation, food preservation, and food, beverage, and industrial process pasteurizers.

29 Feb 2012
2012201211
5 The specific amount of the composition of this invention necessary to inhibit or control the growth of microorganisms and higher aquatic life forms in a locus depends upon the particular locus to be protected. Typically, the amount of the composition of the present invention to control the growth of microorganisms in a locus is sufficient if it provides from 0.1 to 1,000 ppm of the isothiazoline ingredient of the composition in the locus. It is preferred that the isothiazolone ingredients of the composition be present in the locus in an amount of at least 0.5 ppm, more preferably at least 4 ppm and most preferably at least 10 ppm. It is preferred that the isothiazolone ingredients of the composition be present in the locus in an amount of no more than 1000 ppm, more preferably no more than 500 ppm, and most preferably no more than 200 ppm.

0 In one embodiment of the invention, the composition is substantially free of enzymatic biocides. Preferably, when BBIT or MBIT and either methylparaben or ethylparaben are combined, the composition is substantially
5 free of enzymatic biocides. Enzymatic biocides are enzymes having activity against microbes, as defined, e.g., in U.S. Pat. App. Pub. No. 2002/0028754.

EXAMPLES

Materials and Methods

0 The synergism of the combination of the present invention was demonstrated by testing a wide range of concentrations and ratios of the compounds.

One measure of synergism is the industrially accepted method described by Kull, F.C.; Eisman, P.C.; Sylwestrowicz, H.D. and Mayer, R.L., in Applied
25 Microbiology 9:538-541 (1961), using the ratio determined by the formula:

$$Q_a/Q_A + Q_b/Q_B = \text{Synergy Index ("SI")}$$

wherein:

30 Q_A = concentration of compound A (first component) in ppm, acting alone, which produced an end point (MIC of Compound A).

Q_a = concentration of compound A in ppm, in the mixture, which produced an end point.

Q_B = concentration of compound B (second component) in ppm, acting alone, which produced an end point (MIC of Compound B).

Q_b = concentration of compound B in ppm, in the mixture, which produced an end point.

When the sum of Q_a/Q_A and Q_b/Q_B is greater than one, antagonism is indicated. When the sum is equal to one, additivity is indicated, and when less than one, synergism is demonstrated. The lower the SI, the greater the synergy shown by that particular mixture. The minimum inhibitory concentration (MIC) of a microbicide is the lowest concentration tested under a specific set of conditions that prevents the growth of added microorganisms.

Synergy tests were conducted using standard microtiter plate assays with media designed for optimal growth of the test microorganism. Minimal salt medium supplemented with 0.2% glucose and 0.1% yeast extract (M9GY medium) was used for testing bacteria; Potato Dextrose Broth (PDB medium) was used for testing yeast and mold. In this method, a wide range of combinations of microbicides and other personal care raw materials was tested by conducting high resolution MIC assays in the presence of various concentrations of BBIT or MBIT. High resolution MICs were determined by adding varying amounts of microbicide to one column of a microtitre plate and doing subsequent ten-fold dilutions using an automated liquid handling system to obtain a series of endpoints ranging from 2 ppm to 10,000 ppm active ingredient.

The synergy of the combinations of the present invention was determined against a bacterium, *Escherichia coli* (*E. coli* - ATCC #8739), a yeast, *Candida albicans* (*C. albicans* - ATCC 10231), and a mold, *Aspergillus niger* (*A. niger* - ATCC 16404). The bacteria were used at a concentration of about 5×10^6 bacteria per mL and the yeast and mold at 5×10^5 fungi per mL. These microorganisms are representative of natural contaminants in many consumer and industrial applications. The plates were visually evaluated for microbial growth (turbidity) to determine the MIC after various incubation times at 25° C (yeast and mold) or 30° C (bacteria).

2012201211 29 Feb 2012

The test results for demonstration of synergy of the BBIT combinations of the present invention are shown below in Tables 1 through 26. In each test, First Component (A) was BBIT and the Second Component (B) was the other commercial microbicide. Each table shows the specific combinations of BBIT and the second component; results against the microorganisms tested with incubation times; the end-point activity in ppm measured by the MIC for BBIT alone (Q_A), for the second component alone (Q_B), for BBIT in the mixture (Q_a) and for second component in the mixture (Q_b); the calculated SI value; and the range of synergistic ratios for each combination tested (BBIT/second component or A/B).

The test results for demonstration of synergy of the MBIT combinations of the present invention are shown below in Tables 27 through 54. In each test, First Component (A) was MBIT and the Second Component (B) was the other commercial microbicide. Each table shows the specific combinations of MBIT and the second component; results against the microorganisms tested with incubation times; the end-point activity in ppm measured by the MIC for MBIT alone (Q_A), for the second component alone (Q_B), for MBIT in the mixture (Q_a) and for second component in the mixture (Q_b); the calculated SI value; and the range of synergistic ratios for each combination tested (MBIT/second component or A/B).

Table 1

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)

5 Second Component (B) = Benzalkonium chloride (Hyamine 3500)

Microorganism	Q_a	Q_b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	3	1.00	-----
(48 hours)	16.667	40	1.00	1/2
	10	50	1.00	1/5
	12.5	60	1.00	1/5
	100	0	1.00	-----
Microorganism	Q_a	Q_b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	60	1.00	-----
(24 hours)	2	4	0.40	1/2
	2	5	0.42	1/3

Microorganism	Q _a	Q _b	SI	A/B
	2	6	0.43	1/3
	2	8	0.47	1/4
	2	10	0.50	1/5
	2	20	0.67	1/10
	2	30	0.83	1/15
	4	1	0.68	1/0.3
	4	2	0.70	1/0.5
	4	3	0.72	1/0.8
	4	4	0.73	1/1
	4	5	0.75	1/1
	4	6	0.77	1/2
	4	8	0.80	1/2
	4	10	0.83	1/3
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	200	1.00	-----
(7 days)	5	60	0.55	1/12
	5	80	0.65	1/16
	5	100	0.75	1/20
	10	50	0.75	1/5
	10	60	0.80	1/6
	10	80	0.90	1/8
	15	4	0.77	1/0.3
	15	5	0.78	1/0.3
	15	6	0.78	1/0.4
	15	8	0.79	1/0.5
	15	10	0.80	1/0.7
	15	20	0.85	1/1
	15	30	0.90	1/2
	20	0	1.00	-----

The ratios of BBIT/benzalkonium chloride tested ranged from 1/0.02 to 1/500.

Table 2

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = Benzethonium chloride (Hyamine 1622)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	20	1.00	-----
(72 hours)	40	6	0.70	1/0.2
	40	8	0.80	1/0.2
	40	10	0.90	1/0.3
	50	3	0.65	1/0.06
	50	4	0.70	1/0.08
	50	5	0.75	1/0.1
	50	6	0.80	1/0.1
	50	8	0.90	1/0.2
	60	1	0.65	1/0.02
	60	2	0.70	1/0.03
	60	3	0.75	1/0.05
	60	4	0.80	1/0.07
	60	5	0.85	1/0.08
	60	6	0.90	1/0.1
	70	4	0.90	1/0.06
	100	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	4	1.00	-----
(24 hours)	4	1	0.92	1/0.3
	6	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	10	1.00	-----
(7 days)	5	4	0.65	1/0.8
	5	5	0.75	1/0.8
	5	6	0.85	1/1
	10	4	0.90	1/0.4
	20	0	1.00	-----

The ratios of BBIT/benzethonium chloride tested ranged from 1/0.02 to 1/500.

Table 3

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = Benzyl alcohol

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	2000	1.00	-----
(24 hours)	100	2000	0.90	1/20
	200	0	1.00	----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	3000	1.00	-----
(48 hours)	2.5	600	0.45	1/240
	2.5	800	0.52	1/320
	2.5	1000	0.58	1/400
	2.5	2000	0.92	1/800
	5	400	0.63	1/80
	5	500	0.67	1/100
	5	600	0.70	1/120
	5	800	0.77	1/160
	5	1000	0.83	1/200
	7.5	20	0.76	1/3
	7.5	30	0.76	1/4
	7.5	40	0.76	1/5
	7.5	50	0.77	1/7
	7.5	60	0.77	1/8
	7.5	80	0.78	1/11
	7.5	100	0.78	1/13
	7.5	200	0.82	1/27
	7.5	300	0.85	1/40
	7.5	400	0.88	1/53
	7.5	500	0.92	1/67
	7.5	600	0.95	1/80
	10	0	1.00	----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	5000	1.00	-----
(3 days)	5	5000	1.13	1/1000
	10	5000	1.25	1/500
	15	4000	1.18	1/267
	20	5000	1.50	1/250

Microorganism	Q _a	Q _b	SI	A/B
	40	0	1.00	----

The ratios of BBIT/benzyl alcohol tested ranged from 1/0.2 to 1/5000.

Table 4

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = Benzisothiazolinone (BIT)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	8	1.00	-----
(72 hours)	40	1	0.53	1/0.03
	40	2	0.65	1/0.05
	40	3	0.78	1/0.08
	40	4	0.90	1/0.1
	50	1	0.63	1/0.02
	50	2	0.75	1/0.4
	50	3	0.88	1/0.06
	60	1	0.73	1/0.02
	60	2	0.85	1/0.03
	100	0	1.00	----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	30	1.00	-----
(48 hours)	2.5	8	0.60	1/3
	2.5	10	0.67	1/4
	5	3	0.77	1/0.6
	5	4	0.80	1/0.8
	5	5	0.83	1/1
	5	6	0.87	1/1.2
	5	8	0.93	1/1.6
	7.5	0	1.00	----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	100	1.00	-----
(7 days)	5	80	0.86	1/16
	10	20	0.33	1/2
	10	30	0.43	1/3
	10	40	0.53	1/4
	10	50	0.63	1/5
	10	60	0.73	1/6

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
	10	80	0.93	1/8
	15	8	0.27	1/0.5
	15	10	0.29	1/0.7
	15	20	0.39	1/1
	15	30	0.49	1/2
	15	40	0.59	1/3
	15	50	0.69	1/3
	15	60	0.79	1/4
	20	10	0.35	1/0.5
	20	20	0.45	1/1
	20	30	0.55	1/2
	20	40	0.65	1/2
	20	50	0.75	1/3
	20	60	0.85	1/3
	40	2	0.52	1/0.05
	40	3	0.53	1/0.08
	40	4	0.54	1/0.1
	40	5	0.55	1/0.1
	40	6	0.56	1/0.1
	40	8	0.58	1/0.2
	40	10	0.60	1/0.3
	40	20	0.70	1/0.5
	40	30	0.80	1/0.8
	40	40	0.90	1/1
	80	0	1.00	----

The ratios of BBIT/benzisothiazolinone tested ranged from 1/0.02 to 1/500.

5

Table 5

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = 2-bromo-2-nitro-propane-1,3-diol (BNPD)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	10	1.00	-----
(48 hours)	40	1	0.50	1/0.03
	40	2	0.60	1/0.05
	40	3	0.70	1/0.08
	40	4	0.80	1/0.1
	40	5	0.90	1/0.1

Microorganism	Q _a	Q _b	SI	A/B
	50	1	0.60	1/0.02
	50	2	0.70	1/0.04
	50	3	0.80	1/0.06
	50	4	0.90	1/0.08
	70	1	0.80	1/0.01
	70	2	0.90	1/0.03
	80	1	0.90	1/0.01
	100	0	1.00	----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	2000	1.00	-----
(48 hours)	2	1000	0.75	1/500
	4	20	0.51	1/5
	4	30	0.52	1/8
	4	40	0.52	1/10
	4	50	0.53	1/13
	4	60	0.53	1/15
	4	80	0.54	1/20
	4	100	0.55	1/25
	4	200	0.60	1/50
	4	300	0.65	1/75
	4	400	0.70	1/100
	4	500	0.75	1/125
	4	600	0.80	1/150
	4	800	0.90	1/200
	6	1	0.75	1/0.2
	6	2	0.75	1/0.3
	6	3	0.75	1/0.5
	6	4	0.75	1/0.7
	6	5	0.75	1/0.8
	6	6	0.75	1/1
	6	8	0.75	1/1
	6	10	0.76	1/2
	6	20	0.76	1/3
	6	30	0.77	1/5
	6	40	0.77	1/7
	6	50	0.78	1/8
	6	60	0.78	1/10
	6	80	0.79	1/13

Microorganism	Q _a	Q _b	SI	A/B
	6	100	0.80	1/17
	6	200	0.85	1/33
	6	300	0.90	1/50
	8	0	1.00	----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	2000	1.00	-----
(7 days)	5	1000	0.75	1/200
	10	8	0.50	1/0.8
	10	10	0.51	1/1
	10	20	0.51	1/2
	10	30	0.52	1/3
	10	40	0.52	1/4
	10	50	0.53	1/5
	10	60	0.53	1/6
	10	80	0.54	1/8
	10	100	0.55	1/10
	10	200	0.60	1/20
	10	300	0.65	1/30
	10	400	0.70	1/40
	10	500	0.75	1/50
	10	600	0.80	1/60
	10	800	0.90	1/80
	20	0	1.00	----

The ratios of BBIT/2-bromo-2-nitro-propane-1,3-diol tested ranged from 1/0.01 to 1/500.

5

Table 6

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
 Second Component (B) = Butylene glycol

10

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	20000	1.00	-----
(48 hours)	40	5000	0.65	1/125
	40	6000	0.70	1/150
	40	8000	0.80	1/200
	40	10000	0.90	1/250
	50	2000	0.60	1/40
	50	3000	0.65	1/60
	50	4000	0.70	1/80

Microorganism	Q _a	Q _b	SI	A/B
	50	5000	0.75	1/100
	50	6000	0.80	1/120
	50	8000	0.90	1/160
	70	2000	0.80	1/29
	70	3000	0.85	1/43
	70	4000	0.90	1/57
	80	200	0.81	1/3
	80	300	0.82	1/4
	80	400	0.82	1/5
	80	500	0.83	1/6
	80	600	0.83	1/8
	80	800	0.84	1/10
	80	1000	0.85	1/13
	80	2000	0.90	1/25
	100	0	1.00	----
Microorganism	Q _a	Q _b	SI	A/B
C. albicans 10231 · PDB	0	20000	1.00	----
(24 hours)	2	10000	0.83	1/5000
	4	200	0.68	1/50
	4	300	0.68	1/75
	4	400	0.69	1/100
	4	500	0.69	1/125
	4	600	0.70	1/150
	4	800	0.71	1/200
	4	1000	0.72	1/250
	4	2000	0.77	1/500
	4	3000	0.82	1/750
	4	4000	0.87	1/1000
	4	5000	0.92	1/1250
	6	0	1.00	----
Microorganism	Q _a	Q _b	SI	A/B
A. niger 16404 · PDB	0	10000	1.00	----
(3 days)	5	5000	0.83	
	5	6000	0.93	
	15	0	1.00	----

The ratios of BBIT/butylene glycol tested ranged from 1/0.2 to 1/5000.

Table 7

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)

Second Component (B) = Caprylyl glycol

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	2000	1.00	-----
(24 hours)	40	400	0.60	1/10
	40	500	0.65	1/13
	40	600	0.70	1/15
	40	800	0.80	1/20
	40	1000	0.90	1/25
	50	200	0.60	1/4
	50	600	0.80	1/12
	50	800	0.90	1/16
	70	200	0.80	1/3
	70	300	0.85	1/4
	70	400	0.90	1/6
	80	200	0.90	1/3
	100	0	1.00	----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	3000	1.00	-----
(24 hours)	2	2000	0.92	1/1000
	4	500	0.67	1/125
	4	600	0.70	1/150
	4	800	0.77	1/200
	4	1000	0.83	1/250
	6	40	0.76	1/7
	6	50	0.77	1/8
	6	60	0.77	1/10
	6	80	0.78	1/13
	6	100	0.78	1/17
	6	200	0.82	1/33
	6	300	0.85	1/50
	6	400	0.88	1/67
	6	500	0.92	1/83
	8	0	1.00	-----
<i>A. niger</i> 16404 - PDB	0	3000	1.00	-----
(7 days)	5	1000	0.58	1/200

2012201211 29 Feb 2012

5

Microorganism	Q _a	Q _b	SI	A/B
	5	2000	0.92	1/400
	10	1000	0.83	1/100
	20	0	1.00	-----

The ratios of BBIT/caprylyl glycol tested ranged from 1/0.2 to 1/5000.

Table 8

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)

Second Component (B) = Chlorphenesin

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	2000	1.00	-----
(24 hours)	40	500	0.65	1/13
	40	600	0.70	1/115
	40	800	0.80	1/20
	40	1000	0.90	1/25
	50	300	0.65	1/6
	50	400	0.70	1/8
	50	500	0.75	1/10
	50	600	0.80	1/12
	50	800	0.90	1/16
	70	200	0.80	1/3
	70	300	0.85	1/4
	70	400	0.90	1/6
	80	100	0.85	1/1
	80	200	0.90	1/3
	100	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	2000	1.00	-----
(24 hours)	2	800	0.65	1/400
	2	1000	0.75	1/500
	4	300	0.65	1/75
	4	400	0.70	1/100
	4	500	0.75	1/125
	4	600	0.80	1/150
	4	800	0.90	1/200
	6	10	0.76	1/2
	6	20	0.76	1/3

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
	6	30	0.77	1/5
	6	40	0.77	1/7
	6	50	0.78	1/8
	6	60	0.78	1/10
	6	80	0.79	1/13
	6	100	0.80	1/17
	6	200	0.85	1/33
	6	300	0.90	1/50
	8	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	2000	1.00	-----
(4 days)	5	1000	0.83	1/200
	15	0	1.00	-----

The ratios of BBIT/chlorphenesin tested ranged from 1/0.2 to 1/5000.

5

Table 9

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = DMDM Hydantoin (DMDMH)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	200	1.00	-----
(72 hours)	40	8	0.44	1/0.2
	40	10	0.45	1/0.3
	40	20	0.50	1/0.5
	40	30	0.55	1/0.8
	40	40	0.60	1/1
	40	50	0.65	1/1
	40	60	0.70	1/2
	40	80	0.80	1/2
	40	100	0.90	1/3
	50	3	0.52	1/0.06
	50	4	0.52	1/0.08
	50	5	0.53	1/0.1
	50	6	0.53	1/0.1
	50	8	0.54	1/0.2
	50	10	0.55	1/0.2
	50	20	0.60	1/0.4

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
	50	30	0.65	1/0.6
	50	40	0.70	1/0.8
	50	50	0.75	1/1
	50	60	0.80	1/1.2
	50	80	0.90	1/1.6
	60	6	0.63	1/0.1
	60	8	0.64	1/0.1
	60	10	0.65	1/0.2
	60	20	0.70	1/0.3
	60	30	0.75	1/0.5
	60	40	0.80	1/0.7
	60	50	0.85	1/0.8
	60	60	0.90	1/1
	80	8	0.84	1/0.1
	80	10	0.85	1/0.1
	80	20	0.90	1/0.3
	100	0	1.00	-----
<i>C. albicans</i> 10231 · PDB	0	3000	1.00	-----
(24 hours)	5	600	0.87	1/120
	5	800	0.93	1/160
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	1000	1.00	-----
(7 days)	20	80	0.33	1/4
	20	100	0.35	1/5
	20	200	0.45	1/10
	20	300	0.55	1/15
	20	400	0.65	1/20
	20	500	0.75	1/25
	20	600	0.85	1/30
	40	100	0.60	1/3
	40	200	0.70	1/5
	40	300	0.80	1/8
	40	400	0.90	1/10
	80	0	1.00	-----

The ratios of BBIT/DMDM Hydantoin tested ranged from 1/0.2 to 1/5000.

Table 10

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
 Second Component (B) = dithio-2, 2'-bis(N-methylbenzamide) (DTBMA)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	60	1.00	-----
(48 hours)	40	6	0.50	1/0.2
	40	8	0.53	1/0.2
	40	10	0.57	1/0.3
	40	20	0.73	1/0.5
	40	30	0.90	1/0.8
	50	3	0.55	1/1.06
	50	4	0.57	1/0.08
	50	5	0.58	1/0.1
	50	6	0.60	1/0.1
	50	8	0.63	1/1.2
	50	10	0.67	1/0.2
	50	20	0.83	1/0.4
	60	3	0.65	1/0.05
	60	4	0.67	1/0.07
	60	5	0.68	1/0.08
	60	6	0.70	1/0.1
	60	8	0.73	1/0.1
	60	10	0.77	1/0.2
	60	20	0.93	1/0.3
	70	3	0.75	1/0.04
	70	4	0.77	1/0.06
	70	5	0.78	1/0.07
	70	6	0.80	1/0.09
	70	7	0.82	1/0.1
	70	8	0.83	1/0.1
	70	10	0.87	1/0.1
	80	1	0.82	1/0.01
	80	2	0.83	1/0.03
	80	3	0.85	1/0.04
	80	4	0.87	1/0.05
	80	5	0.88	1/0.06
	80	6	0.90	1/0.08

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
	80	8	0.93	1/0.1
	100	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	2000	1.00	-----
(24 hours)	8	40	0.55	1/5
	8	50	0.56	1/6
	8	60	0.56	1/8
	8	80	0.57	1/10
	8	100	0.58	1/13
	8	200	0.63	1/13
	8	300	0.68	1/38
	8	400	0.73	1/50
	8	500	0.78	1/63
	8	600	0.83	1/75
	8	800	0.93	1/100
	10	1	0.67	1/0.1
	10	2	0.67	1/0.2
	10	3	0.67	1/0.3
	10	4	0.67	1/0.4
	10	5	0.67	1/0.5
	10	6	0.67	1/0.6
	10	8	0.67	1/0.8
	10	10	0.67	1/1
	10	20	0.68	1/2
	10	30	0.68	1/3
	10	40	0.69	1/4
	10	50	0.69	1/5
	10	60	0.70	1/6
	10	80	0.71	1/8
	10	100	0.72	1/10
	10	200	0.77	1/20
	10	300	0.82	1/30
	10	400	0.87	1/40
	10	500	0.92	1/50
	15	0	1.00	-----
<i>A. niger</i> 16404 - PDB	0	1000	1.00	-----
(7 days)	5	800	0.93	1/160
	40	0	1.00	-----

The ratios of BBIT/ dithio-2, 2'-bis(N-methylbenzamide) tested ranged from 1/0.02 to 1/500.

Table 11

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = Ethylenediamine tetraacetic acid (EDTA)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	40000	1.00	-----
(72 hours)	40	800	0.42	1/20
	40	1000	0.43	1/25
	40	1200	0.43	1/30
	40	1600	0.44	1/40
	40	2000	0.45	1/50
	40	4000	0.50	1/100
	40	6000	0.55	1/150
	40	8000	0.60	1/200
	40	10000	0.65	1/250
	40	12000	0.70	1/300
	40	16000	0.80	1/400
	40	20000	0.90	1/500
	50	400	0.51	1/8
	50	600	0.52	1/12
	50	800	0.52	1/16
	50	1000	0.53	1/20
	50	1200	0.53	1/24
	50	1600	0.54	1/32
	50	2000	0.55	1/40
	50	4000	0.60	1/80
	50	6000	0.65	1/120
	50	8000	0.70	1/160
	50	10000	0.75	1/200
	50	12000	0.80	1/240
	50	16000	0.90	1/320
	60	400	0.61	1/7
	60	600	0.62	1/10
	60	800	0.62	1/13
	60	1000	0.63	1/17
	60	1200	0.63	1/20
	60	1600	0.64	1/27
	60	2000	0.65	1/33

Microorganism	Q _a	Q _b	SI	A/B
	60	4000	0.70	1/67
	60	6000	0.75	1/100
	60	8000	0.80	1/133
	60	10000	0.85	1/167
	60	12000	0.90	1/200
	70	200	0.71	1/3
	70	400	0.71	1/6
	70	600	0.72	1/9
	70	800	0.72	1/11
	70	1000	0.73	1/14
	70	1200	0.73	1/17
	70	1600	0.74	1/23
	70	2000	0.75	1/29
	70	4000	0.80	1/57
	70	6000	0.85	1/86
	70	8000	0.90	1/114
	70	10000	0.95	1/143
	80	400	0.81	1/5
	80	600	0.82	1/8
	80	800	0.82	1/40
	80	1000	0.83	1/13
	80	1200	0.83	1/15
	80	1600	0.84	1/20
	80	2000	0.85	1/25
	80	4000	0.90	1/50
	80	6000	0.95	1/75
	100	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
C. albicans 10231 · PDB	0	600	1.00	-----
(24 hours)	2.5	60	0.43	1/24
	2.5	80	0.47	1/32
	2.5	100	0.50	1/40
	2.5	120	0.53	1/48
	2.5	160	0.60	1/64
	2.5	200	0.67	1/80
	5	60	0.77	1/12
	5	80	0.80	1/16
	5	100	0.83	1/20
	5	120	0.87	1/24
	5	160	0.93	1/32

Microorganism	Q _a	Q _b	SI	A/B
	7.5	0	1.00	-----
<i>A. niger</i> 16404 - PDB	0	1600	1.00	-----
(7 days)	5	1000	0.75	1/200
	5	1200	0.88	1/240
	10	400	0.50	1/40
	10	600	0.63	1/60
	10	800	0.75	1/80
	10	1000	0.88	1/100
	15	400	0.63	1/27
	15	600	0.75	1/40
	15	800	0.88	1/53
	20	200	0.63	1/10
	20	400	0.75	1/20
	20	600	0.88	1/30
	40	0	1.00	-----

The ratios of BBIT/ethylenediamine tetraacetic acid tested ranged from 1/0.4 to 1/10000.

5

Table 12

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = Ethylparaben

0

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	2000	1.00	-----
(24 hours)	60	1000	0.80	1/17
	80	800	0.80	1/10
	80	1000	0.90	1/13
	100	200	0.60	1/2
	100	300	0.65	1/3
	100	400	0.70	1/4
	100	500	0.75	1/5
	100	600	0.80	1/6
	100	800	0.90	1/8
	200	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	800	1.00	-----
(24 hours)	2.5	80	0.43	1/32
	2.5	100	0.46	1/40

	2.5	200	0.58	1/80
	2.5	300	0.71	1/120
	2.5	400	0.83	1/160
	2.5	500	0.96	1/200
	5	30	0.70	1/6
	5	40	0.72	1/8
	5	50	0.73	1/10
	5	60	0.74	1/12
	5	80	0.77	1/16
	5	100	0.79	1/20
	5	200	0.92	1/40
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	800	1.00	-----
(7 days)	15	600	0.94	1/40
	80	0	1.00	-----

The ratios of BBIT/ethylparaben tested ranged from 1/0.2 to 1/5000.

Table 13

5

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)

Second Component (B) = Hexamidine diisethionate

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	100	1.00	-----
(48 hours)	40	1	0.57	1/0.03
	40	2	0.73	1/0.05
	40	3	0.90	1/0.08
	50	1	0.67	1/0.02
	50	2	0.83	1/0.04
	60	1	0.77	1/0.02
	60	2	0.93	1/0.03
	70	1	0.87	1/0.01
	80	1	0.97	1/0.01
	6	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	10	1.00	-----
(24 hours)	2	4	0.60	1/2
	2	5	0.70	1/2.5

2	6	0.80	1/3
4	2	0.60	1/0.5
4	3	0.70	1/0.75
4	4	0.80	1/1
4	5	0.90	1/1.3
6	1	0.70	1/0.2
6	2	0.80	1/0.3
6	3	0.90	1/0.5
8	1	0.90	1/1.1
10	0	1.00	-----

The ratios of BBIT/hexamidine diisethionate tested ranged from 1/0.02 to 1/500.

Table 14

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = Hexetidine

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	200	1.00	-----
(72 hours)	40	10	0.33	1/0.3
	40	20	0.45	1/1.5
	40	30	0.58	1/0.8
	40	40	0.70	1/1
	40	50	0.83	1/1.3
	40	60	0.95	1/1.5
	50	10	0.38	1/0.2
	50	20	0.50	1/0.4
	50	30	0.63	1/0.6
	50	40	0.75	1/0.8
	50	50	0.88	1/1
	60	4	0.35	1/0.07
	60	5	0.36	1/0.08
	60	6	0.38	1/0.1
	60	8	0.40	1/0.1
	60	10	0.43	1/0.2
	60	20	0.55	1/0.3
	60	30	0.68	1/0.5
	60	40	0.80	1/0.7

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
	60	50	0.93	1/0.8
	70	5	0.41	1/0.07
	70	6	0.43	1/0.09
	70	8	0.45	1/0.1
	70	10	0.48	1/0.1
	70	20	0.60	1/0.3
	70	30	0.73	1/0.4
	70	40	0.85	1/0.6
	70	50	0.98	1/0.7
	80	4	0.45	1/0.05
	80	5	0.46	1/0.06
	80	6	0.48	1/0.08
	80	8	0.50	1/0.1
	80	10	0.53	1/0.1
	80	20	0.65	1/0.3
	80	30	0.78	1/0.4
	80	40	0.90	1/0.5
	100	1	0.51	1/0.01
	100	2	0.53	1/0.02
	100	3	0.54	1/0.03
	100	4	0.55	1/0.04
	100	5	0.56	1/0.05
	100	6	0.58	1/0.06
	100	8	0.60	1/0.08
	100	10	0.63	1/0.1
	100	20	0.75	1/0.2
	100	30	0.88	1/0.3
	80	0	1.00	-----

Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB (72 hours)	0	100	1.00	-----
	2	40	0.53	1/20
	2	50	0.63	1/25
	2	60	0.73	1/30
	2	80	0.93	1/40
	4	30	0.57	1/8
	4	40	0.67	1/10
	4	50	0.77	1/13
	4	60	0.87	1/15
	6	20	0.60	1/3

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
	6	30	0.70	1/5
	6	40	0.80	1/7
	6	50	0.90	1/9
	8	5	0.58	1/0.6
	8	6	0.59	1/0.8
	8	8	0.61	1/1
	8	10	0.63	1/1
	8	20	0.73	1/3
	8	30	0.83	1/4
	8	40	0.93	1/5
	10	1	0.68	1/0.1
	10	2	0.69	1/0.2
	10	3	0.70	1/0.3
	10	4	0.71	1/0.4
	10	5	0.72	1/0.5
	10	6	0.73	1/0.6
	10	8	0.75	1/0.8
	10	10	0.77	1/1
	10	20	0.87	1/2
	10	30	0.97	1/3
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	100	1.00	-----
(4 days)	5	50	0.75	1/10
	5	60	0.85	1/12
	10	40	0.90	1/4
	20	0	1.00	-----

The ratios of BBIT/hexetidine tested ranged from 1/0.02 to 1/500.

5

Table 15

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = Hexylene glycol

10

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	20000	1.00	-----
(24 hours)	40	20000	1.40	1/500
	50	20000	1.50	1/400
	60	20000	1.60	1/333

Microorganism	Q _a	Q _b	SI	A/B
	80	20000	1.80	1/250
	100	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	20000	1.00	-----
(24 hours)	2	20000	1.20	1/10000
	4	20000	1.40	1/5000
	6	20000	1.60	1/3333
	8	20000	1.80	1/2800
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	20000	1.00	-----
(3 days)	5	20000	1.33	1/4000
	10	20000	1.67	1/2000
	15	0	1.00	-----

The ratios of BBIT/hexylene glycol tested ranged from 1/0.2 to 1/5000. No synergy was observed between BBIT and hexylene glycol.

Table 16

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = Methylparaben

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	2000	1.00	-----
(72 hours)	40	400	0.60	1/10
	40	500	0.65	1/13
	40	600	0.70	1/15
	40	800	0.80	1/20
	40	1000	0.90	1/25
	50	100	0.55	1/2
	50	200	0.60	1/4
	50	300	0.65	1/6
	50	400	0.70	1/8
	50	500	0.75	1/10
	50	600	0.80	1/12
	50	800	0.90	1/16
	60	30	0.62	1/0.5
	60	40	0.62	1/0.7

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
	60	50	0.63	1/0.8
	60	60	0.63	1/1
	60	80	0.64	1/1
	60	100	0.65	1/2
	60	200	0.70	1/3
	60	300	0.75	1/5
	60	400	0.80	1/7
	60	500	0.85	1/8
	60	600	0.90	1/10
	70	300	0.85	1/4
	70	400	0.90	1/6
	70	500	0.95	1/7
	80	300	0.95	1/4
	100	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	1000	1.00	-----
(24 hours)	2.5	100	0.43	1/40
	2.5	200	0.53	1/80
	2.5	300	0.63	1/120
	2.5	400	0.73	1/160
	2.5	500	0.83	1/200
	2.5	600	0.93	1/240
	5	40	0.71	1/8
	5	50	0.72	1/10
	5	60	0.73	1/12
	5	80	0.75	1/16
	5	100	0.77	1/20
	5	200	0.87	1/40
	5	300	0.97	1/60
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	1000	1.00	-----
(3 days)	5	600	0.73	1/120
	5	800	0.93	1/160
	40	0	1.00	-----

The ratios of BBIT/methylparaben tested ranged from 1/0.2 to 1/5000.

Table 17

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = Pentylene glycol

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	20000	1.00	-----
(24 hours)	40	20000	1.40	1/500
	60	20000	1.60	1/333
	80	20000	1.80	1/250
	100	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	20000	1.00	-----
(24 hours)	2	20000	1.02	1/10000
	4	20000	1.04	1/5000
	8	20000	1.08	
	10	20000	1.10	
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	20000	1.00	-----
(4 days)	5	20000	1.05	
	10	20000	1.10	
	15	20000	1.15	
	20	0	1.00	-----

The ratios of BBIT/pentylene glycol tested ranged from 1/0.2 to 1/5000. No synergy was observed between BBIT and pentylene glycol.

Table 18

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = Phenoxyethanol

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	5000	1.00	-----
(72 hours)	50	800	0.66	1/16
	50	1000	0.70	1/20
	50	2000	0.90	1/40
	60	60	0.61	1/1
	60	80	0.62	1/1

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
	60	100	0.62	1/2
	60	200	0.64	1/3
	60	300	0.66	1/5
	60	400	0.68	1/7
	60	500	0.70	1/8
	60	600	0.72	1/10
	60	800	0.76	1/13
	60	1000	0.80	1/17
	70	1000	0.90	1/14
	100	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	4000	1.00	-----
(24 hours)	2.5	1000	0.58	1/400
	2.5	2000	0.83	1/800
	5	500	0.79	1/100
	5	600	0.82	1/120
	5	800	0.87	1/160
	5	1000	0.92	1/200
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	4000	1.00	-----
(7 days)	5	3000	0.88	1/600
	40	0	1.00	-----

The ratios of BBIT/phenoxyethanol tested ranged from 1/0.2 to 1/5000.

5

Table 19

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
 Second Component (B) = linoleamidopropyl PG-dimonium chloride phosphate
 (Phospholipid EFA)

10

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	600	1.00	-----
(48 hours)	40	400	0.87	1/10
	50	400	0.92	1/8
	200	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	2000	1.00	-----

Microorganism	Q _a	Q _b	SI	A/B
(24 hours)	2	800	0.60	1/400
	2	1000	0.70	1/500
	4	600	0.70	1/150
	4	800	0.80	1/200
	4	1000	0.90	1/250
	6	500	0.85	1/83
	6	600	0.90	1/100
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	2000	1.00	-----
(4 days)	5	600	0.55	1/120
	5	800	0.65	1/160
	5	1000	0.75	1/200
	10	600	0.80	1/60
	10	800	0.90	1/80
	20	0	1.00	-----

The ratios of BBIT/linoleamidopropyl PG-dimonium chloride phosphate tested ranged from 1/0.02 to 1/500.

Table 20

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
 Second Component (B) = cocamidopropyl PG-dimonium chloride phosphate
 (Phospholipid PTC)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	80	1.00	-----
(72 hours)	40	20	0.45	1/0.5
	40	30	0.58	1/0.75
	40	40	0.70	1/1
	40	50	0.83	1/0.3
	40	60	0.95	1/2
	50	20	0.50	1/0.4
	50	30	0.63	1/0.6
	50	40	0.75	1/0.8
	50	50	0.88	1/1
	60	8	0.40	1/0.1

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
	60	10	0.43	1/0.2
	60	20	0.55	1/0.3
	60	30	0.68	1/0.5
	60	40	0.80	1/0.7
	60	50	0.93	1/0.8
	70	10	0.48	1/0.1
	70	20	0.60	1/0.3
	70	30	0.73	1/0.4
	70	40	0.85	1/0.6
	70	50	0.98	1/0.7
	80	10	0.53	1/0.1
	80	20	0.65	1/0.3
	80	30	0.78	1/0.4
	80	40	0.90	1/0.5
	100	3	0.54	1/0.3
	100	4	0.55	1/0.4
	100	5	0.56	1/0.5
	100	6	0.58	1/0.6
	100	8	0.60	1/0.8
	100	10	0.63	1/0.1
	100	20	0.75	1/0.2
	100	30	0.88	1/0.3
	200	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	1000	1.00	-----
(24 hours)	2	600	0.73	1/300
	2	800	0.93	1/400
	4	400	0.67	1/400
	4	500	0.77	1/125
	4	600	0.87	1/150
	6	300	0.70	1/50
	6	400	0.80	1/67
	6	500	0.90	1/83
	8	4	0.54	1/0.5
	8	5	0.54	1/0.6
	8	6	0.54	1/0.8

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
	8	8	0.54	1/1
	8	10	0.54	1/1
	8	20	0.55	1/6
	8	30	0.56	1/4
	8	40	0.57	1/5
	8	50	0.58	1/6
	8	60	0.59	1/8
	8	80	0.61	1/10
	8	100	0.63	1/13
	8	200	0.73	1/25
	8	300	0.83	1/38
	8	400	0.93	1/50
	10	3	0.67	1/0.3
	10	4	0.67	1/0.4
	10	5	0.67	1/0.5
	10	6	0.67	1/0.6
	10	8	0.67	1/0.8
	10	10	0.68	1/1
	10	20	0.69	1/2
	10	30	0.70	1/3
	10	40	0.71	1/4
	10	50	0.72	1/5
	10	60	0.73	1/6
	10	80	0.75	1/8
	10	100	0.77	1/10
	10	200	0.87	1/20
	10	300	0.97	1/30
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	1000	1.00	-----
(3 days)	5	500	0.83	1/100
	5	600	0.93	1/120
	10	200	0.87	1/20
	10	300	0.97	1/30
	15	0	1.00	-----

The ratios of **BBIT**/cocamidopropyl PG-dimonium chloride phosphate tested ranged from 1/0.02 to 1/500.

Table 21

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = Potassium sorbate

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	20000	1.00	-----
(24 hours)	20	20000	1.10	1/1000
	40	20000	1.20	1/500
	60	20000	1.30	1/333
	80	20000	1.40	1/250
	100	20000	1.50	1/200
	200	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	400	1.00	-----
(24 hours)	2.5	100	0.58	1/40
	2.5	200	0.83	1/80
	5	30	0.74	1/6
	5	60	0.82	1/12
	5	80	0.87	1/16
	5	100	0.92	1/20
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	8000	1.00	-----
(3 days)	5	6000	0.88	1/1200
	10	4000	0.75	1/400
	10	5000	0.88	1/500
	15	4000	0.88	1/267
	20	600	0.58	1/30
	20	800	0.60	1/40
	20	1000	0.63	1/50
	20	2000	0.75	1/100
	20	3000	0.88	1/150
	40	0	1.00	-----

The ratios of **BBIT**/potassium sorbate tested ranged from 1/0.2 to 1/5000.

Table 22

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
 Second Component (B) = Propylparaben

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	2000	1.00	-----
(24 hours)	20	2000	1.10	1/100
	40	2000	1.20	1/50
	60	2000	1.30	1/33
	80	2000	1.40	1/25
	100	2000	1.50	1/25
	200	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	2000	1.00	-----
(24 hours)	2.5	30	0.35	1/12
	2.5	40	0.35	1/16
	2.5	50	0.36	1/20
	2.5	60	0.36	1/24
	2.5	80	0.37	1/32
	2.5	100	0.38	1/40
	2.5	200	0.43	1/80
	2.5	300	0.48	1/120
	2.5	400	0.53	1/160
	2.5	500	0.58	1/200
	2.5	600	0.63	1/240
	2.5	800	0.73	1/320
	2.5	1000	0.83	1/400
	5	20	0.68	1/4
	5	30	0.68	1/6
	5	40	0.69	1/8
	5	50	0.69	1/10
	5	60	0.70	1/12
	5	80	0.71	1/16
	5	100	0.72	1/20
	5	200	0.77	1/40
	5	300	0.82	1/60
	5	400	0.87	1/80
	5	500	0.92	1/100

Microorganism	Q _a	Q _b	SI	A/B
	5	600	0.97	1/120
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	2000	1.00	-----
(7 days)	10	1000	0.63	1/100
	20	800	0.65	1/40
	20	1000	0.75	1/80
	40	800	0.90	1/20
	80	0	1.00	-----

The ratios of BBIT/propylparaben tested ranged from 1/0.2 to 1/5000.

Table 23

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
 Second Component (B) = Quaternium-15 (active - 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	200	1.00	-----
(72 hours)	40	50	0.43	1/1
	40	60	0.43	1/1
	40	80	0.44	1/2
	40	100	0.45	1/3
	40	200	0.50	1/5
	40	300	0.55	1/8
	40	400	0.60	1/10
	40	500	0.65	1/13
	40	600	0.70	1/15
	40	800	0.80	1/20
	200	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	600	1.00	-----
(24 hours)	5	50	0.75	1/10
	5	60	0.77	1/12
	5	80	0.80	1/16
	5	100	0.83	1/20
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B

Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	3000	1.00	-----
(4 days)	10	1000	0.58	1/100
	10	2000	0.92	1/200
	20	1000	0.83	1/50
	40	0	1.00	-----

The ratios of BBIT/1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride tested ranged from 1/0.02 to 1/5000.

Table 24

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)

Second Component (B) = dehydroacetic acid, sodium salt (SDHA)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	20000	1.00	-----
(24 hours)	20	20000	1.10	1/1000
	40	20000	1.20	1/500
	60	20000	1.30	1/333
	80	20000	1.40	1/250
	100	20000	1.50	1/200
	200	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	40	1.00	-----
(24 hours)	4	10	0.75	1/3
	6	4	0.85	1/0.7
	6	5	0.88	1/0.8
	6	6	0.90	1/1
	6	8	0.95	1/1
	8	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	200	1.00	-----
(7 days)	20	80	0.65	1/4
	20	100	0.75	1/5
	40	50	0.75	1/3
	40	60	0.80	1/2
	40	80	0.90	1/2
	80	0	1.00	-----

The ratios of BBIT/dehydroacetic acid, sodium salt tested ranged from 1/0.02 to 1/5000.

Table 25

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
Second Component (B) = Sodium benzoate

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	20000	1.00	-----
(24 hours)	20	20000	1.10	1/1000
	40	20000	1.20	1/500
	60	20000	1.30	1/333
	80	20000	1.40	1/250
	100	20000	1.50	1/200
	200	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	2000	1.00	-----
(48 hours)	2.5	400	0.53	1/160
	2.5	500	0.58	1/200
	2.5	600	0.63	1/240
	2.5	800	0.73	1/320
	2.5	1000	0.83	1/400
	5	300	0.82	1/60
	5	400	0.87	1/80
	5	500	0.92	1/100
	5	600	0.97	1/120
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	10000	1.00	-----
(7 days)	5	8000	0.86	1/1600
	10	3000	0.43	1/300
	10	4000	0.53	1/400
	10	5000	0.63	1/500
	10	6000	0.73	1/600
	10	8000	0.93	1/800
	15	3000	0.49	1/200
	15	4000	0.59	1/267
	15	5000	0.69	1/333
	15	6000	0.79	1/400

Microorganism	Q _a	Q _b	SI	A/B
	15	7000	0.89	1/467
	15	8000	0.99	1/533
	20	1000	0.35	1/50
	20	2000	0.45	1/100
	20	3000	0.55	1/150
	20	4000	0.65	1/200
	20	5000	0.75	1/250
	20	6000	0.85	1/300
	20	7000	0.95	1/350
	40	500	0.55	1/13
	40	600	0.56	1/15
	40	800	0.58	1/20
	40	1000	0.60	1/25
	40	2000	0.70	1/50
	40	3000	0.80	1/75
	40	4000	0.90	1/100
	80	0	1.00	-----

The ratios of BBIT/sodium benzoate tested ranged from 1/0.2 to 1/5000.

5

Table 26

First Component (A) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)
 Second Component (B) = Sodium citrate

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	20000	1.00	-----
(24 hours)	20	20000	1.10	1/1000
	40	20000	1.20	1/500
	60	20000	1.30	1/333
	80	20000	1.40	1/250
	100	20000	1.50	1/200
	200	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	10000	1.00	-----
(48 hours)	2.5	3000	0.55	1/1200
	2.5	4000	0.65	1/1600
	2.5	5000	0.75	1/2000
	2.5	6000	0.85	1/2400

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
	5	600	0.56	1/120
	5	800	0.58	1/160
	5	1000	0.60	1/200
	5	2000	0.70	1/400
	5	3000	0.80	1/600
	5	4000	0.90	1/800
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	10000	1.00	-----
(3 days)	5	8000	0.93	1/1600
	10	5000	0.75	1/500
	10	6000	0.85	1/600
	15	3000	0.68	1/200
	15	4000	0.78	1/267
	15	5000	0.88	1/333
	15	6000	0.98	1/400
	20	1000	0.60	1/50
	20	2000	0.70	1/100
	20	3000	0.80	1/150
	20	4000	0.90	1/200
	40	0	1.00	-----

The ratios of **BBIT**/sodium citrate tested ranged from 1/0.2 to 1/5000.

5

Table 27

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)

Second Component (B) = N-(n-butyl)-1,2-benzisothiazolinone (BBIT)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	100	1.00	-----
(24 hours)	2.5	50	0.75	1/20
	2.5	60	0.85	1/24
	5	40	0.90	1/8
	7.5	10	0.85	1/1
	7.5	20	0.95	1/3
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	6	1.00	-----

Microorganism	Q _a	Q _b	SI	A/B
(48 hours)	2.5	3	0.75	1/1
	2.5	4	0.92	1/2
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	40	1.00	-----
(4 days)	10	5	0.38	1/0.5
	10	6	0.40	1/0.6
	10	8	0.45	1/0.8
	10	10	0.50	1/1
	10	20	0.75	1/2
	20	6	0.65	1/0.3
	20	8	0.70	1/0.4
	20	10	0.75	1/0.5
	40	0	1.00	-----

The ratios of MBIT/BBIT tested ranged from 1/0.01 to 1/400.

5

Table 28

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)

Second Component (B) = Benzalkonium chloride (Hyamine 3500)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	10	1.00	-----
(24 hours)	2.5	4	1.25	1/2
	5	3	1.25	1/0.6
	7.5	3	1.50	1/0.4
	100	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	60	1.00	-----
(24 hours)	2	10	0.42	1/5
	2	20	0.58	1/10
	2	30	0.75	1/15
	2	40	0.92	1/20
	4	3	0.55	1/0.8
	4	4	0.57	1/1
	4	5	0.58	1/1
	4	6	0.60	1/2

Microorganism	Q _a	Q _b	SI	A/B
	4	8	0.63	1/2
	4	10	0.67	1/3
	4	20	0.83	1/5
	6	3	0.80	1/0.5
	6	4	0.82	1/0.7
	6	5	0.83	1/0.9
	6	6	0.85	1/1
	6	8	0.88	1/1
	6	10	0.92	1/2
	8	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	200	1.00	-----
(4 days)	10	30	0.48	1/3
	10	40	0.53	1/4
	10	50	0.58	1/5
	10	60	0.63	1/6
	10	80	0.73	1/8
	10	100	0.83	1/10
	20	4	0.69	1/0.2
	20	5	0.69	1/0.3
	20	6	0.70	1/0.3
	20	8	0.71	1/0.4
	20	10	0.72	1/0.5
	20	20	0.77	1/4
	20	30	0.82	1/1.5
	20	40	0.87	1/2
	20	50	0.92	1/3
	20	60	0.97	1/3
	30	0	1.00	-----

The ratios of MBIT/benzalkonium chloride tested ranged from 1/0.01 to 1/400.

5

Table 29

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
 Second Component (B) = Benzethonium chloride (Hyamine 1622)

10

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	20	1.00	-----

Microorganism	Q _a	Q _b	SI	A/B
(24 hours)	2.5	20	1.25	1/8
	5	20	1.50	1/4
	7.5	5	1.00	1/0.7
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	4	1.00	-----
(24 hours)	2	3	1.00	1/2
	4	3	1.25	1/0.8
	6	1	1.00	1/0.2
	8	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	10	1.00	-----
(4 days)	10	4	0.73	1/0.4
	10	5	0.83	1/0.5
	10	6	0.93	1/0.6
	30	0	1.00	-----

The ratios of MBIT/benzethonium chloride tested ranged from 1/0.01 to 1/400.

Table 30

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
Second Component (B) = Benzyl alcohol

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	5000	1.00	-----
(24 hours)	2.5	5000	1.25	1/2000
	5	4000	1.30	1/800
	7.5	4000	1.55	1/533
	10	0	1.00	----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	4000	1.00	-----
(48 hours)	2.5	2000	0.75	1/800
	5	500	0.63	1/100
	5	1000	0.75	1/200
	7.5	40	0.76	1/5
	7.5	50	0.76	1/7
	7.5	60	0.77	1/8

	7.5	80	0.77	1/11
	7.5	100	0.78	1/13
	7.5	200	0.80	1/27
	7.5	300	0.08	1/40
	7.5	400	0.85	1/53
	7.5	500	0.88	1/67
	7.5	600	0.90	1/80
	7.5	800	0.95	1/107
	10	0	1.00	-----
Microorganism	Q_a	Q_b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	6000	1.00	-----
(4 days)	10	6000	1.20	1/600
	20	6000	1.40	1/300
	30	4000	1.27	1/133
	50	0	1.00	----

The ratios of MBIT/benzyl alcohol tested ranged from 1/0.1 to 1/4000.

5

Table 31

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)

Second Component (B) = Benzisothiazolinone (BIT)

Microorganism	Q_a	Q_b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	20	1.00	-----
(24 hours)	2.5	10	0.75	1/4
	5	5	0.75	1/1
	5	6	0.80	1/1
	5	8	0.90	1/2
	7.5	3	0.90	1/0.4
	7.5	4	0.95	1/0.5
	10	0	1.00	-----
Microorganism	Q_a	Q_b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	30	1.00	-----
(48 hours)	2.5	10	0.58	1/4
	2.5	20	0.92	1/8
	5	4	0.63	1/0.8
	5	5	0.67	1/1
	5	6	0.70	1/1

Microorganism	Q _a	Q _b	SI	A/B
	5	8	0.77	1/2
	5	10	0.83	1/2
	7.5	2	0.82	1/0.3
	7.5	3	0.85	1/0.4
	7.5	4	0.88	1/0.5
	7.5	5	0.92	1/0.7
	7.5	6	0.95	1/0.8
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB (7 days)	0	300	1.00	-----
	10	100	0.53	1/10
	10	200	0.87	1/20
	20	30	0.50	1/2
	20	40	0.53	1/2
	20	50	0.57	1/3
	20	60	0.60	1/3
	20	80	0.67	1/4
	20	100	0.73	1/5
	30	20	0.67	1/0.7
	30	30	0.70	1/1
	30	40	0.73	1/1
	30	50	0.77	1/2
	30	60	0.80	1/2
	30	80	0.87	1/3
	30	100	0.93	1/3
	50	0	1.00	-----

The ratios of MBIT/benzisothiazolinone tested ranged from 1/0.01 to 1/400.

5

Table 32

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)

Second Component (B) = 2-bromo-2-nitro-propane-1, 3-diol (BNPD)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	8	1.00	-----
(24 hours)	2.5	5	0.88	1/2
	5	1	0.63	1/0.2
	5	2	0.75	1/0.4

Microorganism	Q _a	Q _b	SI	A/B
	5	3	0.88	1/0.6
	7.5	1	0.88	1/0.1
	7.5	2	1.00	1/0.3
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	1000	1.00	-----
(24 hours)	4	100	0.60	1/25
	4	200	0.70	1/50
	4	300	0.80	1/75
	4	400	0.90	1/100
	6	4	0.75	1/0.7
	6	5	0.76	1/0.8
	6	6	0.76	1/1
	6	8	0.76	1/1
	6	10	0.76	1/2
	6	20	0.77	1/3
	6	30	0.78	1/5
	6	40	0.79	1/7
	6	50	0.80	1/8
	6	60	0.81	1/10
	6	80	0.83	1/13
	6	100	0.85	1/17
	6	200	0.95	1/33
	8	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	2000	1.00	-----
(4 days)	10	800	0.60	0/80
	10	1000	0.70	1/100
	20	5	0.40	1/0.3
	20	6	0.40	1/0.3
	20	8	0.40	1/0.4
	20	10	0.41	1/0.5
	20	20	0.41	1/1
	20	30	0.42	1/2
	20	40	0.42	1/2
	20	50	0.43	1/3
	20	60	0.43	1/3
	20	80	0.44	1/4

Microorganism	Q _a	Q _b	SI	A/B
	20	100	0.45	1/5
	20	200	0.50	1/10
	20	300	0.55	1/15
	20	400	0.60	1/20
	20	500	0.65	1/25
	20	600	0.70	1/30
	20	800	0.80	1/40
	20	1000	0.90	1/50
	30	0	1.00	-----

The ratios of MBIT/2-bromo-2-nitro-propane-1, 3-diol tested ranged from 1/0.01 to 1/400.

Table 33

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
Second Component (B) = Butylene glycol

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	20000	1.00	-----
(24 hours)	2.5	20000	1.33	1/8000
	5	20000	1.67	1/4000
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	20000	1.00	-----
(72 hours)	8	100	0.81	1/13
	8	200	0.81	1/25
	8	300	0.82	1/38
	8	400	0.82	1/50
	8	500	0.83	1/63
	8	600	0.83	1/75
	8	800	0.84	1/100
	8	1000	0.85	1/125
	8	2000	0.90	1/250
	8	3000	0.95	1/375
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	10000	1.00	----
(7 days)	10	10000	1.25	1/1000

Microorganism	Q _a	Q _b	SI	A/B
	20	10000	1.50	1/500
	30	10000	1.75	1/333
	40	0	1.00	----

The ratios of MBIT/butylene glycol tested ranged from 1/0.1 to 1/4000.

Table 34

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
Second Component (B) = Caprylyl glycol

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	2000	1.00	-----
(24 hours)	2.5	2000	1.25	1/800
	5	2000	1.50	1/400
	7.5	2000	1.75	1/267
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	4000	1.00	-----
(24 hours)	2	2000	0.70	1/1000
	2	3000	0.95	1/1500
	4	2000	0.90	1/500
	6	600	0.75	1/100
	6	800	0.80	1/133
	6	1000	0.85	1/167
	8	200	0.85	1/25
	8	300	0.88	1/38
	8	400	0.90	1/50
	8	500	0.93	1/63
	8	600	0.95	1/75
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	1000	1.00	-----
(4 days)	10	1000	0.58	1/100
	10	2000	0.83	1/200
	20	1000	0.92	1/50
	20	0	1.00	-----

The ratios of MBIT/caprylyl glycol tested ranged from 1/0.1 to 1/4000.

Table 35

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
Second Component (B) = Chlorphenesin

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	2000	1.00	-----
(24 hours)	7.5	800	0.90	1/107
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	2000	1.00	-----
(72 hours)	4	200	0.50	1/50
	4	300	0.55	1/75
	4	400	0.60	1/100
	4	500	0.65	1/125
	4	600	0.70	1/150
	4	800	0.80	1/200
	4	1000	0.90	1/250
	6	80	0.64	1/13
	6	100	0.65	1/17
	6	200	0.70	1/33
	6	300	0.75	1/50
	6	400	0.80	1/67
	6	500	0.85	1/83
	6	600	0.90	1/100
	8	40	0.82	1/5
	8	50	0.83	1/6
	8	60	0.83	1/8
	8	80	0.84	1/10
	8	100	0.85	1/13
	8	200	0.90	1/25
	8	300	0.95	1/38
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	2000	1.00	-----
(3 days)	10	2000	1.50	1/200
	20	0	1.00	-----

The ratios of MBIT/chlorphenesin tested ranged from 1/0.1 to 1/4000.

Table 36

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
Second Component (B) = DMDM Hydantoin (DMDMH)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	50	1.00	-----
(24 hours)	2.5	60	1.45	1/24
	5	30	1.10	1/6
	7.5	10	0.95	1/1
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	3000	1.00	-----
(24 hours)	2.5	3000	1.33	1/1200
	5	1000	1.00	1/200
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	1000	1.00	-----
(3 days)	10	2000	2.25	1/200
	20	2000	2.50	1/100
	40	0	1.00	-----

The ratios of MBIT/DMDM Hydantoin tested ranged from 1/0.1 to 1/4000.
No synergy was observed between MBIT and DMDM Hydantoin.

Table 37

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
Second Component (B) = dithio-2, 2'-bis(N-methylbenzamide) (DTBMA)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	80	1.00	-----
(24 hours)	2.5	20	0.42	1/8
	2.5	30	0.54	1/12
	2.5	40	0.67	1/16
	2.5	50	0.79	1/20
	2.5	60	0.92	1/24
	5	20	0.58	1/4

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
	5	30	0.71	1/6
	5	40	0.83	1/8
	5	50	0.96	1/10
	7.5	3	0.54	1/0.4
	7.5	4	0.55	1/0.5
	7.5	5	0.56	1/0.7
	7.5	6	0.58	1/0.8
	7.5	8	0.60	1/1
	7.5	10	0.63	1/1
	7.5	20	0.75	1/3
	7.5	30	0.88	1/4
	10	1	0.68	1/0.1
	10	2	0.69	1/0.2
	10	3	0.70	1/0.3
	10	4	0.72	1/0.4
	10	5	0.73	1/0.5
	10	6	0.74	1/0.6
	10	8	0.77	1/0.8
	10	10	0.79	1/1
	10	20	0.92	1/2
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	2000	1.00	-----
(24 hours)	6	60	0.63	1/10
	6	80	0.64	1/13
	6	100	0.65	1/17
	6	200	0.70	1/33
	6	300	0.75	1/50
	6	400	0.80	1/67
	6	500	0.85	1/83
	6	600	0.90	1/100
	8	10	0.81	1/1
	8	20	0.81	1/3
	8	30	0.82	1/4
	8	40	0.82	1/5
	8	50	0.83	1/6
	8	60	0.83	1/8
	8	80	0.84	1/10
	8	100	0.85	1/13

Microorganism	Q _a	Q _b	SI	A/B
	8	200	0.90	1/25
	8	300	0.95	1/38
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	800	1.00	-----
(3 days)	10	500	0.96	1/50
	30	0	1.00	-----

The ratios of MBIT/ dithio-2, 2'-bis(N-methylbenzamide) tested ranged from 1/0.01 to 1/400.

Table 38

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)

Second Component (B) = Ethylenediamine tetraacetic acid (EDTA)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	40000	1.00	-----
(24 hours)	5	1600	0.54	1/320
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	1000	1.00	-----
(48 hours)	2.5	60	0.43	1/24
	2.5	80	0.47	1/32
	2.5	100	0.50	1/40
	2.5	120	0.53	1/48
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	1600	1.00	-----
(3 days)	10	800	0.75	1/80
	10	1000	0.88	1/100
	20	200	0.63	1/10
	20	400	0.75	1/20
	20	600	0.88	1/30
	40	0	1.00	-----

The ratios of MBIT/ethylenediamine tetraacetic acid tested ranged from 1/0.3 to 1/8000.

Table 39

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
Second Component (B) = Ethylparaben

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	2000	1.00	-----
(24 hours)	2.5	2000	1.25	1/800
	5	2000	1.50	1/400
	7.5	400	0.95	1/53
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	2000	1.00	-----
(24 hours)	2.5	600	0.63	1/240
	2.5	800	0.73	1/320
	2.5	1000	0.83	1/400
	2.5	2000	1.33	1/800
	5	50	0.69	1/10
	5	60	0.70	1/12
	5	80	0.71	1/16
	5	100	0.72	1/20
	5	200	0.77	1/40
	5	300	0.82	1/60
	5	400	0.87	1/80
	5	500	0.92	1/100
	5	600	0.97	1/120
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	800	1.00	-----
(7 days)	30	200	0.85	1/7
	30	300	0.98	1/10
	50	0	1.00	-----

The ratios of MBIT/ethylparaben tested ranged from 1/0.1 to 1/4000.

Table 40

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)

Second Component (B) = Hexamidine diisethionate

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	8	1.00	-----
(24 hours)	2.5	8	1.17	1/3
	5	8	1.33	1/2
	7.5	5	1.13	1/0.7
	10	3	1.04	1/0.3
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	10	1.00	-----
(24 hours)	2	5	0.70	1/3
	2	6	0.80	1/3
	4	4	0.80	1/1
	4	5	0.90	1/1
	8	1	0.90	1/0.1
	10	0	1.00	-----

The ratios of MBIT/hexamidine diisethionate tested ranged from 1/0.01 to 1/400.

Table 41

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)

Second Component (B) = Hexetidine

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	80	1.00	-----
(72 hours)	5	40	0.83	1/8
	5	50	0.96	1/10
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	80	1.00	-----
(72 hours)	2	40	0.53	1/20
	2	50	0.63	1/25
	2	60	0.73	1/30
	2	80	0.93	1/40

Microorganism	Q _a	Q _b	SI	A/B
	4	30	0.57	1/8
	4	40	0.67	1/10
	4	50	0.77	1/13
	4	60	0.87	1/15
	6	20	0.60	1/3
	6	30	0.70	1/5
	6	40	0.80	1/7
	6	50	0.90	1/8
	8	5	0.58	1/0.6
	8	6	0.59	1/0.7
	8	8	0.61	1/1
	8	10	0.63	1/1
	8	20	0.73	1/3
	8	30	0.83	1/4
	8	40	0.93	1/5
	10	1	0.68	1/0.1
	10	2	0.69	1/0.2
	10	3	0.70	1/0.3
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	200	1.00	-----
(3 days)	10	40	0.53	1/4
	10	50	0.58	1/5
	10	60	0.63	1/6
	10	80	0.73	1/8
	10	100	0.83	1/10
	20	20	0.77	1/1
	20	30	0.82	1/2
	20	40	0.87	1/2
	20	50	0.92	1/3
	20	60	0.97	1/3
	20	0	1.00	-----

The ratios of MBIT/hexetidine tested ranged from 1/0.01 to 1/400.

Table 42

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
Second Component (B) = Hexylene glycol

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	20000	1.00	-----
(24 hours)	2.5	20000	1.17	1/8000
	5	20000	1.33	1/4000
	7.5	20000	1.50	1/2667
	10	20000	1.67	1/2000
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	20000	1.00	-----
(24 hours)	2	20000	1.13	1/10000
	4	20000	1.27	1/5000
	6	20000	1.40	1/3333
	8	20000	1.53	1/2500
	10	20000	1.67	1/2000
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	20000	1.00	-----
(3 days)	10	20000	1.33	1/2000
	20	20000	1.67	1/1000
	30	0	1.00	-----

The ratios of MBIT/hexylene glycol tested ranged from 1/0.1 to 1/10000.
No synergy was observed between MBIT and hexylene glycol.

Table 43

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
Second Component (B) = Methylparaben

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	2000	1.00	-----
(24 hours)	2.5	2000	1.25	1/800
	5	2000	1.50	1/400
	7.5	800	1.15	1/107
	10	0	1.00	-----

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	2000	1.00	-----
(48 hours)	2.5	1000	0.83	1/400
	5	400	0.87	1/80
	5	500	0.92	1/100
	5	600	0.97	1/120
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	800	1.00	-----
(7 days)	30	50	0.66	1/2
	30	60	0.68	1/2
	30	80	0.70	1/3
	30	100	0.73	1/6
	30	200	0.85	1/7
	30	300	0.98	1/10
	50	0	1.00	-----

The ratios of MBIT/methylparaben tested ranged from 1/0.1 to 1/4000.

5

Table 44

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
Second Component (B) = Pentylene glycol

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	20000	1.00	-----
(24 hours)	2.5	20000	1.17	1/8000
	5	20000	1.33	1/4000
	7.5	20000	1.50	1/2667
	10	20000	1.67	1/2000
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	20000	1.00	-----
(24 hours)	2	20000	1.13	1/10000
	4	20000	1.27	1/5000
	8	20000	1.53	1/2500
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B

Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	20000	1.00	-----
(3 days)	10	20000	1.67	1/2000
	20	20000	2.33	1/1000
	30	0	1.00	-----

The ratios of MBIT/pentylene glycol tested ranged from 1/0.1 to 1/10000. No synergy was observed between MBIT and pentylene glycol.

Table 45

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
Second Component (B) = Phenoxyethanol

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	4000	1.00	-----
(24 hours)	2.5	4000	1.17	1/1600
	5	4000	1.33	1/800
	7.5	4000	1.50	1/533
	10	3000	1.42	1/300
	40	3000	3.42	1/75
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	3000	1.00	-----
(24 hours)	5	600	0.87	1/120
	5	800	0.93	1/160
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	4000	1.00	-----
(3 days)	10	4000	1.25	1/400
	20	2000	1.00	1/100
	40	0	1.00	-----

The ratios of MBIT/phenoxyethanol tested ranged from 1/0.1 to 1/4000.

Table 46

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
 Second Component (B) = linoleamidopropyl PG-dimonium chloride phosphate
 (Phospholipid EFA)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	500	1.00	-----
(48 hours)	15	40	0.83	1/3
	15	50	0.85	1/3
	15	60	0.87	1/4
	15	80	0.91	1/5
	15	100	0.95	1/7
	20	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	1000	1.00	-----
(24 hours)	2	800	0.93	1/400
	4	600	0.87	1/150
	6	500	0.90	1/83
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	2000	1.00	-----
(3 days)	10	800	0.73	1/80
	10	1000	0.83	1/100
	20	300	0.82	1/15
	20	400	0.87	1/20
	20	500	0.92	1/25
	20	600	0.97	1/30
	30	0	1.00	-----

The ratios of MBIT/linoleamidopropyl PG-dimonium chloride phosphate tested ranged from 1/0.01 to 1/400.

Table 47

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
 Second Component (B) = cocamidopropyl PG-dimonium chloride phosphate
 (Phospholipid PTC)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	100	1.00	-----
(48 hours)	2.5	60	0.73	1/24
	2.5	80	0.93	1/32
	10	30	0.80	1/3
	10	40	0.90	1/4
	15	10	0.85	1/0.7
	15	20	0.95	1/1
	20	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	1000	1.00	-----
(24 hours)	4	500	0.90	1/125
	6	300	0.90	1/50
	8	3	0.80	1/0.4
	8	4	0.80	1/0.5
	8	5	0.81	1/0.7
	8	6	0.81	1/0.8
	8	8	0.81	1/1
	8	10	0.81	1/1
	8	20	0.82	1/3
	8	30	0.83	1/4
	8	40	0.84	1/5
	8	50	0.85	1/6
	8	60	0.86	1/8
	8	80	0.88	1/10
	8	100	0.90	1/13
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	1000	1.00	-----
(4 days)	10	500	0.83	1/50
	10	600	0.93	1/60
	30	0	1.00	-----

The ratios of MBIT/cocamidopropyl PG-dimonium chloride phosphate tested ranged from 1/0.01 to 1/400.

Table 48

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)

Second Component (B) = Potassium sorbate

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 · M9GY	0	20000	1.00	-----
(48 hours)	10	1000	0.72	1/100
	10	2000	0.77	1/200
	10	3000	0.82	1/300
	10	4000	0.87	1/400
	10	5000	0.92	1/500
	10	6000	0.97	1/600
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 · PDB	0	400	1.00	-----
(24 hours)	2.5	200	0.83	1/80
	5	40	0.77	1/8
	5	50	0.79	1/10
	5	60	0.82	1/12
	5	80	0.87	1/16
	5	100	0.92	1/20
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 · PDB	0	10000	1.00	-----
(4 days)	10	6000	0.85	1/600
	20	1000	0.60	1/50
	20	2000	0.70	1/100
	20	3000	0.80	1/150
	20	4000	0.90	1/200
	40	0	1.00	-----

The ratios of MBIT/potassium sorbate tested ranged from 1/0.1 to 1/4000.

Table 49

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
Second Component (B) = Propylparaben

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	4000	1.00	-----
(48 hours)	2.5	2000	0.67	1/800
	2.5	3000	0.92	1/1200
	5	2000	0.83	1/400
	15	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	2000	1.00	-----
(24 hours)	2.5	800	0.73	1/320
	2.5	1000	0.83	1/400
	5	400	0.87	1/80
	5	500	0.92	1/100
	5	600	0.97	1/120
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	2000	1.00	-----
(7 days)	10	800	0.60	1/80
	10	1000	0.70	1/100
	20	1000	0.90	1/50
	30	300	0.75	1/10
	30	400	0.80	1/13
	30	500	0.85	1/17
	30	600	0.90	1/20
	50	0	1.00	-----

The ratios of MBIT/propylparaben tested ranged from 1/0.1 to 1/4000.

2012201211 29 Feb 2012

5

Table 50

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)

Second Component (B) = Quaternium-15 (active - 1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	50	1.00	-----
(24 hours)	5	20	0.90	1/400
	7.5	2	0.79	1/0.3
	7.5	3	0.81	1/0.4
	7.5	4	0.83	1/0.5
	7.5	5	0.85	1/0.7
	7.5	6	0.87	1/0.8
	7.5	8	0.91	1/1
	7.5	10	0.95	1/1
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	500	1.00	-----
(24 hours)	2.5	500	1.33	1/200
	5	300	1.27	1/60
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	3000	1.00	-----
(3 days)	10	1000	0.58	1/100
	10	2000	0.92	1/200
	20	1000	0.83	1/50
	40	0	1.00	-----

The ratios of MBIT/1-(3-chloroallyl)-3,5,7-triaza-1-azoniaadamantane chloride tested ranged from 1/0.01 to 1/4000.

Table 51

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)

Second Component (B) = Dehydroacetic acid, sodium salt (SDHA)

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	20000	1.00	-----
(24 hours)	2.5	20000	1.25	1/8000

Microorganism	Q _a	Q _b	SI	A/B
	5	20000	1.50	1/4000
	7.5	20000	1.75	1/2667
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	40	1.00	-----
(24 hours)	2	40	1.25	1/20
	4	30	1.25	1/8
	6	8	0.95	1/1
	8	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	200	1.00	-----
(7 days)	20	80	0.80	1/4
	20	100	0.90	1/5
	30	50	0.85	1/2
	30	60	0.90	1/2
	50	0	1.00	-----

The ratios of MBIT/dehydroacetic acid, sodium salt tested ranged from 1/0.01 to 1/4000.

Table 52

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
Second Component (B) = Sodium benzoate

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	10000	1.00	-----
(24 hours)	2.5	10000	1.25	1/4000
	5	10000	1.50	1/2000
	7.5	10000	1.75	1/1333
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	2000	1.00	-----
(48 hours)	2.5	800	0.65	1/320
	2.5	1000	0.75	1/400
	5	300	0.65	1/60
	5	400	0.70	1/80
	5	500	0.75	1/100

Microorganism	Q _a	Q _b	SI	A/B
	5	600	0.80	1/120
	5	800	0.90	1/160
	7.5	30	0.77	1/4
	7.5	40	0.77	1/5
	7.5	50	0.78	1/7
	7.5	60	0.78	1/8
	7.5	80	0.79	1/11
	7.5	100	0.80	1/13
	7.5	200	0.85	1/27
	7.5	300	0.90	1/40
	7.5	400	0.95	1/53
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	10000	1.00	-----
(7 days)	10	6000	0.77	1/600
	10	8000	0.97	1/800
	20	6000	0.93	1/300
	30	2000	0.70	1/67
	30	3000	0.80	1/100
	30	4000	0.90	1/133
	40	2000	0.87	1/50
	40	3000	0.97	1/75
	60	0	1.00	-----

The ratios of MBIT/sodium benzoate tested ranged from 1/0.1 to 1/4000.

5

Table 53

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)
 Second Component (B) = Sodium citrate

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	20000	1.00	-----
(24 hours)	2.5	20000	1.25	1/8000
	5	20000	1.50	1/4000
	7.5	20000	1.75	1/2667
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B

2012201211 29 Feb 2012

Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB	0	10000	1.00	-----
(24 hours)	2.5	3000	0.63	1/1200
	2.5	4000	0.73	1/1600
	2.5	5000	0.83	1/2000
	2.5	6000	0.93	1/2400
	5	500	0.72	1/100
	5	600	0.73	1/120
	5	800	0.75	1/160
	5	1000	0.77	1/200
	5	2000	0.87	1/400
	5	3000	0.97	1/600
	7.5	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>A. niger</i> 16404 - PDB	0	10000	1.00	-----
(7 days)	30	1000	0.70	1/33
	30	2000	0.80	1/67
	30	3000	0.90	1/100
	50	0	1.00	-----

The ratios of MBIT/sodium citrate tested ranged from 1/0.1 to 1/4000.

5

Table 54

First Component (A) = N-methyl-1,2-benzisothiazolin-3-one (MBIT)

Second Component (B) = Zinc pyrithione

Microorganism	Q _a	Q _b	SI	A/B
<i>E. coli</i> 8739 - M9GY	0	2	1.00	-----
(48 hours)	2.5	0.4	0.45	1/0.2
	2.5	0.5	0.50	1/0.2
	2.5	0.6	0.55	1/0.2
	2.5	0.8	0.65	1/0.3
	2.5	1	0.75	1/0.4
	5	0.3	0.65	1/0.06
	5	0.4	0.70	1/0.08
	5	0.5	0.75	1/0.1
	5	0.6	0.80	1/0.1
	5	0.8	0.90	1/0.2
	7.5	0.3	0.90	1/0.04

Microorganism	Q _a	Q _b	SI	A/B
	7.5	0.4	0.95	1/0.05
	10	0	1.00	-----
Microorganism	Q _a	Q _b	SI	A/B
<i>C. albicans</i> 10231 - PDB (72 hours)	0	30	1.00	-----
	2.5	4	0.47	1/2
	2.5	5	0.50	1/2
	2.5	6	0.53	1/2
	2.5	8	0.60	1/3
	2.5	10	0.67	1/4
	5	2	0.73	1/0.4
	5	3	0.77	1/0.6
	5	4	0.80	1/0.8
	5	5	0.83	1/1
	5	6	0.87	1/1
	5	8	0.93	1/2
	7.5	0	1.00	-----
<i>A. niger</i> 16404 - PDB (3 days)	0	20	1.00	-----
	20	5	0.75	1/3
	20	6	0.80	1/0.3
	20	8	0.90	1/0.4
	40	0	1.00	-----

The ratios of MBIT/zinc pyrithione tested ranged from 1/0.001 to 1/400.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A microbicidal composition comprising:
(a) N-(n-butyl)-1,2-benzisothiazolin-3-one; and
(b) hexamidine diisethionate.
2. A microbicidal composition comprising:
(a) N-methyl-1,2-benzisothiazolin-3-one; and
(b) hexamidine diisethionate.
3. The composition according to claim 1 or claim 2, substantially as
hereinbefore described with reference to any one of the Examples.