



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2370123 C 2010/09/14

(11)(21) **2 370 123**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2000/05/03
(87) Date publication PCT/PCT Publication Date: 2000/11/09
(45) Date de délivrance/Issue Date: 2010/09/14
(85) Entrée phase nationale/National Entry: 2001/11/02
(86) N° demande PCT/PCT Application No.: US 2000/012243
(87) N° publication PCT/PCT Publication No.: 2000/066786
(30) Priorité/Priority: 1999/05/03 (US60/132,412)

(51) Cl.Int./Int.Cl. *C12Q 1/68* (2006.01)
(72) Inventeurs/Inventors:
HOGAN, JAMES J., US;
GORDON, PATRICIA, US
(73) Propriétaire/Owner:
GEN-PROBE INCORPORATED, US
(74) Agent: SMART & BIGGAR

(54) Titre : **SONDES POLYNUCLEOTIDIQUES DESTINEES A LA DETECTION ET A LA QUANTIFICATION
D'ACTINOMYCETES**
(54) Title: **POLYNUCLEOTIDE PROBES FOR DETECTION AND QUANTITATION OF ACTINOMYCETES**

(57) **Abrégé/Abstract:**

Polynucleotide probes and accessory helper oligonucleotides useful for detecting the subset of High (G+C) subset of Gram-positive bacteria known as the "Actinomycetes." The hybridization probes are highly specific for the Actinomycetes and do not cross-hybridize with the rRNA or rDNA of numerous other bacterial and fungal species.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
9 November 2000 (09.11.2000)

PCT

(10) International Publication Number
WO 00/66786 A3(51) International Patent Classification⁷: **C12Q 1/68**

(21) International Application Number: PCT/US00/12243

(22) International Filing Date: 3 May 2000 (03.05.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/132,412 3 May 1999 (03.05.1999) US(71) Applicant (for all designated States except US):
GEN-PROBE INCORPORATED [US/US]; Patent
Department, 10210 Genetic Center Drive, San Diego, CA
92121-4362 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HOGAN, James, J.**
[US/US]; 1038 Olive Lane, Coronado, CA 92118 (US).
GORDON, Patricia [US/US]; 3501 Saddle Drive, San
Diego, CA 92077 (US).(74) Agent: **GILLY, Michael, J.**; Gen-Probe Incorporated,
Patent Department, 10210 Genetic Center Drive, San
Diego, CA 92121-4362 (US).(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ,
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,
TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GW, ML, MR, NE, SN, TD, TG).**Published:**

— With international search report.

(88) Date of publication of the international search report:
12 July 2001For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: POLYNUCLEOTIDE PROBES FOR DETECTION AND QUANTITATION OF ACTINOMYCETES

(57) Abstract: Polynucleotide probes and accessory helper oligonucleotides useful for detecting the subset of High (G+C) subset of Gram-positive bacteria known as the "Actinomycetes." The hybridization probes are highly specific for the Actinomycetes and do not cross-hybridize with the rRNA or rDNA of numerous other bacterial and fungal species.

**WO 00/66786 A3**

POLYNUCLEOTIDE PROBES FOR DETECTION AND QUANTITATION OF ACTINOMYCETES

5

Field of the Invention

The present invention relates to nucleic acid detection systems. More specifically, the invention relates to polynucleotide probes having binding specificity
10 for rRNA or rDNA of the subset of Gram-positive bacteria known as the "Actinomycetes."

Background of the Invention

The Actinomycetes, or the "High (G+C)" subset of Gram-positive (Gram⁽⁺⁾) bacteria, are a distinct evolutionary lineage within the eubacteria. The Actinomycetes
15 exhibit highly unusual phenotypic features as a reflection of a characteristically high mutation rate. Many members of this group of bacteria produce antibiotics and are commonly found in soil. Actinomycetes are responsible for a variety of significant animal diseases including tuberculosis, leprosy, diphtheria and periodontal diseases. Notably, immunodeficient individuals are particularly susceptible to infection by the
20 *Mycobacteria avium*, *Mycobacteria intracellulare*, and *Mycobacteria scrofulaceum*, all of which are individual species among the Actinomycetes. Additionally, the Actinomycetes are responsible for a variety of economically important plant diseases.

It is well established that two single strands of deoxyribonucleic acid ("DNA") or ribonucleic acid ("RNA") can associate or "hybridize" with one another to form a
25 double-stranded structure having two strands held together by hydrogen bonds between complementary base pairs. The individual strands of nucleic acid are formed from nucleotides that comprise the bases: adenine (A), cytosine (C), thymine (T), guanine (G), uracil (U) and inosine (I). In the double helical structure of nucleic acids, the base adenine hydrogen bonds with the base thymine or uracil, the base guanine hydrogen
30 bonds with the base cytosine and the base inosine hydrogen bonds with adenine, cytosine or uracil. At any point along the chain, therefore, one may find the classical

"Watson-Crick" base pairs A:T or A:U, T:A or U:A, and G:C or C:G. However, one may also find A:G, G:U and other "wobble" or mismatched base pairs in addition to the traditional ("canonical") base pairs.

A double-stranded nucleic acid hybrid will result if a first single-stranded
5 polynucleotide is contacted under hybridization-promoting conditions with a second single-stranded polynucleotide having a sufficient number of contiguous bases complementary to the sequence of the first polynucleotide. DNA/DNA, RNA/DNA or RNA/RNA hybrids may be formed under appropriate conditions.

Generally, a probe is a single-stranded polynucleotide having some degree of
10 complementarity with the nucleic acid sequence that is to be detected ("target sequence"). Probes commonly are labeled with a detectable moiety such as a radioisotope, an antigen or a chemiluminescent moiety.

Descriptions of nucleic acid hybridization as a procedure for detecting particular nucleic acid sequences are given by Kohne in U.S. Patent No. 4,851,330, and by Hogan
15 et al., in U.S. Patent Nos. 5,541,308 and 5,681,698. These references also describe methods for determining the presence of RNA-containing organisms in a sample which might contain such organisms. These procedures require probes that are sufficiently complementary to the ribosomal RNA (rRNA) of one or more non-viral organisms or groups of non-viral organisms. According to the method, nucleic acids from a sample
20 to be tested and an appropriate probe are first mixed and then incubated under specified hybridization conditions. Conventionally, but not necessarily, the probe will be labeled with a detectable label. The resulting hybridization reaction is then assayed to detect and quantitate the amount of labeled probe that has formed duplex structures in order to detect the presence of rRNA contained in the test sample.

25 With the exception of viruses, all prokaryotic organisms contain rRNA genes encoding homologs of the procaryotic 5S, 16S and 23S rRNA molecules. In eucaryotes, these rRNA molecules are the 5S rRNA, 5.8S rRNA, 18S rRNA and 28S rRNA which are substantially similar to the prokaryotic molecules. Probes for detecting specifically targeted rRNA subsequences in particular organisms or groups of
30 organisms in a sample have been described previously. These highly specific probe sequences advantageously do not cross react with nucleic acids from any other bacterial

species or infectious agent under appropriate stringency conditions.

The present invention provides polynucleotide probes that can be used to detect the Actinomycetes in a highly specific manner.

Summary of the Invention

5 Various embodiments of this invention provide an oligonucleotide probe that specifically hybridizes to an Actinomycetes nucleic acid target region corresponding to *E. coli* 23S rRNA nucleotide positions 1986 - 2064 under a high stringency hybridization condition provided by a solution comprising 0.48 M sodium phosphate or 0.6 M LiCl to form a detectable probe:target duplex, said oligonucleotide probe having a length of up to
10 100 nucleotides and comprising at least 17 contiguous nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof.

Other embodiments of this invention provide an oligonucleotide probe for detecting nucleic acids of Actinomycetes bacteria, wherein the oligonucleotide probe hybridizes under a high stringency condition provided by a solution comprising 0.48 M sodium
15 phosphate or 0.6 M LiCl to an Actinomycetes nucleic acid target region corresponding to *E. coli* 23S rRNA nucleotide positions 1986 - 2064 to form a detectable target:probe duplex, wherein said oligonucleotide probe has a length of up to 100 nucleotide bases and comprises at least 17 contiguous nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof, and wherein under said hybridization condition said
20 oligonucleotide probe specifically hybridizes to nucleic acids present in *Corynebacterium aquaticum*, *Corynebacterium jeikeium*, *Corynebacterium xerosis*, *Micrococcus luteus*, *Propionibacterium acnes*, *Mycobacterium chelonae*, *Mycobacterium terrae*, *Mycobacterium intracellulare*, *Mycobacterium simiae*, *Mycobacterium avium*, *Mycobacterium scrofulaceum*, *Mycobacterium gordonae*, *Mycobacterium kansasii*, *Mycobacterium smegatis*, *Mycobacterium fortuitum*, *Mycobacterium gastri*, *Mycobacterium xenopi*,
25 *Mycobacterium marinum* and *Mycobacterium phlei*.

Other embodiments of this invention provide a method for detecting the presence of Actinomycetes in a test sample, comprising the steps of: (a) providing to said test sample an oligonucleotide probe that hybridizes under a high stringency condition provided by a
30 solution comprising 0.48 M sodium phosphate or 0.6 M LiCl to an Actinomycetes nucleic acid target region corresponding to *E. coli* 23S rRNA nucleotide positions 1986 - 2064 to form a detectable target:probe duplex, said oligonucleotide probe having a length of up to

3a

100 nucleotide bases and comprising at least 25 contiguous nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof, and wherein under said hybridization condition said oligonucleotide probe specifically hybridizes to nucleic acids present in *Corynebacterium aquaticum*, *Corynebacterium jeikieum*, *Corynebacterium xerosis*, *Micrococcus luteus*, *Propionibacterium acnes*, *Mycobacterium chelonae*,
 5 *Mycobacterium terrae*, *Mycobacterium intracellulare*, *Mycobacterium simiae*, *Mycobacterium avium*, *Mycobacterium scrofulaceum*, *Mycobacterium gordonae*, *Mycobacterium kansasii*, *Mycobacterium smegatis*, *Mycobacterium fortuitum*, *Mycobacterium gastri*, *Mycobacterium xenopi*, *Mycobacterium marinum* and
 10 *Mycobacterium phlei*; (b) hybridizing under high stringency conditions any Actinomycetes nucleic acid that may be present in the test sample with said probe to form a probe:target duplex; and (c) detecting said probe:target duplex as an indicator of the presence of Actinomycetes in the test sample.

Other embodiments of this invention provide a kit for detecting the presence of
 15 Actinomycetes nucleic acids in a test sample, comprising: an oligonucleotide probe that hybridizes under a high stringency condition provided by a solution comprising 0.48 M sodium phosphate or 0.6 M LiCl to an Actinomycetes nucleic acid target region corresponding to *E. coli* 23S rRNA nucleotide positions 1986 - 2064 to form a detectable target:probe duplex, said oligonucleotide probe having a length of up to 100 nucleotide
 20 bases and comprising at least 25 contiguous nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof, and wherein under said hybridization condition said oligonucleotide probe specifically hybridizes to nucleic acids present in *Corynebacterium aquaticum*, *Corynebacterium jeikieum*, *Corynebacterium xerosis*, *Micrococcus luteus*, *Propionibacterium acnes*, *Mycobacterium chelonae*, *Mycobacterium terrae*, *Mycobacterium intracellulare*, *Mycobacterium simiae*, *Mycobacterium avium*,
 25 *Mycobacterium scrofulaceum*, *Mycobacterium gordonae*, *Mycobacterium kansasii*, *Mycobacterium smegatis*, *Mycobacterium fortuitum*, *Mycobacterium gastri*, *Mycobacterium xenopi*, *Mycobacterium marinum* and *Mycobacterium phlei*; and printed instructions specifying, in order of implementation, the steps to be followed for detecting said
 30 Actinomycetes nucleic acid by detecting a complex between the oligonucleotide probe and an Actinomycetes nucleic acid target, wherein said probe and said printed instructions are in packaged combination.

3b

One aspect of the present invention relates to an oligonucleotide probe that specifically hybridizes under high stringency hybridization conditions to a nucleic acid target region characteristic of Actinomycetes bacteria to form a detectable probe:target duplex. This target region corresponds to *E. coli* rRNA nucleotide positions 1986 - 2064. The invented oligonucleotide probe has a length of up to 100 nucleotides and includes at least 17 contiguous nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof. In certain embodiments, the oligonucleotide probe includes at least 25 contiguous nucleotides, and may include at least 29 contiguous nucleotides contained within the sequence of SEQ ID NO:10. The high stringency hybridization condition may be provided either by 0.48 M sodium phosphate buffer, 0.1% sodium dodecyl sulfate and 1 mM each of EDTA and EGTA, or by 0.6 M LiCl, 1% lithium lauryl sulfate, 60 mM lithium succinate and 10 mM each of EDTA and EGTA. The oligonucleotide probe may be made of DNA, but also may include at least one nucleotide analog. For example, the probe may include at least one nucleotide analog that has a methoxy group at the 2' position of a ribose moiety. In one embodiment the invented oligonucleotide probe has a sequence that is any one of SEQ ID NO:1 or the complement thereof, SEQ ID NO:2 or the complement thereof, or SEQ ID NO:3 or the complement thereof. In a preferred embodiment, the sequence of the oligonucleotide is given by SEQ ID NO:2 or SEQ ID NO:3, and the oligonucleotide is a helper oligonucleotide. Any of the disclosed oligonucleotides can include a detectable label. Particular examples of detectable labels include chemiluminescent labels and radiolabels. In another preferred embodiment, the oligonucleotide has a sequence given by SEQ ID NO:1. This oligonucleotide is particularly useful as a hybridization probe, and may include a detectable label. A highly preferred detectable label for the hybridization probe is an acridinium ester.

Another aspect of the present invention relates to a probe composition for detecting nucleic acids of Actinomycetes bacteria. This composition includes an

oligonucleotide probe that hybridizes under a high stringency condition to an Actinomycetes nucleic acid target region corresponding to *E. coli* 23S rRNA nucleotide positions 1986 - 2064 to form a detectable target:probe duplex. This oligonucleotide probe has a length of up to 100 nucleotide bases and includes at least 17 contiguous

5 nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof. Under high stringency hybridization conditions the oligonucleotide probe specifically hybridizes nucleic acids present in *Corynebacterium aquaticum*, *Corynebacterium jeikeium*, *Corynebacterium xerosis*, *Micrococcus luteus*, *Propionibacterium acnes*, *Mycobacterium chelonae*, *Mycobacterium terrae*,

10 *Mycobacterium intracellulare*, *Mycobacterium simiae*, *Mycobacterium avium*, *Mycobacterium scrofulaceum*, *Mycobacterium gordonae*, *Mycobacterium kansasii*, *Mycobacterium smegatis*, *Mycobacterium fortuitum*, *Mycobacterium gastri*, *Mycobacterium xenopi*, *Mycobacterium marinum* and *Mycobacterium phlei*. It is preferred that the oligonucleotide probe have a length of up to 100 nucleotide bases and

15 include at least 25 contiguous nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof. In certain embodiments, the invented oligonucleotide probe is made of DNA. Examples of useful high stringency hybridization conditions are alternatively provided by 0.48 M sodium phosphate buffer, 0.1% sodium dodecyl sulfate, and 1 mM each of EDTA and EGTA, or by 0.6 M LiCl,

20 1% lithium lauryl sulfate, 60 mM lithium succinate and 10 mM each of EDTA and EGTA. In a highly preferred embodiment, the oligonucleotide probe has the sequence of SEQ ID NO:1 or the complement thereof. In a preferred embodiment, the length of the oligonucleotide probe may be up to 60 bases. A highly preferred oligonucleotide probe has the length and sequence of SEQ ID NO:1. Certain embodiments of the

25 invented probe composition include a detectable label on the oligonucleotide probe. For example, when the oligonucleotide probe has a length of up to 60 nucleotides, the oligonucleotide probe can include a detectable label. Alternatively, when the oligonucleotide probe has the sequence of SEQ ID NO:1 there can be included a detectable label. Regardless of whether the probe composition includes a labeled

30 oligonucleotide probe of from 17-100 nucleotides in length, or from 17-60 nucleotides in length, or having the sequence of SEQ ID NO:1, the detectable label may be either a

chemiluminescent label (such as an acridinium ester) or a radiolabel. It is preferred that these labeled oligonucleotide probes are used in combination with at least one helper oligonucleotide that facilitates formation of the detectable probe:target duplex under high stringency hybridization conditions. At least one of the helper oligonucleotides
5 can include at least one nucleotide analog. An example of a highly preferred nucleotide analog would be one in which a ribose moiety has a methoxy group disposed at the 2' position. In a highly preferred embodiment of the invention, the labeled oligonucleotide probe, regardless of its length, is used in combination with at least one helper oligonucleotide having the sequence of SEQ ID NO:2 or SEQ ID NO:3.

10 Yet another aspect of the invention relates to a method for detecting the presence of Actinomycetes bacteria in a test sample. This method involves steps for providing to said test sample a probe composition comprising an oligonucleotide probe that hybridizes under a high stringency condition to an Actinomycetes nucleic acid target region corresponding to *E. coli* 23S rRNA nucleotide positions 1986 - 2064 to
15 form a detectable target:probe duplex, said oligonucleotide probe having a length of up to 100 nucleotide bases and comprising at least 25 contiguous nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof, and wherein under said hybridization condition said oligonucleotide probe specifically hybridizes nucleic acids present in *Corynebacterium aquaticum*, *Corynebacterium jeikeium*,
20 *Corynebacterium xerosis*, *Micrococcus luteus*, *Propionibacterium acnes*, *Mycobacterium chelonae*, *Mycobacterium terrae*, *Mycobacterium intracellulare*, *Mycobacterium simiae*, *Mycobacterium avium*, *Mycobacterium scrofulaceum*, *Mycobacterium gordonae*, *Mycobacterium kansasii*, *Mycobacterium smegatis*, *Mycobacterium fortuitum*, *Mycobacterium gastri*, *Mycobacterium xenopi*,
25 *Mycobacterium marinum* and *Mycobacterium phlei*. Thereafter, the resulting mixture is hybridized under high stringency conditions so that any Actinomycetes nucleic acid that may be present in the test sample combines with the probe composition to form a probe:target duplex. Finally, the method involves detecting the probe:target duplex as an indicator of the presence of Actinomycetes bacteria in the test sample. In one
30 embodiment of the invented method the test sample includes bacteria, and there is conducted a preliminary step for releasing nucleic acid from any bacteria that may be

present in said test sample. In a different embodiment of the method the test sample is a lysate. In general, the high stringency hybridization conditions can be provided either by: (a) 0.48 M sodium phosphate buffer, 0.1% sodium dodecyl sulfate, and 1 mM each of EDTA and EGTA, or (b) 0.6 M LiCl, 1% lithium lauryl sulfate, 60 mM lithium succinate and 10 mM each of EDTA and EGTA. However, it is to be understood that other high stringency hybridization conditions can give good results. In a preferred embodiment, the oligonucleotide probe used in the invented method has the length and sequence of SEQ ID NO:1. When this is the case, the oligonucleotide probe may include a detectable label. In a highly preferred embodiment the labeled oligonucleotide probe has the sequence of SEQ ID NO:1, the label on the probe is an acridinium ester, and the detecting step in the invented method involves performing luminometry to detect probe:target duplexs. In instances wherein the labeled oligonucleotide probe has the sequence of SEQ ID NO:1, the probe composition further may include at least one helper oligonucleotide that facilitates formation of probe:target duplexs. Exemplary helper oligonucleotide have the sequences of SEQ ID NO:2 and SEQ ID NO:3.

Still yet another aspect of the invention relates to a kit that can be used for detecting the presence of Actinomycetes nucleic acids in a test sample. The kit contains a probe composition that includes an oligonucleotide probe that hybridizes under a high stringency condition to an Actinomycetes nucleic acid target region corresponding to *E. coli* 23S rRNA nucleotide positions 1986 - 2064 to form a detectable target:probe duplex. the oligonucleotide probe has a length of up to 100 nucleotide bases and includes at least 25 contiguous nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof. Under the high stringency hybridization condition the oligonucleotide probe specifically hybridizes nucleic acids present in *Corynebacterium aquaticum*, *Corynebacterium jeikeium*, *Corynebacterium xerosis*, *Micrococcus luteus*, *Propionibacterium acnes*, *Mycobacterium chelonae*, *Mycobacterium terrae*, *Mycobacterium intracellulare*, *Mycobacterium simiae*, *Mycobacterium avium*, *Mycobacterium scrofulaceum*, *Mycobacterium gordonae*, *Mycobacterium kansasii*, *Mycobacterium smegatis*, *Mycobacterium fortuitum*, *Mycobacterium gastri*, *Mycobacterium xenopi*, *Mycobacterium marinum* and

Mycobacterium phlei. Additionally, the invented kit includes printed instructions specifying, in order of implementation, the steps to be followed for detecting the Actinomycetes nucleic acid by detecting a complex between the oligonucleotide probe and an Actinomycetes nucleic acid target. Both the probe composition and the printed instructions are in packaged combination with each other.

Definitions

As used herein, the following terms have the given meanings unless expressly stated to the contrary.

A “nucleotide” is a subunit of a nucleic acid consisting of a phosphate group, a 5-carbon sugar and a nitrogenous base. The 5-carbon sugar found in RNA is ribose. In DNA, the 5-carbon sugar is 2'-deoxyribose. The sugar of a 5'-nucleotide contains a hydroxyl group (-OH) at the 5'-carbon-5 position. The term also includes analogs of naturally occurring nucleotides and particularly includes analogs having a methoxy group at the 2' position of the ribose (OMe). As used herein, methoxy oligonucleotides containing “T” residues have a methoxy group at the 2' position of the ribose moiety, and a uracil at the base position of the nucleotide.

A “non-nucleotide unit” is a unit which does not significantly participate in hybridization of a polymer. Such units must not, for example, participate in any significant hydrogen bonding with a nucleotide, and would exclude units having as a component one of the five nucleotide bases or analogs thereof.

An “oligonucleotide” is a nucleotide polymer having two or more nucleotide subunits covalently joined together. Oligonucleotides are generally about 10 to about 100 nucleotides in length. The sugar groups of the nucleotide subunits may be ribose, deoxyribose, or modified derivatives thereof such as OMe. The nucleotide subunits may be joined by linkages such as phosphodiester linkages, modified linkages or by non-nucleotide moieties that do not prevent hybridization of the oligonucleotide to its complementary target nucleotide sequence. Modified linkages include those in which a standard phosphodiester linkage is replaced with a different linkage, such as a phosphorothioate linkage, a methylphosphonate linkage, or a neutral peptide linkage. Nitrogenous base analogs also may be components of oligonucleotides in accordance with the invention.

A "target nucleic acid" is a nucleic acid comprising a target nucleic acid sequence.

A "target nucleic acid sequence," "target nucleotide sequence" or "target sequence" is a specific deoxyribonucleotide or ribonucleotide sequence that can be
5 hybridized by an oligonucleotide.

An "oligonucleotide probe" is an oligonucleotide having a nucleotide sequence sufficiently complementary to its target nucleic acid sequence to be able to form a detectable hybrid probe:target duplex under high stringency hybridization conditions. An oligonucleotide probe is an isolated chemical species and may include additional
10 nucleotides outside of the targeted region as long as such nucleotides do not prevent hybridization under high stringency hybridization conditions. Non-complementary sequences, such as promotor sequences, restriction endonuclease recognition sites, or sequences that confer a desired secondary or tertiary structure such as a catalytic active site can be used to facilitate detection using the invented probes. An oligonucleotide
15 probe optionally may be labeled with a detectable moiety such as a radioisotope, a fluorescent moiety, a chemiluminescent moiety, an enzyme or a ligand, which can be used to detect or confirm probe hybridization to its target sequence. Oligonucleotide probes are preferred to be in the size range of from 10 to 100 nucleotides in length.

A "detectable moiety" is a molecule attached to, or synthesized as part of, a
20 nucleic acid probe. This molecule should be uniquely detectable and will allow the probe to be detected as a result. These detectable moieties are often radioisotopes, chemiluminescent molecules, enzymes, haptens, or even unique oligonucleotide sequences.

A "hybrid" or a "duplex" is a complex formed between two single-stranded
25 nucleic acid sequences by Watson-Crick base pairings or non-canonical base pairings between the complementary bases.

"Hybridization" is the process by which two complementary strands of nucleic acid combine to form a double-stranded structure ("hybrid" or "duplex").

"Complementarity" is a property conferred by the base sequence of a single
30 strand of DNA or RNA which may form a hybrid or double-stranded DNA:DNA, RNA:RNA or DNA:RNA through hydrogen bonding between Watson-Crick base pairs

on the respective strands. Adenine (A) ordinarily complements thymine (T) or uracil (U), while guanine (G) ordinarily complements cytosine (C).

“Mismatch” refers to any pairing, in a hybrid, of two nucleotides which do not form canonical Watson-Crick hydrogen bonds. In addition, for the purposes of the following discussions, a mismatch can include an insertion or deletion in one strand of the hybrid which results in an unpaired nucleotide(s).

The term “stringency” is used to describe the temperature and solvent composition existing during hybridization and the subsequent processing steps. Under high stringency conditions only highly complementary nucleic acid hybrids will form; hybrids without a sufficient degree of complementarity will not form. Accordingly, the stringency of the assay conditions determines the amount of complementarity needed between two nucleic acid strands forming a hybrid. Stringency conditions are chosen to maximize the difference in stability between the hybrid formed with the target and the non-target nucleic acid. Exemplary high stringency conditions are provided in the working Examples.

The term “probe specificity” refers to a characteristic of a probe which describes its ability to distinguish between target and non-target sequences.

The term “variable region” refers to a nucleotide polymer which differs by at least one base between the target organism and non-target organisms contained in a sample.

A “conserved region” is a nucleic acid subsequence which is not variable between at least two different polynucleotides.

“Bacteria” are members of the phylogenetic group eubacteria, which is considered one of the three primary kingdoms.

The term “sequence divergence” refers to a process by which nucleotide polymers become less similar during evolution.

The term “sequence convergence” refers to a process by which nucleotide polymers become more similar during evolution.

“T_m” refers to the temperature at which 50% of the probe is converted from the hybridized to the unhybridized form.

A "helper oligonucleotide" is an oligonucleotide that binds a region of a target nucleic acid other than the region that is bound by an oligonucleotide probe. Helper oligonucleotides impose new secondary and tertiary structures on the targeted region of the single-stranded nucleic acid so that the rate of binding of the oligonucleotide probe is accelerated. Although helper oligonucleotides are not labeled with a detectable label when used in conjunction with labeled oligonucleotide probes, they facilitate binding of labeled probes and so indirectly enhance hybridization signals.

The phrases "consist essentially of" or "consisting essentially of" means that the oligonucleotide has a nucleotide sequence substantially similar to a specified nucleotide sequence. Any additions or deletions are non-material variations of the specified nucleotide sequence which do not prevent the oligonucleotide from having its claimed property, such as being able to preferentially hybridize under high stringency hybridization conditions to its target nucleic acid over non-target nucleic acids.

One skilled in the art will understand that substantially corresponding probes of the invention can vary from the referred-to sequence and still hybridize to the same target nucleic acid sequence. This variation from the nucleic acid may be stated in terms of a percentage of identical bases within the sequence or the percentage of perfectly complementary bases between the probe and its target sequence. Probes of the present invention substantially correspond to a nucleic acid sequence if these percentages are from 100% to 80% or from 0 base mismatches in a 10 nucleotide target sequence to 2 bases mismatched in a 10 nucleotide target sequence. In preferred embodiments, the percentage is from 100% to 85%. In more preferred embodiments, this percentage is from 90% to 100%; in other preferred embodiments, this percentage is from 95% to 100%.

By "sufficiently complementary" or "substantially complementary" is meant nucleic acids having a sufficient amount of contiguous complementary nucleotides to form, under high stringency hybridization conditions, a hybrid that is stable for detection.

By "nucleic acid hybrid" or "probe:target duplex" is meant a structure that is a double-stranded, hydrogen-bonded structure, preferably 10 to 100 nucleotides in length, more preferably 14 to 50 nucleotides in length. The structure is sufficiently stable to be

detected by means such as chemiluminescent or fluorescent light detection, autoradiography, electrochemical analysis or gel electrophoresis. Such hybrids include RNA:RNA, RNA:DNA, or DNA:DNA duplex molecules.

By "negative sense" is meant a nucleic acid molecule perfectly complementary
5 to a reference (i.e., sense) nucleic acid molecule.

"RNA and DNA equivalents" refer to RNA and DNA molecules having the same complementary base pair hybridization properties. RNA and DNA equivalents have different sugar groups (i.e., ribose versus deoxyribose), and may differ by the presence of uracil in RNA and thymine in DNA. The difference between RNA and
10 DNA equivalents do not contribute to differences in substantially corresponding nucleic acid sequences because the equivalents have the same degree of complementarity to a particular sequence.

By "preferentially hybridize" is meant that under high stringency hybridization conditions oligonucleotide probes can hybridize their target nucleic acids to form stable
15 probe:target hybrids (thereby indicating the presence of the target nucleic acids) without forming stable probe:non-target hybrids (that would indicate the presence of non-target nucleic acids from other organisms). Thus, the probe hybridizes to target nucleic acid to a sufficiently greater extent than to non-target nucleic acid to enable one skilled in the art to accurately detect the presence of Actinomycetes bacteria and distinguish their
20 presence from that of other organisms. Preferential hybridization can be measured using techniques known in the art and described herein. For example, when compared with hybridization to *C. albicans* nucleic acids, oligonucleotide probes of the invention preferentially hybridize Actinomycetes nucleic acids by about 100 - 2,000 fold.

An Actinomycetes "target nucleic acid sequence region" refers to a nucleic acid
25 sequence present in the nucleic acid of Actinomycetes bacteria or a sequence complementary thereto, which is not present in the nucleic acids of other species. Nucleic acids having nucleotide sequences complementary to a target sequence may be generated by target amplification techniques such as polymerase chain reaction (PCR) or transcription mediated amplification (e.g., Kacian and Fultz, Nucleic Acid Sequence
30 Amplification Methods, U.S. Patent No. 5,824,518).

Brief Description of the Drawings

Figure 1 shows an alignment of one oligonucleotide probe and two helper oligonucleotides with a collection of positively reactive and non-reactive target sequences.

Detailed Description of the Invention

5 Herein we disclose preferred target nucleotide sequences for oligonucleotide probes and helper oligonucleotides that can be used to detect and identify rRNA or rDNA of Actinomycetes bacteria. Highly preferred polynucleotide probes and accessory helper oligonucleotides that are useful for specifically detecting
10 Actinomycetes are particularly disclosed. The probes, which are complementary to particular rRNA sequences of the 23S rRNA, advantageously are capable of distinguishing Actinomycetes from the known phylogenetically nearest neighbors.

In addition to having nucleic acid sequences that permit hybridization to the ribosomal RNA (rRNA) or DNA (rDNA) sequences of Actinomycetes bacteria, the
15 oligonucleotide probes of the invention are at least 90% complementary, preferably perfectly complementary, to at least a portion of the described target sequence region identified by SEQ ID NO:11. The portion is preferably at least 17 nucleotides in length, more preferably at least 25 nucleotides in length, and still more preferably at least 29 nucleotides in length.

20 As indicated above, the invented oligonucleotides are targeted to nucleic acid sequences of Actinomycetes bacteria. These oligonucleotides can be used as probes that preferentially hybridize to an Actinomycetes nucleic acid target region to form a detectable duplex that indicates the presence of Actinomycetes bacteria. Alternatively, the invented oligonucleotides can be used as Helper oligonucleotides that hybridize to
25 an Actinomycetes nucleic acid target region under high stringency hybridization conditions, and that can enhance the formation of a duplex between a labeled oligonucleotide probe and its complementary target nucleic acid.

In preferred embodiments, the oligonucleotide probes described herein selectively hybridize nucleic acids from Actinomycetes bacteria over those from other
30 organisms under high stringency hybridization conditions. In some embodiments of the present invention, the oligonucleotide probe comprises a detectable moiety, such as an

acridinium ester or a radioisotope.

Preferred methods for detecting the presence of Actinomycetes bacteria include the step of contacting a test sample under high stringency hybridization conditions with an oligonucleotide probe that preferentially hybridizes to an Actinomycetes target
5 nucleic acid sequence over a nucleic acid sequence of other organisms. Preferably the target nucleic acid sequence is fully complementary to a sequence of at least 17 contiguous nucleotides, more preferably at least 25 contiguous nucleotides, still more preferably at least 29 contiguous nucleotides contained in the sequence given by SEQ ID NO:11. Thus, oligonucleotides useful in connection with the invention preferably
10 are up to 100 nucleotides in length and have at least 17 contiguous nucleotides, more preferably at least 25 contiguous nucleotides, still more preferably at least 29 contiguous nucleotides contained in the sequence given by
GGTCTTTCCGTCCTGCCGCGCGTAACGAGCATCTTTACTCGTAGTGCAATTT
CGCCGAGTCTGTGGTTGAGACAGTGGG (SEQ ID NO:10) or the complement
15 thereof. The oligonucleotides may be RNA and DNA equivalents, and may contain nucleotide analogs.

Introduction and Background

In the development of the invention, rRNA sequences from a collection of related and unrelated organisms were aligned to identify candidate conserved sequences
20 present in the 23S rRNA that could be used to distinguish Actinomycetes organisms from other bacterial as well as eukaryotic organisms. The rRNA or rDNA sequences of Actinomycetes and distant phylogenetic neighbors were aligned to reveal areas of maximum homology. Homologous regions were examined for sequence variation in order to identify rRNA sequences that were conserved among the Actinomycetes and
25 that showed mismatches with other closely and distantly related genera. The sequences deduced as candidate probes according to the methods described below finally were tested against a panel of rRNA standards and bacterial lysates to verify their utility as probes under laboratory conditions.

Polynucleotide sequences of rRNAs are most conveniently determined using a
30 dideoxynucleotide sequencing procedure. In this procedure, oligonucleotide primers of about 10-100 bases in length and complementary to conserved regions of rRNA from

any of the 5S, 16S or 23S ribosome subunits can be extended by reverse transcriptase. The resulting DNA extension products can then be sequenced either by chemical degradation or by dideoxynucleotide sequencing (Lane et al., *Proc. Natl. Acad. Sci. USA* 82: 6955 (1985)). According to another preferred method, genomic sequences
5 encoding the rRNA can also be determined.

The strong interdependence of secondary structure and function of the rRNA molecules is well known. Indeed, evolutionary changes in the primary sequence of the rRNA are effectively restricted such that secondary structure of the molecule will be maintained. For example, if a base is changed on one side of a helix of a rRNA
10 molecule, then a compensating change will be made on the other side of the helix to preserve complementarity (this is referred to as covariance). This relationship allows two very different rRNA sequences to be "aligned" based on conserved primary sequence and conserved elements of the secondary structure. Once the sequences have been aligned, it becomes possible to identify conserved and variable regions of the
15 rRNA sequence.

Variable regions of rRNAs were identified by comparative analysis using published rRNA sequences and sequences that were determined during the development of the present invention. Commercially available software can be used or adapted for the purposes disclosed herein. Since the sequence evolution at each of the
20 variable regions (for example, spanning a minimum of 10 nucleotides) of rRNA is, for the most part, divergent and not convergent, we can confidently design probes based on a few rRNA sequences which differ between the target organism and its phylogenetically closest relatives. Indeed, we have detected sufficient variation between the rRNA sequences of numerous target organisms and their closest
25 phylogenetic relatives in a single sample to permit the design of a probe that can be used according to the methods described below.

Probe Selection Guidelines

The following general guidelines can be used for designing probes having desirable characteristics in accordance with the present invention. Manipulation of one
30 or more of the many factors that influence the extent and specificity of a hybridization reaction can determine the sensitivity and specificity of a particular probe. This is true

whether or not the probe is perfectly complementary over the full length of its target polynucleotide sequence. Guidelines for preparing probes useful in connection with the invention now follow.

First, the stability of the probe:target nucleic acid hybrid should be chosen to be compatible with the assay conditions. This may be accomplished by avoiding long A and T rich sequences, by terminating the hybrids with G:C base pairs and by designing the probe in such a way that the T_m will be appropriate for standard conditions to be employed in the assay. The nucleotide sequence of the probe should be chosen so that the length and %G and %C result in a probe having a T_m about 2-10°C higher than the temperature at which the final assay will be performed. The base composition of the probe is significant because G:C base pairs exhibit greater thermal stability when compared with A:T or A:U base pairs. Thus, hybrids involving complementary nucleic acids having a high G:C content will be stable at higher temperatures when compared with hybrids having a lower G:C content.

Ionic strength and temperature conditions at which a hybridization reaction will be conducted also should be considered when designing a probe having a negatively charged backbone, such as would be provided by phosphodiester linkages between nucleotides. It is generally known that hybridization rate increases as ionic strength of the reaction mixture increases. Similarly, the thermal stability of hybrids increases with increasing ionic strength. Conversely, hydrogen bond-disrupting reagents such as formamide, urea, DMSO and alcohols increase the stringency of hybridization. Destabilization of hydrogen bonds by reagents in this class can greatly reduce the T_m . In general, optimal hybridization for synthetic oligonucleotide probes of about 10-50 bases in length occurs approximately 5° C below the melting temperature for a given duplex. Hybridization reactions conducted below the temperature optimum may allow mismatched base sequences to hybridize and can result in reduced probe specificity.

Second, the position at which the probe binds its target polynucleotide should be chosen to minimize the stability of hybrids formed between probe:non-target polynucleotides. This may be accomplished by minimizing the length of perfect complementarity with polynucleotides of non-target organisms, by avoiding G:C rich regions of homology with non-target sequences, and by positioning the probe to span as

many destabilizing mismatches as possible. Whether a probe sequence will be useful for detecting only a specific type of organism depends largely on thermal stability differences between probe:target hybrids and probe:non-target hybrids. The differences in T_m should be as large as possible to produce highly specific probes.

5 The length of the target nucleic acid sequence and the corresponding length of the probe sequence also are important factors to be considered when designing a probe useful for specifically detecting Actinomycetes. While it is possible for polynucleotides that are not perfectly complementary to hybridize to each other, the longest stretch of perfectly homologous base sequence will ordinarily be the primary
10 determinant of hybrid stability.

Third, regions of the rRNA which are known to form strong internal structures inhibitory to hybridization of a probe are less preferred as targets. Probes having extensive self-complementarity also should be avoided. As indicated above, hybridization is the association of two single strands of complementary nucleic acid to
15 form a hydrogen bonded double-stranded structure. If one of the two strands is wholly or partially double-stranded, then it will be less able to participate in the formation of a new hybrid. Significantly, all rRNA molecules form very stable intramolecular hybrids.

The rate and extent of hybridization between a probe and its target can be increased substantially by designing the probe such that a substantial portion of the
20 sequence of interest is single-stranded. If the target nucleic acid to be detected is a genomic sequence encoding a rRNA, then that target will naturally occur in a double-stranded form. This is also the case with products of the polymerase chain reaction (PCR). These double-stranded targets are naturally inhibitory to hybridization with a probe. Finally, undesirable intramolecular and intermolecular hybrids can form within
25 a single probe molecule or between different probe molecules if there is sufficient self-complementarity. Thus, extensive self-complementarity in a probe sequence should be avoided.

Preferably, probes useful for carrying out the procedures described below will hybridize only under conditions of high stringency. Under these conditions only highly
30 complementary nucleic acid hybrids will form (i.e., those having at least 14 out of 17 bases in a contiguous series of bases being complementary). Hybrids will not form in

the absence of a sufficient degree of complementarity. Accordingly, the stringency of the assay conditions determines the amount of complementarity needed between two nucleic acid strands forming a hybrid. Stringency is chosen to maximize the difference in stability between the hybrid formed with the target and non-target nucleic acid.

- 5 Exemplary high stringency conditions are employed in the Examples presented below.

While oligonucleotide probes of different lengths and base composition may be used for detecting Actinomycetes, preferred probes in this invention have lengths of up to 100 nucleotides, and more preferably 60 nucleotides. Preferred length ranges for the invented oligonucleotides are from 10 to 100 bases in length, or more preferably
10 between 15 and 50 bases in length, and are sufficiently homologous to the target nucleic acid to permit hybridization under high stringency conditions such as those employed in the Examples described below. However, the specific probe sequences described below also may be provided in a nucleic acid cloning vector or transcript or other longer nucleic acid and still can be used for detecting members of the
15 Actinomycetes.

Chemical Structure of Oligonucleotides

All of the oligonucleotides of the present invention may be modified with chemical groups to enhance their performance. Thus, it is to be understood that references to "oligonucleotide probes" or "helper oligonucleotides" or simply
20 "oligonucleotides" embrace polymers of native nucleotides as well as polymers that include at least one nucleotide analog.

Backbone-modified oligonucleotides, such as those having phosphorothioate or methylphosphonate groups, are examples of analogs that can be used in conjunction with oligonucleotides of the present invention. These modifications render the
25 oligonucleotides resistant to the nucleolytic activity of certain polymerases or to nuclease enzymes. Other analogs that can be incorporated into the structures of the oligonucleotides disclosed herein include peptide nucleic acids, or "PNAs." The PNAs are compounds comprising ligands linked to a peptide backbone rather than to a phosphodiester backbone. Representative ligands include either the four main naturally
30 occurring DNA bases (i.e., thymine, cytosine, adenine or guanine) or other naturally occurring nucleobases (e.g., inosine, uracil, 5-methylcytosine or thiouracil) or artificial

bases (e.g., bromothymine, azaadenines or azaguanines, etc.) attached to a peptide backbone through a suitable linker. The PNAs are able to bind complementary ssDNA and RNA strands. Methods for making and using PNAs are disclosed in U.S. Patent No. 5,539,082. Another type of modification that can be used to make oligonucleotides
5 having the sequences described herein involves the use of non-nucleotide linkers (e.g., Arnold, et al., "Non-Nucleotide Linking Reagents for Nucleotide Probes", U.S. Patent No. 6,031,091 incorporated between nucleotides in the nucleic acid chain which do not interfere with hybridization or the elongation of a primer.

10 Nucleic Acid Based Methods of Detecting rRNA or rDNA

A composition that includes an oligonucleotide probe, either alone or in combination with one or more helper oligonucleotides, can be used for detecting rRNA or rDNA of Actinomycetes bacteria in a hybridization assay. Defined oligonucleotides that can be used to practice the invention can be produced by any of several well-known
15 methods, including automated solid-phase chemical synthesis using cyanoethylphosphoramidite precursors (Barone et al., *Nucl Acids Res* 12:4051 (1984)). Other well-known methods for preparing synthetic oligonucleotides also may be employed.

Essentially any labeling and detection system that can be used for monitoring
20 specific nucleic acid hybridization can be used in conjunction with the probes disclosed herein when a labeled probe is desired. Included among the collection of useful labels are: isotopic labels, enzymes, haptens, linked oligonucleotides, chemiluminescent molecules and redox-active moieties that are amenable to electrochemical detection methods. Standard isotopic labels that can be used to produce labeled oligonucleotides
25 include ^3H , ^{35}S , ^{32}P , ^{125}I , ^{57}Co and ^{14}C . When using radiolabeled probes, hybrids can be detected by autoradiography, scintillation counting or gamma counting.

Non-isotopic materials can also be used for labeling oligonucleotide probes. These non-isotopic labels can be positioned internally or at a terminus of the oligonucleotide probe. Modified nucleotides may be incorporated enzymatically or
30 chemically with modifications of the probe being performed during or after probe synthesis, for example, by the use of non-nucleotide linker groups. Non-isotopic labels

include fluorescent molecules, chemiluminescent molecules, enzymes, cofactors, enzyme substrates, haptens or other ligands. Acridinium esters are particularly preferred non-isotopic labels for detecting probe hybrids.

Indeed, any number of different non-isotopic labels can be used for preparing
5 labeled oligonucleotides in accordance with the invention. Preferred chemiluminescent molecules include acridinium esters of the type disclosed by Arnold et al., in U.S. Patent No. 5,283,174 for use in connection with homogenous protection assays, and of the type disclosed by Woodhead et al., in U.S. Patent No. 5,656,207 for use in connection with assays that quantify multiple targets in a single reaction.

10

U.S. Patent 5,998,135 discloses yet another method that can be used for labeling and detecting the probes of the present invention using fluorimetry to detect fluorescence emission from lanthanide metal labels disposed on probes, where the emission from these labels becomes enhanced when it is in close proximity to an energy transfer
15 partner. Preferred electrochemical labeling and detection approaches are disclosed in U.S. Patent Nos. 5,591,578 and 5,770,369, and the published International Patent Application No. PCT/US98/12082.

Redox active moieties useful as electrochemical labels in the present invention include transition metals such as Cd, Mg, Cu, Co, Pd, Zn, Fe and Ru.

20

Those having an ordinary level of skill in the art will appreciate that alternative procedures for detecting nucleic acids of Actinomycetes bacteria using the invented probes can be carried out using either labeled probes or unlabeled probes. For example, hybridization assay methods that do not rely on the use of a labeled probe are disclosed in U.S. Patent No. 5,945,286 which describes immobilization of unlabeled probes made
25 of peptide nucleic acids (PNAs), and detectably labeled intercalating molecules which can bind double-stranded PNA probe/target nucleic acid duplexes. In these procedures, as well as in certain electrochemical detection procedures, such as those disclosed in published International Patent Application No. PCT/US98/12082 entitled "Detection of Analytes Using Reorganization Energy," published International Patent Application
30 No. PCT/US98/12430 entitled "Electronic Methods for the Detection of Analytes," and in published International Patent Application No. PCT/US97/20014 entitled

“Electrodes Linked Via Conductive Oligomers to Nucleic Acids” the oligonucleotide probe is not required to harbor a detectable label.

Acceptability of the final product following synthesis and purification of an oligonucleotide may be verified by any of several procedures. First, polyacrylamide gel electrophoresis can be used to determine the size and purity of the oligonucleotide according to standard laboratory methods (see Molecular Cloning: A Laboratory Manual, Sambrook et al., eds. Cold Spring Harbor Lab Publ., 11.51, (1989)). Alternatively, High Pressure Liquid Chromatography (“HPLC”) procedures can be used for this same purpose.

Hybridization between the labeled oligonucleotide probe and target nucleic acid in the procedures described below can be enhanced through the use of unlabeled “helper oligonucleotides” according to the procedure disclosed by Hogan et al., in U.S. Patent No. 5,030,557 entitled, “Means and Methods for Enhancing Nucleic Acid Hybridization.” As indicated above, helper oligonucleotides bind a region of the target nucleic acid other than the region that is bound by the assay probe. This binding imposes new secondary and tertiary structures on the targeted region of the single-stranded nucleic acid and accelerates the rate of probe binding. Helper oligonucleotides which can be used in combination with labeled oligonucleotide probes of the present invention are preferably 17 to 100 nucleotides in length and have a sequence that includes at least 17 contiguous nucleotides contained within the sequence of SEQ ID NO:10. Other preferred helper oligonucleotides have lengths of up to 100 nucleotides and include at least 25 contiguous nucleotides contained within the sequence of SEQ ID NO:10.

Those having ordinary skill in the art will appreciate that factors affecting the thermal stability of a probe:target hybrid also can influence probe specificity. Accordingly, the melting profile, including the melting temperature (T_m) of probe:target hybrids, should be empirically determined for each probe:target combination. A preferred method for making this determination is described by Arnold et al., in U.S. Patent No. 5,283,174, entitled “Homogeneous Protection Assay.”

One approach for measuring the T_m of a probe:target hybrid involves conducting a hybridization protection assay. According to the method of this assay, a

probe:target hybrid is formed under conditions of target excess in a lithium succinate buffered solution containing lithium lauryl sulfate. Aliquots of the "preformed" hybrids are diluted in the hybridization buffer and incubated for five minutes at various temperatures starting below the anticipated T_m (typically 55°C) and increasing in 2-5
5 degree increments. This solution is then diluted with a mildly alkaline borate buffer and incubated at a lower temperature (for example 50°C) for ten minutes. An acridinium ester (AE) linked to a single-stranded probe will be hydrolyzed under these conditions while an acridinium ester linked to a hybridized probe will be relatively "protected." This procedure is referred to as the hybridization protection assay
10 ("HPA"). The amount of chemiluminescence remaining is proportional to the amount of hybrid and is measured in a luminometer by addition of hydrogen peroxide followed by alkali. The data is plotted as percent of maximum signal (usually from the lowest temperature) versus temperature. The T_m is defined as the point at which 50% of the maximum signal remains.

15 In an alternative approach, the T_m of a probe:target hybrid can be determined using an isotopically labeled probe. In all cases, the T_m for a given hybrid will vary depending on the concentration of salts, detergents and other solutes contained in the hybridization solution. All of these factors influence relative hybrid stability during thermal denaturation (Molecular Cloning: A Laboratory Manual Sambrook et al., eds.
20 Cold Spring Harbor Lab Publ., 9.51 (1989)).

The rate at which a probe hybridizes to its target is a measure of the thermal stability of the target secondary structure in the probe region, and can be determined using $C_0t_{1/2}$ measurements. These kinetic measurements of hybridization rate have units of (moles of nucleotide per liter) x (seconds). Expressed more simply, the $C_0t_{1/2}$ value
25 is the concentration of probe times the half-life of hybridization at that concentration. This value can be determined by hybridizing various amounts of probe to a constant amount of target nucleic acid for a fixed time. For example, 0.05 pmol of target is incubated with 0.012, 0.025, 0.05, 0.1 and 0.2 pmol of probe for 30 minutes. The $C_0t_{1/2}$ may also be determined by hybridizing the target and probe under conditions of target
30 excess and then measuring the increase of duplex formation over time. The amount of hybrid present can be measured using the above-described HPA procedure or by

scintillation counting if a radiolabeled probe is used in the procedure. The measured signal, when using AE labeled probe, is then plotted as the log of the percent of maximum Relative Light Units ("RLU") from the highest probe concentration versus probe concentration (moles of nucleotide per liter). The $C_0t_{1/2}$ is graphically
5 determined from the concentration corresponding to 50% of maximum hybridization multiplied by the hybridization time in seconds. These values range from 9×10^{-6} to 9×10^{-5} with the preferred values being less than 3.5×10^{-5} . Similar values may be obtained by measuring radioactivity and plotting % hybridization at a given time point vs maximum extent.

10 In a preferred method of determining whether a biological sample contains rRNA or rDNA that would indicate the presence of members of the Actinomycetes, nucleic acids may be released from bacterial cells by sonic disruption, for example according to the method disclosed by Murphy et al., in U.S. Patent No. 5,374,522. Other known methods for disrupting cells include the use of enzymes, osmotic shock,
15 chemical treatment, and vortexing with glass beads. Other methods suitable for liberating from microorganisms the nucleic acids that can be subjected to the hybridization methods disclosed herein have been described by Clark et al., in U.S. Patent No. 5,837,452 and by Kacian et al., in U.S. Patent No. 5,536,763. Following or concurrent with the release of rRNA, labeled probe may be added in the presence of
20 accelerating agents and incubated at the optimal hybridization temperature for a period of time necessary to achieve significant hybridization reaction.

The following polynucleotide sequence was characterized by the criteria of length, T_m and nucleotide sequence and was found to be specific for the rRNA of Actinomycetes, CGAGCATCTTTACTCGTAGTGCAATTTCG (SEQ ID NO:1). This
25 polynucleotide, referred to herein as MtuB2011, is complementary to a unique segment found in the 23S rRNA of all Actinomycetes. A representative list of Actinomycetes bacteria can be found in Table 2. The probe is 29 bases in length, has an RXL linker between 11 and 12 nucleotides from the 5' end and has a T_m of 65.5°C , and hybridized rRNA of *Mycobacterium avium* in a region corresponding to bases 2011- 2040 of *E.*
30 *coli* 23S rRNA.

This probe is one illustration of an oligonucleotide that: (1) hybridizes the target nucleic acid under high stringency hybridization conditions, (2) has a length of up to 100 nucleotide bases, and (3) includes at least 17 contiguous nucleotides falling within the 1986 - 2064 target region identified by SEQ ID NO:10 or its complement. Other
5 oligonucleotides having these properties are contemplated for use as hybridization assay detection probes and are embraced by the invention.

Similarly, oligonucleotides having the sequences of SEQ ID NOs:2 and 3 are disclosed herein as illustrations of useful helper oligonucleotides. Like the helper oligonucleotides employed in the working Examples herein, other helper
10 oligonucleotides embraced by the invention also have sequences of up to 100 nucleotides in length and further have at least 17, or more preferably at least 25, contiguous nucleotides contained within the target region identified by SEQ ID NO:10 or its complement.

As indicated below, the MtuB2011 probe hybridized Actinomycetes rRNA in a
15 manner that was promoted by the use of helper oligonucleotides. According to the procedure used to make this determination, single-stranded probe oligonucleotide radiolabeled at the 5'-end was contacted with rRNA from *Mycobacterium avium* in the presence or absence of helper oligonucleotides. Probe molecules hybridizing the rRNA to form double-stranded hybrids were separated from single-stranded probe molecules
20 by hydroxyapatite capture. The double-stranded hybrids bound to the hydroxyapatite and were detected and quantitated by scintillation counting. The extent of hybridization was then calculated as a percentage. As indicated below, the T_m of the probe:target hybrid advantageously was significantly increased in the presence of one or more helper oligonucleotides.

25 The following Example describes the methods used to demonstrate that the MtuB2011 probe hybridized rRNA from *Mycobacterium avium* and that this interaction was facilitated by including helper oligonucleotides in the hybridization mixture.

Example 1

T_m Determination for Probe:Target Hybrids

30 T_m values for probe:target and helper:target hybrids were determined using an end-labeled probe having the sequence of MtuB2011 and end-labeled helper

oligonucleotides selected from the group: (A) OMeFraB1986 and (B) OMeFraB2040. The sequence of MtuB2011 is CGAGCATCTTTACTCGTAGTGCAATTTTCG (SEQ ID NO:1), the sequence of OMeFraB1986 is CCGAGUCUGUGGUUGAGACAGUGGG (SEQ ID NO:2) and the sequence of OMeFraB2040 is GGUCUUUCCGUCCUGCCGCGCGUAA (SEQ ID NO:3). Helper oligonucleotides A and B were selected to bind the rRNA of Actinomycetes in regions of the molecule immediately adjacent to the probe, helper A binding in about the 1986-2010 region of the 23S rRNA, and helper B binding in the 2040-2064 region of the 23S rRNA. The probe and helper oligonucleotides were 5'-end labeled using [γ - ^{32} P]ATP as a phosphate donor and T4 polynucleotide kinase to catalyze the phosphate transfer reaction essentially as described in Molecular Cloning: A Laboratory Manual (Sambrook et al., eds. Cold Spring Harbor Lab Publ. 10.59 (1989)). End-labeled helper oligonucleotides were separately combined with purified rRNA from *Mycobacterium gordonae* and the probe oligonucleotide was combined with purified rRNA from *Mycobacterium avium* to provide conditions of target excess. In trials that included both the probe and helper oligonucleotides, only the probe was end-labeled and each helper oligonucleotide was present in at least a 10 fold molar excess over the *Mycobacterium avium* rRNA that served as a target. All mixtures were hybridized to completion in a solution that included 0.48 M sodium phosphate buffer, 0.1% sodium dodecyl sulfate, 1 mM EDTA and 1 mM EGTA. As negative controls, the probe and/or helper oligonucleotides were hybridized in the absence of the nucleic acid target. At the conclusion of the hybridization procedure, mixtures were diluted and passed over a hydroxyapatite column to separate single-stranded nucleic acids from double-stranded hybrids. The amount of radioactivity in the column flow-through represented single-stranded probe and was measured by scintillation counting. The amount of radioactivity bound to the hydroxyapatite was separately measured by scintillation counting. The extent of hybrid formation, expressed as a percentage, was calculated by dividing the amount of probe (measured in cpm) bound to the hydroxyapatite by the total amount of probe (in cpm) that was applied to the column. Results of these procedures are presented in Table 1.

TABLE 1

Hybridization of the MtuB2011 Probe with Target rRNA

Oligonucleotide(s)	% Hybridization	T _m (°C)
MtuB2011 (Probe)	84	65.5
helper A, OMeFraB1986	96	78
helper B, OMeFraB2040	96	≈80
Probe + helper A	98	68
Probe + helper B	97	67
Probe + helper A + helper B	96	69.5

The results from this procedure confirmed that the end-labeled probe hybridized rRNA from *Mycobacterium avium* and/or *Mycobacterium gordonae* and indicated this interaction advantageously was facilitated by helper oligonucleotides. We particularly observed that the T_m of the probe:target complex could be increased from 65.5 to 69.5°C when both helper oligonucleotides were included in the hybridization reaction. Although the probe can be used either alone or in combination with one or more helper oligonucleotides for hybridizing Actinomycetes rRNA, the below-described experiments to characterize the probe were conducted using the probe in combination with helper oligonucleotides having the sequences of OMeFraB1986 and OMeFraB2040. Combinations of probe and helper oligonucleotides useful in the procedures described herein preferably have probe:target T_m values in the range of from about 65-70°C under the conditions described above.

Probe specificity was confirmed by demonstrating positive hybridization to rRNAs from a specificity panel. The collection of organisms used as sources of target nucleic acids in this procedure represented a broad taxonomic cross-section of organisms and a nearest-neighbor group. In the following procedure, quantitative results using the AE-labeled hybridization probe were compared to the amount of bacterial rRNA present in each sample using a positive control probe. This positive control probe, which hybridized rRNA from all species of bacteria, was particularly useful for confirming the presence of bacterial rRNA in samples that failed to hybridize

the MtuB2011 probe. In such an event, the positive control probe provided confirmation for the presence of hybridizable rRNA and so validated the negative results. In the case of fungal rRNA targets, a broadly reactive fungal rRNA hybridization probe served as the positive control.

- 5 The following Example describes the methods used to demonstrate that the MtuB2011 probe hybridized rRNAs from a panel of Actinomycetes organisms.

Example 2

Verification of Probe Specificity

- Bacterial lysates or purified RNA were used as nucleic acid targets for
- 10 hybridization of a probe having the sequence of
CGAGCATCTTTACTCGTAGTGCAATTTCG (MtuB2011) (SEQ ID NO:1) together
with helper oligonucleotides having the sequences of
CCGAGUCUGUGGUUGAGACAGUGGG (OMeFraB1986) (SEQ ID NO:2) and
GGUCUUUCCGUCCUGCCGCGCGUAA (OMeFraB2040) (SEQ ID NO:3).
- 15 Organisms employed as sources of rRNA in this procedure were either typed clinical
isolates or obtained from the American Type Culture Collection (ATCC). All samples
are identified in Table 2 by master log numbers for Gen-Probe Incorporated. Most
samples were obtained from the ATCC. Parallel samples of each rRNA were
hybridized with a labeled positive control probe having the sequence
- 20 CGACAAGGAAUUUCGC (OMe Eco B 1933) (SEQ ID NO:4) and unlabeled helper
oligonucleotides having the sequences UACCUUAGGACCGUUAU (OMe Eco B
1916) (SEQ ID NO:5) and CAGGUCGGAACUUACC (OMe Eco B 1949a) (SEQ ID
NO:6). The hybridization solution contained 0.6M LiCl, 1% lithium lauryl sulfate, 60
mM lithium succinate and 10mM each of EDTA and EGTA, pH 5.5. Both the
- 25 MtuB2011 probe and the positive control probe were labeled with acridinium ester
essentially according to the method disclosed in U.S. Patent No. 5,185,439, entitled
"Acridinium Ester Labeling and Purification of Nucleotide Probes." At the conclusion
of the hybridization reaction, acridinium ester linked to unhybridized probe was
rendered non-chemiluminescent under mild alkaline conditions, while acridinium ester
- 30 attached to hybridized probe remained resistant to the inactivation. Conditions for the
hydrolysis and detection of hybridized probe labeled with acridinium ester are

described by Arnold et al., in *Clin. Chem.* **35**:1588 (1989)). The magnitudes of probe hybridization in these procedures were quantitated by luminometry using procedures familiar to those having ordinary skill in the art. The magnitude of the Actinomycetes probe signal was then divided by the magnitude of the bacterial positive control signal to quantitatively normalize results in the study. Samples having MtuB2011 probe signals that were greater than 30% of the positive control signal indicated specific hybridization with the MtuB2011 probe with helpers, while lower values indicated negative results for the assay format. Assay results are shown in Table 2.

TABLE 2

Hybridization of the MtuB2011 Probe and rRNA-Containing Lysates from a Collection of Actinomycetes

*GP #	rRNA Source (Organism)	Hybridization Results		
		Pan-Bacterial Probe (RLU)	Actinomycetes Probe (RLU)	Fractional Hybridization (%)
236	<i>Corynebacterium aquaticum</i>	19450	18950	97
239	<i>Corynebacterium jeikeium</i>	9258	4404	48
237	<i>Corynebacterium xerosis</i>	9207	9624	105
240	<i>Micrococcus luteus</i>	2143	2301	107
190	<i>Propionibacterium acnes</i>	5680	4804	85
159	<i>Mycobacterium chelonae</i>	256985	106195	41
543	<i>Mycobacterium terrae</i>	43323	58933	136
87	<i>Mycobacterium intracellulare</i>	80497	123777	154
84	<i>Mycobacterium simiae</i>	29865	39705	133
PN 101676	<i>Mycobacterium avium</i>	1690	1551	92
79	<i>Mycobacterium scrofulaceum</i>	7601	10373	136
85	<i>Mycobacterium gordonae</i>	22560	22180	98
82	<i>Mycobacterium kansasii</i>	35457	44997	127
86	<i>Mycobacterium smegmatis</i>	108778	137648	127
160	<i>Mycobacterium fortuitum</i>	71278	127108	178
15	<i>Mycobacterium gastri</i>	5406	7274	135
74	<i>Mycobacterium xenopi</i>	36080	27110	75
83	<i>Mycobacterium marinum</i>	268900	330700	123
78	<i>Mycobacterium phlei</i>	12999	19780	152

* "GP#" entries indicate master log numbers for Gen-Probe Incorporated.

The results presented in Table 2 confirmed that the probe directed against Actinomycetes rRNA efficiently hybridized rRNA samples from numerous Actinomycetes species.

Specificity of the Actinomycetes-specific probe was further investigated by hybridizing labeled probe with a collection of species representing a broad spectrum of phylogenetically diverse organisms. In this procedure, AE-labeled probe was separately mixed with individual rRNA containing lysates from organisms that were only distant
5 phylogenetic relatives of the Actinomycetes. Positive hybridization results obtained using the positive control probe and negative results obtained using the MtuB2011 probe in the following procedure further indicated that the MtuB2011 probe advantageously was highly specific for Actinomycetes organisms.

The following Example describes additional methods used to demonstrate
10 specificity of the probe. More particularly, the following procedures showed that the MtuB2011 probe did not cross hybridize with lysates from non-phylogenetically related organisms.

Example 3

Absence of Cross Hybridization with Phylogenetically Unrelated Organisms

15 Hybridization assays were conducted using the AE-labeled probes and helper oligonucleotides according to the procedures described in the previous Example except that lysates containing rRNA isolated from numerous diverse species served as target nucleic acids. Results of the procedure are presented in Table 3. A pan-fungal probe having the sequence GTCTGGACCTGGTGAGTTTCCC (SEQ ID NO:7), and
20 methoxy helper oligonucleotides having the sequences CGUGUUGAGUCAAAUUAAGCCGC (SEQ ID NO:8) and GCUCUCAAUCUGUCAAUCCUUAUUGU (SEQ ID NO:9) were used as positive controls to detect fungal rRNAs. Results obtained in hybridization procedures that employed rRNA targets from fungal organisms are presented in Table 4.

TABLE 3
Hybridization of the MtuB2011 Probe with rRNA from a Collection of
Phylogenetically Non-Related Organisms

5			Hybridization Results		
	*GP#	rRNA Source (Organism)	Pan-Bacterial Probe (RLU)	Actinomycetes Probe (RLU)	Fractional Hybridization (%)
	#234	<i>Acinetobacter calcoaceticus</i>	4215	629	15
	#233	<i>Acinetobacter lwoffii</i>	4345	551	13
	#13	<i>Bacillus brevis</i>	9293	818	9
10	#11	<i>Bacillus subtilis</i>	5184	860	17
	#212	<i>Bacteriodes fragilis</i>	7837	771	10
	#226	<i>Bacteroides ovatus</i>	4107	563	14
	#225	<i>Bacteroides thetaiotamicron</i>	33410	553	2
	#152	<i>Citrobacter diversus</i>	6835	744	11
15	#150	<i>Citrobacter freundii</i>	9188	704	8
	#192	<i>Clostridium perfringens</i>	7816	956	12
	#156	<i>Enterobact gergoviae</i>	3175	598	19
	#153	<i>Enterobacter aerogenes</i>	8306	733	9
	#154	<i>Enterobacter agglomerans</i>	8753	752	9
20	#155	<i>Enterobacter cloacae</i>	8150	732	9
	#215	<i>Enterobacter fragilis</i>	7139	724	10
	#46	<i>Enterococcus avium</i>	9813	858	9
	#27	<i>Enterococcus casseliflavus</i>	10339	667	6
	#7	<i>Enterococcus cecorum</i>	5911	901	15
25	#15	<i>Enterococcus dispar</i>	9170	772	8
	#85	<i>Enterococcus durans</i>	8839	877	10
	#82	<i>Enterococcus faecalis</i>	7910	879	11
	#79	<i>Enterococcus faecium</i>	7696	926	12
	#23	<i>Enterococcus faecium V1</i>	5607	713	13
30	#17	<i>Enterococcus faecium V6</i>	5761	755	13
	#89	<i>Enterococcus gallinarum</i>	8682	777	9
	#81	<i>Enterococcus hirae</i>	6138	793	13
	#45	<i>Enterococcus malodoratus</i>	15407	737	5
	#25	<i>Enterococcus mundtii</i>	9482	817	9
35	#26	<i>Enterococcus pseudoavium</i>	10042	703	7
	#33	<i>Enterococcus raffinosus</i>	6662	675	10
	#47	<i>Enterococcus sacchrolyticus</i>	9017	844	9
	#159	<i>Escherichia coli</i>	5587	598	11
	#161	<i>Escherichiafergusonii</i>	4598	585	13
40	#162	<i>Escherichia hermanii</i>	3267	601	18
	#217	<i>Haemophilus influenzae</i>	4157	635	15
	#219	<i>Haemophilus influenzae A</i>	4182	596	14
	#220	<i>Haemophilus influenzae B</i>	4247	570	13
	#222	<i>Haemophilus parainfluenzae</i>	25100	557	2

		Hybridization Results			
	*GP#	rRNA Source (Organism)	Pan-Bacterial Probe (RLU)	Actinomycetes Probe (RLU)	Fractional Hybridization (%)
5	#188	<i>Hafnia alvei</i>	47770	941	2
	#163	<i>Klebsiella oxytoca</i>	3088	560	18
	#164	<i>Klebsiella ozaenae</i>	5252	555	11
	#176	<i>Klebsiella pneumoniae</i>	4631	441	10
	#178	<i>Klebsiella rhinoscleromatis</i>	3567	474	13
10	#36	<i>Lactobacillus acidophilus</i>	20730	704	3
	#56	<i>Lactobacillus jensenii</i>	6541	783	12
	#9	<i>Lactococcus lactis</i>	36850	1022	3
	#41	<i>Listeria grayi</i>	2956	834	28
	#72	<i>Listeria ivanovii</i>	6355	908	14
15	#31	<i>Listeria monocytogenes 1/2b</i>	10102	677	7
	#28	<i>Listeria monocytogenes 4b</i>	8225	685	8
	#73	<i>Listeria seeligeri</i>	7001	849	12
	#40	<i>Listeria welshimeri</i>	5288	862	16
	#184	<i>Morganella morganii</i>	3852	440	11
20	#196	<i>Neisseria gonorrhoea</i>	6824	872	13
	#198	<i>Neisseria meningitidis</i>	20430	766	4
	#191	<i>Peptostreptococcus anaerobius</i>	6985	824	12
	#179	<i>Proteus mirabilis</i>	5115	424	8
	#183	<i>Proteus penneri</i>	4285	927	22
25	#181	<i>Proteus vulgaris</i>	4883	433	9
	#186	<i>Providencia alcalifaciens</i>	44220	947	2
	#187	<i>Providencia rettgeri</i>	31350	923	3
	#185	<i>Providencia stuartii</i>	13926	437	3
	#200	<i>Pseudomonas aeruginosa</i>	5771	832	14
30	#203	<i>Pseudomonas cepacia</i>	25590	982	4
	#205	<i>Pseudomonas fluorescens</i>	8676	773	9
	#206	<i>Pseudomonas maltophilia</i>	7703	759	10
	#209	<i>Pseudomonas mendocina</i>	17980	816	5
	#208	<i>Pseudomonas picketti</i>	7954	991	12
35	#210	<i>Pseudomonas putida A</i>	8216	788	10
	#211	<i>Pseudomonas stutzeri</i>	7772	786	10
	#189	<i>Salmonella enteritidis</i>	20900	900	4
	#216	<i>Salmonella paratyphi</i>	4253	560	13
	#165	<i>Salmonella typhi</i>	7967	1019	13
40	#166	<i>Salmonella typhimurium</i>	5565	507	9
	#170	<i>Serratia liquifaciens</i>	5801	551	9
	#171	<i>Serratia marcescens</i>	5093	471	9
	#168	<i>Shigella dysenterae</i>	5233	764	15
	#169	<i>Shigella sonnei</i>	5121	581	11
	#49	<i>Staphylococcus aureus</i>	5964	838	14

		Hybridization Results			
	*GP#	rRNA Source (Organism)	Pan-Bacterial Probe (RLU)	Actinomycetes Probe (RLU)	Fractional Hybridization (%)
5	#63	<i>Staphylococcus cohnii</i>	6253	1066	17
	#6	<i>Staphylococcus delphini</i>	6595	857	13
	#50	<i>Staphylococcus epidermidis</i>	8432	806	10
	#62	<i>Staphylococcus haemolyticus</i>	5354	985	18
	#61	<i>Staphylococcus hominis</i>	5593	1011	18
10	#69	<i>Staphylococcus hyicus</i>	6193	877	14
	#60	<i>Staphylococcus intermedius</i>	7501	791	11
	#59	<i>Staphylococcus saprophyticus</i>	4162	774	19
	#39	<i>Staphylococcus stimulants</i>	6568	926	14
	#67	<i>Staphylococcus warner</i>	4274	967	23
15	#53	<i>Streptococcus agalactiae</i>	7295	845	12
	#32	<i>Streptococcus agalactiae Ia</i>	10144	691	7
	#43	<i>Streptococcus anginosus</i>	8861	891	10
	#16	<i>Streptococcus avium</i>	6366	687	11
	#34	<i>Streptococcus bovis</i>	9298	664	7
20	#54	<i>Streptococcus dysgalactiae</i>	7095	744	10
	#51	<i>Streptococcus equi</i>	12933	904	7
	#80	<i>Streptococcus equinus</i>	5259	888	17
	#37	<i>Streptococcus equisimilis</i>	8123	774	10
	#97	<i>Streptococcus grp C</i>	9261	796	9
25	#98	<i>Streptococcus grp G</i>	7478	770	10
	#44	<i>Streptococcus mutans</i>	6874	939	14
	#42	<i>Streptococcus pneumoniae</i>	6153	869	14
	#92	<i>Streptococcus pyogenes</i>	6723	886	13
	#38	<i>Streptococcus salivarius</i>	8044	977	12
30	#35	<i>Streptococcus sanguis</i>	11918	705	6
	#66	<i>Streptococcus sp gp F2</i>	6921	894	13
	#3	<i>Streptococcus</i> sp. <i>Gp. B, II</i>	154	915	15
	#5	<i>Streptococcus uberis</i>	6067	879	14
	#173	<i>Yersinia enterocolitica</i>	6165	521	8
	#175	<i>Yersinia intermedia</i>	5246	460	9
	#174	<i>Yersinia pseudotuberculosis</i>	5564	418	8

* "GP#" entries indicate master log numbers for Gen-Probe Incorporated.

TABLE 4
Hybridization of the MtuB2011 Probe with rRNA from a
Collection of Fungal Organisms

	GP#*	rRNA Source (Organism)	Pan-Bacterial Probe (RLU)	MtuB2011 Probe (RLU)	Fungal Probe (RLU)	Hybrid (%)
5	F-932	<i>Arachnoitus flavoluteus</i>	481	417	232076	0.2
	F-906	<i>Arachnoitus flavus</i>	364	395	348387	0.1
	F-899	<i>Aspergillus fumigatus</i>	382	456	419831	0.1
	F-907	<i>Aspergillus niger</i>	194	411	650747	0.1
	F-930	<i>Auxarthron thaxteri</i>	301	399	494055	0.1
10	F-1022	<i>Blastomyces dermatitidis</i>	296	405	422465	0.1
	715	<i>Candida albicans</i>	369	333	327951	0.1
	1123	<i>Candida glabrata</i>	1419	1398	45039	3.1
	717	<i>Candida parapsillosis</i>	352	374	312482	0.1
	1091	<i>Candida tropicalis</i>	1566	1437	24023	6
15	F-1399	<i>Coccidioides immitis</i>	303	329	141956	0.2
	F-900	<i>Cryptococcus neoformans</i>	316	302	452943	0.1
	F-965	<i>Gymnoascus dugwayensis</i>	317	317	506033	0.1
	F-968	<i>Histoplasma capsulatum</i>	254	255	346283	0.1
	F-933	<i>Myxotrichum deflexum</i>	267	319	366688	0.1
20	F-934	<i>Oidiodendron echinulatum</i>	238	262	322685	0.1
	716	<i>Candida krusei</i>	669	432	71371	0.6
	1087	<i>Candida pseudotropicalis</i>	243	35	79868	0
	384	<i>Saccharomyces cerevisiae</i>	116	76	75954	0.1
	1080	<i>Candida guilliermondii</i>	320	203	65011	0.3

25 * "GP#" identifies organisms by master log numbers for Gen-Probe Incorporated.

The results presented in Table 3 and Table 4 confirmed that the MtuB2011 probe did not cross hybridize with rRNA from numerous unrelated bacterial and fungal species. Taken together with the positive hybridization results presented in the Table 2, it was
 30 clear that the probe was highly specific for rRNA of the Actinomycetes.

These results confirmed that the novel probes disclosed herein were capable of detecting the Actinomycetes. Moreover, the probes were capable of distinguishing Actinomycetes from organisms that were phylogenetically closely related.

This invention has been described with reference to a number of specific
 35 examples and embodiments thereof. Of course, a number of different embodiments of the present invention will suggest themselves to those having ordinary skill in the art upon review of the foregoing detailed description. Thus, the true scope of the present invention is to be determined upon reference to the appended claims.

32a

SEQUENCE LISTING

<110> Gen-Probe Incorporated

<120> Polynucleotide Probes for Detection and
Quantitation of Actinomycetes

<130> 82022-12

<140> PCT/US00/12243

<141> 2000-05-03

<150> 60/132,412

<151> 1999-05-03

<160> 19

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Probe for hybridizing ribosomal nucleic acids of
Actinomycetes bacteria

<400> 1

cgagcatctt tactcgtagt gcaatttcg

29

<210> 2

<211> 25

<212> RNA

<213> Artificial Sequence

<220>

<223> Helper oligonucleotide for hybridizing ribosomal
nucleic acids of Actinomycetes bacteria

<400> 2

ccgagucugu gguugagaca guggg

25

<210> 3

<211> 25

<212> RNA

<213> Artificial Sequence

<220>

<223> Helper oligonucleotide for hybridizing ribosomal
nucleic acids of Actinomycetes bacteria

<400> 3

ggucuuuccg uccugccgcg cguaa

25

<210> 4

<211> 16

32b

<212> RNA
 <213> Artificial Sequence

 <220>
 <223> Probe for hybridizing ribosomal nucleic acids of
 pan-bacterial organisms

 <400> 4
 cgacaaggaa uuucgc 16

 <210> 5
 <211> 17
 <212> RNA
 <213> Artificial Sequence

 <220>
 <223> Helper oligonucleotide for hybridizing ribosomal
 nucleic acids of pan-bacterial organisms

 <400> 5
 uaccuuagga ccguuau 17

 <210> 6
 <211> 16
 <212> RNA
 <213> Artificial Sequence

 <220>
 <223> Helper oligonucleotide for hybridizing ribosomal
 nucleic acids of pan-bacterial organisms

 <400> 6
 caggucggaa cuuacc 16

 <210> 7
 <211> 22
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Probe for hybridizing ribosomal nucleic acids of
 pan-fungal organisms

 <400> 7
 gtctggacct ggtgagtttc cc 22

 <210> 8
 <211> 23
 <212> RNA
 <213> Artificial Sequence

 <220>
 <223> Helper oligonucleotide for hybridizing ribosomal
 nucleic acids of pan-fungal organisms

 <400> 8
 cguguugagu caaauuaagc cgc 23

32c

<210> 9
 <211> 26
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Helper oligonucleotide for hybridizing ribosomal
 nucleic acids of pan-fungal organisms

<400> 9
 gcucucaauc ugucaauccu uauugu 26

<210> 10
 <211> 79
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Contiguous probe sequence for hybridizing
 ribosomal nucleic acids of Actinomycetes bacteria

<400> 10
 ggtctttccg tcctgccgcg cgtaacgagc atctttactc gtagtgcaat ttcgccgagt 60
 ctgtggttga gacagtggg 79

<210> 11
 <211> 79
 <212> RNA
 <213> Artificial Sequence

<220>
 <223> Contiguous ribosomal nucleic acid target sequence
 for Actinomycetes bacteria

<400> 11
 cccacugucu caaccacaga cucggcgaaa uugcacuacg aguaaagaug cucguuacgc 60
 gcggcaggac ggaaagacc 79

<210> 12
 <211> 86
 <212> RNA
 <213> Staphylococcus aureus

<220>
 <221> misc_feature
 <222> (1)...(86)
 <223> Segment of rRNA

<400> 12
 uugggcacug ucucaacgag agacucggug aaaucauagu accugugaag augcagguua 60
 cccgcgacag gacggaaaga ccccg 86

<210> 13
 <211> 86
 <212> RNA
 <213> Escherichia coli

<220>

32d

<221> misc_feature
 <222> (1)...(86)
 <223> Segment of rRNA

<400> 13

ggccaggcug ucuccacccg agacucagug aaauugaacu cgcugugaag augcagugua 60
 cccgcggcaa gacggaaaga ccccg 86

<210> 14
 <211> 85
 <212> RNA
 <213> Frankia sp

<220>
 <221> misc_feature
 <222> (1)...(85)
 <223> Segment of rRNA

<300>
 <301> Normand et al.
 <302> Analysis of a ribosomal RNA operon...
 <303> Gene
 <304> 111
 <306> 119-124

<400> 14

uuucccacug ucucaaccac agacucggcg aaauugcauu acgaguaaag augcucguua 60
 cgcgcggcag gacggaaaga ccccg 85

<210> 15
 <211> 83
 <212> RNA
 <213> Mycobacterium tuberculosis

<220>
 <221> misc_feature
 <222> (1)...(83)
 <223> n=a, g, c, u

<221> misc_feature
 <222> (1)...(83)
 <223> Segment of rRNA

<400> 15

uucucaacug ucucaaccac agacucggcg aaauugcacu acgaguaaag augcucgnac 60
 gcgcggcagg ncgaaaanac ccc 83

<210> 16
 <211> 61
 <212> RNA
 <213> Corynebacterium xerosis

<220>
 <221> misc_feature
 <222> (1)...(61)
 <223> n=a, g, c, u

<221> misc_feature

32e

<222> (1)...(61)
 <223> Segment of rRNA

<400> 16

cccggcgaaa uugcgauacg aguaaagaug cunnuacgcg cggcaggacg aaaagacccc 60
 g 61

<210> 17
 <211> 86
 <212> RNA
 <213> Streptomyces griseus

<220>
 <221> misc_feature
 <222> (1)...(86)
 <223> Segment of rRNA

<400> 17

uucucgacug ucucaaccu aggcccgug aaauugcacu acgaguaaag augcucguuu 60
 cgcgcgagcag gacggaaaga ccccg 86

<210> 18
 <211> 85
 <212> RNA
 <213> Mycobacterium avium

<220>
 <221> misc_feature
 <222> (1)...(85)
 <223> n=a, g, c, u

<221> misc_feature
 <222> (1)...(85)
 <223> Segment of rRNA

<400> 18

uucccaacug ucucaaccu agacucggcg aaauugcacu acgaguaaag augcucgnua 60
 cgcgcgagcag gncgaaaaga ccccg 85

<210> 19
 <211> 85
 <212> RNA
 <213> Nocardia asteroides

<220>
 <221> misc_feature
 <222> (1)...(85)
 <223> Segment of rRNA

<400> 19

uucucugcug ucucaaccu agacucggcg aaauugcauu acgaguaaag augcucguua 60
 cgcgcgagcag gacgaaaaga ccccg 85

WHAT IS CLAIMED IS:

1. An oligonucleotide probe that specifically hybridizes to an Actinomycetes nucleic acid target region corresponding to *E. coli* 23S rRNA nucleotide positions 1986 - 2064 under a high stringency hybridization condition provided by a solution comprising 0.48 M sodium phosphate or 0.6 M LiCl to form a detectable probe:target duplex, said oligonucleotide probe having a length of up to 100 nucleotides and comprising at least 17 contiguous nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof.
2. The oligonucleotide probe of Claim 1, wherein said probe comprises at least 25 contiguous nucleotides contained within the sequence of SEQ ID NO:10.
3. The oligonucleotide probe of Claim 2, wherein said probe comprises at least 29 contiguous nucleotides contained within the sequence of SEQ ID NO:10.
4. The oligonucleotide probe of Claim 1, wherein said oligonucleotide probe has a length of up to 100 nucleotide bases and comprises at least 25 contiguous nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof.
5. The oligonucleotide probe of Claim 1, 2 or 3, wherein the high stringency hybridization condition is provided by 0.48 M sodium phosphate buffer, 0.1% sodium dodecyl sulfate, 1 mM each of EDTA and EGTA.
6. The oligonucleotide probe of Claim 1, 2 or 3, wherein the high stringency hybridization condition is provided by 0.6 M LiCl, 1% lithium lauryl sulfate, 60 mM lithium succinate and 10 mM each of EDTA and EGTA.
7. The oligonucleotide probe of any one of Claims 1 to 5, wherein said oligonucleotide probe comprises DNA.
8. The oligonucleotide probe of any one of Claims 1 to 6, wherein said oligonucleotide probe comprises at least one nucleotide analog.
9. The oligonucleotide probe of Claim 7, wherein said at least one nucleotide analog comprises a methoxy group at the 2' position of a ribose moiety.
10. The oligonucleotide probe of any one of Claims 1 to 9, further comprising a detectable label.

11. The oligonucleotide probe of Claim 1, wherein said oligonucleotide probe has a sequence selected from the group consisting of SEQ ID NO:1 or the complement thereof, SEQ ID NO:2 or the complement thereof, and SEQ ID NO:3 or the complement thereof.

12. The oligonucleotide of Claim 11, wherein said sequence is given by SEQ ID NO:2 or SEQ ID NO:3, said oligonucleotide being a helper oligonucleotide.

13. The oligonucleotide probe of Claim 11 or 12, further comprising a detectable label.

14. The oligonucleotide probe of Claim 13, wherein the detectable label is a chemiluminescent label or a radiolabel.

15. The oligonucleotide probe of Claim 11, wherein said sequence is given by SEQ ID NO:1.

16. The oligonucleotide probe of Claim 15, wherein said oligonucleotide probe further comprises a detectable label.

17. The oligonucleotide probe of Claim 16, wherein the detectable label is an acridinium ester.

18. An oligonucleotide probe for detecting nucleic acids of Actinomycetes bacteria, wherein the oligonucleotide probe hybridizes under a high stringency condition provided by a solution comprising 0.48 M sodium phosphate or 0.6 M LiCl to an Actinomycetes nucleic acid target region corresponding to *E. coli* 23S rRNA nucleotide positions 1986 - 2064 to form a detectable target:probe duplex,

wherein said oligonucleotide probe has a length of up to 100 nucleotide bases and comprises at least 17 contiguous nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof, and

wherein under said hybridization condition said oligonucleotide probe specifically hybridizes to nucleic acids present in *Corynebacterium aquaticum*, *Corynebacterium jeikeium*, *Corynebacterium xerosis*, *Micrococcus luteus*, *Propionibacterium acnes*, *Mycobacterium chelonae*, *Mycobacterium terrae*, *Mycobacterium intracellulare*, *Mycobacterium simiae*, *Mycobacterium avium*, *Mycobacterium scrofulaceum*, *Mycobacterium gordonae*, *Mycobacterium kansasii*, *Mycobacterium smegatis*,

Mycobacterium fortuitum, *Mycobacterium gastri*, *Mycobacterium xenopi*, *Mycobacterium marinum* and *Mycobacterium phlei*.

19. The oligonucleotide probe of Claim 18, wherein the oligonucleotide probe comprises DNA.

20. The oligonucleotide probe of Claim 18 or 19, wherein said high stringency condition is provided by 0.48 M sodium phosphate buffer, 0.1% sodium dodecyl sulfate, 1 mM each of EDTA and EGTA.

21. The oligonucleotide probe of Claim 18 or 19, wherein said high stringency condition is provided by 0.6 M LiCl, 1% lithium lauryl sulfate, 60 mM lithium succinate and 10 mM each of EDTA and EGTA.

22. The oligonucleotide probe of any one of Claims 18 to 21, wherein said oligonucleotide probe comprises the sequence of SEQ ID NO:1 or the complement thereof.

23. The oligonucleotide probe of any one of Claims 18 to 22, wherein the length of said oligonucleotide probe is up to 60 bases.

24. The oligonucleotide probe of any one of Claims 18 to 21, wherein said oligonucleotide probe has the length and sequence of SEQ ID NO:1.

25. The oligonucleotide probe of any one of Claims 18 to 24, wherein said oligonucleotide probe further comprises a detectable label.

26. The oligonucleotide probe of Claim 25, wherein the detectable label is a chemiluminescent label or a radiolabel.

27. The oligonucleotide probe of Claim 26, wherein the chemiluminescent label is an acridinium ester.

28. The oligonucleotide probe of Claim 25, 26 or 27, further comprising at least one helper oligonucleotide that facilitates formation of the detectable probe:target duplex under said hybridization conditions.

29. The oligonucleotide probe of Claim 28, wherein said at least one helper oligonucleotide includes at least one nucleotide analog.

30. The oligonucleotide probe of Claim 29, wherein said at least one nucleotide analog comprises a ribose moiety having a methoxy group disposed at the 2' position.

31. The oligonucleotide probe of Claim 28, wherein said at least one helper oligonucleotide has a sequence selected from the group consisting of SEQ ID NO:2 and SEQ ID NO:3.

32. A method for detecting the presence of Actinomycetes in a test sample, comprising the steps of:

(a) providing to said test sample an oligonucleotide probe that hybridizes under a high stringency condition provided by a solution comprising 0.48 M sodium phosphate or 0.6 M LiCl to an Actinomycetes nucleic acid target region corresponding to *E. coli* 23S rRNA nucleotide positions 1986 - 2064 to form a detectable target:probe duplex, said oligonucleotide probe having a length of up to 100 nucleotide bases and comprising at least 25 contiguous nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof, and wherein under said hybridization condition said oligonucleotide probe specifically hybridizes to nucleic acids present in *Corynebacterium aquaticum*, *Corynebacterium jeikeium*, *Corynebacterium xerosis*, *Micrococcus luteus*, *Propionibacterium acnes*, *Mycobacterium chelonae*, *Mycobacterium terrae*, *Mycobacterium intracellulare*, *Mycobacterium simiae*, *Mycobacterium avium*, *Mycobacterium scrofulaceum*, *Mycobacterium gordonae*, *Mycobacterium kansasii*, *Mycobacterium smegatis*, *Mycobacterium fortuitum*, *Mycobacterium gastri*, *Mycobacterium xenopi*, *Mycobacterium marinum* and *Mycobacterium phlei*;

(b) hybridizing under high stringency conditions any Actinomycetes nucleic acid that may be present in the test sample with said probe to form a probe:target duplex; and

(c) detecting said probe:target duplex as an indicator of the presence of Actinomycetes in the test sample.

33. The method of Claim 32, wherein said test sample may comprise bacteria, and wherein before step (a) there is a step for releasing nucleic acid from any bacteria that may be present in said test sample.

34. The method of Claim 32, wherein said test sample is a lysate.

35. The method of Claim 32, 33 or 34, wherein said high stringency hybridization conditions are provided by 0.48 M sodium phosphate buffer, 0.1% sodium dodecyl sulfate, 1 mM each of EDTA and EGTA.

36. The method of Claim 32, 33 or 34, wherein said high stringency hybridization conditions are provided by 0.6 M LiCl, 1% lithium lauryl sulfate, 60 mM lithium succinate and 10 mM each of EDTA and EGTA.

37. The method of any one of Claims 32 to 36, wherein the oligonucleotide probe has the length and sequence of SEQ ID NO:1.

38. The method of any one of Claims 32 to 36, wherein said probe further comprises at least one helper oligonucleotide that facilitates formation of the probe:target duplex.

39. The method of Claim 38, wherein said at least one helper oligonucleotide is selected from the group consisting of SEQ ID NO:2 and SEQ ID NO:3.

40. The method of any one of Claims 32 to 39, wherein the oligonucleotide probe comprises a detectable label.

41. The method of Claim 40, wherein the detectable label is an acridinium ester, and wherein the detecting step comprises performing luminometry to detect any of said probe:target duplex.

42. A kit for detecting the presence of Actinomycetes nucleic acids in a test sample, comprising:

an oligonucleotide probe that hybridizes under a high stringency condition provided by a solution comprising 0.48 M sodium phosphate or 0.6 M LiCl to an Actinomycetes nucleic acid target region corresponding to *E. coli* 23S rRNA nucleotide positions 1986 - 2064 to form a detectable target:probe duplex, said oligonucleotide probe having a length of up to 100 nucleotide bases and comprising at least 25 contiguous nucleotides contained within the sequence of SEQ ID NO:10 or the complement thereof, and wherein under said hybridization condition said oligonucleotide probe specifically hybridizes to nucleic acids present in *Corynebacterium aquaticum*, *Corynebacterium jeikeium*, *Corynebacterium xerosis*, *Micrococcus luteus*, *Propionibacterium acnes*, *Mycobacterium chelonae*, *Mycobacterium terrae*, *Mycobacterium intracellulare*, *Mycobacterium simiae*, *Mycobacterium avium*, *Mycobacterium scrofulaceum*, *Mycobacterium gordonae*, *Mycobacterium kansasii*, *Mycobacterium smegatis*, *Mycobacterium fortuitum*, *Mycobacterium gastri*, *Mycobacterium xenopi*, *Mycobacterium marinum* and *Mycobacterium phlei*; and

printed instructions specifying, in order of implementation, the steps to be followed for detecting said Actinomycetes nucleic acid by detecting a complex between the oligonucleotide probe and an Actinomycetes nucleic acid target, wherein said probe and said printed instructions are in packaged combination.

HIGH.G+C

S.aureus

SEQ ID NO: 12

E.coli

SEQ ID NO: 13

OMeFraB2040

SEQ ID NO: 3

MtuB2011

SEQ ID NO: 1

OMeFraB1986

SEQ ID NO: 2

Frankia .sp

SEQ ID NO: 14

M.tuberculosis

SEQ ID NO: 15

C.xerosis

SEQ ID NO: 16

S.firseus

SEQ ID NO: 17

M.avium

SEQ ID NO: 18

N.asteroides

SEQ ID NO: 19

UUGGGACUGUCUCAACGAGAGACUCGGUGAAUCAUAGUACCUUGAAGAUGCAGGUUACCCGCGACAGGACGGAAAGACCCC
GGCCAGGCUGUCUCCACCCGAGACUCAGUGAAAUUGAACUCGCUGUGAAGAUGCAGUGUACCCGCGCAAGACGGAAAGACCCC
(3 ') AAUGCGCGCCGUCUCCUGCCUUUCUGG (5 ')
(3 ') GCTTTAACGTGATGCTCATTTTCTACGAGC (5 ')
(3 ') GGGUGACAGAGUUGGUGUCUGAGCC (5 ')
UUUCCCAACUGUCUCAACCAACACAGACUCGGCGGAAAUUGCAUUACGAGUAAAGAUGCUCGUUACGCGCGGACGGAAAGACCCC
UUCUCAACUGUCUCAACCAUAGACUCGGCGAAAUUGCACUACGAGUAAAGAUGCUCGN - ACGCGCGGACGNCGAAANACCCC
CCCCGGCGAAAUUGCGAUACGAGUAAAGAUGCUNN - UACGCGCGGACGACGAAAGACCCC
UUCUCGACUGUCUCAACCAUAGGCCCGGUGAAAUUGCACUACGAGUAAAGAUGCUGUUCGCGCAGCAGGACGGAAAGACCCC
UUCCCAACUGUCUCAACCAUAGACUCGGCGAAAUUGCACUACGAGUAAAGAUGCUGNUACGCGCGCAGNCGAAAGACCCC
UUCUCUGCUGUCUCAACCAACAGACUCGGCGAAAUUGCAUUACGAGUAAAGAUGCUGUACGCGCGGACGACGAAAGACCCC

FIG. 1