



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12Q 1/68	A2	(11) International Publication Number: WO 00/61795 (43) International Publication Date: 19 October 2000 (19.10.00)
(21) International Application Number: PCT/EP00/02998 (22) International Filing Date: 5 April 2000 (05.04.00) (30) Priority Data: 99870068.6 9 April 1999 (09.04.99) EP 60/138,614 11 June 1999 (11.06.99) US (71) Applicant (for all designated States except US): INNOGENETICS N.V. [BE/BE]; Industriepark Zwijnaarde 7, Box 4, B-9052 Ghent (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): DE CANCK, Ilse [BE/BE]; Kruishofstraat 146, Bus 105, B-2020 Antwerpen (BE). ROMBOUT, Annelies [BE/BE]; Hospitaalstraat 3, B-9080 Beervelde (BE). ROSSAU, Rudi [BE/BE]; Wilgehoevestraat 45, B-2180 Ekeren (BE). (74) Common Representative: INNOGENETICS N.V.; Intellectual Property Department, Industriepark Zwijnaarde 7, Box 4, B-9052 Ghent (BE).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHOD FOR THE AMPLIFICATION OF HLA CLASS I ALLELES		
(57) Abstract <p>The present invention relates to a method and to specific primers for the locus-specific, separate amplification of exon 2, exon 3 and/or exon 4 of HLA-A, HLA-B or HLA-C alleles, making use of at least one primer set wherein: for the amplification of exon 2, the reverse primer specifically hybridizes to a locus-specific target sequence in intron 2 of respectively HLA-A, HLA-B or HLA-C; for the amplification of exon 3, the forward primer specifically hybridizes to a locus-specific target sequence in intron 2 of respectively HLA-A, HLA-B or HLA-C and/or the reverse primer specifically hybridizes to a locus-specific target sequence in intron 3 of respectively HLA-A, HLA-B or HLA-C; for the amplification of exon 4, the forward primer specifically hybridizes to a locus-specific target sequence in intron 3 of respectively HLA-A, HLA-B or HLA-C. In accordance, the present invention provides an improved method for the typing or subtyping of HLA Class I alleles making use of the amplification method of the invention.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHOD FOR THE AMPLIFICATION OF HLA CLASS I ALLELES

5 FIELD OF THE INVENTION

The present invention relates to a method for the typing or subtyping of HLA-A, HLA-B or HLA-C. More specifically, the present invention relates to a method for the locus-specific, separate amplification of exon 2, exon 3 and/or exon 4 of HLA-A,
10 HLA-B or HLA-C alleles.

BACKGROUND OF THE INVENTION

15 The human major histocompatibility complex (MHC) is contained within about 4 Mbp of DNA on the short arm of chromosome 6 at 6p21.3 (Campbell and Trowsdale, 1993). The human MHC is divided into class I, class II and class III regions. The genes of class I and class II encode highly polymorphic cell-surface molecules that bind and present processed antigens in the form of peptides to T-lymphocytes,
20 initiating both cellular and humoral immune responses.

The class I molecules of the human MHC, HLA-A, -B, and -C, are found on most nucleated cells. They are cell-surface glycoproteins that bind and present processed peptides derived from endogenously synthesized proteins to CD8+ T-cells. These heterodimers consist of an HLA-encoded α -chain associated with a non-MHC
25 encoded monomorphic polypeptide, β_2 -microglobulin (Townsend and Bodmer, 1989; Spencer and Parham, 1996). The class II molecules of the human MHC are encoded in the HLA-D region. These cell-surface glycoproteins consist of HLA-encoded α -, and β -chains, associated as heterodimers on the cell surface of antigen-presenting cells such as B-cells and macrophages. Class II molecules serve as receptors for
30 processed peptides. However, these peptides are derived predominantly from membrane and extracellular proteins and are presented to CD4+ T-cells. The HLA-D region contains several class II genes and has three main subregions: HLA-DR, -DQ, and -DP. Both the HLA-DQ and -DP regions contain one functional gene for each of

their α - and β -chains. The HLA-DR subregion contains one functional gene for the α -chain; the number of functional genes for the β -chain varies from one to two according to the haplotype (Andersson et al., 1987; Apple and Erlich, 1996).

- 5 Extensive polymorphism exists at most loci. In view of the biological and medical importance of these antigens, a highly sensitive and rapid technique for HLA typing is required. A variety of techniques are currently used to detect HLA polymorphism, including serological, biochemical, T-cell recognition and, most recently, molecular biological methods.
- 10 Serology remains the mainstay method for HLA typing - especially for class I - for many routine histocompatibility laboratories. The micro-lymphocytotoxicity assay (Kissmeyer et al., 1969; Terasaki and McClelland, 1964) is the standard approach: viable peripheral blood mononuclear cells (class I) or separate B-cells (class II) are mixed with antisera (polyclonal or monoclonal) of known HLA specificity.
- 15 Detection of polymorphism can be achieved by looking at the different amino acid composition of HLA molecules through biochemical techniques such as one-dimensional isoelectric focusing (IEF; Yang, 1987). This method relies on amino acid substitutions contributing to changes in charge of the HLA molecule.
- Another HLA typing method is the mixed lymphocyte reaction (MLR). Concurrent to
20 observations being made using HLA-specific antisera, it was noted that lymphocytes from two unrelated sources, when mixed in culture, would proliferate (Hirschorn et al., 1963).
- Analysis of HLA specificities from DNA provided a new approach to defining their polymorphic differences. Rather than looking at differences in the expressed
25 molecule, polymorphism is characterized at the nucleotide level.
- An important and powerful development in the field of molecular biology has been the polymerase chain reaction (PCR; Mullis et al., 1986; Mullis and Faloona, 1987). In tissue typing, PCR is used to amplify the polymorphic regions of HLA genes. This HLA PCR product can then be analyzed for its polymorphic differences, to establish
30 the tissue type. A number of such approaches have been developed, including hetero duplex analysis of PCR products (Clay et al., 1994), single-stranded conformational polymorphism analysis of the PCR product (PCR-SSCP; Yoshida *et al.*, 1992), sequence-based typing (SBT; Santamaria et al., 1992 and 1993), the use of sequence

specific primers in PCR reaction (PCR-SSP; Olerup and Zetterquist, 1991), the use of PCR in combination with sequence-specific oligonucleotide probing (PCR-SSOP; Saiki et al., 1986) or probing by reverse dot-blot (Saiki et al., 1989). These approaches, used singly or in combination, have all been applied as DNA-based methods for tissue-typing of class I and class II HLA specificities. DNA typing methods should be preferred over serological methods provided that an easy, rapid and reliable DNA typing method is available. Some differences at the subtype level which are detectable by DNA methods might go undetected by current serological typing methods, although these differences might provoke allograft rejection (Fleischhauer et al., 1990).

The HLA system is the most polymorphic human genetic system yet known. HLA class I genes share a similar structure (from 5' to 3'): a 5' untranslated flanking region, a first exon (exon 1) having a length of approximately 73 base pairs, a first intron (intron 1) having a length of approximately 130 base pairs, a second exon (exon 2), having a length of approximately 250 base pairs, a second intron (intron 2), having a length of approximately 272 base pairs, a third exon (exon 3), having a length of approximately 276 base pairs, a third intron (intron 3), having a length of approximately 588 base pairs and a fourth exon (exon 4), having a length of approximately 276 base pairs. Polymorphic substitutions within HLA class I alleles are mostly located in both exon 2 and exon 3, encoding the peptide binding groove of the class I molecule. These polymorphisms make differentiation between alleles achievable through a variety of molecular biological techniques such as sequencing or hybridization with relevant probes. In the current diagnostic kits exon 2 and exon 3 are amplified together, resulting in amplicons of about 1 kb, consisting at least of exon 2, intron 2 and exon 3. Locus-specific primers are available for the amplification of these 1 kb amplicons. However, such large amplicons are difficult to amplify and show secondary structure formation resulting in inefficient hybridization of some probes. In addition, due to the emergence of new HLA-Class I alleles, certain allele combinations cannot be distinguished anymore by the detection of polymorphism's only in exon 2 and exon 3 and additional typing in exon 4 is required. This raises the need for the additional amplification of exon 4, resulting in an even larger amplicon. Therefore, a separate amplification of exon 2, exon 3 and/or exon 4 would be desired resulting in amplification products that enable a more efficient typing of HLA class I

alleles. However, as locus-specific primer annealing sites are scarce and cannot be found in exon 2, exon 3 or exon 4, the separate and locus-specific amplification of exon 2, exon 3 and/or exon 4 of HLA-A, HLA-B or HLA-C is not that evident.

AIMS OF THE INVENTION

It is an aim of the present invention to provide a method for the locus-specific and separate amplification of exon 2, exon 3 and/or exon 4 of HLA-A, HLA-B or HLA-C alleles.

It is a more specific aim of the present invention to provide a method for a one-step, locus-specific, separate amplification of both exon 2 and exon 3 of HLA-A, HLA-B or HLA-C alleles.

It is a more specific aim of the present invention to provide a method for a one-step, locus-specific, separate amplification of both exon 2 and exon 4 of HLA-A, HLA-B or HLA-C alleles.

It is a more specific aim of the present invention to provide a method for a one-step, locus-specific, separate amplification of both exon 3 and exon 4 of HLA-A, HLA-B or HLA-C alleles.

It is a more specific aim of the present invention to provide a method for a one-step, locus-specific, separate amplification of all three exons, exon 2, exon 3 and exon 4, of HLA-A, HLA-B or HLA-C alleles.

It is another aim of the present invention to provide a primer for use in a method for the locus-specific, separate amplification of exon 2 of HLA-A, HLA-B or HLA-C alleles.

It is another aim of the present invention to provide a primer for use in a method for the locus-specific, separate amplification of exon 3 of HLA-A, HLA-B or HLA-C alleles.

It is another aim of the present invention to provide a primer for use in a method for the locus-specific, separate amplification of exon 4 of HLA-A, HLA-B or HLA-C alleles.

It is a more specific aim of the present invention to provide a primer set for use in a method for the locus-specific, separate amplification of exon 2 of HLA-A, HLA-B or HLA-C alleles.

It is another more specific aim of the present invention to provide a primer set for use in a method for the locus-specific, separate amplification of exon 3 of HLA-A, HLA-B or HLA-C alleles.

It is another more specific aim of the present invention to provide a primer set for use in a method for the locus-specific, separate amplification of exon 4 of HLA-A, HLA-B or HLA-C alleles.

5 It is another more specific aim of the present invention to provide a multiplex primer mix for use in a method for the one step, locus-specific, separate amplification of exon 2 and exon 3 of HLA-A, HLA-B or HLA-C.

It is another more specific aim of the present invention to provide a multiplex primer mix for use in a method for the one step, locus-specific, separate amplification of exon 2 and exon 4 of HLA-A, HLA-B or HLA-C.

10 It is another more specific aim of the present invention to provide a multiplex primer mix for use in a method for the one step, locus-specific, separate amplification of exon 3 and exon 4 of HLA-A, HLA-B or HLA-C.

It is another more specific aim of the present invention to provide a multiplex primer mix for use in a method for the one step, locus-specific, separate amplification of
15 exon 2, exon 3 and exon 4 of HLA-A, HLA-B or HLA-C.

It is another aim of the present invention to provide an improved method for the typing or subtyping of one or more HLA-A, HLA-B or HLA-C alleles in a sample.

It is another aim of the present invention to provide an improved diagnostic kit for the typing or subtyping of one or more HLA-A, HLA-B or HLA-C alleles in a sample.

20 It is a further aim of the present invention to provide an improved Line Probe Assay for the typing or subtyping of one or more HLA-A, HLA-B or HLA-C alleles in a sample.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to a method for the locus-specific, separate amplification of exon 2, exon 3 and/or exon 4 of HLA-A, HLA-B or HLA-C alleles, making use of

5 at least one primer set wherein:

- for the amplification of exon 2, the reverse primer specifically hybridizes to a locus-specific target sequence in intron 2 of respectively HLA-A, HLA-B or HLA-C;
- 10 - for the amplification of exon 3, the forward primer specifically hybridizes to a locus-specific target sequence in intron 2 of respectively HLA-A, HLA-B or HLA-C and/or the reverse primer specifically hybridizes to a locus-specific target sequence in intron 3 of respectively HLA-A, HLA-B or HLA-C;
- 15 - for the amplification of exon 4, the forward primer specifically hybridizes to a locus-specific target sequence in intron 3 of respectively HLA-A, HLA-B or HLA-C.

The amplification method of the invention thus, makes use of a primer set of which at least one of the primers is hybridizing to a locus-specific target sequence in intron 2 or in intron 3 of the HLA Class I gene in order to obtain the separate amplification of
20 exon 2 (without exon 3 or exon 4), exon 3 (without exon 2 or exon 4) and/or exon 4 (without exon 2 or exon 3).

In the case where exon 2 is amplified, the reverse primer hybridizes to a locus-specific target sequence in intron 2 while the forward primer might be chosen to hybridize to a region upstream of exon 2 (e.g. in intron 1 or in exon 1) or to the initial nucleotides of
25 exon 2. In a preferred embodiment, the forward primer hybridizes to a locus-specific region upstream of exon 2 (e.g. in intron 1 or in exon 1) or to locus-specific initial nucleotides of exon 2 and the reverse primer hybridizes to a locus-specific target sequence in intron 2.

In the case where exon 3 is amplified, the forward primer is hybridizing to a locus-specific target sequence in intron 2 while the reverse primer might be chosen to
30 hybridize to the final nucleotides of exon 3 or downstream of exon 3 (e.g. in intron 3) or the reverse primer is hybridizing to a locus-specific target sequence in intron 3 while the forward primer might be chosen to hybridize to a region upstream to exon 3 (e.g. intron 2) or to the initial nucleotides of exon 3. In a preferred embodiment, exon

3 is amplified by use of a primer set of which the forward primer is hybridizing to a locus-specific target sequence in intron 2 and the reverse primer is hybridizing to a locus-specific target sequence in intron 3.

In the case where exon 4 is amplified, the forward primer is hybridizing to a locus-specific target sequence in intron 3 while the reverse primer might be chosen to hybridize to the final nucleotides of exon 4 or downstream of exon 4 (e.g. in intron 4). In a preferred embodiment, the forward primer is hybridizing to a locus-specific target sequence in intron 3 while the reverse primer hybridizes to locus-specific target sequences at the end of exon 4 or downstream of exon 4 (e.g. in intron 4).

10 This new amplification method will result in the amplification of shorter DNA fragments, containing only exon 2, only exon 3 or only exon 4, which are much easier to amplify and much easier for use in different typing methods such as sequencing or hybridization with different allele specific probes. From the example section it is clear that primer sets containing one primer hybridizing to a target sequence in intron 2 or
15 intron 3 provide a much better and easier amplification of exon 2, exon 3 or exon 4 of HLA Class I alleles and a more clear and pronounced hybridization pattern with the allele typing probes. It has been the merit of the present inventors to define specific primers that enable this separate amplification of exon 2, exon 3 or exon 4 of only one HLA locus, while said exons of other HLA loci (i.e. other classical HLA genes, non-classical HLA genes or pseudogenes) are not co-amplified. As a-specific amplicons in the amplification product might result in a fault typing, the amplification method of the invention will certainly improve the security of the present typing methods.

The term "locus-specific" thus means that exon 2, exon 3 and/or exon 4 of only one locus (i.e. HLA-A, HLA-B or HLA-C) is amplified while exon 2, exon 3 and/or exon
25 4 of the other HLA loci (i.e. other classical HLA genes, non-classical HLA genes or pseudogenes) is not amplified. Thus, when primers specific for the amplification of exon 2, exon 3 and/or exon 4 of HLA-A are used, exon 2, exon 3 and/or exon 4 of HLA-B, HLA-C or of other HLA loci are not amplified. Similarly, when primers specific for the amplification of exon 2, exon 3 and/or exon 4 of HLA-B are used,
30 exon 2, exon 3 and/or exon 4 of HLA-A, HLA-C or of other HLA loci are not amplified. Similarly, when primers specific for the amplification of exon 2, exon 3 and/or exon 4 of HLA-C are used, exon 2, exon 3 and/or exon 4 of HLA-A, HLA-B or of other HLA loci are not amplified. From this it is clear that exon 2, exon 3 and/or exon 4 of 2, 3 or more different HLA loci can never be amplified together. Thus, exon

2, exon 3 and/or exon 4 of HLA-A can never be amplified in the same reaction tube with exon 2, exon 3 and/or exon 4 of HLA-B. Similarly, exon 2, exon 3 and/or exon 4 of HLA-A can never be amplified in the same reaction tube with exon 2, exon 3 and/or exon 4 of HLA-C. Similarly, exon 2, exon 3 and/or exon 4 of HLA-B can never be amplified in the same reaction tube with exon 2, exon 3 and/or exon 4 of HLA-C. Similarly, exon 2, exon 3 and/or exon 4 of HLA-A can never be amplified in the same reaction tube with exon 2, exon 3 and/or exon 4 of HLA-B and exon 2, exon 3 and/or exon 4 of HLA-C. As a consequence, the present invention relates to a method for the locus-specific, separate amplification of exon 2, exon 3 and/or exon 4 of HLA-A, HLA-B or HLA-C alleles. This means that the invention relates to a method for the separate amplification of exon 2, exon 3 and/or exon 4 of HLA-A, the invention relates to a method for the separate amplification of exon 2, exon 3 and/or exon 4 of HLA-B and the invention relates to a method for the separate amplification of exon 2, exon 3 and/or exon 4 of HLA-C.

The term “primer” refers to a single stranded oligonucleotide sequence capable of acting as a point of initiation for the synthesis of a primer extension product that is complementary to the nucleic acid strand to be copied. The length and the sequence of the primer must be such that they allow to prime the synthesis of the extension products. In a specific embodiment, the length of the primer is about 5-50 nucleotides. In another specific embodiment, the length of the primer is about 10-30 nucleotides. In another specific embodiment, the length of the primers is about 20-25 nucleotides. Specific length and sequence will depend on the complexity of the required DNA or RNA target, as well as on the conditions at which the primer is used, such as temperature and ionic strength.

The expression “primer set” or “primer pair” refers to a pair of primers allowing the amplification of part or all of exon 2, exon 3 or exon 4 of HLA-A, HLA-B or HLA-C. A primer set always consists of a forward primer (or 5' primer) and a reverse primer (or 3' primer).

The terms “to hybridize specifically” and “specifically hybridizes” mean that, during the amplification step, said primer forms a duplex with part of its target sequence or with the entire target sequence under the experimental conditions used, and that under those conditions said primer does not form a duplex with other sequences of the polynucleic acids present in the sample to be analysed. It should be understood that primers that are designed to specifically hybridize to a target sequence of a nucleic

acid, may fall within said target sequence or may to a large extent overlap with said target sequence (i.e. form a duplex with nucleotides outside as well as within said target sequence).

The term "target sequence" of a primer according to the present invention is a sequence within intron 2 or intron 3 of the HLA Class I alleles to which the primer is completely complementary or partially complementary (i.e. with up to 20%, 15%, 10% or 5% mismatches). It is to be understood that the complement of said target sequence is also a suitable target sequence in some cases. The fact that amplification primers do not have to match exactly with the corresponding target sequence in the template to warrant proper amplification is amply documented in the literature (Kwok et al., 1990). However, when the primers are not completely complementary to their target sequence, it should be taken into account that the amplified fragments will have the sequence of the primers and not of the target sequence.

In one embodiment, the amplification is performed in a reaction tube with a primer set for the amplification of exon 2 as described above (reverse primer hybridizes with a locus-specific target sequence in intron 2) and consequently only exon 2 is amplified.

In another embodiment, the amplification is performed in a reaction tube with a primer set for the amplification of exon 3 as described above (forward primer hybridizes with a locus-specific target sequence in intron 2) and consequently only exon 3 is amplified.

In another embodiment, the amplification is performed in a reaction tube with a primer set for the amplification of exon 3 as described above (reverse primer hybridizes with a locus-specific target sequence in intron 3) and consequently only exon 3 is amplified.

In another embodiment, the amplification is performed in a reaction tube with a primer set for the amplification of exon 3 as described above (forward primer hybridizes with a locus-specific target sequence in intron 2 and reverse primer hybridizes with a locus-specific target sequence in intron 3) and consequently only exon 3 is amplified.

In another embodiment, the amplification is performed in a reaction tube with a primer set for the amplification of exon 4 as described above (forward primer hybridizes with a locus-specific target sequence in intron 3) and consequently only exon 4 is amplified.

Accordingly, the present invention relates to a method for the separate amplification of exon 2, exon 3 or exon 4 of HLA Class I alleles.

In another embodiment, the different primer sets involved in the amplification of exon 2 and the amplification of exon 3 are mixed and the separate amplification of both
5 exon 2 and exon 3 is performed in a single reaction tube. Thus, the present invention also relates to a method as described above further characterized that both exon 2 and exon 3 of HLA-A, HLA-B or HLA-C are amplified by use of a multiplex primer mix containing at least one primer pair for amplification of exon 2 and at least one primer pair for amplification of exon 3. Accordingly, the present invention relates to a
10 method for the separate amplification of exon 2 and exon 3 of HLA Class I alleles.

In another embodiment, the different primer sets involved in the amplification of exon 2 and the amplification of exon 4 are mixed and the separate amplification of both exon 2 and exon 4 is performed in a single reaction tube. Thus, the present invention also relates to a method as described above further characterized that both exon 2 and
15 exon 4 of HLA-A, HLA-B or HLA-C are amplified by use of a multiplex primer mix containing at least one primer pair for amplification of exon 2 and at least one primer pair for amplification of exon 4. Accordingly, the present invention relates to a method for the separate amplification of exon 2 and exon 4 of HLA Class I alleles.

In another embodiment, the different primer sets involved in the amplification of exon 3 and the amplification of exon 4 are mixed and the separate amplification of both
20 exon 3 and exon 4 is performed in a single reaction tube. Thus, the present invention also relates to a method as described above further characterized that both exon 3 and exon 4 of HLA-A, HLA-B or HLA-C are amplified by use of a multiplex primer mix containing at least one primer pair for amplification of exon 3 and at least one primer pair for amplification of exon 4. Accordingly, the present invention relates to a
25 method for the separate amplification of exon 3 and exon 4 of HLA Class I alleles.

In another embodiment, the different primer sets involved in the amplification of exon 2, the amplification of exon 3 and the amplification of exon 4 are mixed and the separate amplification of all three exons, exon 2, exon 3 and exon 4, is performed in a
30 single reaction tube. Thus, the present invention also relates to a method as described above further characterized that all three exons, exon 2, exon 3 and exon 4, of HLA-A, HLA-B or HLA-C are amplified by use of a multiplex primer mix containing at least one primer pair for amplification of exon 2, at least one primer pair for the amplification of exon 3 and at least one primer pair for amplification of exon 4.

Accordingly, the present invention relates to a method for the separate amplification of exon 2, exon 3 and exon 4 of HLA Class I alleles.

The amplification method used can be either polymerase chain reaction (PCR; Saiki et al., 1988), ligase chain reaction (LCR; Landgren et al., 1988; Wu and Wallace, 1989; 5 Barany, 1991), nucleic acid sequence-based amplification (NASBA; Guatelli et al., 1990; Compton, 1991), transcription-based amplification system (TAS; Kwoh et al., 1989), strand displacement amplification (SDA; Duck, 1990) or amplification by means of Q β replicase (Lomeli et al., 1989) or any other suitable method to amplify nucleic acid molecules known in the art. Also TMA (Guatelli et al., 1990) or bDNA 10 (Sanchez-Pescador et al., 1988; Urdea et al., 1991) techniques can be used in the method of the present invention. In a specific embodiment, exon 2, exon 3 and/or exon 4 of HLA-A, HLA-B or HLA-C are amplified by PCR.

In a specific embodiment, the present invention relates to a method as described above, further characterized that the locus-specific target sequence is situated at one 15 of the following positions:

- 67, 96, 109, 110, 118, 123, 131 or 181 of the HLA-A intron 2 (Fig. 1) and/or 32, 50, 62, 73, 83, 86, 118, 130, 150, 501, 525, 561 or 571 (Fig. 4) of the HLA-A intron 3; or
- 35 or 170 of the HLA-B intron 2 (Fig. 2) and/or 42, 46, 65, 68, 96, 438, 20 502, 524, 547 or 571 of the HLA-B intron 3 (Fig. 5); or
- 84, 107 or 142 of the HLA-C intron 2 (Fig. 3) and/or 461, 477, 527, 545 or 561 of the HLA-C intron 3 (Fig. 6).

As these positions all contain locus-specific nucleotides, these position are particularly suitable for designing an efficient primer for the locus-specific 25 amplification of exon 2, exon 3 or exon 4. The primers can be of different length. In a specific embodiment, the length of the primer is about 5-50 nucleotides. In another specific embodiment, the length of the primer is about 10-30 nucleotides. In another specific embodiment, the length of the primers is about 20-25 nucleotides. Specific length and sequence will depend on the complexity of the required DNA or RNA 30 target, as well as on the conditions at which the primer is used, such as temperature and ionic strength.

In a specific embodiment, the present invention relates to a method as described above further characterized that said positions constitute the 3' end of the primer that is used for the amplification of exon 2, exon 3 or exon 4. Thus, from the above

described positions, forward as well as reverse primers can be designed that have their 3' end in these specific positions. The reverse primer having its 3' end in one of the above-mentioned specific positions of intron 2 will enable the locus-specific amplification of exon 2 of the respective HLA-A, HLA-B or HLA-C allele. The forward primer having its 3' end in one of the above-mentioned specific positions of intron 2 will enable the locus-specific amplification of exon 3 of the respective HLA-A, HLA-B or HLA-C allele. The reverse primer having its 3' end in one of the above-mentioned specific positions of intron 3 will enable the locus-specific amplification of exon 3 of the respective HLA-A, HLA-B or HLA-C allele. The forward primer having its 3' end in one of the above-mentioned specific positions of intron 3 will enable the locus-specific amplification of exon 4 of the respective HLA-A, HLA-B or HLA-C allele. The primers can be of different length. In a specific embodiment, the length of the primer is about 5-50 nucleotides. In another specific embodiment, the length of the primer is about 10-30 nucleotides. In another specific embodiment, the length of the primers is about 20-25 nucleotides. Specific length and sequence will depend on the complexity of the required DNA or RNA target, as well as on the conditions at which the primer is used, such as temperature and ionic strength.

In another embodiment, the present invention relates to a method as described above, further characterized that the primer is chosen from the following list:

- 20 - for the amplification of exon 2 of HLA-A (table 1):
 - 5'ATCTCGGACCCGGAGACTGT3' (SEQ ID NO 1)
 - 5'GATCTCGGACCCGGAGACTGT3' (SEQ ID NO 2)
 - 5'GGATCTCGGACCCGGAGACTGT3' (SEQ ID NO 3)
 - 5'YGGATCTCGGACCCGGAGACTGT3' (SEQ ID NO 4)
 - 25 5'GYGGATCTCGGACCCGGAGACTGT3' (SEQ ID NO 5)
 - 5'GGYGGATCTCGGACCCGGAGACTGT3' (SEQ ID NO 6)
 - 5'GGTCTCGGRGTCCCGCGGCT3' (SEQ ID NO 7)
 - 5'GGGTCTCGGRGTCCCGCGGCT3' (SEQ ID NO 8)
 - 5'AGGGTCTCGGRGTCCCGCGGCT3' (SEQ ID NO 9)
 - 30 5'AAGGGTCTCGGRGTCCCGCGGCT3' (SEQ ID NO 10)
 - 5'CAAGGGTCTCGGRGTCCCGCGGCT3' (SEQ ID NO 11)
 - 5'CTCCCGGGDCAAGGGTCTCG3' (SEQ ID NO 12)
 - 5'TCTCCCGGGDCAAGGGTCTCG3' (SEQ ID NO 13)
 - 5'CTCTCCCGGGDCAAGGGTCTCG3' (SEQ ID NO 14)

5'CCTCTCCCGGGDCAAGGGTCTCG3' (SEQ ID NO 15)
5'GCCTCTCCCGGGDCAAGGGTCTCG3' (SEQ ID NO 16)
5'GGCCTCTCCCGGGDCAAGGGTCTCG3' (SEQ ID NO 17)
5'TCTCCCGGGDCAAGGGTCTC3' (SEQ ID NO 18)
5'CTCTCCCGGGDCAAGGGTCTC3' (SEQ ID NO 19)
5'CCTCTCCCGGGDCAAGGGTCTC3' (SEQ ID NO 20)
5'GCCTCTCCCGGGDCAAGGGTCTC3' (SEQ ID NO 21)
5'GGCCTCTCCCGGGDCAAGGGTCTC3' (SEQ ID NO 22)
5'GGGCCTCTCCCGGGDCAAGGGTCTC3' (SEQ ID NO 23)
5'CCTGGGCCTCTCCCGGGDCA3' (SEQ ID NO 30)
5'GCCTGGGCCTCTCCCGGGDCA3' (SEQ ID NO 31)
5'CGCCTGGGCCTCTCCCGGGDCA3' (SEQ ID NO 32)
5'GCGCCTGGGCCTCTCCCGGGDCA3' (SEQ ID NO 33)
5'GGCGCCTGGGCCTCTCCCGGGDCA3' (SEQ ID NO 34)
5'AGGCGCCTGGGCCTCTCCCGGGDCA3' (SEQ ID NO 35)
5'AGGCGCCTGGGCCTCTCCCG3' (SEQ ID NO 36)
5'AAGGCGCCTGGGCCTCTCCCG3' (SEQ ID NO 37)
5'WAAGGCGCCTGGGCCTCTCCCG3' (SEQ ID NO 38)
5'TWAAGGCGCCTGGGCCTCTCCCG3' (SEQ ID NO 39)
5'GTWAAGGCGCCTGGGCCTCTCCCG3' (SEQ ID NO 40)
5'GGTWAAGGCGCCTGGGCCTCTCCCG3' (SEQ ID NO 41)
5'CCGGGTWAAGGCGCCTGGGC3' (SEQ ID NO 42)
5'ACCGGGTWAAGGCGCCTGGGC3' (SEQ ID NO 43)
5'AACCGGGTWAAGGCGCCTGGGC3' (SEQ ID NO 44)
5'AAACCGGGTWAAGGCGCCTGGGC3' (SEQ ID NO 45)
5'GAAACCGGGTWAAGGCGCCTGGGC3' (SEQ ID NO 46)
5'TGAAACCGGGTWAAGGCGCCTGGGC3' (SEQ ID NO 47)
5'YCCVGGCCCCGACCAACCYGG3' (SEQ ID NO 48)
5'GYCCVGGCCCCGACCAACCYGG3' (SEQ ID NO 49)
5'YGYCCVGGCCCCGACCAACCYGG3' (SEQ ID NO 50)
5'CYGYCCVGGCCCCGACCAACCYGG3' (SEQ ID NO 51)
5'CCYGYCCVGGCCCCGACCAACCYGG3' (SEQ ID NO 52)
5'CCCYGYCCVGGCCCCGACCAACCYGG3' (SEQ ID NO 53)

- for the amplification of exon 3 of HLA-A (table 2; table 3):

5'CGGACGGGCCRGGTSRCCCA3' (SEQ ID NO 54)
 5'ACGGACGGGCCRGGTSRCCCA3' (SEQ ID NO 55)
 5'CACGGACGGGCCRGGTSRCCCA3' (SEQ ID NO 56)
 5'CCACGGACGGGCCRGGTSRCCCA3' (SEQ ID NO 57)
 5'CCCACGGACGGGCCRGGTSRCCCA3' (SEQ ID NO 58)
 5'CCCCACGGACGGGCCRGGTSRCCCA3' (SEQ ID NO 59)
 5'GGTCCGAGATCCRCCCCGAA3' (SEQ ID NO 60)
 5'GGGTCCGAGATCCRCCCCGAA3' (SEQ ID NO 61)
 5'CGGGTCCGAGATCCRCCCCGAA3' (SEQ ID NO 62)
 5'CCGGGTCCGAGATCCRCCCCGAA3' (SEQ ID NO 63)
 5'TCCGGGTCCGAGATCCRCCCCGAA3' (SEQ ID NO 64)
 5'CTCCGGGTCCGAGATCCRCCCCGAA3' (SEQ ID NO 65)
 5'CCCCGAAGCCGCGGGACYCC3' (SEQ ID NO 66)
 5'RCCCCGAAGCCGCGGGACYCC3' (SEQ ID NO 67)
 5'CRCCCCGAAGCCGCGGGACYCC3' (SEQ ID NO 68)
 5'CCRCCCCGAAGCCGCGGGACYCC3' (SEQ ID NO 69)
 5'TCCRCCCCGAAGCCGCGGGACYCC3' (SEQ ID NO 70)
 5'ATCCRCCCCGAAGCCGCGGGACYCC3' (SEQ ID NO 71)
 5'CCCGAAGCCGCGGGACYCCG3' (SEQ ID NO 72)
 5'CCCCGAAGCCGCGGGACYCCG3' (SEQ ID NO 73)
 5'RCCCCGAAGCCGCGGGACYCCG3' (SEQ ID NO 74)
 5'CRCCCCGAAGCCGCGGGACYCCG3' (SEQ ID NO 75)
 5'CCRCCCCGAAGCCGCGGGACYCCG3' (SEQ ID NO 76)
 5'TCCRCCCCGAAGCCGCGGGACYCCG3' (SEQ ID NO 77)
 5'CGCGGGACYCCGAGACCCTT3' (SEQ ID NO 84)
 5'CCGCGGGACYCCGAGACCCTT3' (SEQ ID NO 85)
 5'GCCGCGGGACYCCGAGACCCTT3' (SEQ ID NO 86)
 5'AGCCGCGGGACYCCGAGACCCTT3' (SEQ ID NO 87)
 5'AAGCCGCGGGACYCCGAGACCCTT3' (SEQ ID NO 88)
 5'GAAGCCGCGGGACYCCGAGACCCTT3' (SEQ ID NO 89)
 5'GACYCCGAGACCCTTGDCCC3' (SEQ ID NO 90)
 5'GGACYCCGAGACCCTTGDCCC3' (SEQ ID NO 91)
 5'GGGACYCCGAGACCCTTGDCCC3' (SEQ ID NO 92)

5'CGGGACYCCGAGACCCTTGDCCC3' (SEQ ID NO 93)
5'GCGGGACYCCGAGACCCTTGDCCC3' (SEQ ID NO 94)
5'CGCGGGACYCCGAGACCCTTGDCCC3' (SEQ ID NO 95)
5'GACCCTTGDCCCGGGAGAGG3' (SEQ ID NO 96)
5 5'AGACCCTTGDCCCGGGAGAGG3' (SEQ ID NO 97)
5'GAGACCCTTGDCCCGGGAGAGG3' (SEQ ID NO 98)
5'CGAGACCCTTGDCCCGGGAGAGG3' (SEQ ID NO 99)
5'CCGAGACCCTTGDCCCGGGAGAGG3' (SEQ ID NO 100)
5'YCCGAGACCCTTGDCCCGGGAGAGG3' (SEQ ID NO 101)
10 5'GTTTAGGCCAAAAATCCCCC3' (SEQ ID NO 102)
5'AGTTTAGGCCAAAAATCCCCC3' (SEQ ID NO 103)
5'CAGTTTAGGCCAAAAATCCCCC3' (SEQ ID NO 104)
5'TCAGTTTAGGCCAAAAATCCCCC3' (SEQ ID NO 105)
5'TTCAGTTTAGGCCAAAAATCCCCC3' (SEQ ID NO 106)
15 5'TTTCAGTTTAGGCCAAAAATCCCCC3' (SEQ ID NO 107)
5'AGCCCGGGAGATCTAYAGGC3' (SEQ ID NO 150)
5'CAGCCCGGGAGATCTAYAGGC3' (SEQ ID NO 151)
5'CCAGCCCGGGAGATCTAYAGGC3' (SEQ ID NO 152)
5'GCCAGCCCGGGAGATCTAYAGGC3' (SEQ ID NO 153)
20 5'GGCCAGCCCGGGAGATCTAYAGGC3' (SEQ ID NO 154)
5'AGGCCAGCCCGGGAGATCTAYAGGC3' (SEQ ID NO 155)
5'CCCTCCTTGTGGGAGGCCAG3' (SEQ ID NO 156)
5'CCCCTCCTTGTGGGAGGCCAG3' (SEQ ID NO 157)
5'TCCCCTCCTTGTGGGAGGCCAG3' (SEQ ID NO 158)
25 5'CTCCCCTCCTTGTGGGAGGCCAG3' (SEQ ID NO 159)
5'TCTCCCCTCCTTGTGGGAGGCCAG3' (SEQ ID NO 160)
5'GTCTCCCCTCCTTGTGGGAGGCCAG3' (SEQ ID NO 161)
5'CCCAAWTGTCTCCCCTCCTT3' (SEQ ID NO 162)
5'TCCCAAWTGTCTCCCCTCCTT3' (SEQ ID NO 163)
30 5'GTCCCAAWTGTCTCCCCTCCTT3' (SEQ ID NO 164)
5'GGTCCCAAWTGTCTCCCCTCCTT3' (SEQ ID NO 165)
5'TGGTCCCAAWTGTCTCCCCTCCTT3' (SEQ ID NO 166)
5'TTGGTCCCAAWTGTCTCCCCTCCTT3' (SEQ ID NO 167)
5'CTAGTGTTGGTCCCAAWTGT3' (SEQ ID NO 168)

5'TCTAGTGTTGGTCCCAAWTGT3' (SEQ ID NO 169)
5'TTCTAGTGTTGGTCCCAAWTGT3' (SEQ ID NO 170)
5'ATTCTAGTGTTGGTCCCAAWTGT3' (SEQ ID NO 171)
5'TATTCTAGTGTTGGTCCCAAWTGT3' (SEQ ID NO 172)
5 ATATTCTAGTGTTGGTCCCAAWTGT3' (SEQ ID NO 173)
5'GGGYGATATTCTAGTGTTGG3' (SEQ ID NO 174)
5'AGGGYGATATTCTAGTGTTGG3' (SEQ ID NO 175)
5'GAGGGYGATATTCTAGTGTTGG3' (SEQ ID NO 176)
5'GGAGGGYGATATTCTAGTGTTGG3' (SEQ ID NO 177)
10 5'GGGAGGGYGATATTCTAGTGTTGG3' (SEQ ID NO 178)
5'AGGGAGGGYGATATTCTAGTGTTGG3' (SEQ ID NO 179)
5'GGAGGGYGATATTCTAGTGT3' (SEQ ID NO 180)
5'GGGAGGGYGATATTCTAGTGT3' (SEQ ID NO 181)
5'AGGGAGGGYGATATTCTAGTGT3' (SEQ ID NO 182)
15 5'GAGGGAGGGYGATATTCTAGTGT3' (SEQ ID NO 183)
5'AGAGGGAGGGYGATATTCTAGTGT3' (SEQ ID NO 184)
5'CAGAGGGAGGGYGATATTCTAGTGT3' (SEQ ID NO 185)
5'CCCAGGAGGAKTCCTCTCCC3' (SEQ ID NO 186)
5'ACCCAGGAGGAKTCCTCTCCC3' (SEQ ID NO 187)
20 5'AACCCAGGAGGAKTCCTCTCCC3' (SEQ ID NO 188)
5'AAACCCAGGAGGAKTCCTCTCCC3' (SEQ ID NO 189)
5'GAAACCCAGGAGGAKTCCTCTCCC3' (SEQ ID NO 190)
5'GGAAACCCAGGAGGAKTCCTCTCCC3' (SEQ ID NO 191)
5'AGGATCTGGAAACCCAGGAG3' (SEQ ID NO 192)
25 5'CAGGATCTGGAAACCCAGGAG3' (SEQ ID NO 193)
5'ACAGGATCTGGAAACCCAGGAG3' (SEQ ID NO 194)
5'TACAGGATCTGGAAACCCAGGAG3' (SEQ ID NO 195)
5'GTACAGGATCTGGAAACCCAGGAG3' (SEQ ID NO 196)
5'GGTACAGGATCTGGAAACCCAGGAG3' (SEQ ID NO 197)
30 5'TCAGAGTCACTCTCTGGTAC3' (SEQ ID NO 198)
5'CTCAGAGTCACTCTCTGGTAC3' (SEQ ID NO 199)
5'CCTCAGAGTCACTCTCTGGTAC3' (SEQ ID NO 200)
5'ACCTCAGAGTCACTCTCTGGTAC3' (SEQ ID NO 201)
5'AACCTCAGAGTCACTCTCTGGTAC3' (SEQ ID NO 202)

5'GAACCTCAGAGTCACTCTCTGGTAC3' (SEQ ID NO 203)

- for the amplification of exon 4 of HLA-A (table 4):

5' TTCTGTGCTCYCTTCCCCAT3' (SEQ ID NO 204)
 5' GTTCTGTGCTCYCTTCCCCAT3' (SEQ ID NO 205)
 5' GGTTCCTGTGCTCYCTTCCCCAT3' (SEQ ID NO 206)
 5' GGGTTCCTGTGCTCYCTTCCCCAT3' (SEQ ID NO 207)
 5' TGGGTTCCTGTGCTCYCTTCCCCAT3' (SEQ ID NO 208)
 5' CTGGGTTCCTGTGCTCYCTTCCCCAT3' (SEQ ID NO 209)
 5' GGTGTCCTGTCCATTCTCAA3' (SEQ ID NO 24)
 5' RGGTGTCTGTCCATTCTCAA3' (SEQ ID NO 25)
 5' CRGGTGTCTGTCCATTCTCAA3' (SEQ ID NO 26)
 5' CCRGGTGTCTGTCCATTCTCAA3' (SEQ ID NO 27)
 5' CCCRGGTGTCTGTCCATTCTCAA3' (SEQ ID NO 28)
 5' TCCCRGGTGTCTGTCCATTCTCAA3' (SEQ ID NO 29)
 5' CTGGWGGAGTGTCCCATKAC3' (SEQ ID NO 210)
 5' GCTGGWGGAGTGTCCCATKAC3' (SEQ ID NO 211)
 5' TGCTGGWGGAGTGTCCCATKAC3' (SEQ ID NO 212)
 5' RTGCTGGWGGAGTGTCCCATKAC3' (SEQ ID NO 213)
 5' YRTGCTGGWGGAGTGTCCCATKAC3' (SEQ ID NO 214)
 5' GYRTGCTGGWGGAGTGTCCCATKAC3' (SEQ ID NO 215)
 5' GTCCCATKACAGATRCMMAA3' (SEQ ID NO 216)
 5' TGTCCCATKACAGATRCMMAA3' (SEQ ID NO 217)
 5' GTGTCCCATKACAGATRCMMAA3' (SEQ ID NO 218)
 5' AGTGTCCCATKACAGATRCMMAA3' (SEQ ID NO 219)
 5' GAGTGTCCCATKACAGATRCMMAA3' (SEQ ID NO 220)
 5' GGAGTGTCCCATKACAGATRCMMAA3' (SEQ ID NO 221)

- for the amplification of exon 2 of HLA-B (table 5):

5' ACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 108)
 5' AACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 109)
 5' CAACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 110)
 5' CCAACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 111)
 5' MCCAACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 112)

5'GMCCAACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 113)

5'YGMCCAACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 314)

- for the amplification of exon 3 of HLA-B (table 6; table 7):

5 5'CYGGGGCGSAGGTCACGACT3' (SEQ ID NO 114)
 5'CCYGGGGCGSAGGTCACGACT3' (SEQ ID NO 115)
 5'GCCYGGGGCGSAGGTCACGACT3' (SEQ ID NO 116)
 5'GGCCYGGGGCGSAGGTCACGACT3' (SEQ ID NO 117)
 5'CGGCCYGGGGCGSAGGTCACGACT3' (SEQ ID NO 118)
 10 5'CCGGCCYGGGGCGSAGGTCACGACT3' (SEQ ID NO 119)
 5'CCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 120)
 5'ACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 121)
 5'TACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 122)
 5'TTACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 123)
 15 5'TTTACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 124)
 5'GTTTACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 125)
 5'CGTTTACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 222)
 5'GCGTTTACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 223)
 5'CGCGTTTACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 224)
 20 5'CGTKGGAGSCCATCCCCGSC3' (SEQ ID NO 225)
 5'TCGTKGGAGSCCATCCCCGSC3' (SEQ ID NO 226)
 5'CTCGTKGGAGSCCATCCCCGSC3' (SEQ ID NO 227)
 5'TCTCGTKGGAGSCCATCCCCGSC3' (SEQ ID NO 228)
 5'TTCTCGTKGGAGSCCATCCCCGSC3' (SEQ ID NO 229)
 25 5'CTTCTCGTKGGAGSCCATCCCCGSC3' (SEQ ID NO 230)
 5'TCTCGTKGGAGSCCATCCCC3' (SEQ ID NO 231)
 5'TTCTCGTKGGAGSCCATCCCC3' (SEQ ID NO 232)
 5'CTTCTCGTKGGAGSCCATCCCC3' (SEQ ID NO 233)
 5'TCTTCTCGTKGGAGSCCATCCCC3' (SEQ ID NO 234)
 30 5'YTCTTCTCGTKGGAGSCCATCCCC3' (SEQ ID NO 235)
 5'CYTCTTCTCGTKGGAGSCCATCCCC3' (SEQ ID NO 236)
 5'GATCCCATTTTCCTCYTCTT3' (SEQ ID NO 237)
 5'TGATCCCATTTTCCTCYTCTT3' (SEQ ID NO 238)
 5'CTGATCCCATTTTCCTCYTCTT3' (SEQ ID NO 239)

5'GCTGATCCCATTTTCCTCYTCTT3' (SEQ ID NO 240)
 5'CGCTGATCCCATTTTCCTCYTCTT3' (SEQ ID NO 241)
 5'GCGCTGATCCCATTTTCCTCYTCTT3' (SEQ ID NO 242)
 5'GCTGATCCCATTTTCCTCYT3' (SEQ ID NO 243)
 5'CGCTGATCCCATTTTCCTCYT3' (SEQ ID NO 244)
 5'GCGCTGATCCCATTTTCCTCYT3' (SEQ ID NO 245)
 5'AGCGCTGATCCCATTTTCCTCYT3' (SEQ ID NO 246)
 5'TAGCGCTGATCCCATTTTCCTCYT3' (SEQ ID NO 247)
 5'CTAGCGCTGATCCCATTTTCCTCYT3' (SEQ ID NO 248)
 5'TCCATTCAAGGGAGGGCGAC3' (SEQ ID NO 249)
 5'CTCCATTCAAGGGAGGGCGAC3' (SEQ ID NO 250)
 5'TCTCCATTCAAGGGAGGGCGAC3' (SEQ ID NO 251)
 5'TTCTCCATTCAAGGGAGGGCGAC3' (SEQ ID NO 252)
 5'ATTCTCCATTCAAGGGAGGGCGAC3' (SEQ ID NO 253)
 5'CATTCTCCATTCAAGGGAGGGCGAC3' (SEQ ID NO 254)

- for the amplification of exon 4 of HLA-B (table 8):

5'AGATTATCCCAGGTGCCTGC3' (SEQ ID NO 255)
 5'GAGATTATCCCAGGTGCCTGC3' (SEQ ID NO 256)
 5'GGAGATTATCCCAGGTGCCTGC3' (SEQ ID NO 257)
 5'AGGAGATTATCCCAGGTGCCTGC3' (SEQ ID NO 258)
 5'TAGGAGATTATCCCAGGTGCCTGC3' (SEQ ID NO 259)
 5'ATAGGAGATTATCCCAGGTGCCTGC3' (SEQ ID NO 260)
 5'TGTCCTGYCCATTCTCAGKC3' (SEQ ID NO 261)
 5'GTGTCCTGYCCATTCTCAGKC3' (SEQ ID NO 262)
 5'GGTGTCTCCTGYCCATTCTCAGKC3' (SEQ ID NO 263)
 5'AGGTGTCTCCTGYCCATTCTCAGKC3' (SEQ ID NO 264)
 5'CAGGTGTCTCCTGYCCATTCTCAGKC3' (SEQ ID NO 265)
 5'CCAGGTGTCTCCTGYCCATTCTCAGKC3' (SEQ ID NO 266)
 5'TCACATGGGTGGTCCTAGG3' (SEQ ID NO 267)
 5'GTCACATGGGTGGTCCTAGG3' (SEQ ID NO 268)
 5'GGTCACATGGGTGGTCCTAGG3' (SEQ ID NO 269)
 5'TGGTCACATGGGTGGTCCTAGG3' (SEQ ID NO 270)
 5'CTGGTCACATGGGTGGTCCTAGG3' (SEQ ID NO 271)

- 5'KCTGGTCACATGGGTGGTCCTAGG3' (SEQ ID NO 272)
 5'GKCTGGTCACATGGGTGGTCCTAGG3' (SEQ ID NO 273)
 5'TSCCATGARAGATGCMAAGC3' (SEQ ID NO 274)
 5'GTSCCATGARAGATGCMAAGC3' (SEQ ID NO 275)
 5 5'TGTSCCATGARAGATGCMAAGC3' (SEQ ID NO 276)
 5'GTGTSCCATGARAGATGCMAAGC3' (SEQ ID NO 277)
 5'GGTGTSCCATGARAGATGCMAAGC3' (SEQ ID NO 278)
 5'GGGTGTSCCATGARAGATGCMAAGC3' (SEQ ID NO 279)
 5'GAWTTTCTGACTCTTCCCA3' (SEQ ID NO 280)
 10 5'TGAWTTTCTGACTCTTCCCA3' (SEQ ID NO 281)
 5'CTGAWTTTCTGACTCTTCCCA3' (SEQ ID NO 282)
 5'CCTGAWTTTCTGACTCTTCCCA3' (SEQ ID NO 283)
 5'GCCTGAWTTTCTGACTCTTCCCA3' (SEQ ID NO 284)
 5'CGCCTGAWTTTCTGACTCTTCCCA3' (SEQ ID NO 285)
 15
- for the amplification of exon 2 of HLA-C (table 9):
 5'GTCGAGGGTCTGGGCGGGTT3' (SEQ ID NO 126)
 5'GGTCGAGGGTCTGGGCGGGTT3' (SEQ ID NO 127)
 5'CGGTCGAGGGTCTGGGCGGGTT3' (SEQ ID NO 128)
 20 5'CCGGTCGAGGGTCTGGGCGGGTT3' (SEQ ID NO 129)
 5'YCCGGTCGAGGGTCTGGGCGGGTT3' (SEQ ID NO 130)
 5'CYCCGGTCGAGGGTCTGGGCGGGTT3' (SEQ ID NO 131)
 - for the amplification of exon 3 of HLA-C (table 10):
 25 5'CGCCCCRAGTCTCCSSGTCT3' (SEQ ID NO 132)
 5'TCGCCCCRAGTCTCCSSGTCT3' (SEQ ID NO 133)
 5'GTCGCCCCRAGTCTCCSSGTCT3' (SEQ ID NO 134)
 5'GGTCGCCCCRAGTCTCCSSGTCT3' (SEQ ID NO 135)
 5'GGGTCGCCCCRAGTCTCCSSGTCT3' (SEQ ID NO 136)
 30 5'CGGGTCGCCCCRAGTCTCCSSGTCT3' (SEQ ID NO 137)
 5'CGRCCGGRGAGAGCCCCAGT3' (SEQ ID NO 138)
 5'TCGRCCGGRGAGAGCCCCAGT3' (SEQ ID NO 139)
 5'CTCGRCCGGRGAGAGCCCCAGT3' (SEQ ID NO 140)
 5'CCTCGRCCGGRGAGAGCCCCAGT3' (SEQ ID NO 141)

5'CCCTCGRCCGGRGAGAGCCCCAGT3' (SEQ ID NO 142)

5'ACCCTCGRCCGGRGAGAGCCCCAGT3' (SEQ ID NO 143)

- for the amplification of exon 4 of HLA-C (table 11):

5 5'GTGCCTGTGTCCAGGCTGGC3' (SEQ ID NO 286)
 5'GGTGCCTGTGTCCAGGCTGGC3' (SEQ ID NO 287)
 5'AGGTGCCTGTGTCCAGGCTGGC3' (SEQ ID NO 288)
 5'CAGGTGCCTGTGTCCAGGCTGGC3' (SEQ ID NO 289)
 5'CCAGGTGCCTGTGTCCAGGCTGGC3' (SEQ ID NO 290)
 10 5'CCCAGGTGCCTGTGTCCAGGCTGGC3' (SEQ ID NO 291)
 5'TGGCGTCTGGGTTCTGTGCC3' (SEQ ID NO 292)
 5'CTGGCGTCTGGGTTCTGTGCC3' (SEQ ID NO 293)
 5'GCTGGCGTCTGGGTTCTGTGCC3' (SEQ ID NO 294)
 5'GGCTGGCGTCTGGGTTCTGTGCC3' (SEQ ID NO 295)
 15 5'AGGCTGGCGTCTGGGTTCTGTGCC3' (SEQ ID NO 296)
 5'CAGGCTGGCGTCTGGGTTCTGTGCC3' (SEQ ID NO 297)
 5'CTCAGGATRGTCACATGGSC3' (SEQ ID NO 298)
 5'TCTCAGGATRGTCACATGGSC3' (SEQ ID NO 299)
 5'TTCTCAGGATRGTCACATGGSC3' (SEQ ID NO 300)
 20 5'RTTCTCAGGATRGTCACATGGSC3' (SEQ ID NO 301)
 5'CRTTCTCAGGATRGTCACATGGSC3' (SEQ ID NO 302)
 5'CCRTTCTCAGGATRGTCACATGGSC3' (SEQ ID NO 303)
 5'GCGCTGTTGGAGTGTCGCAA3' (SEQ ID NO 78)
 5'GGCGCTGTTGGAGTGTCGCAA3' (SEQ ID NO 79)
 25 5'GGGCGCTGTTGGAGTGTCGCAA3' (SEQ ID NO 80)
 5'TGGGCGCTGTTGGAGTGTCGCAA3' (SEQ ID NO 81)
 5'ATGGGCGCTGTTGGAGTGTCGCAA3' (SEQ ID NO 82)
 5'CATGGGCGCTGTTGGAGTGTCGCAA3' (SEQ ID NO 83)
 5'SCAAGAGAGAWRCAAAGTGT3' (SEQ ID NO 304)
 30 5'CSCAAGAGAGAWRCAAAGTGT3' (SEQ ID NO 305)
 5'TCSCAAGAGAGAWRCAAAGTGT3' (SEQ ID NO 306)
 5'GTCSCAAGAGAGAWRCAAAGTGT3' (SEQ ID NO 307)
 5'TGTCSCAAGAGAGAWRCAAAGTGT3' (SEQ ID NO 308)
 5'GTGTCSCAAGAGAGAWRCAAAGTGT3' (SEQ ID NO 309)

In another preferred embodiment, the present invention relates to any method as described above, further characterized that the amplification of exon 2 is carried out with at least one of the following forward primers:

- 5 • for HLA-A: 5APBio: B-TTCTCCCCAGACGCCGAGGATGGCC (SEQ ID NO 144);
- for HLA-B: IBPin1: B-GGGAGGAGCGAGGGGACCSCAG (SEQ ID NO 145);
- 10 • for HLA-C: 5CIN1: B-AGCGAGGGGCCCCGCCCGGCGA (SEQ ID NO 146).

In another preferred embodiment, the present invention relates to any method as described above, further characterized that the amplification of exon 4 is carried out with at least one of the following reverse primers:

- 15 • for HLA-A: 3ex4APbio: B-TTGGGCAGACCCTCATGCTGC (SEQ ID NO 311);
- for HLA-B: 3ex4IBbio: B-TCGGCAGCCCCCTCATGCTGT (SEQ ID NO 312);
- for HLA-C: 3ex4ICbio: B-CATCTCAGGGTGMRRGGGCTT (SEQ ID NO 313).

20 In a very specific embodiment, the present invention relates to any method as described above, further characterized that:

- the amplification of exon 2 is carried out with at least one of the following primers sets:

- 25 • for HLA-A: 5APbio (B-TTCTCCCCAGACGCCGAGGATGGCC; SEQ ID NO 144) and 3ex2APbio (B-ATCTCGGACCCGGAGACTGT; SEQ ID NO 1);
- for HLA-B: IBPin1 (B-GGGAGGAGCGAGGGGACCSCAG; SEQ ID NO 145) and IB3Pin2bio (B-AACCCGCGGGGATTTTGGCCTC; SEQ ID NO 109);
- 30 • for HLA-C: 5CIN1 (B-AGCGAGGGGCCCCGCCCGGCGA; SEQ ID NO 146) and IC3Pin2bio (B-GGTCGAGGGTCTGGGCGGGTT; SEQ ID NO 127);

- the amplification of exon 3 is carried out with at least one of the following primer sets:
 - for HLA-A: 5ex3APbio (B-CAGTTTtagGCCAAAAATCCCCC; SEQ ID NO 104) and 3ex3APbio (B-CCCTCCTTGTGGGAGGCCAG; SEQ ID NO 156);
 - for HLA-B: IB5Pin 2bio (B-CGCGTTTACCCGGTTTCATTTTCAGTTG; SEQ ID NO 224) and IB3Pin3bio (B-TCTTCTCGTKGGAGSCCATCCCC; SEQ ID NO 234);
 - for HLA-C: IC5Pin2bio (B-TCGRCCGGRGAGAGCCCCAGT; SEQ ID NO 139) and 3CIN3 (B-GGAGATGGGGAAGGCTCCCCACT; SEQ ID NO 149);
- the amplification of exon 4 is carried out with at least one of the following primer sets:
 - for HLA-A: 5ex4APbio (B-GTTCTGTGCTCYCTTCCCCAT; SEQ ID NO 205) and 3ex4APbio (B-TTGGGCAGACCCTCATGCTGC ; SEQ ID NO 311);
 - for HLA-B: 5ex4IBbio (B-TCACATGGGTGGTCCTAGG; SEQ ID NO 267) and 3ex4IBbio (B-TCGGCAGCCCTCATGCTGT; SEQ ID NO 312);
 - for HLA-C: 5ex4ICbio (B-TCTCAGGATRGTCACATGGSC; SEQ ID NO 299) and 3ex4ICbio (B-CATCTCAGGGTGMRGGGCTT; SEQ ID NO 313).

The skilled man will understand that these primers (SEQ ID NOs 1 to 314) may be adapted by addition or deletion of one or more nucleotides at their extremities. Such adaptations may be required, for instance, if the conditions of amplification are changed, if the amplified material is RNA instead of DNA, as is the case, for example, in the NASBA system.

The present invention further relates to a primer as described above, for use in the amplification of exon 2 of HLA-A, HLA-B or HLA-C alleles, said primer specifically hybridizing to a locus-specific target sequence in intron 2 of respectively HLA-A, HLA-B or HLA-C; more specifically, said primer specifically hybridizing to a locus-specific target sequence situated at position:

- 67, 96, 109, 110, 118, 123, 131 or 181 of the HLA-A intron 2 (Fig. 1); or
- 35 or 170 of the HLA-B intron 2 (Fig. 2); or
- 84, 107 or 142 of the HLA-C intron 2 (Fig. 3);

more specifically, said primer being chosen from table 1 (for the amplification of
 5 exon 2 of HLA-A), from table 5 (for the amplification of exon 2 of HLA-B) or from
 table 9 (for the amplification of exon 2 of HLA-C).

The present invention further also relates to a primer as described above, for use in the
 amplification of exon 3 of HLA-A, HLA-B or HLA-C alleles, said primer specifically
 hybridizing to a locus-specific target sequence in intron 2 (forward primer) or to a
 10 locus-specific target sequence in intron 3 (reverse primer) of respectively HLA-A,
 HLA-B or HLA-C; more specifically, said primer specifically hybridizing to a locus-
 specific target sequence situated at position:

- 67, 96, 109, 110, 118, 123, 131 or 181 of the HLA-A intron 2 (Fig. 1) or
 32, 50, 62, 73, 83, 86, 118, 130, 150 of the HLA-A intron 3 (Fig. 4); or
- 15 - 35 or 170 of the HLA-B intron 2 (Fig. 2) or 42, 46, 65, 68, 96, of the
 HLA-B intron 3 (Fig. 5); or
- 84, 107 or 142 of the HLA-C intron 2 (Fig. 3).

more specifically, said primer being chosen from table 2 or table 3 (for the
 amplification of exon 3 of HLA-A) or from table 6 or table 7 (for the amplification of
 20 exon 3 of HLA-B).

The present invention further also relates to a primer as described above, for use in the
 amplification of exon 4 of HLA-A, HLA-B or HLA-C alleles, said primer specifically
 hybridizing to a locus-specific target sequence in intron 3 of respectively HLA-A,
 HLA-B or HLA-C; more specifically, said primer specifically hybridizing to a locus-
 25 specific target sequence situated at position:

- 501, 525, 561 or 571 of the HLA-A intron 3 (Fig. 4); or
- 438, 502, 524, 547 or 571 of the HLA-B intron 3 (Fig. 5); or
- 461, 477, 527, 545 or 561 of the HLA-C intron 3 (Fig. 6).

more specifically, said primer being chosen from table 4 (for the amplification of
 30 exon 4 of HLA-A), from table 8 (for the amplification of exon 4 of HLA-B) or from
 table 11 (for the amplification of exon 4 of HLA-C).

The present invention further relates to a primer set consisting of a combination of a
 forward and a reverse primer as defined above, for use in the amplification of exon 2
 of HLA-A, HLA-B or HLA-C alleles. In a specific embodiment, the present invention

relates to the combination of the forward primer 5APBio (SEQ ID NO 144) for HLA-A, IBPin1 (SEQ ID NO 145) for HLA-B or 5CIN1 (SEQ ID NO 146) for HLA-C and a reverse primer specifically hybridizing to a locus-specific target sequence in intron 2 of respectively HLA-A, HLA-B or HLA-C; more specifically, said reverse primer

5 specifically hybridizing to a locus-specific target sequence situated at position:

- 67, 96, 109, 110, 118, 123, 131 or 181 of the HLA-A intron 2; or
- 35 or 170 of the HLA-B intron 2; or
- 84, 107 or 142 of the HLA-C intron 2;

more specifically, said reverse primer being chosen from table 1 (for the amplification of exon 2 of HLA-A), from table 5 (for the amplification of exon 2 of HLA-B) or

10 from table 9 (for the amplification of exon 2 of HLA-C).

The present invention also relates to a primer set consisting of a combination of a forward and a reverse primer as described above, for use in the amplification of exon 3 of HLA-A, HLA-B or HLA-C alleles. In a specific embodiment, the present

15 invention relates to the combination of a forward primer specifically hybridizing to a locus-specific target sequence in intron 2 of respectively HLA-A, HLA-B or HLA-C; more specifically, said forward primer specifically hybridizing to a locus-specific target sequences situated at position:

- 67, 96, 109, 110, 118, 123, 131 or 181 of the HLA-A intron 2; or
- 20 - 35 or 170 of the HLA-B intron 2; or
- 84, 107 or 142 of the HLA-C intron 2;

more specifically, said forward primer being chosen from table 2 (for the amplification of exon 3 of HLA-A), from table 6 (for the amplification of exon 3 of HLA-B) or from table 10 (for the amplification of exon 3 of HLA-C) and a reverse

25 primer specifically hybridizing to a locus-specific target sequence in intron 3 of respectively HLA-A, HLA-B or HLA-C; more specifically, said reverse primer specifically hybridizing to a locus-specific target sequences situated at position:

- 32, 50, 62, 73, 83, 86, 118, 130, 150 (Fig. 4) of the HLA-A intron 3; or
- 42, 46, 65, 68, 96, of the HLA-B intron 3 (Fig. 5);

30 more specifically, said reverse primer being chosen from table 3 (for the amplification of exon 3 of HLA-A), from table 7 (for the amplification of exon 3 of HLA-B) or from 3CIN3 (SEQ ID NO 149) (for the amplification of exon 3 of HLA-C).

The present invention also relates to a primer set consisting of a combination of a forward and a reverse primer as described above, for use in the amplification of exon

4 of HLA-A, HLA-B or HLA-C alleles. In a specific embodiment, the present invention relates to the combination of a forward primer specifically hybridizing to a locus-specific target sequence in intron 3 of respectively HLA-A, HLA-B or HLA-C; more specifically, said forward primer specifically hybridizing to a locus-specific target sequences situated at position:

- 501, 525, 561 or 571 of the HLA-A intron 3 (Fig. 4); or
- 438, 502, 524, 547 or 571 of the HLA-B intron 3 (Fig. 5); or
- 461, 477, 527, 545 or 561 of the HLA-C intron 3 (Fig. 6).

more specifically, said forward primer being chosen from table 4 (for the amplification of exon 4 of HLA-A), from table 8 (for the amplification of exon 4 of HLA-B) or from table 11 (for the amplification of exon 4 of HLA-C) and a reverse primer 3ex4APbio (SEQ ID NO 311) for HLA-A, 3ex4IBbio (SEQ ID NO 312) for HLA-B or 3ex4ICbio (SEQ ID NO 313) for HLA-C.

In a specific embodiment, the primers are used in a mix that allows the separate amplification of exon 2 and exon 3. Accordingly, the present invention relates to a multiplex primer mix containing at least one primer pair as described above for the amplification of exon 2 and one primer pair as described above for the amplification of exon 3.

In a specific embodiment, the multiplex primer mix for the separate amplification of exon 2 and exon 3 comprises the following primer sets:

- for the separate amplification of exon 2 and exon 3 of HLA-A:

Amplicon	Primerset	Sequence (5'-3')	SEQ ID
Exon 1-2	5APbio (5')	B-TTCTCCCCAGACGCCGAGGATGGCC	144
	3ex2APbio (3')	B-ATCTCGGACCCGGAGACTGT	1
Exon 3	5ex3APbio (5')	B-CAGTTTAGGCCAAAAATCCCCC	104
	3ex3APbio (3')	B-CCCTCCTTGTGGGAGGCCAG	156

- for the separate amplification of exon 2 and exon 3 of HLA-B:

Amplicon	Primer set	Sequence (5'-3')	SEQ ID
Exon 2	IBPin1bio (5')	B-GGGAGGAGCGAGGGGACCSCAG	145
	IB3Pin2bio (3')	B-AACCCGCGGGGATTTTGGCCTC	109
Exon 3	IB5Pin2bio (5')	B-CGCGTTTACCCGGTTTCATTTTCAGTTG	224
	IB3Pin3bio (3')	B-TCTTCTCGTKGGAGSCCATCCCC	234

- for the separate amplification of exon 2 and exon 3 of HLA-C:

5

Amplicon	Primer set	Sequence (5'-3')	SEQ ID
Exon 2	5CIN1(5')	B-AGCGAGGGGCCCCGCGGCGA	146
	IC3Pin2bio(3')	B-GGTCGAGGGTCTGGGCGGGTT	127
Exon 3	IC5Pin2bio(5')	B-TCGRCCGGRGAGAGCCCCAGT	139
	3CIN3 (3')	B-GGAGATGGGGAAGGCTCCCCACT	149

In a specific embodiment, the primers are used in a mix that allows the separate amplification of exon 2, exon 3 and exon 4. Accordingly, the present invention relates to a multiplex primer mix containing at least one primer pair as described above for amplification of exon 2, one primer pair as described above for the amplification of exon 3 and one primer pair as described above for the amplification of exon 4.

In a specific embodiment, the multiplex primer mix for the separate amplification of exon 2, exon 3 and exon 4 comprises the following primer sets:

15 - for the separate amplification of exon 2, exon 3 and exon 4 of HLA-A:

Amplicon	Primer set	Sequence (5'-3')	SEQ ID
Exon 1-2	5APbio (5')	B-TTCTCCCCAGACGCCGAGGATGGCC	144
	3ex2APbio (3')	B-ATCTCGGACCCGGAGACTGT	1
Exon 3	5ex3APbio (5')	B-CAGTTTAGGCCAAAAATCCCCC	104
	3ex3APbio (3')	B-CCCTCCTTGTGGGAGGCCAG	156
Exon 4	5ex4APbio (5')	B-GTTCTGTGCTCYCTTCCCCAT	205
	3ex4APbio (3')	B-TTGGGCAGACCCTCATGCTGC	311

- for the separate amplification of exon 2, exon 3 and exon 4 of HLA-B:

Amplicon	Primerset	Sequence (5'-3')	SEQ ID
Exon 2	IBPin1bio (5')	B-GGGAGGAGCGAGGGGACCSCAG	145
	IB3Pin2bio (3')	B-AACCCGCGGGGATTTTGGCCTC	109
Exon 3	IB5Pin2bio (5')	B-CGCGTTTACCCGGTTTCATTTTCAGTTG	224
	IB3Pin3bio (3')	B-TCTTCTCGTKGGAGSCCATCCCC	234
Exon 4	5ex4IBbio (5')	B-TCACATGGGTGGTCCTAGG	267
	3ex4IBbio (3')	B-TCGGCAGCCCCCTCATGCTGT	312

- for the separate amplification of exon 2, exon 3 and exon 4 of HLA-C:

5

Amplicon	Primerset	Sequence (5'-3')	SEQ ID
Exon 2	5CIN1(5')	B-AGCGAGGGGCCCCGCCCCGGCGA	146
	IC3Pin2bio(3')	B-GGTCGAGGGTCTGGGCGGGTT	127
Exon 3	IC5Pin2bio(5')	B-TCGRCCGGRGAGAGCCCCAGT	139
	3CIN3 (3')	B-GGAGATGGGGAAGGCTCCCCACT	149
Exon 4	5ex4ICbio(5')	B-TCTCAGGATRGTCACATGGSC	299
	3ex4ICbio(3')	B-CATCTCAGGGTGMRGGGCTT	313

The primers of the invention may be labeled. Labeling may be carried out by any method known to the person skilled in the art. The nature of the label may be isotopic (^{32}P , ^{35}S , etc.) or non-isotopic (biotin, digoxigenin, etc.).

- 10 The oligonucleotides used as primers may also comprise nucleotide analogues such as phosphorothiates (Matsukura et al., 1987), alkylphosphorothiates (Miller et al., 1979) or peptide nucleic acids (Nielsen et al., 1991; Nielsen et al., 1993) or may contain intercalating agents (Asseline et al., 1984). As most other variations or modifications introduced into the original DNA sequences of the invention, these variations will
- 15 necessitate adaptations with respect to the conditions under which the oligonucleotide should be used to obtain the required specificity and sensitivity. However, the eventual results of hybridization will be essentially the same as those obtained with the unmodified oligonucleotides. The introduction of these modifications may be advantageous in order to positively influence characteristics such as hybridization

kinetics, reversibility of the hybrid-formation, biological stability of the oligonucleotide molecules, etc.

The present invention also relates to the use of the primers, the primer sets and/or the primer mixes of the invention in a method for the locus-specific and separate
5 amplification of exon 2, exon 3 and/or exon 4 of HLA-A, HLA-B or HLA-C.

The present invention also relates to a method for typing or subtyping of one or more HLA-A, HLA-B or HLA-C alleles in a sample comprising the following steps:

- (i) if needed, release, isolation and/or concentration of the nucleic acids present in said sample;
- 10 (ii) amplification of the nucleic acids according to the invention;
- (iii) typing of the specific HLA-A, HLA-B or HLA-C alleles present in said sample.

Release, concentration and isolation of the nucleic acids from the sample can be done by any method known in the art. Currently, various commercial kits are available such
15 as the QIAamp Blood Kit from Qiagen (Hilden, Germany) for the isolation of nucleic acids from blood samples or the 'High pure PCR Template Preparation Kit' (Roche Diagnostics, Brussels, Belgium). Other well-known procedures for the isolation of DNA or RNA from a biological sample are also available (Sambrook et al., 1989).

The nucleic acids are subsequently amplified by the method of the invention
20 described above. The products of this amplification step are then ideally suited for typing of the specific allele present in the sample. Currently 169 different alleles of HLA-A, 332 different alleles of HLA-B and 87 different alleles of HLA-C are known. Typing of these alleles can be done by any method known in the art, such as duplex analysis of the PCR products (Clay et al., 1994), single-stranded conformational
25 polymorphism analysis of the PCR product (PCR-SSCP; Yoshida *et al.*, 1992), sequence-based typing (SBT; Santamaria et al., 1992 and 1993), the use of sequence specific primers in PCR reaction (PCR-SSP; Olerup and Zetterquist, 1991), the use of PCR in combination with sequence-specific oligonucleotide probing (PCR-SSOP; Saiki et al., 1986), conventional dot-blot, Southern blot, sandwich or probing by
30 reverse dot-blot (Saiki et al., 1989). In order to obtain fast and easy results if a multitude of probes is involved, a reverse hybridization format may be convenient. Accordingly, in a preferred embodiment the selected probes are immobilized to certain locations on a solid support and the amplified polynucleic acids are labeled in order to enable the detection of the hybrids formed. The term "solid support" can refer

- to any substrate to which an oligonucleotide probe can be coupled, provided that it retains its hybridization characteristics and provided that the background level of hybridization remains low. Usually the solid substrate will be a microtiter plate (e.g. in the DEIA technique), a membrane (e.g. nylon or nitrocellulose) or a microsphere (bead) or a chip. Prior to application to the membrane or fixation, it may be convenient to modify the nucleic acid probe in order to facilitate fixation or improve the hybridization efficiency. Such modifications may encompass homopolymer tailing, coupling with different reactive groups such as aliphatic groups, NH₂ groups, SH groups, carboxylic groups, or coupling with biotin, haptens or proteins.
- 10 The present invention further relates to a diagnostic kit for the typing or subtyping of one or more HLA-A, HLA-B or HLA-C alleles in a sample comprising the following components:
- (i) when appropriate, a means for releasing, isolating or concentrating the nucleic acids present in said sample;
 - 15 (ii) a primer set or a primer mix according to the invention;
 - (iv) a means for typing of the specific HLA-A, HLA-B or HLA-C alleles present in said sample.

- A specific and very user-friendly diagnostic kit is the a Line Probe Assay for the typing or subtyping of one or more HLA-A, HLA-B or HLA-C alleles in a sample comprising the following components:
- (i) when appropriate, a means for releasing, isolating or concentrating the nucleic acids present in said sample;
 - (ii) a primer pair or a primer mix according to the invention;
 - (iii) at least one probe that specifically hybridizes with one of the HLA-A, HLA-B or HLA-C alleles, fixed to a solid support;
 - 25 (iv) a hybridization buffer, or components necessary for producing said buffer;
 - (v) a wash solution, or components necessary for producing said solution;
 - (vi) when appropriate, a means for detecting the hybrids resulting from the preceding hybridization.

- 30 In this embodiment, the selected set of probes is immobilized to a membrane strip in a line fashion. Said probes may be immobilized individually or as mixtures to the delineated locations. The amplified HLA-A, HLA-B or HLA-C polynucleic acids can be labelled with biotine, and the hybrid can then, via a biotine-streptavidine coupling, be detected with a non-radioactive colour developing system.

The term "hybridization buffer" means a buffer allowing a hybridization reaction between the probes and the polynucleic acids present in the sample, or the amplified products, under the appropriate stringency conditions.

- 5 The term "wash solution" means a solution enabling washing of the hybrids formed under the appropriate stringency conditions.

The present invention also relates to the use of the primers, the primer sets and/or the primer mixes of the invention for the manufacturing of a diagnostic kit or Line Probe Assay for HLA Class I typing.

- 10 Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of stated integers or steps but not to the exclusion of any other integer or step or group of integers or steps.

FIGURE LEGENDS

Figure 1. Alignment of 29 HLA-A intron 2 sequences. The consensus sequence is shown on top of the figure. Nucleotides conform to the consensus sequence are indicated with a vertical line. Nucleotides that differ from the consensus sequence are indicated. N indicates the absence of a nucleotide at that position. Sequencing was carried out according to example 1.

Figure 2. Alignment of 38 HLA-B intron 2 sequences. The consensus sequence is shown on top of the figure. Nucleotides conform to the consensus sequence are indicated with a vertical line. Nucleotides that differ from the consensus sequence are indicated. N indicates the absence of a nucleotide at that position. Sequencing was carried out according to example 1.

Figure 3. Alignment of 13 HLA-C intron 2 sequences. The consensus sequence is shown on top of the figure. Nucleotides conform to the consensus sequence are indicated with a vertical line. Nucleotides that differ from the consensus sequence are indicated. N indicates the absence of a nucleotide at that position. Sequencing was carried out according to example 1.

Figure 4. Alignment of 12 HLA-A intron 3 sequences. The consensus sequence is shown on top of the figure. Nucleotides conform to the consensus sequence are indicated with a vertical line. Nucleotides that differ from the consensus sequence are indicated. N indicates the absence of a nucleotide at that position.

Figure 5. Alignment of 22 HLA-B intron 3 sequences. The consensus sequence is shown on top of the figure. Nucleotides conform to the consensus sequence are indicated with a vertical line. Nucleotides that differ from the consensus sequence are indicated. N indicates the absence of a nucleotide at that position.

Figure 6. Alignment of 12 HLA-C intron 3 sequences. The consensus sequence is shown on top of the figure. Nucleotides conform to the consensus sequence are indicated with a vertical line. Nucleotides that differ from the consensus sequence are indicated. N indicates the absence of a nucleotide at that position.

Figure 7. Results of a Line Probe Assay for the typing of HLA-A as indicated in example 5. Nucleic acids were amplified by use of: (A) 5APBio (forward primer; SEQ ID NO 144) and 3APBio (reverse primer; SEQ ID NO 147) for the amplification of exon 2 and exon 3 of HLA-A in one single amplicon; (B) the multiplex primer mix for the separate amplification of exon 2, exon 3 and exon 4 of HLA-A. The probes with numbers 1, 4, 9, 10, 12 and 13 clearly show a stronger signal after the hybridization with the amplicons obtained by use of the multiplex primer mix (strip B).

10

Figure 8. Results of a Line Probe Assay for the typing of HLA-B as indicated in example 6. Nucleic acids were amplified by use of: (B) IBPin1 (forward primer; SEQ ID NO 145) and IBPin3 (reverse primer; SEQ ID NO 148) for the amplification of exon 2 and exon 3 of HLA-B in one single amplicon; (B) the multiplex primer mix for the separate amplification of exon 2, exon 3 and exon 4 of HLA-B. The probes with numbers 9, 10, 18, 19, 20, 34 and 35 clearly show a stronger signal after the hybridization with the amplicons obtained by use of the multiplex primer mix (strips B).

15

TABLES

Table 1. Reverse primers used for the amplification of HLA-A exon 2.

3' position in intron 2	Sequence (5'-3')	SEQ ID NO
67	ATCTCGGACCCGGAGACTGT	1
	GATCTCGGACCCGGAGACTGT	2
	GGATCTCGGACCCGGAGACTGT	3
	YGGATCTCGGACCCGGAGACTGT	4
	GYGGATCTCGGACCCGGAGACTGT	5
	GGYGGATCTCGGACCCGGAGACTGT	6
96	GGTCTCGGRGTCCCGCGGCT	7
	GGGTCTCGGRGTCCCGCGGCT	8
	AGGGTCTCGGRGTCCCGCGGCT	9
	AAGGGTCTCGGRGTCCCGCGGCT	10
	CAAGGGTCTCGGRGTCCCGCGGCT	11
109	CTCCCGGGDCAAGGGTCTCG	12
	TCTCCCGGGDCAAGGGTCTCG	13
	CTCTCCCGGGDCAAGGGTCTCG	14
	CCTCTCCCGGGDCAAGGGTCTCG	15
	GCCTCTCCCGGGDCAAGGGTCTCG	16
	GGCCTCTCCCGGGDCAAGGGTCTCG	17
110	TCTCCCGGGDCAAGGGTCTC	18
	CTCTCCCGGGDCAAGGGTCTC	19
	CCTCTCCCGGGDCAAGGGTCTC	20
	GCCTCTCCCGGGDCAAGGGTCTC	21
	GGCCTCTCCCGGGDCAAGGGTCTC	22
	GGGCCTCTCCCGGGDCAAGGGTCTC	23
118	CCTGGGCCTCTCCCGGGDCA	30
	GCCTGGGCCTCTCCCGGGDCA	31
	CGCCTGGGCCTCTCCCGGGDCA	32
	GCGCCTGGGCCTCTCCCGGGDCA	33

	GGCGCCTGGGCCTCTCCCGGGDCA	34
	AGGCGCCTGGGCCTCTCCCGGGDCA	35
123	AGGCGCCTGGGCCTCTCCCG	36
	AAGGCGCCTGGGCCTCTCCCG	37
	WAAGGCGCCTGGGCCTCTCCCG	38
	TWAAGGCGCCTGGGCCTCTCCCG	39
	GTWAAGGCGCCTGGGCCTCTCCCG	40
	GGTWAAGGCGCCTGGGCCTCTCCCG	41
131	CCGGGTWAAGGCGCCTGGGC	42
	ACCGGGTWAAGGCGCCTGGGC	43
	AACCGGGTWAAGGCGCCTGGGC	44
	AAACCGGGTWAAGGCGCCTGGGC	45
	GAAACCGGGTWAAGGCGCCTGGGC	46
	TGAAACCGGGTWAAGGCGCCTGGGC	47
181	YCCVGCCCCGACCAACCYGG	48
	GYCCVGCCCCGACCAACCYGG	49
	YGYCCVGCCCCGACCAACCYGG	50
	CYGYCCVGCCCCGACCAACCYGG	51
	CCYGYCCVGCCCCGACCAACCYGG	52
	CCCYGYCCVGCCCCGACCAACCYGG	53

Table 2. Forward primers used for the amplification of HLA-A exon 3.

3' position in intron 2	Sequence (5'-3')	SEQ ID NO
67	CGGACGGGCCRGGTSRCCCA	54
	ACGGACGGGCCRGGTSRCCCA	55
	CACGGACGGGCCRGGTSRCCCA	56
	CCACGGACGGGCCRGGTSRCCCA	57
	CCCACGGACGGGCCRGGTSRCCCA	58
	CCCCACGGACGGGCCRGGTSRCCCA	59
96	GGTCCGAGATCCRCCCCGAA	60
	GGGTCCGAGATCCRCCCCGAA	61
	CGGGTCCGAGATCCRCCCCGAA	62
	CCGGGTCCGAGATCCRCCCCGAA	63
	TCCGGGTCCGAGATCCRCCCCGAA	64
	CTCCGGGTCCGAGATCCRCCCCGAA	65
109	CCCCGAAGCCGCGGGACYCC	66
	RCCCCGAAGCCGCGGGACYCC	67
	CRCCCCGAAGCCGCGGGACYCC	68
	CCRCCCCGAAGCCGCGGGACYCC	69
	TCCRCCCCGAAGCCGCGGGACYCC	70
	ATCCRCCCCGAAGCCGCGGGACYCC	71
110	CCCGAAGCCGCGGGACYCCG	72
	CCCCGAAGCCGCGGGACYCCG	73
	RCCCCGAAGCCGCGGGACYCCG	74
	CRCCCCGAAGCCGCGGGACYCCG	75
	CCRCCCCGAAGCCGCGGGACYCCG	76
	TCCRCCCCGAAGCCGCGGGACYCCG	77
118	CGCGGGACYCCGAGACCCTT	84
	CCGCGGGACYCCGAGACCCTT	85
	GCCGCGGGACYCCGAGACCCTT	86
	AGCCGCGGGACYCCGAGACCCTT	87
	AAGCCGCGGGACYCCGAGACCCTT	88

	GAAGCCGCGGGACYCCGAGACCCTT	89
123	GACYCCGAGACCCTTGDCCC	90
	GGACYCCGAGACCCTTGDCCC	91
	GGGACYCCGAGACCCTTGDCCC	92
	CGGGACYCCGAGACCCTTGDCCC	93
	GCGGGACYCCGAGACCCTTGDCCC	94
	CGCGGGACYCCGAGACCCTTGDCCC	95
131	GACCCTTGDCCCGGGAGAGG	96
	AGACCCTTGDCCCGGGAGAGG	97
	GAGACCCTTGDCCCGGGAGAGG	98
	CGAGACCCTTGDCCCGGGAGAGG	99
	CCGAGACCCTTGDCCCGGGAGAGG	100
	YCCGAGACCCTTGDCCCGGGAGAGG	101
181	GTTTAGGCCAAAAATCCCCC	102
	AGTTTAGGCCAAAAATCCCCC	103
	CAGTTTAGGCCAAAAATCCCCC	104
	TCAGTTTAGGCCAAAAATCCCCC	105
	TTCAGTTTAGGCCAAAAATCCCCC	106
	TTTCAGTTTAGGCCAAAAATCCCCC	107

Table 3. Reverse primers used for the amplification of HLA-A exon 3.

3' position in intron 3	Sequence (5'-3')	SEQ ID NO
32	AGCCCGGGAGATCTAYAGGC	150
	CAGCCCGGGAGATCTAYAGGC	151
	CCAGCCCGGGAGATCTAYAGGC	152
	GCCAGCCCGGGAGATCTAYAGGC	153
	GGCCAGCCCGGGAGATCTAYAGGC	154
	AGGCCAGCCCGGGAGATCTAYAGGC	155
50	CCCTCCTTGTGGGAGGCCAG	156
	CCCCTCCTTGTGGGAGGCCAG	157
	TCCCCTCCTTGTGGGAGGCCAG	158
	CTCCCCTCCTTGTGGGAGGCCAG	159
	TCTCCCCTCCTTGTGGGAGGCCAG	160
	GTCTCCCCTCCTTGTGGGAGGCCAG	161
62	CCCAAWTGTCTCCCCTCCTT	162
	TCCCAAWTGTCTCCCCTCCTT	163
	GTCCCAAWTGTCTCCCCTCCTT	164
	GGTCCCAAWTGTCTCCCCTCCTT	165
	TGGTCCCAAWTGTCTCCCCTCCTT	166
	TTGGTCCCAAWTGTCTCCCCTCCTT	167
73	CTAGTGTTGGTCCCAAWTGT	168
	TCTAGTGTTGGTCCCAAWTGT	169
	TTCTAGTGTTGGTCCCAAWTGT	170
	ATTCTAGTGTTGGTCCCAAWTGT	171
	TATTCTAGTGTTGGTCCCAAWTGT	172
	ATATTCTAGTGTTGGTCCCAAWTGT	173
83	GGGYGATATTCTAGTGTTGG	174
	AGGGYGATATTCTAGTGTTGG	175
	GAGGGYGATATTCTAGTGTTGG	176
	GGAGGGYGATATTCTAGTGTTGG	177
	GGGAGGGYGATATTCTAGTGTTGG	178

	AGGGAGGGYGATATTCTAGTGTTGG	179
86	GGAGGGYGATATTCTAGTGT	180
	GGGAGGGYGATATTCTAGTGT	181
	AGGGAGGGYGATATTCTAGTGT	182
	GAGGGAGGGYGATATTCTAGTGT	183
	AGAGGGAGGGYGATATTCTAGTGT	184
	CAGAGGGAGGGYGATATTCTAGTGT	185
118	CCCAGGAGGAKTCCTCTCCC	186
	ACCCAGGAGGAKTCCTCTCCC	187
	AACCCAGGAGGAKTCCTCTCCC	188
	AAACCCAGGAGGAKTCCTCTCCC	189
	GAAACCCAGGAGGAKTCCTCTCCC	190
	GGAAACCCAGGAGGAKTCCTCTCCC	191
130	AGGATCTGGAAACCCAGGAG	192
	CAGGATCTGGAAACCCAGGAG	193
	ACAGGATCTGGAAACCCAGGAG	194
	TACAGGATCTGGAAACCCAGGAG	195
	GTACAGGATCTGGAAACCCAGGAG	196
	GGTACAGGATCTGGAAACCCAGGAG	197
150	TCAGAGTCACTCTCTGGTAC	198
	CTCAGAGTCACTCTCTGGTAC	199
	CCTCAGAGTCACTCTCTGGTAC	200
	ACCTCAGAGTCACTCTCTGGTAC	201
	AACCTCAGAGTCACTCTCTGGTAC	202
	GAACCTCAGAGTCACTCTCTGGTAC	203

Table 4. Forward primers used for the amplification of HLA-A exon 4.

3' position in intron 3	Sequence (5'-3')	SEQ ID NO
501	TTCTGTGCTCYCTTCCCCAT	204
	GTTCTGTGCTCYCTTCCCCAT	205
	GGTTCTGTGCTCYCTTCCCCAT	206
	GGGTTCTGTGCTCYCTTCCCCAT	207
	TGGGTTCTGTGCTCYCTTCCCCAT	208
	CTGGGTTCTGTGCTCYCTTCCCCAT	209
525	GGTGTCTGTCCATTCTCAA	24
	RGGTGTCTGTCCATTCTCAA	25
	CRGGTGTCTGTCCATTCTCAA	26
	CCRGGTGTCTGTCCATTCTCAA	27
	CCCRGGTGTCTGTCCATTCTCAA	28
	TCCCRGGTGTCTGTCCATTCTCAA	29
561	CTGGWGGAGTGTCCCATKAC	201
	GCTGGWGGAGTGTCCCATKAC	211
	TGCTGGWGGAGTGTCCCATKAC	212
	RTGCTGGWGGAGTGTCCCATKAC	213
	YRTGCTGGWGGAGTGTCCCATKAC	214
	GYRTGCTGGWGGAGTGTCCCATKAC	215
571	GTCCCATKACAGATRCMMAA	216
	TGTCCCATKACAGATRCMMAA	217
	GTGTCCCATKACAGATRCMMAA	218
	AGTGTCCCATKACAGATRCMMAA	219
	GAGTGTCCCATKACAGATRCMMAA	220
	GGAGTGTCCCATKACAGATRCMMAA	221

Table 5. Reverse primers used for the amplification of HLA-B exon 2.

3' position in intron 2	Sequence (5'-3')	SEQ ID NO
170	ACCCGCGGGGATTTTGGCCTC	108
	AACCCGCGGGGATTTTGGCCTC	109
	CAACCCGCGGGGATTTTGGCCTC	110
	CCAACCCGCGGGGATTTTGGCCTC	111
	MCCAACCCGCGGGGATTTTGGCCTC	112
	GMCCAACCCGCGGGGATTTTGGCCTC	113
	YGMCCAACCCGCGGGGATTTTGGCCTC	314

5 Table 6. Forward primers used for the amplification of HLA-B exon 3.

3' position in intron 2	Sequence (5'-3')	SEQ ID NO
35	CYGGGGCGSAGGTCACGACT	114
	CCYGGGGCGSAGGTCACGACT	115
	GCCYGGGGCGSAGGTCACGACT	116
	GGCCYGGGGCGSAGGTCACGACT	117
	CGGCCYGGGGCGSAGGTCACGACT	118
	CCGGCCYGGGGCGSAGGTCACGACT	119
170	CCCGGTTTCATTTTCAGTTG	120
	ACCCGGTTTCATTTTCAGTTG	121
	TACCCGGTTTCATTTTCAGTTG	122
	TTACCCGGTTTCATTTTCAGTTG	123
	TTTACCCGGTTTCATTTTCAGTTG	124
	GTTTACCCGGTTTCATTTTCAGTTG	125
	CGTTTACCCGGTTTCATTTTCAGTTG	222
	GCGTTTACCCGGTTTCATTTTCAGTTG	223
	CGCGTTTACCCGGTTTCATTTTCAGTTG	224

Table 7. Reverse primers used for the amplification of HLA-B exon 3.

3' position in intron 3	Sequence (5'-3')	SEQ ID NO
42	CGTKGGAGSCCATCCCCGSC	225
	TCGTKGGAGSCCATCCCCGSC	226
	CTCGTKGGAGSCCATCCCCGSC	227
	TCTCGTKGGAGSCCATCCCCGSC	228
	TTCTCGTKGGAGSCCATCCCCGSC	229
	CTTCTCGTKGGAGSCCATCCCCGSC	230
46	TCTCGTKGGAGSCCATCCCC	231
	TTCTCGTKGGAGSCCATCCCC	232
	CTTCTCGTKGGAGSCCATCCCC	233
	TCTTCTCGTKGGAGSCCATCCCC	234
	YTCTTCTCGTKGGAGSCCATCCCC	235
	CYTCTTCTCGTKGGAGSCCATCCCC	236
65	GATCCCATTTTCCTCYTCTT	237
	TGATCCCATTTTCCTCYTCTT	238
	CTGATCCCATTTTCCTCYTCTT	239
	GCTGATCCCATTTTCCTCYTCTT	240
	CGCTGATCCCATTTTCCTCYTCTT	241
	GCGCTGATCCCATTTTCCTCYTCTT	242
68	GCTGATCCCATTTTCCTCYT	243
	CGCTGATCCCATTTTCCTCYT	244
	GCGCTGATCCCATTTTCCTCYT	245
	AGCGCTGATCCCATTTTCCTCYT	246
	TAGCGCTGATCCCATTTTCCTCYT	247
	CTAGCGCTGATCCCATTTTCCTCYT	248
96	TCCATTCAAGGGAGGGCGAC	249
	CTCCATTCAAGGGAGGGCGAC	250
	TCTCCATTCAAGGGAGGGCGAC	251
	TTCTCCATTCAAGGGAGGGCGAC	252
	ATTCTCCATTCAAGGGAGGGCGAC	253

	CATTCTCCATTCAAGGGAGGGCGAC	254
--	---------------------------	-----

Table 8. Forward primers used for the amplification of HLA-B exon 4.

3' position in intron 3	Sequence (5'-3')	SEQ ID NO
438	AGATTATCCCAGGTGCCTGC	255
	GAGATTATCCCAGGTGCCTGC	256
	GGAGATTATCCCAGGTGCCTGC	257
	AGGAGATTATCCCAGGTGCCTGC	258
	TAGGAGATTATCCCAGGTGCCTGC	259
	ATAGGAGATTATCCCAGGTGCCTGC	260
502	TGTCCTGYCCATTCTCAGKC	261
	GTGTCCTGYCCATTCTCAGKC	262
	GGTGTCTGYCCATTCTCAGKC	263
	AGGTGTCTGYCCATTCTCAGKC	264
	CAGGTGTCTGYCCATTCTCAGKC	265
	CCAGGTGTCTGYCCATTCTCAGKC	266
524	TCACATGGGTGGTCCTAGG	267
	GTCACATGGGTGGTCCTAGG	268
	GGTCACATGGGTGGTCCTAGG	269
	TGGTCACATGGGTGGTCCTAGG	270
	CTGGTCACATGGGTGGTCCTAGG	271
	KCTGGTCACATGGGTGGTCCTAGG	272
	GKCTGGTCACATGGGTGGTCCTAGG	273
547	TSCCATGARAGATGCMAAGC	274
	GTSCCATGARAGATGCMAAGC	275
	TGTSCCATGARAGATGCMAAGC	276
	GTGTSCCATGARAGATGCMAAGC	277
	GGTGTSCCATGARAGATGCMAAGC	278
	GGGTGTSCCATGARAGATGCMAAGC	279
571	GWAWTTTCTGACTCTTCCCA	280
	TGWAWTTTCTGACTCTTCCCA	281

	CTGWA WTTTCTGACTCTTCCCA	282
	CCTGWA WTTTCTGACTCTTCCCA	283
	GCCTGWA WTTTCTGACTCTTCCCA	284
	CGCCTGWA WTTTCTGACTCTTCCCA	285

Table 9. Reverse primers used for the amplification of HLA-C exon 2.

3' position in intron 2	Sequence (5'-3')	SEQ ID NO
107	GTCGAGGGTCTGGGCGGGTT	126
	GGTCGAGGGTCTGGGCGGGTT	127
	CGGTCGAGGGTCTGGGCGGGTT	128
	CCGGTCGAGGGTCTGGGCGGGTT	129
	YCCGGTCGAGGGTCTGGGCGGGTT	130
	CYCCGGTCGAGGGTCTGGGCGGGTT	131

Table 10. Forward primers used for the amplification of HLA-C exon 3.

3' position in intron 2	Sequence (5'-3')	SEQ ID NO
84	CGCCCCRAGTCTCCSSGTCT	132
	TCGCCCCRAGTCTCCSSGTCT	133
	GTCGCCCCRAGTCTCCSSGTCT	134
	GGTCGCCCCRAGTCTCCSSGTCT	135
	GGGTCGCCCCRAGTCTCCSSGTCT	136
	CGGGTCGCCCCRAGTCTCCSSGTCT	137
142	CGRCCGGRGAGAGCCCCAGT	138
	TCGRCCGGRGAGAGCCCCAGT	139
	CTCGRCCGGRGAGAGCCCCAGT	140
	CCTCGRCCGGRGAGAGCCCCAGT	141
	CCCTCGRCCGGRGAGAGCCCCAGT	142
	ACCCTCGRCCGGRGAGAGCCCCAGT	143

5 Table 11. Forward primers used for the amplification of HLA-C exon 4.

3' position in intron 3	Sequence (5'-3')	SEQ ID NO
461	GTGCCTGTGTCCAGGCTGGC	286
	GGTGCCTGTGTCCAGGCTGGC	287
	AGGTGCCTGTGTCCAGGCTGGC	288
	CAGGTGCCTGTGTCCAGGCTGGC	289
	CCAGGTGCCTGTGTCCAGGCTGGC	290
	CCCAGGTGCCTGTGTCCAGGCTGGC	291
477	TGGCGTCTGGGTTCTGTGCC	292
	CTGGCGTCTGGGTTCTGTGCC	293
	GCTGGCGTCTGGGTTCTGTGCC	294
	GGCTGGCGTCTGGGTTCTGTGCC	295
	AGGCTGGCGTCTGGGTTCTGTGCC	296

	CAGGCTGGCGTCTGGGTTCTGTGCC	297
527	CTCAGGATRGTCACATGGSC	298
	TCTCAGGATRGTCACATGGSC	299
	TTCTCAGGATRGTCACATGGSC	300
	RTTCTCAGGATRGTCACATGGSC	301
	CRTTCTCAGGATRGTCACATGGSC	302
	CCRTTCTCAGGATRGTCACATGGSC	303
545	GCGCTGTTGGAGTGTGCGAA	78
	GGCGCTGTTGGAGTGTGCGAA	79
	GGGCGCTGTTGGAGTGTGCGAA	80
	TGGGCGCTGTTGGAGTGTGCGAA	81
	ATGGGCGCTGTTGGAGTGTGCGAA	82
	CATGGGCGCTGTTGGAGTGTGCGAA	83
561	SCAAGAGAGAWRCAAAGTGT	304
	CSCAAGAGAGAWRCAAAGTGT	305
	TCSCAAGAGAGAWRCAAAGTGT	306
	GTCSCAAGAGAGAWRCAAAGTGT	307
	TGTCSCAAGAGAGAWRCAAAGTGT	308
	GTGTCSCAAGAGAGAWRCAAAGTGT	309

EXAMPLES**Example 1: Sequence determination of intron 2 of various HLA-A, HLA-B and HLA-C alleles**

5

Nucleic acids were prepared from different blood samples by use of the QIAamp Blood Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol.

Part of exon 1, intron 2, exon 2, intron 2 and exon 3 of HLA-A were amplified by use of the following primer set:

10

Primer	Sequence (5'-3')	SEQ ID NO
5APBio (5')	B-TTCTCCCCAGACGCCGAGGATGGCC	144
3APBio (3')	B-CCGTGCGCTGCAGCGTCTCCTTCCCG	147

B = biotine.

Exon 2, intron 2 and exon 3 of HLA-B were amplified by use of the following primer set:

15

Primer	Sequence (5'-3')	SEQ ID NO
IBPin1 (5')	B-GGGAGGAGCGAGGGGACCSCAG	145
IBPin3 (3')	B-GGAGGCCATCCCCGGCGACCTAT	148

B = biotine.

Exon 2, intron 2 and exon 3 of HLA-C were amplified by use of the following primer set:

20

Primer	Sequence (5'-3')	SEQ ID NO
5CIN1 (5')	B-AGCGAGGGGCCCCGGCGA	146
3CIN3 (3')	B-GGAGATGGGGAAGGCTCCCCACT	149

B = biotine.

The PCR reaction cycle was composed of the following steps:

1 min at 96°C;

- 5 times (30 s at 96°C; 50 s at 64°C; 50 s at 72°C);
 5 times (30 s at 96°C; 50 s at 62°C; 50 s at 72°C);
 10 times (30 s at 96°C; 50 s at 60°C; 50 s at 72°C);
 15 times (30 s at 96°C; 50 s at 55°C; 50 s at 72°C);
 5 min at 72°C.

The amplification reaction was carried out in 50 mM Tris-HCl pH 9.2, 16 mM (NH₄)₂SO₄, 200 μM dNTPs, 2.5 U Taq polymerase, 1.5 mM MgCl₂, 15 pmol of each primer and 0.1 to 0.5 μg DNA.

10

The resulting amplicon was cloned in the pGEMt-vector (Promega, Madison, WI, USA). Nucleotide sequence analysis was performed by use of an automated DNA sequencer Model 373A (Applied Biosystems, Foster City, CA, USA) with fluorescence-labelled dideoxy nucleotides (Prism™ Ready Reaction Dye Terminator Cycle Sequencing Kit; Applied Biosystems, Foster City, CA, USA). The primers used for the sequencing reaction were the same as for the amplification step. 29 intron 2 sequences were obtained for HLA-A, 38 for HLA-B and 13 for HLA-C. The sequences are shown in Figures 1, 2 and 3, respectively.

20 **Example 2: Amplification of exon 2 and exon 3 of HLA-A**

Nucleic acids were prepared from different blood samples by use of the QIAamp Blood Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol. Based on the sequence alignment of the HLA-A intron 2 sequences (Figure 1), a reverse and a forward, locus-specific primer was designed for the specific amplification of the HLA-A exon 2 and exon 3, respectively. With these primers, a primer mix was constructed for the separate amplification of exon 2 and exon 3 of HLA-A consisting of the following 2 primer sets:

- for exon 2: 5APBio (SEQ ID NO 144) as forward primer and
 5'ATCTCGGACCCGGAGACTGT3' (SEQ ID NO 1) as reverse primer;
- for exon 3: 5'CAGTTTAGGCCAAAAATCCCCC3' (SEQ ID NO 104) as
 forward primer and 3APBio (SEQ ID NO 147) as reverse primer.

The PCR reaction cycle was composed of the following steps:

5 min at 96°C;

35 times (30 s at 96°C; 20 s at 58°C; 30 s at 72°C);

10 min at 72°C.

- 5 The PCR reaction was carried out in 10 mM Tris.HCl pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.001% (w/v) gelatine, 200µM dNTP's (dATP, dGTP, dCTP, dTTP), 20 pmol of each primer and 1U AmpliTaq (Applied Biosystems, Foster City, CA, USA). The length of the obtained PCR products was verified on an agarose gel according to Sambrook et al. (1989).

10

Example 3: Amplification of exon 2 and exon 3 of HLA-B

Nucleic acids were prepared from different blood samples by use of the QIAamp Blood Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol.

- 15 Based on the sequence alignment of the HLA-B intron 2 sequences (Figure 2), a reverse and a forward, locus-specific primer was designed for the specific amplification of the HLA-B exon 2 and exon 3, respectively. With these primers, a primer mix was constructed for the separate amplification of exon 2 and exon 3 of HLA-B consisting of the following 2 primer sets:

- 20 - for exon 2: IBPin1 (SEQ ID NO 145) as forward primer and 5'ACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 310) as reverse primer;
- for exon 3: 5'ACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 121) as forward primer and IBPin3 (SEQ ID NO 148) as reverse primer.

25

The PCR reaction cycle was composed of the following steps:

5 min at 96°C;

35 times (30 s at 96°C; 20 s at 58°C; 30 s at 72°C);

10 min at 72°C.

30

The PCR reaction was carried out in 10 mM Tris.HCl pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.001% (w/v) gelatine, 200µM dNTP's (dATP, dGTP, dCTP, dTTP), 20 pmol of each primer and 1U AmpliTaq (Applied Biosystems, Foster City, CA, USA). The

length of the obtained PCR products was verified on an agarose gel according to Sambrook et al. (1989).

Example 4: Amplification of exon 2 and exon 3 of HLA-C

5

Nucleic acids were prepared from different blood samples by use of the QIAamp Blood Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol. Based on the sequence alignment of the HLA-C intron 2 sequences (Figure 3), a reverse and a forward, locus-specific primer was designed for the specific amplification of the HLA-C exon 2 and exon 3, respectively. With these primers, a primer mix was constructed for the separate amplification of exon 2 and exon 3 of HLA-C consisting of the following 2 primer sets:

- for exon 2: 5CIN1 (SEQ ID NO 146) as forward primer and 5'GGTCGAGGGTCTGGGCGGGTT3' (SEQ ID NO 127) as reverse primer;
- for exon 3: 5'TCGRCCGGRGAGAGCCCCAGT3' (SEQ ID NO 139) as forward primer and 3CIN3 (SEQ ID NO 149) as reverse primer.

The PCR reaction cycle was composed of the following steps:

- 5 min at 96°C;
- 35 times (30 s at 96°C; 20 s at 58°C; 30 s at 72°C);
- 10 min at 72°C.

The PCR reaction was carried out in 10 mM Tris.HCl pH 8.3? 50 mM KCl, 1.5 mM MgCl₂, 0.001% (w/v) gelatine, 200µM DNTP's (dATP, dGTP, dCTP, dTTP), 20 pmol of each primer and 1U AmpliTaq (Applied Biosystems, Foster City, CA, USA). The length of the obtained PCR products was verified on an agarose gel according to Sambrook et al. (1989).

Example 5: Amplification exon 2, exon 3 and exon 4 of HLA-A

Blood samples were collected from a Caucasian donor at B.A.R.C. (Gent, Belgium). Nucleic acids were prepared from the blood samples by use of the QIAamp Blood Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol.

5.1 Primer mix for the separate amplification of exon 2, exon 3 and exon 4 of HLA-A

- 5 A primer mix was used for the separate amplification of exon 2, exon 3 and exon 4 of HLA-A consisting of the following primer sets:

Amplicon	Primerset	Sequence (5'-3')	SEQ ID NO
Exon 1-2	5APbio (5')	B-TTCTCCCCAGACGCCGAGGATGGCC	144
	3ex2APbio (3')	B-ATCTCGGACCCGGAGACTGT	1
Exon 3	5ex3APbio (5')	B-CAGTTTAGGCCAAAAATCCCCC	104
	3ex3APbio (3')	B-CCCTCCTTGTGGGAGGCCAG	156
Exon 4	5ex4APbio (5')	B-GTTCTGTGCTCYCTTCCCCAT	205
	3ex4APbio (3')	B-TTGGGCAGACCCTCATGCTGC	311

B = biotine.

- 10 The PCR reaction cycle was composed of the following steps:

1 min 96 °C;

5 times (30 s at 96 °C; 50 s at 64 °C; 50 s at 72 °C);

5 times (30 s at 96 °C; 50 s at 62 °C; 50 s at 72 °C);

10 times (30 s at 96 °C; 50 s at 60 °C; 50 s at 72 °C);

- 15 15 times (30 s at 96 °C; 50 s at 55 °C; 50 s at 72 °C);

5 min 72 °C;

4°C.

- 20 The PCR reaction was carried out in 50 mM Tris.HCl pH 9.2, 16 mM (NH₄)₂SO₄, 1.5 mM MgCl₂, 200µM dNTP's (dATP, dGTP, dCTP, dTTP), 2.5 U Taq polymerase (Perkin Elmer, Roche Molecular Systems, Branchburg, New Jersey, US), 20 pmol primer 5APbio, 20 pmol primer 3ex2APbio, 40 pmol primer 5ex3APbio, 40 pmol primer 3ex3APbio, 15 pmol primer 5ex4APbio, 15 pmol primer 3ex4APbio and 0.1-0.5 µg DNA.

25

The length of the obtained amplification products was verified on a 3% agarose gel according to Sambrook et al. (1989). 3 bands of different length were obtained

relating respectively to exon 2 (559 bp), exon 3 (439 bp) and exon 4 (380 bp) of HLA-A. By use of this protocol, the HLA-A exons were amplified strong enough for subsequent typing in a hybridization assay.

5 5.2 Testing of the obtained amplification products in a HLA-A typing assay

The HLA-A amplicons were subsequently typed in a reverse hybridization assay based on the LiPA technology (Stuyver et al., 1993). After the amplification step as described above, the amplified nucleic acids were hybridized to a panel of 36 probes by use of the LiPA HLA-A kit (Innogenetics, Gent, Belgium) according to the manufacturer's instructions. Results of this reverse hybridization are shown in Figure 7.

For comparison, also an amplification product comprising exon 2 and exon 3 in one single amplicon was obtained by use of 5APBio (forward primer; SEQ ID NO 144) and 3APBio (reverse primer; SEQ ID NO 147). The hybridization of this larger amplicon to the probes on the LiPA strip is also shown in Figure 7. Figure 7 clearly illustrates that the amplicons obtained by separate amplification of exon 2 and exon 3 enable a more clear and pronounced typing than the larger amplicon obtained by amplification of exon 2 and exon 3 in one single amplicon.

Example 6: Amplification exon 2, exon 3 and exon 4 of HLA-B

Blood samples were collected from a Caucasian donor at B.A.R.C. (Gent, Belgium). Nucleic acids were prepared from the blood samples by use of the QIAamp Blood Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol.

6.1 Primer mix for the separate amplification of exon 2, exon 3 and exon 4 of HLA-B

A primer mix was used for the separate amplification of exon 2, exon 3 and exon 4 of HLA-B consisting of the following primer sets:

Amplicon	Primer set	Sequence (5'-3')	SEQ ID NO
Exon 2	IBPin1bio (5')	B-GGGAGGAGCGAGGGGACCSCAG	145
	IB3Pin2bio (3')	B-AACCCGCGGGGATTTTGGCCTC	109
Exon 3	IB5Pin2bio (5')	B-CGCGTTTACCCGGTTTCATTTTCAGTTG	224
	IB3Pin3bio (3')	B-TCTTCTCGTKGGAGSCCATCCCC	234
Exon 4	5ex4IBbio (5')	B-TCACATGGGTGGTCCTAGG	267
	3ex4IBbio (3')	B-TCGGCAGCCCTCATGCTGT	312

B = biotine.

The PCR reaction cycle was composed of the following steps:

- 1 min 96 °C;
- 5 5 times (30 s at 96 °C; 50 s at 64 °C; 50 s at 72 °C);
- 5 times (30 s at 96 °C; 50 s at 62 °C; 50 s at 72 °C);
- 10 times (30 s at 96 °C; 50 s at 60 °C; 50 s at 72 °C);
- 15 times (30 s at 96 °C; 50 s at 55 °C; 50 s at 72 °C);
- 5 min 72 °C;
- 10 4°C

The PCR reaction was carried out in 50 mM Tris.HCl pH 9.2, 16 mM (NH₄)₂SO₄, 1.5 mM MgCl₂, 200µM dNTP's (dATP, dGTP, dCTP, dTTP), 2.5 U Taq polymerase (Perkin Elmer, Roche Molecular Systems, Branchburg, New Jersey, US), 35 pmol primer IBPin1bio, 35 pmol primer IBPin2bio, 50 pmol primer IB5Pin2bio, 50 pmol primer IB3Pin3bio, 10 pmol primer 5ex4IBbio, 10 pmol primer 3ex4IBbio and 0.1-0.5 µg DNA.

The length of the obtained amplification products was verified on a 3% agarose gel according to Sambrook et al. (1989). 3 bands of different length were obtained relating respectively to HLA-B exon 2 (555 bp), exon 3 (446 bp) and exon 4 (323 bp). By use of this protocol, the HLA-B exons were amplified strong enough for subsequent typing in a Line Probe Assay.

6.2 Testing of the obtained amplification products in a HLA-B typing assay

The HLA-B amplicons were subsequently typed in a reverse hybridization assay based on the LiPA technology (Stuyver et al., 1993). After the amplification step as described above, the amplified nucleic acids were hybridized to a panel of 60 probes by use of the LiPA HLA-B kit (Innogenetics, Gent, Belgium) according to the manufacturer's instructions. Results of this reverse hybridization are shown in Figure 8.

For comparison, also an amplification product comprising exon 2 and exon 3 in one single amplicon was obtained by use of IBPin1 (forward primer; SEQ ID NO 145) and IBPin3 (reverse primer; SEQ ID NO 148). The hybridization of this larger amplicon to the probes on the LiPA strip is also shown in Figure 8. Figure 8 clearly illustrates that the amplicons obtained by separate amplification of exon 2 and exon 3 enable a more clear and pronounced typing than the larger amplicon obtained by amplification of exon 2 and exon 3 in one single amplicon.

Example 7: Amplification exon 2, exon 3 and exon 4 of HLA-C

Nucleic acids are prepared from the blood samples by use of the QIAamp Blood Kit (Qiagen, Hilden, Germany) according to the manufacturer's protocol. A primer mix is used for the separate amplification of exon 2, exon 3 and exon 4 of HLA-C consisting of the following primer sets:

Amplicon	Primerset	Sequence (5'-3')	SEQ ID NO
Exon 2	5CIN1(5')	B-AGCGAGGGGCCCCGCCCCGGCGA	146
	IC3Pin2bio(3')	B-GGTCGAGGGTCTGGGCGGGTT	127
Exon 3	IC5Pin2bio(5')	B-TCGRCCGGRGAGAGCCCCAGT	139
	3CIN3 (3')	B-GGAGATGGGGAAGGCTCCCCACT	149
Exon 4	5ex4ICbio(5')	B-TCTCAGGATRGTCACATGGSC	299
	3ex4ICbio(3')	B-CATCTCAGGGTGMRRGGGCTT	313

B = biotine.

The PCR reaction cycle is composed of the following steps:

- 1 min 96 °C;
5 times (30 s at 96 °C; 50 s at 64 °C; 50 s at 72 °C);
5 times (30 s at 96 °C; 50 s at 62 °C; 50 s at 72 °C);
10 times (30 s at 96 °C; 50 s at 60 °C; 50 s at 72 °C);
5 15 times (30 s at 96 °C; 50 s at 55 °C; 50 s at 72 °C);
5 min 72 °C;
4°C.

- The PCR reaction is carried out in 50 mM Tris.HCl pH 9.2, 16 mM (NH₄)₂SO₄, 1.5
10 mM MgCl₂, 200µM dNTP's (dATP, dGTP, dCTP, dTTP), 2.5 U Taq polymerase
(Perkin Elmer, Roche Molecular Systems, Branchburg, New Jersey, US), 20 pmol of
each primer and 0.1-0.5 µg DNA.

- The length of the obtained amplification products is verified on a 3% agarose gel
15 according to Sambrook et al. (1989). 3 bands of different length are obtained relating
respectively to HLA-C exon 2, exon 3 and exon 4. By use of this protocol, the HLA-C
exons are amplified strong enough for subsequent typing in a Line Probe Assay.

REFERENCES

- Andersson G, Larhammar D, Widmark E, Servenius B, Peterson PA and Rask L (1987) Class II genes of the human major histocompatibility complex. Organization and evolutionary relationship of the DRbeta genes. J Biol Chem 262: 8748-8758.
- Apple RJ and Erlich HA (1996) HLA classII genes: structure and diversity. Chapter 5 HLA and MHC: genes, molecules and function. BIOS Scientific Publishers Ltd., Oxford, UK.
- Asseline U, Delarue M, Lancelot G, Toulme F and Thuong N (1984) Nucleic acid-binding molecules with high affinity and base sequence specificity: intercalating agents covalently linked to oligodeoxynucleotides. Proc Natl Acad Sci USA 81: 3297-3301.
- Barany F (1991) The ligase chain reaction in a PCR world. PCR Methods Appl 1: 5-16.
- Campbell RD and Trowsdale J (1993) Map of the human MHC. Immunology Today 14: 349-352.
- Clay TM, Culpan D, Howell WM, Sage DA, Bradley BA and Bidwell JL (1994) UHG crossmatching. A comparison with PCR-SSO typing in the selection of HLA-DPB-compatible bone marrow donors. Transplantation 58: 200-207.
- Compton J (1991) Nucleic acid sequence-based amplification. Nature 350: 91-92.
- Duck P (1990) Probe amplifier system based on chimeric cycling oligonucleotides. Biotechniques 9: 142-147.
- Fleischhauer K, Kernan NA, O'Reilly RJ, Dupont B and Yang SY (1990) Bone marrow-allograft rejection by T lymphocytes recognizing a single amino acid difference in HLA-B44. New Eng J Med 323: 1818-1822.

Guatelli J, Whitfield K, Kwoh D, Barringer K, Richman D and Gengeras T (1990) Isothermal, in vitro amplification of nucleic acids by a multienzyme reaction modeled after retroviral replication. *Proc Natl Acad Sci USA* 87: 1874-1878.

- 5 Hirschorn K, Bach F, Kolodny RL and Firschen IL (1963) Immune response and mitosis of human peripheral blood lymphocytes in vitro. *Science* 142: 1185-1187.

Kissmeyer NF, Svejgaard A and Hauge M (1969) The HLA system defined with lymphocytotoxic and platelet antibodies in relation to kidney transplantation.

- 10 Transplant Proc 1: 357-361.

Kwoh D, Davis G, Whitfield K, Chappelle H, Dimichele L and Gengeras T (1989) Transcription-based amplification system and detection of amplified human immunodeficiency virus type 1 with a bead-based sandwich hybridization format.

- 15 Proc Natl Acad Sci USA 86: 1173-1177.

Kwok S, Kellogg DE, McKinney N, Spasic D, Goda L, Levenson C and Sninsky JJ (1990) Effects of primer-template mismatches on the polymerase chain reaction: human immunodeficiency virus type 1 model studies. *Nucleic Acids Res.* 18: 999-

- 20 1005.

Landgren U, Kaiser R, Sanders J and Hood L (1988) A ligase-mediated gene detection technique. *Science* 241: 1077-1080.

- 25 Lomeli H, Tyagi S, Pritchard C, Lisardi P and Kramer F (1989) Quantitative assays based on the use of replicatable hybridization probes. *Clin Chem* 35: 1826-1831.

Matsukura M, Shinozuka K, Zon G, Mitsuya H, Reitz M, Cohen J and Broder S (1987) Phosphorothioate analogs of oligodeoxynucleotides: inhibitors of replication and cytopathic effects of human immunodeficiency virus. *Proc Natl Acad Sci USA* 84: 7706-7710.

- 30

- Miller P, Yano J, Yano E, Carroll C, Jayaram K and Ts'o P (1979) Nonionic nucleic acid analogues. Synthesis and characterization of dideoxyribonucleoside methylphosphonates. *Biochemistry* 18: 5134-5143.
- 5 Mullis K, Faloona F, Scharf S, Saiki R, Horn G and Erlich H (1986) Specific enzymatic amplification of DNA in vitro: the polymerase chain reaction. *Gold Spring Harb Symp Quant Biol* 1: 263-273.
- Mullis KB and Faloona FA (1987) Specific synthesis of DNA in vitro via a
10 polymerase-catalyzed chain reaction. *Methods Enzymol* 155: 335-350.
- Nielsen P, Egholm M, Berg R and Buchardt O (1991) Sequence-selective recognition of DNA by strand displacement with a thymine-substituted polyamide. *Science* 254: 1497-1500.
- 15 Nielsen P, Egholm M, Berg R and Buchardt O (1993) Sequence specific inhibition of DNA restriction enzyme cleavage by PNA. *Nucleic-Acids-Res* 21: 197-200.
- Olerup O and Zetterquist H (1991) HLA-DRB1*01 subtyping by allele-specific PCR
20 amplification: a sensitive, specific and rapid technique. *Tissue Antigens* 37: 197-204.
- Saiki RK, Bugawan TL, Horn GT, Mullis KB and Erlich HA (1986) Analysis of enzymatically amplified beta-globin and HLA-DQ alpha DNA with allele-specific oligonucleotide probes. *Nature* 324: 163-166.
- 25 Saiki RK, Gelfand DH, Stoffel S, Scharf SJ, Higuchi R, Horn GT, Mullis KB and Erlich HA (1988) Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239: 487-491.
- 30 Saiki RK, Walsh PS, Levenson CH and Erlich HA (1989) Genetic analysis of amplified DNA with immobilizes sequence-specific oligonucleotide probes. *Proc Natl Acad Sci USA* 86: 6230-6234.

Sambrook J, Fritsch E and Maniatis T (1989) Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA.

- 5 Santamaria P, Boyce-Jacino MT, Lindstrom AL, Barbosa JJ, Faras AJ and Rich SS (1992) HLA class II "typing": direct sequencing of DRB, DQB and DQA genes. Hum Immunol 33: 69-81.

- 10 Santamaria P, Lindstrom AL, Boyce JM, Jacino MT, Myster SH, Barbosa JJ, Faras AJ and Rich SS (1993) HLA class I sequence-based typing. Hum Immunol 37: 39-50.

Sanchez-Pescador R, Stempien MS and Urdea MS (1988) Rapid chemiluminescent nucleic acid assays for detection of TEM-1 beta-lactamase-mediated penicillin resistance in *Neisseria gonorrhoeae* and other bacteria. J Clin Microbiol 26: 1934-1938.

- 15 Spencer WR and Parham P (1996) HLA class I genes: structure and diversity. Chapter 4. HLA and MHC: genes, molecules and function. BIOS Scientific Publishers Ltd., Oxford, UK.

- 20 Stuyver L, Rossau R, Wyseur A, Duhamel M, Vanderborght B, Van Heuverswyn H and Maertens G (1993) Typing of hepatitis C virus isolates and characterization of new subtypes using a line probe assay. J Gen Virol 74: 1093-1102.

- 25 Terasaki PH and McClelland JD (1964) Microdot assay of human serum cytotoxins. Nature 204: 998-1007.

Townsend A and Bodmer H (1989) Antigen recognition by Class I-restricted T lymphocytes. Annu Rev Immunol 7: 601-624.

- 30 Urdea MS, Horn T, Fultz TJ, Anderson M, Running JA, Hamren S, Ahle D and Chang CA (1991) Branched DNA amplification multimers for the sensitive, direct detection of human hepatitis viruses. Nucleic Acids Symp Ser 24:197-200.

Wu D and Wallace B (1989) The ligation amplification reaction (LAR) - amplification of specific DNA sequences using sequential rounds of template-dependent ligation. *Genomics* 4: 560-569.

- 5 Yang SY (1987) A standardised method for detection of HLA-A and HLA-B alleles by one-dimensional isoelectric focusing (IEF) gel electrophoresis. *Immunobiology of HLA. Histocompatibility testing* (ed. B Dupont). Springer-Verlag, New York, USA. pp. 332-335.
- 10 Yoshida M, Kimura A, Numano F and Sasazuki T (1992) Polymerase-chain-reaction-based analysis of polymorphism in the HLA-B gene. *Hum Immunol* 34: 257-266.

CLAIMS

1. Method for the locus-specific, separate amplification of exon 2, exon 3 and/or exon 4 of HLA-A, HLA-B or HLA-C alleles, making use of at least one primer set wherein:
 - for the amplification of exon 2, the reverse primer specifically hybridizes to a locus-specific target sequence in intron 2 of respectively HLA-A, HLA-B or HLA-C;
 - for the amplification of exon 3, the forward primer specifically hybridizes to a locus-specific target sequence in intron 2 of respectively HLA-A, HLA-B or HLA-C and/or the reverse primer specifically hybridizes to a locus-specific target sequence in intron 3 of respectively HLA-A, HLA-B or HLA-C;
 - for the amplification of exon 4, the forward primer specifically hybridizes to a locus-specific target sequence in intron 3 of respectively HLA-A, HLA-B or HLA-C.
2. Method according to claim 1 further characterized that the locus-specific target sequence is situated at:
 - position 67, 96, 109, 110, 118, 123, 131 or 181 of the HLA-A intron 2 (Fig. 1) and/or position 32, 50, 62, 73, 83, 86, 118, 130, 150, 501, 525, 561 or 571 of the HLA-A intron 3 (Fig. 4); or
 - position 35 or 170 of the HLA-B intron 2 (Fig. 2) and/or position 42, 46, 65, 68, 96, 438, 502, 524, 547 or 571 of the HLA-B intron 3 (Fig. 5); or
 - position 84, 107 or 142 of the HLA-C intron 2 (Fig. 3) and/or position 461, 477, 527, 545 or 561 of the HLA-C intron 3 (Fig. 6).
3. Method according to claim 2 further characterized that said positions constitute the 3' end of the primer that is used for the amplification of exon 2, exon 3 or exon 4.
4. Method according to claim 3 further characterized that the primer is chosen from the following list:
 - for the amplification of exon 2 of HLA-A (table 1):

- 5'ATCTCGGACCCGGAGACTGT3' (SEQ ID NO 1)
5'GATCTCGGACCCGGAGACTGT3' (SEQ ID NO 2)
5'GGATCTCGGACCCGGAGACTGT3' (SEQ ID NO 3)
5'YGGATCTCGGACCCGGAGACTGT3' (SEQ ID NO 4)
5 5'GYGGATCTCGGACCCGGAGACTGT3' (SEQ ID NO 5)
5'GGYGGATCTCGGACCCGGAGACTGT3' (SEQ ID NO 6)
5'GGTCTCGGRGTCCCGCGGCT3' (SEQ ID NO 7)
5'GGGTCTCGGRGTCCCGCGGCT3' (SEQ ID NO 8)
5'AGGGTCTCGGRGTCCCGCGGCT3' (SEQ ID NO 9)
10 5'AAGGGTCTCGGRGTCCCGCGGCT3' (SEQ ID NO 10)
5'CAAGGGTCTCGGRGTCCCGCGGCT3' (SEQ ID NO 11)
5'CTCCCGGGDCAAGGGTCTCG3' (SEQ ID NO 12)
5'TCTCCCGGGDCAAGGGTCTCG3' (SEQ ID NO 13)
5'CTCTCCCGGGDCAAGGGTCTCG3' (SEQ ID NO 14)
15 5'CCTCTCCCGGGDCAAGGGTCTCG3' (SEQ ID NO 15)
5'GCCTCTCCCGGGDCAAGGGTCTCG3' (SEQ ID NO 16)
5'GGCCTCTCCCGGGDCAAGGGTCTCG3' (SEQ ID NO 17)
5'TCTCCCGGGDCAAGGGTCTC3' (SEQ ID NO 18)
5'CTCTCCCGGGDCAAGGGTCTC3' (SEQ ID NO 19)
20 5'CCTCTCCCGGGDCAAGGGTCTC3' (SEQ ID NO 20)
5'GCCTCTCCCGGGDCAAGGGTCTC3' (SEQ ID NO 21)
5'GGCCTCTCCCGGGDCAAGGGTCTC3' (SEQ ID NO 22)
5'GGGCCTCTCCCGGGDCAAGGGTCTC3' (SEQ ID NO 23)
5'CCTGGGCCTCTCCCGGGDCA3' (SEQ ID NO 30)
25 5'GCCTGGGCCTCTCCCGGGDCA3' (SEQ ID NO 31)
5'CGCCTGGGCCTCTCCCGGGDCA3' (SEQ ID NO 32)
5'GCGCCTGGGCCTCTCCCGGGDCA3' (SEQ ID NO 33)
5'GGCGCCTGGGCCTCTCCCGGGDCA3' (SEQ ID NO 34)
5'AGGCGCCTGGGCCTCTCCCGGGDCA3' (SEQ ID NO 35)
30 5'AGGCGCCTGGGCCTCTCCCG3' (SEQ ID NO 36)
5'AAGGCGCCTGGGCCTCTCCCG3' (SEQ ID NO 37)
5'WAAGGCGCCTGGGCCTCTCCCG3' (SEQ ID NO 38)
5'TWAAGGCGCCTGGGCCTCTCCCG3' (SEQ ID NO 39)
5'GTWAAGGCGCCTGGGCCTCTCCCG3' (SEQ ID NO 40)

- 5'GGTWAAGGCGCCTGGGCCTCTCCCG3' (SEQ ID NO 41)
 5'CCGGGTWAAGGCGCCTGGGC3' (SEQ ID NO 42)
 5'ACCGGGTWAAGGCGCCTGGGC3' (SEQ ID NO 43)
 5'AACCGGGTWAAGGCGCCTGGGC3' (SEQ ID NO 44)
 5 5'AAACCGGGTWAAGGCGCCTGGGC3' (SEQ ID NO 45)
 5'GAAACCGGGTWAAGGCGCCTGGGC3' (SEQ ID NO 46)
 5'TGAAACCGGGTWAAGGCGCCTGGGC3' (SEQ ID NO 47)
 5'YCCVGCCCCGACCAACCYGG3' (SEQ ID NO 48)
 5'GYCCVGCCCCGACCAACCYGG3' (SEQ ID NO 49)
 10 5'YGYCCVGCCCCGACCAACCYGG3' (SEQ ID NO 50)
 5'CYGYCCVGCCCCGACCAACCYGG3' (SEQ ID NO 51)
 5'CCYGYCCVGCCCCGACCAACCYGG3' (SEQ ID NO 52)
 5'CCCYGYCCVGCCCCGACCAACCYGG3' (SEQ ID NO 53)
- 15 - for the amplification of exon 3 of HLA-A (table 2; table 3):
 5'CGGACGGGCCRGGTSRCCCA3' (SEQ ID NO 54)
 5'ACGGACGGGCCRGGTSRCCCA3' (SEQ ID NO 55)
 5'CACGGACGGGCCRGGTSRCCCA3' (SEQ ID NO 56)
 5'CCACGGACGGGCCRGGTSRCCCA3' (SEQ ID NO 57)
 20 5'CCCACGGACGGGCCRGGTSRCCCA3' (SEQ ID NO 58)
 5'CCCCACGGACGGGCCRGGTSRCCCA3' (SEQ ID NO 59)
 5'GGTCCGAGATCCRCCCCGAA3' (SEQ ID NO 60)
 5'GGGTCCGAGATCCRCCCCGAA3' (SEQ ID NO 61)
 5'CGGGTCCGAGATCCRCCCCGAA3' (SEQ ID NO 62)
 25 5'CCGGGTCCGAGATCCRCCCCGAA3' (SEQ ID NO 63)
 5'TCCGGGTCCGAGATCCRCCCCGAA3' (SEQ ID NO 64)
 5'CTCCGGGTCCGAGATCCRCCCCGAA3' (SEQ ID NO 65)
 5'CCCCGAAGCCGCGGGACYCC3' (SEQ ID NO 66)
 5'RCCCCGAAGCCGCGGGACYCC3' (SEQ ID NO 67)
 30 5'CRCCCCGAAGCCGCGGGACYCC3' (SEQ ID NO 68)
 5'CCRCCCCGAAGCCGCGGGACYCC3' (SEQ ID NO 69)
 5'TCCRCCCCGAAGCCGCGGGACYCC3' (SEQ ID NO 70)
 5'ATCCRCCCCGAAGCCGCGGGACYCC3' (SEQ ID NO 71)
 5'CCCGAAGCCGCGGGACYCCG3' (SEQ ID NO 72)

5'CCCCGAAGCCGCGGGACYCCG3' (SEQ ID NO 73)
5'RCCCCGAAGCCGCGGGACYCCG3' (SEQ ID NO 74)
5'CRCCCCGAAGCCGCGGGACYCCG3' (SEQ ID NO 75)
5'CCRCCCCGAAGCCGCGGGACYCCG3' (SEQ ID NO 76)
5 5'TCCRCCCCGAAGCCGCGGGACYCCG3' (SEQ ID NO 77)
5'CGCGGGACYCCGAGACCCTT3' (SEQ ID NO 84)
5'CCGCGGGACYCCGAGACCCTT3' (SEQ ID NO 85)
5'GCCGCGGGACYCCGAGACCCTT3' (SEQ ID NO 86)
5'AGCCGCGGGACYCCGAGACCCTT3' (SEQ ID NO 87)
10 5'AAGCCGCGGGACYCCGAGACCCTT3' (SEQ ID NO 88)
5'GAAGCCGCGGGACYCCGAGACCCTT3' (SEQ ID NO 89)
5'GACYCCGAGACCCTTGDCCC3' (SEQ ID NO 90)
5'GGACYCCGAGACCCTTGDCCC3' (SEQ ID NO 91)
5'GGGACYCCGAGACCCTTGDCCC3' (SEQ ID NO 92)
15 5'CGGGACYCCGAGACCCTTGDCCC3' (SEQ ID NO 93)
5'GCGGGACYCCGAGACCCTTGDCCC3' (SEQ ID NO 94)
5'CGCGGGACYCCGAGACCCTTGDCCC3' (SEQ ID NO 95)
5'GACCCTTGDCCCCGGGAGAGG3' (SEQ ID NO 96)
5'AGACCCTTGDCCCCGGGAGAGG3' (SEQ ID NO 97)
20 5'GAGACCCTTGDCCCCGGGAGAGG3' (SEQ ID NO 98)
5'CGAGACCCTTGDCCCCGGGAGAGG3' (SEQ ID NO 99)
5'CCGAGACCCTTGDCCCCGGGAGAGG3' (SEQ ID NO 100)
5'YCCGAGACCCTTGDCCCCGGGAGAGG3' (SEQ ID NO 101)
5'GTTTAGGCCAAAAATCCCCC3' (SEQ ID NO 102)
25 5'AGTTTAGGCCAAAAATCCCCC3' (SEQ ID NO 103)
5'CAGTTTAGGCCAAAAATCCCCC3' (SEQ ID NO 104)
5'TCAGTTTAGGCCAAAAATCCCCC3' (SEQ ID NO 105)
5'TTCAGTTTAGGCCAAAAATCCCCC3' (SEQ ID NO 106)
5'TTTCAGTTTAGGCCAAAAATCCCCC3' (SEQ ID NO 107)
30 5'AGCCCGGGAGATCTAYAGGC3' (SEQ ID NO 150)
5'CAGCCCGGGAGATCTAYAGGC3' (SEQ ID NO 151)
5'CCAGCCCGGGAGATCTAYAGGC3' (SEQ ID NO 152)
5'GCCAGCCCGGGAGATCTAYAGGC3' (SEQ ID NO 153)
5'GGCCAGCCCGGGAGATCTAYAGGC3' (SEQ ID NO 154)

5'AGGCCAGCCCGGGAGATCTAYAGGC3' (SEQ ID NO 155)
5'CCCTCCTTGTGGGAGGCCAG3' (SEQ ID NO 156)
5'CCCCTCCTTGTGGGAGGCCAG3' (SEQ ID NO 157)
5'TCCCCTCCTTGTGGGAGGCCAG3' (SEQ ID NO 158)
5'CTCCCCTCCTTGTGGGAGGCCAG3' (SEQ ID NO 159)
5'TCTCCCCTCCTTGTGGGAGGCCAG3' (SEQ ID NO 160)
5'GTCTCCCCTCCTTGTGGGAGGCCAG3' (SEQ ID NO 161)
5'CCCAAWTGTCTCCCCTCCTT3' (SEQ ID NO 162)
5'TCCCAAWTGTCTCCCCTCCTT3' (SEQ ID NO 163)
5'GTCCCAAWTGTCTCCCCTCCTT3' (SEQ ID NO 164)
5'GGTCCCAAWTGTCTCCCCTCCTT3' (SEQ ID NO 165)
5'TGGTCCCAAWTGTCTCCCCTCCTT3' (SEQ ID NO 166)
5'TTGGTCCCAAWTGTCTCCCCTCCTT3' (SEQ ID NO 167)
5'CTAGTGTTGGTCCCAAWTGT3' (SEQ ID NO 168)
5'TCTAGTGTTGGTCCCAAWTGT3' (SEQ ID NO 169)
5'TTCTAGTGTTGGTCCCAAWTGT3' (SEQ ID NO 170)
5'ATTCTAGTGTTGGTCCCAAWTGT3' (SEQ ID NO 171)
5'TATTCTAGTGTTGGTCCCAAWTGT3' (SEQ ID NO 172)
5'ATATTCTAGTGTTGGTCCCAAWTGT3' (SEQ ID NO 173)
5'GGGYGATATTCTAGTGTTGG3' (SEQ ID NO 174)
5'AGGGYGATATTCTAGTGTTGG3' (SEQ ID NO 175)
5'GAGGGYGATATTCTAGTGTTGG3' (SEQ ID NO 176)
5'GGAGGGYGATATTCTAGTGTTGG3' (SEQ ID NO 177)
5'GGGAGGGYGATATTCTAGTGTTGG3' (SEQ ID NO 178)
5'AGGGAGGGYGATATTCTAGTGTTGG3' (SEQ ID NO 179)
5'GGAGGGYGATATTCTAGTGT3' (SEQ ID NO 180)
5'GGGAGGGYGATATTCTAGTGT3' (SEQ ID NO 181)
5'AGGGAGGGYGATATTCTAGTGT3' (SEQ ID NO 182)
5'GAGGGAGGGYGATATTCTAGTGT3' (SEQ ID NO 183)
5'AGAGGGAGGGYGATATTCTAGTGT3' (SEQ ID NO 184)
5'CAGAGGGAGGGYGATATTCTAGTGT3' (SEQ ID NO 185)
5'CCCAGGAGGAKTCCTCTCCC3' (SEQ ID NO 186)
5'ACCCAGGAGGAKTCCTCTCCC3' (SEQ ID NO 187)
5'AACCCAGGAGGAKTCCTCTCCC3' (SEQ ID NO 188)

5'AAACCCAGGAGGAKTCCTCTCCC3' (SEQ ID NO 189)
 5'GAAACCCAGGAGGAKTCCTCTCCC3' (SEQ ID NO 190)
 5'GGAAACCCAGGAGGAKTCCTCTCCC3' (SEQ ID NO 191)
 5'AGGATCTGGAAACCCAGGAG3' (SEQ ID NO 192)
 5'CAGGATCTGGAAACCCAGGAG3' (SEQ ID NO 193)
 5'ACAGGATCTGGAAACCCAGGAG3' (SEQ ID NO 194)
 5'TACAGGATCTGGAAACCCAGGAG3' (SEQ ID NO 195)
 5'GTACAGGATCTGGAAACCCAGGAG3' (SEQ ID NO 196)
 5'GGTACAGGATCTGGAAACCCAGGAG3' (SEQ ID NO 197)
 5'TCAGAGTCACTCTCTGGTAC3' (SEQ ID NO 198)
 5'CTCAGAGTCACTCTCTGGTAC3' (SEQ ID NO 199)
 5'CCTCAGAGTCACTCTCTGGTAC3' (SEQ ID NO 200)
 5'ACCTCAGAGTCACTCTCTGGTAC3' (SEQ ID NO 201)
 5'AACCTCAGAGTCACTCTCTGGTAC3' (SEQ ID NO 202)
 5'GAACCTCAGAGTCACTCTCTGGTAC3' (SEQ ID NO 203)

- for the amplification of exon 4 of HLA-A (table 4):

5'TTCTGTGCTCYCTTCCCCAT3' (SEQ ID NO 204)
 5'GTTCTGTGCTCYCTTCCCCAT3' (SEQ ID NO 205)
 5'GGTTCTGTGCTCYCTTCCCCAT3' (SEQ ID NO 206)
 5'GGGTTCTGTGCTCYCTTCCCCAT3' (SEQ ID NO 207)
 5'TGGGTTCTGTGCTCYCTTCCCCAT3' (SEQ ID NO 208)
 5'CTGGGTTCTGTGCTCYCTTCCCCAT3' (SEQ ID NO 209)
 5'GGTGTCCCTGTCCATTCTCAA3' (SEQ ID NO 24)
 5'RGGTGTCCTGTCCATTCTCAA3' (SEQ ID NO 25)
 5'CRGGTGTCCCTGTCCATTCTCAA3' (SEQ ID NO 26)
 5'CCRGGTGTCCCTGTCCATTCTCAA3' (SEQ ID NO 27)
 5'CCCRGGTGTCCCTGTCCATTCTCAA3' (SEQ ID NO 28)
 5'TCCCRGGTGTCCCTGTCCATTCTCAA3' (SEQ ID NO 29)
 5'CTGGWGGAGTGTCCCATKAC3' (SEQ ID NO 210)
 5'GCTGGWGGAGTGTCCCATKAC3' (SEQ ID NO 211)
 5'TGCTGGWGGAGTGTCCCATKAC3' (SEQ ID NO 212)
 5'RTGCTGGWGGAGTGTCCCATKAC3' (SEQ ID NO 213)
 5'YRTGCTGGWGGAGTGTCCCATKAC3' (SEQ ID NO 214)

5'GYRTGCTGGWGGAGTGTCCCATKAC3' (SEQ ID NO 215)
 5'GTCCCATKACAGATRCMMAA3' (SEQ ID NO 216)
 5'TGTCCCATKACAGATRCMMAA3' (SEQ ID NO 217)
 5'GTGTCCCATKACAGATRCMMAA3' (SEQ ID NO 218)
 5'AGTGTCCCATKACAGATRCMMAA3' (SEQ ID NO 219)
 5'GAGTGTCCCATKACAGATRCMMAA3' (SEQ ID NO 220)
 5'GGAGTGTCCCATKACAGATRCMMAA3' (SEQ ID NO 221)

- for the amplification of exon 2 of HLA-B (table 5):

5'ACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 108)
 5'AACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 109)
 5'CAACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 110)
 5'CCAACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 111)
 5'MCCAACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 112)
 5'GMCCAACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 113)
 5'YGMCCAACCCGCGGGGATTTTGGCCTC3' (SEQ ID NO 314)

- for the amplification of exon 3 of HLA-B (table 6; table 7):

5'CYGGGGCGSAGGTCACGACT3' (SEQ ID NO 114)
 5'CCYGGGGCGSAGGTCACGACT3' (SEQ ID NO 115)
 5'GCCYGGGGCGSAGGTCACGACT3' (SEQ ID NO 116)
 5'GGCCYGGGGCGSAGGTCACGACT3' (SEQ ID NO 117)
 5'CGGCCYGGGGCGSAGGTCACGACT3' (SEQ ID NO 118)
 5'CCGGCCYGGGGCGSAGGTCACGACT3' (SEQ ID NO 119)
 5'CCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 120)
 5'ACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 121)
 5'TACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 122)
 5'TTACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 123)
 5'TTTACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 124)
 5'GTTTACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 125)
 5'CGTTTACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 222)
 5'GCGTTTACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 223)
 5'CGCGTTTACCCGGTTTCATTTTCAGTTG3' (SEQ ID NO 224)
 5'CGTKGGAGSCCATCCCCGSC3' (SEQ ID NO 225)

5'TCGTKGGAGSCCATCCCCGSC3' (SEQ ID NO 226)
 5'CTCGTKGGAGSCCATCCCCGSC3' (SEQ ID NO 227)
 5'TCTCGTKGGAGSCCATCCCCGSC3' (SEQ ID NO 228)
 5'TTCTCGTKGGAGSCCATCCCCGSC3' (SEQ ID NO 229)
 5'CTTCTCGTKGGAGSCCATCCCCGSC3' (SEQ ID NO 230)
 5'TCTCGTKGGAGSCCATCCCC3' (SEQ ID NO 231)
 5'TTCTCGTKGGAGSCCATCCCC3' (SEQ ID NO 232)
 5'CTTCTCGTKGGAGSCCATCCCC3' (SEQ ID NO 233)
 5'TCTTCTCGTKGGAGSCCATCCCC3' (SEQ ID NO 234)
 5'YTCTTCTCGTKGGAGSCCATCCCC3' (SEQ ID NO 235)
 5'CYTCTTCTCGTKGGAGSCCATCCCC3' (SEQ ID NO 236)
 5'GATCCCATTTTCCTCYTCTT3' (SEQ ID NO 237)
 5'TGATCCCATTTTCCTCYTCTT3' (SEQ ID NO 238)
 5'CTGATCCCATTTTCCTCYTCTT3' (SEQ ID NO 239)
 5'GCTGATCCCATTTTCCTCYTCTT3' (SEQ ID NO 240)
 5'CGCTGATCCCATTTTCCTCYTCTT3' (SEQ ID NO 241)
 5'GCGCTGATCCCATTTTCCTCYTCTT3' (SEQ ID NO 242)
 5'GCTGATCCCATTTTCCTCYT3' (SEQ ID NO 243)
 5'CGCTGATCCCATTTTCCTCYT3' (SEQ ID NO 244)
 5'GCGCTGATCCCATTTTCCTCYT3' (SEQ ID NO 245)
 5'AGCGCTGATCCCATTTTCCTCYT3' (SEQ ID NO 246)
 5'TAGCGCTGATCCCATTTTCCTCYT3' (SEQ ID NO 247)
 5'CTAGCGCTGATCCCATTTTCCTCYT3' (SEQ ID NO 248)
 5'TCCATTCAAGGGAGGGCGAC3' (SEQ ID NO 249)
 5'CTCCATTCAAGGGAGGGCGAC3' (SEQ ID NO 250)
 5'TCTCCATTCAAGGGAGGGCGAC3' (SEQ ID NO 251)
 5'TTCTCCATTCAAGGGAGGGCGAC3' (SEQ ID NO 252)
 5'ATTCTCCATTCAAGGGAGGGCGAC3' (SEQ ID NO 253)
 5'CATTCTCCATTCAAGGGAGGGCGAC3' (SEQ ID NO 254)

30

- for the amplification of exon 4 of HLA-B (table 8):

5'AGATTATCCCAGGTGCCTGC3' (SEQ ID NO 255)
 5'GAGATTATCCCAGGTGCCTGC3' (SEQ ID NO 256)
 5'GGAGATTATCCCAGGTGCCTGC3' (SEQ ID NO 257)

- 5'AGGAGATTATCCCAGGTGCCTGC3' (SEQ ID NO 258)
 5'TAGGAGATTATCCCAGGTGCCTGC3' (SEQ ID NO 259)
 5'ATAGGAGATTATCCCAGGTGCCTGC3' (SEQ ID NO 260)
 5'TGTCCTGYCCATTCTCAGKC3' (SEQ ID NO 261)
 5'GTGTCCTGYCCATTCTCAGKC3' (SEQ ID NO 262)
 5'GGTGTCTGTCCTGYCCATTCTCAGKC3' (SEQ ID NO 263)
 5'AGGTGTCTGTCCTGYCCATTCTCAGKC3' (SEQ ID NO 264)
 5'CAGGTGTCTGTCCTGYCCATTCTCAGKC3' (SEQ ID NO 265)
 5'CCAGGTGTCTGTCCTGYCCATTCTCAGKC3' (SEQ ID NO 266)
 5'TCACATGGGTGGTCCTAGG3' (SEQ ID NO 267)
 5'GTCACATGGGTGGTCCTAGG3' (SEQ ID NO 268)
 5'GGTCACATGGGTGGTCCTAGG3' (SEQ ID NO 269)
 5'TGGTCACATGGGTGGTCCTAGG3' (SEQ ID NO 270)
 5'CTGGTCACATGGGTGGTCCTAGG3' (SEQ ID NO 271)
 5'KCTGGTCACATGGGTGGTCCTAGG3' (SEQ ID NO 272)
 5'GKCTGGTCACATGGGTGGTCCTAGG3' (SEQ ID NO 273)
 5'TSCCATGARAGATGCMAAGC3' (SEQ ID NO 274)
 5'GTSCCATGARAGATGCMAAGC3' (SEQ ID NO 275)
 5'TGTSCCATGARAGATGCMAAGC3' (SEQ ID NO 276)
 5'GTGTSCCATGARAGATGCMAAGC3' (SEQ ID NO 277)
 5'GGTGTSCCATGARAGATGCMAAGC3' (SEQ ID NO 278)
 5'GGGTGTSCCATGARAGATGCMAAGC3' (SEQ ID NO 279)
 5'GAWTTTCTGACTCTTCCCA3' (SEQ ID NO 280)
 5'TGAWTTTCTGACTCTTCCCA3' (SEQ ID NO 281)
 5'CTGAWTTTCTGACTCTTCCCA3' (SEQ ID NO 282)
 5'CCTGAWTTTCTGACTCTTCCCA3' (SEQ ID NO 283)
 5'GCCTGAWTTTCTGACTCTTCCCA3' (SEQ ID NO 284)
 5'CGCCTGAWTTTCTGACTCTTCCCA3' (SEQ ID NO 285)
- 30 - for the amplification of exon 2 of HLA-C (table 9):
 5'GTCGAGGGTCTGGGCGGGTT3' (SEQ ID NO 126)
 5'GGTCGAGGGTCTGGGCGGGTT3' (SEQ ID NO 127)
 5'CGGTCGAGGGTCTGGGCGGGTT3' (SEQ ID NO 128)
 5'CCGGTCGAGGGTCTGGGCGGGTT3' (SEQ ID NO 129)

5'YCCGGTCGAGGGTCTGGGCGGGTT3' (SEQ ID NO 130)

5'CYCCGGTCGAGGGTCTGGGCGGGTT3' (SEQ ID NO 131)

- for the amplification of exon 3 of HLA-C (table 10):

5 5'CGCCCCRAGTCTCCSSGTCT3' (SEQ ID NO 132)

5'TCGCCCCRAGTCTCCSSGTCT3' (SEQ ID NO 133)

5'GTCGCCCCRAGTCTCCSSGTCT3' (SEQ ID NO 134)

5'GGTCGCCCCRAGTCTCCSSGTCT3' (SEQ ID NO 135)

5'GGGTCGCCCCRAGTCTCCSSGTCT3' (SEQ ID NO 136)

10 5'CGGGTCGCCCCRAGTCTCCSSGTCT3' (SEQ ID NO 137)

5'CGRCCGGRGAGAGCCCCAGT3' (SEQ ID NO 138)

5'TCGRCCGGRGAGAGCCCCAGT3' (SEQ ID NO 139)

5'CTCGRCCGGRGAGAGCCCCAGT3' (SEQ ID NO 140)

5'CCTCGRCCGGRGAGAGCCCCAGT3' (SEQ ID NO 141)

15 5'CCCTCGRCCGGRGAGAGCCCCAGT3' (SEQ ID NO 142)

5'ACCCTCGRCCGGRGAGAGCCCCAGT3' (SEQ ID NO 143)

- for the amplification of exon 4 of HLA-C (table 11):

5'GTGCCTGTGTCCAGGCTGGC3' (SEQ ID NO 286)

20 5'GGTGCCTGTGTCCAGGCTGGC3' (SEQ ID NO 287)

5'AGGTGCCTGTGTCCAGGCTGGC3' (SEQ ID NO 288)

5'CAGGTGCCTGTGTCCAGGCTGGC3' (SEQ ID NO 289)

5'CCAGGTGCCTGTGTCCAGGCTGGC3' (SEQ ID NO 290)

5'CCCAGGTGCCTGTGTCCAGGCTGGC3' (SEQ ID NO 291)

25 5'TGGCGTCTGGGTTCTGTGCC3' (SEQ ID NO 292)

5'CTGGCGTCTGGGTTCTGTGCC3' (SEQ ID NO 293)

5'GCTGGCGTCTGGGTTCTGTGCC3' (SEQ ID NO 294)

5'GGCTGGCGTCTGGGTTCTGTGCC3' (SEQ ID NO 295)

5'AGGCTGGCGTCTGGGTTCTGTGCC3' (SEQ ID NO 296)

30 5'CAGGCTGGCGTCTGGGTTCTGTGCC3' (SEQ ID NO 297)

5'CTCAGGATRGTCACATGGSC3' (SEQ ID NO 298)

5'TCTCAGGATRGTCACATGGSC3' (SEQ ID NO 299)

5'TTCTCAGGATRGTCACATGGSC3' (SEQ ID NO 300)

5'RTTCTCAGGATRGTCACATGGSC3' (SEQ ID NO 301)

5'CRTTCTCAGGATRGTCACATGGSC3' (SEQ ID NO 302)
 5'CCRTTCTCAGGATRGTCACATGGSC3' (SEQ ID NO 303)
 5'GCGCTGTTGGAGTGTCGCAA3' (SEQ ID NO 78)
 5'GGCGCTGTTGGAGTGTCGCAA3' (SEQ ID NO 79)
 5 5'GGGCGCTGTTGGAGTGTCGCAA3' (SEQ ID NO 80)
 5'TGGGCGCTGTTGGAGTGTCGCAA3' (SEQ ID NO 81)
 5'ATGGGCGCTGTTGGAGTGTCGCAA3' (SEQ ID NO 82)
 5'CATGGGCGCTGTTGGAGTGTCGCAA3' (SEQ ID NO 83)
 5'SCAAGAGAGAWRCAAAGTGT3' (SEQ ID NO 304)
 10 5'CSCAAGAGAGAWRCAAAGTGT3' (SEQ ID NO 305)
 5'TCSCAAGAGAGAWRCAAAGTGT3' (SEQ ID NO 306)
 5'GTCSCAAGAGAGAWRCAAAGTGT3' (SEQ ID NO 307)
 5'TGTCSCAAGAGAGAWRCAAAGTGT3' (SEQ ID NO 308)
 5'GTGTCSCAAGAGAGAWRCAAAGTGT3' (SEQ ID NO 309)

15

5. Method according to any of claims 1 to 4 further characterized that the amplification of exon 2 is carried out with at least one of the following forward primers:

- for HLA-A: 5APBio: B-TTCTCCCCAGACGCCGAGGATGGCC (SEQ ID NO 144); or
- for HLA-B: IBPin1: B-GGGAGGAGCGAGGGGACCSCAG (SEQ ID NO 145); or
- for HLA-C: 5CIN1: B-AGCGAGGGGCCCCGCCCCGGCGA (SEQ ID NO 146).

25

6. Method according to any of claims 1 to 5 further characterized that the amplification of exon 4 is carried out with at least one of the following reverse primers:

- for HLA-A: 3ex4APbio: B-TTGGGCAGACCCTCATGCTGC (SEQ ID NO 311); or
- for HLA-B: 3ex4IBbio: B-TCGGCAGCCCCTCATGCTGT (SEQ ID NO 312); or

30

- for HLA-C: 3ex4ICbio: B-CATCTCAGGGTGMRRGGGCTT (SEQ ID NO 313).

7. Method according to any of claims 1 to 6 further characterized that:

- 5 - the amplification of exon 2 is carried out with at least one of the following primers sets:
 - for HLA-A: 5APbio (B-TTCTCCCCAGACGCCGAGGATGGCC; SEQ ID NO 144) and 3ex2Apbio (B-ATCTCGGACCCGGAGACTGT; SEQ ID NO 1);
 - 10 • for HLA-B: IBPin1 (B-GGGAGGAGCGAGGGGACCSCAG; SEQ ID NO 145) and IB3Pin2bio (B-AACCCGCGGGGATTTTGGCCTC; SEQ ID NO 109);
 - for HLA-C: 5CIN1 (B-AGCGAGGGGCCCGCCCGGCGA; SEQ ID NO 146) and IC3Pin2bio (B-15 GGTCGAGGGTCTGGGCGGGTT; SEQ ID NO 127);
- the amplification of exon 3 is carried out with at least one of the following primer sets:
 - for HLA-A: 5ex3APbio (B-CAGTTTAGGCCAAAAATCCCCC; SEQ ID NO 104) and 3ex3APbio (B-20 CCCTCCTTGTGGGAGGCCAG; SEQ ID NO 156);
 - for HLA-B: IB5Pin 2bio (B-CGCGTTTACCCGGTTTCATTTTCAGTTG; SEQ ID NO 224) and IB3Pin3bio (B-TCTTCTCGTKGGAGSCCATCCCC; SEQ ID NO 234);
 - 25 • for HLA-C: IC5Pin2bio (B-TCGRCCGGRGAGAGCCCCAGT; SEQ ID NO 139) and 3CIN3 (B-GGAGATGGGGAAGGCTCCCCACT; SEQ ID NO 149);
- the amplification of exon 4 is carried out with at least one of the following primer sets:
 - 30 • for HLA-A: 5ex4APbio (B-GTTCTGTGCTCYCTTCCCCAT; SEQ ID NO 205) and 3ex4APbio (B-TTGGGCAGACCCTCATGCTGC ; SEQ ID NO 311);

- for HLA-B: 5ex4IBbio (B-TCACATGGGTGGTCCTAGG; SEQ ID NO 267) and 3ex4IBbio (B-TCGGCAGCCCCTCATGCTGT; SEQ ID NO 312);
 - for HLA-C: 5ex4ICbio (B-TCTCAGGATRGTCACATGGSC; SEQ ID NO 299) and 3ex4ICbio (B-CATCTCAGGGTGMRGGGCTT; SEQ ID NO 313).
- 5
8. Method according to any of claims 1 to 7 further characterized that both exon 2 and exon 3 of HLA-A, HLA-B or HLA-C are amplified by use of a multiplex primer mix containing at least one primer pair for the amplification of exon 2 and at least one primer pair for the amplification of exon 3.
- 10
9. Method according to any of claims 1 to 7 further characterized that all three exons, exon 2, exon 3 and exon 4, of HLA-A, HLA-B or HLA-C are amplified by use of a multiplex primer mix containing at least one primer pair for the amplification of exon 2, at least one primer pair for the amplification of exon 3 and at least one primer pair for the amplification of exon 4.
- 15
10. A primer as defined by any of claims 1 to 7, for use in the amplification of exon 2 of HLA-A, HLA-B or HLA-C alleles.
- 20
11. A primer as defined by any of claims 1 to 7, for use in the amplification of exon 3 of HLA-A, HLA-B or HLA-C alleles.
- 25
12. A primer as defined by any of claims 1 to 7, for use in the amplification of exon 4 of HLA-A, HLA-B or HLA-C alleles.
- 30
13. A primer set consisting of a combination of a forward and a reverse primer as defined in any of claims 1 to 7, for use in the amplification of exon 2 of HLA-A, HLA-B or HLA-C alleles.

14. A primer set consisting of a combination of a forward and a reverse primer as defined in any of claims 1 to 7, for use in the amplification of exon 3 of HLA-A, HLA-B or HLA-C alleles.
- 5 15. A primer set consisting of a combination of a forward and a reverse primer as defined in any of claims 1 to 7, for use in the amplification of exon 4 of HLA-A, HLA-B or HLA-C alleles.
- 10 16. A multiplex primer mix containing at least one primer set according to claim 13 for the amplification of exon 2 and one primer set according to claim 14 for amplification of exon 3.
- 15 17. A multiplex primer mix containing at least one primer set according to claim 13 for the amplification of exon 2, one primer set according to claim 14 for amplification of exon 3 and one primer set according to claim 15 for the amplification of exon 4.
- 20 18. Method for the typing or subtyping of one or more HLA-A, HLA-B or HLA-C alleles in a sample comprising the following steps:
(i) if needed, release, isolation and/or concentration of the nucleic acids present in said sample;
(ii) amplification of the nucleic acids according to any of claims 1 to 9;
(iii) typing of the specific HLA-A, HLA-B or HLA-C allele present in said sample.
- 25 19. Method according to claim 18 further characterized that the typing step is carried out by hybridization with one or more suitable probes.
- 30 20. A diagnostic kit for the typing or subtyping of one or more HLA-A, HLA-B or HLA-C alleles in a sample comprising the following components:
(i) when appropriate, a means for releasing, isolating or concentrating the nucleic acids present in said sample;
(ii) a primer set or a primer mix according to any of claims 13 to 17;

- (iii) a means for the typing of the specific HLA-A, HLA-B or HLA-C allele present in said sample.
21. A line probe assay for the typing or subtyping of one or more HLA-A, HLA-B or HLA-C alleles in a sample comprising the following components:
- 5 (i) when appropriate, a means for releasing, isolating or concentrating the nucleic acids present in said sample;
- (ii) a primer pair or a primer mix according to any of claims 13 to 17;
- (iii) at least one probe that specifically hybridizes with exon 2, exon 3 or exon 4
- 10 of HLA-A, HLA-B or HLA-C, fixed to a solid support;
- (iv) a hybridization buffer, or components necessary for producing said buffer;
- (v) a wash solution, or components necessary for producing said solution;
- (vi) when appropriate, a means for detecting the hybrids resulting from the preceding hybridization.




Fig 1

41 51 61 71 81
TCCCCCACGGACGGGCCAGGTCTCGCCACAGTCTCCGGGTCCGAGAT




Figure 1 - cont'd 1

CCACCCCGAAGCCGCGGGACCCCGAGACCCTTGCCCCGGGAGAGGCC




Figure 1 - cont'd 2

AGGCGCCTTTACCCGGTTTCATTTTCAGTTTAGGCCAAAAATCCCCCG




Figure 1 - cont'd 3

										Seq. ID	Nos.
GCGGGG ²³¹ TCGGGG ²⁴¹ CCAG											315
						C					316
						C					317
						C					318
						C					319
						C					320
						C					321
						C					322
T						C					323
T						C					324
T						C					325
T						C					326
T						C					327
											328
											329
											330
											331
											332
											333
											334
											335
											336
											337
											338
T											339
T											340
T											341
T											342
T											343
T											344

Figure 1 - cont'd 5




Figure 2

31 41 51 61 71
CACGACNNNTCCCCATCCCCACGTACGGCCCGGGTCGNCCCCGAGT




Figure 2 - cont'd 1

121 131 141 151 161
ACCCTCGACCGGCGAGAGCCCCAGGCGCGTTTACCCGGTTTCATTT




Figure 2 - cont'd 3

171 181 191 201
TCAGTTGAGGCCAAAATCCCCGCGGGTTGGTCGGGGCGGGGCGGGGC




Figure 2 - cont'd of 4

211		221		231		241		Seq. ID Nos.	
TCGGNNNNGGGGACGGGGCTGACCGCGGGGCCGGGGCCAGN									
							T		345
GG	CTCG						T		346
									347
									348
		T							349
			A						350
									351
								G	352
									353
									354
									355
									356
									357
									358
									359
			T						360
			T						361
			T						362
			T						363
		T							364
						G			365
						G			366
						G			367
									368
									369
							T		370
							T		371
							T		372
							T		373
		T		T					374
		T				A		N	375
						N			376
		T	T					A	377
		T				G			378
								A	379
		T				G		G	380
N		N		NN			N		381
		T	N		A	G	T		382

Figure 2 - cont'd 5




Figure 3

41 51 61 71
GACCCCTCCCCATCCCCACGGACGGCCCGGGTCGCCCCGAGTCT




Figure 3- cont'd 1

81 91 101 111
 CCCGGTCTGAGATCCACCCGAGGNNNNNCTGCGGAACCCGCCCAGAC




Figure 3 - cont'd 2

121 131 141 151 161
CCTCGACCGGAGAGAGCCCCAGTCACCTTTACCCGGTTTCATTTT




Figure 3- cont'd 3

171 181 191 201

CAGTTTAGGCCAAAATCCCCGCGGGT TGGTCGGGGCTGGGGCGGGG

[illegible]

Figure 3 - cont'd 4

211	221	231	241	Seq. ID Nos.
CTCGGGGGACGGGGCTGACCACGGGGGCGGGGCCAG				
	C			383
				384
				385
				386
				387
				388
				389
			G	390
			G	391
				392
				393
	T		G	394
C		T		395

Figure 3- cont'd 5




Figure 4




Figure 4- cont'd 1




Figure 4 - cont'd 2




Figure 4 - cont'd 4

321 331 341 351 361
TCTCAGGCCCTTGTTCTCTGCTTCACACTCAATGTGTGGGGGCTGTAGTCCAGCACTTCTGAG

Figure 4- cont'd 5

371
 TCCCTCAGCCTCCACTCAGGTCAGGACCAAGTCTGGTCTCCCTCTTCAGGGACTAGAAATTT
 381
 391
 401
 411
 421
 431
 441
 451
 461
 471
 481
 491
 501
 511
 521
 531
 541
 551
 561
 571
 581
 591
 601
 611
 621
 631
 641
 651
 661
 671
 681
 691
 701
 711
 721
 731
 741
 751
 761
 771
 781
 791
 801
 811
 821
 831
 841
 851
 861
 871
 881
 891
 901
 911
 921
 931
 941
 951
 961
 971
 981
 991
 1001
 1011
 1021
 1031
 1041
 1051
 1061
 1071
 1081
 1091
 1101
 1111
 1121
 1131
 1141
 1151
 1161
 1171
 1181
 1191
 1201
 1211
 1221
 1231
 1241
 1251
 1261
 1271
 1281
 1291
 1301
 1311
 1321
 1331
 1341
 1351
 1361
 1371
 1381
 1391
 1401
 1411
 1421
 1431
 1441
 1451
 1461
 1471
 1481
 1491
 1501
 1511
 1521
 1531
 1541
 1551
 1561
 1571
 1581
 1591
 1601
 1611
 1621
 1631
 1641
 1651
 1661
 1671
 1681
 1691
 1701
 1711
 1721
 1731
 1741
 1751
 1761
 1771
 1781
 1791
 1801
 1811
 1821
 1831
 1841
 1851
 1861
 1871
 1881
 1891
 1901
 1911
 1921
 1931
 1941
 1951
 1961
 1971
 1981
 1991
 2001
 2011
 2021
 2031
 2041
 2051
 2061
 2071
 2081
 2091
 2101
 2111
 2121
 2131
 2141
 2151
 2161
 2171
 2181
 2191
 2201
 2211
 2221
 2231
 2241
 2251
 2261
 2271
 2281
 2291
 2301
 2311
 2321
 2331
 2341
 2351
 2361
 2371
 2381
 2391
 2401
 2411
 2421
 2431
 2441
 2451
 2461
 2471
 2481
 2491
 2501
 2511
 2521
 2531
 2541
 2551
 2561
 2571
 2581
 2591
 2601
 2611
 2621
 2631
 2641
 2651
 2661
 2671
 2681
 2691
 2701
 2711
 2721
 2731
 2741
 2751
 2761
 2771
 2781
 2791
 2801
 2811
 2821
 2831
 2841
 2851
 2861
 2871
 2881
 2891
 2901
 2911
 2921
 2931
 2941
 2951
 2961
 2971
 2981
 2991
 3001
 3011
 3021
 3031
 3041
 3051
 3061
 3071
 3081
 3091
 3101
 3111
 3121
 3131
 3141
 3151
 3161
 3171
 3181
 3191
 3201
 3211
 3221
 3231
 3241
 3251
 3261
 3271
 3281
 3291
 3301
 3311
 3321
 3331
 3341
 3351
 3361
 3371
 3381
 3391
 3401
 3411
 3421
 3431
 3441
 3451
 3461
 3471
 3481
 3491
 3501
 3511
 3521
 3531
 3541
 3551
 3561
 3571
 3581
 3591
 3601
 3611
 3621
 3631
 3641
 3651
 3661
 3671
 3681
 3691
 3701
 3711
 3721
 3731
 3741
 3751
 3761
 3771
 3781
 3791
 3801
 3811
 3821
 3831
 3841
 3851
 3861
 3871
 3881
 3891
 3901
 3911
 3921
 3931
 3941
 3951
 3961
 3971
 3981
 3991
 4001
 4011
 4021
 4031
 4041
 4051
 4061
 4071
 4081
 4091
 4101
 4111
 4121
 4131
 4141
 4151
 4161
 4171
 4181
 4191
 4201
 4211
 4221
 4231
 4241
 4251
 4261
 4271
 4281
 4291
 4301
 4311
 4321
 4331
 4341
 4351
 4361
 4371
 4381
 4391
 4401
 4411
 4421
 4431
 4441
 4451
 4461
 4471
 4481
 4491
 4501
 4511
 4521
 4531
 4541
 4551
 4561
 4571
 4581
 4591
 4601
 4611
 4621
 4631
 4641
 4651
 4661
 4671
 4681
 4691
 4701
 4711
 4721
 4731
 4741
 4751
 4761
 4771
 4781
 4791
 4801
 4811
 4821
 4831
 4841
 4851
 4861
 4871
 4881
 4891
 4901
 4911
 4921
 4931
 4941
 4951
 4961
 4971
 4981
 4991
 5001
 5011
 5021
 5031
 5041
 5051
 5061
 5071
 5081
 5091
 5101
 5111
 5121
 5131
 5141
 5151
 5161
 5171
 5181
 5191
 5201
 5211
 5221
 5231
 5241
 5251
 5261
 5271
 5281
 5291
 5301
 5311
 5321
 5331
 5341
 5351
 5361
 5371
 5381
 5391
 5401
 5411
 5421
 5431
 5441
 5451
 5461
 5471
 5481
 5491
 5501
 5511
 5521
 5531
 5541
 5551
 5561
 5571
 5581
 5591
 5601
 5611
 5621
 5631
 5641
 5651
 5661
 5671
 5681
 5691
 5701
 5711
 5721
 5731
 5741
 5751
 5761
 5771
 5781
 5791
 5801
 5811
 5821
 5831
 5841
 5851
 5861
 5871
 5881
 5891
 5901
 5911
 5921
 5931
 5941
 5951
 5961
 5971
 5981
 5991
 6001
 6011
 6021
 6031
 6041
 6051
 6061
 6071
 6081
 6091
 6101
 6111
 6121
 6131
 6141
 6151
 6161
 6171
 6181
 6191
 6201
 6211
 6221
 6231
 6241
 6251
 6261
 6271
 6281
 6291
 6301
 6311
 6321
 6331
 6341
 6351
 6361
 6371
 6381
 6391
 6401
 6411
 6421
 6431
 6441
 6451
 6461
 6471
 6481
 6491
 6501
 6511
 6521
 6531
 6541
 6551
 6561
 6571
 6581
 6591
 6601
 6611
 6621
 6631
 6641
 6651
 6661
 6671
 6681
 6691
 6701
 6711
 6721
 6731
 6741
 6751
 6761
 6771
 6781
 6791
 6801
 6811
 6821
 6831
 6841
 6851
 6861
 6871
 6881
 6891
 6901
 6911
 6921
 6931
 6941
 6951
 6961
 6971
 6981
 6991
 7001
 7011
 7021
 7031
 7041
 7051
 7061
 7071
 7081
 7091
 7101
 7111
 7121
 7131
 7141
 7151
 7161
 7171
 7181
 7191
 7201
 7211
 7221
 7231
 7241
 7251
 7261
 7271
 7281
 7291
 7301
 7311
 7321
 7331
 7341
 7351
 7361
 7371
 7381
 7391
 7401
 7411
 7421
 7431
 7441
 7451
 7461
 7471
 7481
 7491
 7501
 7511
 7521
 7531
 7541
 7551
 7561
 7571
 7581
 7591
 7601
 7611
 7621
 7631
 7641
 7651
 7661
 7671
 7681
 7691
 7701
 7711
 7721
 7731
 7741
 7751
 7761
 7771
 7781
 7791
 7801
 7811
 7821
 7831
 7841
 7851
 7861
 7871
 7881
 7891
 7901
 7911
 7921
 7931
 7941
 7951
 7961
 7971
 7981
 7991
 8001
 8011
 8021
 8031
 8041
 8051
 8061
 8071
 8081
 8091
 8101
 8111
 8121
 8131
 8141
 8151
 8161
 8171
 8181
 8191
 8201
 8211
 8221
 8231
 8241
 8251
 8261
 8271
 8281
 8291
 8301
 8311
 8321
 8331
 8341
 8351
 8361
 8371
 8381
 8391
 8401
 8411
 8421
 8431
 8441
 8451
 8461
 8471
 8481
 8491
 8501
 8511
 8521
 8531
 8541
 8551
 8561
 8571
 8581
 8591
 8601
 8611
 8621
 8631
 8641
 8651
 8661
 8671
 8681
 8691
 8701
 8711
 8721
 8731
 8741
 8751
 8761
 8771
 8781
 8791
 8801
 8811
 8821
 8831
 8841
 8851
 8861
 8871
 8881
 8891
 8901
 8911
 8921
 8931
 8941
 8951
 8961
 8971
 8981
 8991
 9001
 9011
 9021
 9031
 9041
 9051
 9061
 9071
 9081
 9091
 9101
 9111
 9121
 9131
 9141
 9151
 9161
 9171
 9181
 9191
 9201
 9211
 9221
 9231
 9241
 9251
 9261
 9271
 9281
 9291
 9301
 9311
 9321
 9331
 9341
 9351
 9361
 9371
 9381
 9391
 9401
 9411
 9421
 9431
 9441
 9451
 9461
 9471
 9481
 9491
 9501
 9511
 9521
 9531
 9541
 9551
 9561
 9571
 9581
 9591
 9601
 9611
 9621
 9631
 9641
 9651
 9661
 9671
 9681
 9691
 9701
 9711
 9721
 9731
 9741
 9751
 9761
 9771
 9781
 9791
 9801
 9811
 9821
 9831
 9841
 9851
 9861
 9871
 9881
 9891
 9901
 9911
 9921
 9931
 9941
 9951
 9961
 9971
 9981
 9991
 10001
 10011
 10021
 10031
 10041
 10051
 10061
 10071
 10081
 10091
 10101
 10111
 10121
 10131
 10141
 10151
 10161
 10171
 10181
 10191
 10201
 10211
 10221
 10231
 10241
 10251
 10261
 10271
 10281
 10291
 10301
 10311
 10321
 10331
 10341
 10351
 10361
 10371
 10381
 10391
 10401
 10411
 10421
 10431
 10441
 10451
 10461
 10471
 10481
 10491
 10501
 10511
 10521
 10531
 10541
 10551
 10561
 10571
 10581
 10591
 10601
 10611
 10621
 10631
 10641
 10651
 10661
 10671
 10681
 10691
 10701
 10711
 10721
 10731
 10741
 10751
 10761
 10771
 10781
 10791
 10801
 10811
 10821
 10831
 10841
 10851
 10861
 10871
 10881
 10891
 10901
 10911
 10921
 10931
 10941
 10951
 10961
 10971
 10981
 10991
 11001
 11011
 11021
 11031
 11041
 11051
 11061
 11071
 11081
 11091
 11101
 11111
 11121
 11131
 11141
 11151
 11161
 11171
 11181
 11191
 11201
 11211
 11221
 11231
 11241
 11251
 11261
 11271
 11281
 11291
 11301
 11311
 11321
 11331
 11341
 11351
 11361
 11371
 11381
 11391
 11401
 11411
 11421
 11431
 11441
 11451
 11461
 11471
 11481
 11491
 11501
 11511
 11521
 11531
 11541
 11551
 11561
 11571
 11581
 11591
 11601
 11611
 11621
 11631
 11641
 11651
 11661
 11671
 11681
 11691
 11701
 11711
 11721
 11731
 11741
 11751
 11761
 11771
 11781
 11791
 11801
 11811
 11821
 11831
 11841
 11851
 11861
 11871
 11881
 11891
 11901
 11911
 11921
 11931
 11941
 11951
 11961
 11971
 11981
 11991
 12001
 12011
 12021
 12031
 12041
 12051
 12061
 12071
 12081
 12091
 12101
 12111
 12121
 12131
 12141
 12151
 12161
 12171
 12181
 12191
 12201
 12211
 12221
 12231
 12241
 12251
 12261
 12271
 12281
 12291
 12301
 12311
 12321
 12331
 12341
 12351
 12361
 12371
 12381
 12391
 12401
 12411
 12421
 12431
 12441
 12451
 12461
 12471
 12481
 12491
 12501
 12511
 12521
 12531
 12541
 12551
 12561
 12571
 12581
 12591
 12601
 12611
 12621
 12631
 12641
 12651
 12661
 12671
 12681
 12691
 12701
 12711
 12721
 12731
 12741
 12751
 12761
 12771
 12781
 12791
 12801
 12811
 12821
 12831
 12841
 12851
 12861
 12871
 12881
 12891
 12901
 12911
 12921
 12931
 12941
 12951
 12961
 12971
 12981
 12991
 13001
 13011
 13021
 13031
 13041
 13051
 13061
 13071
 13081
 13091
 13101
 13111
 13121
 13131
 13141
 13151
 13161
 13171
 13181
 13191
 13201
 13211
 13221
 13231
 13241
 13251
 13261
 13271
 13281
 13291
 13301
 13311
 13321
 13331
 13341
 13351
 13361
 13371
 13381
 13391
 13401
 13411
 13421
 13431
 13441
 13451
 13461
 13471
 13481
 13491
 13501
 13511
 13521
 13531
 13541
 13551
 13561
 13




Figure 5




Figure 5- cont'd 1

NGANGAATGGCATGAGTTTCTCCTGAGTTTCCCTCTGAGGGCCCCCTTCTCTCTCTAGGACAA

121


131

141

151

161

171



A
G




Figure 5- cont'd 3




Figure 5- cont'd 4




Figure 5- cont'd 5

361 371 381 391 401 411
ACTCAGATCAGGAGCAGAAAGTCCCTGTTCCCGCTCAGAGACTCGAACTTCCCAATGAA




Figure 5-cont'd 6




Figure 5- cont'd 8




Figure 6




Figure 6 - cont'd 1




Figure 6 - cont'd 2




Figure 6 - cont'd 4




Figure 6 - cont'd 5




Figure 6 - cont'd 6

⁴⁸¹TCCAA⁴⁸¹TAGGAG⁴⁸¹ATTATCC⁴⁸¹AGGTG⁴⁸¹CTG⁴⁸¹TCCAGG⁴⁸¹CTGG⁴⁸¹CTGG⁴⁸¹GTCT⁴⁸¹GG⁴⁸¹GTCT⁴⁸¹GTGCC




Figure 6 - cont'd 4

481 491 501 511 521 531
 CCCTTCCCCACCCCAAGGTGTCCTGTCCTCAATCTCAGGATGGTCACATGGGCGCTGTTGGA




Figure 6 - cont'd 8

49/50




Fig 7

50/50




Fig 8