



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, C07K 14/47, C12N 5/10, C07K 16/18, A61K 38/17, C12Q 1/68	A1	(11) International Publication Number: WO 00/05368 (43) International Publication Date: 3 February 2000 (03.02.00)
(21) International Application Number: PCT/US99/12385 (22) International Filing Date: 19 July 1999 (19.07.99) (30) Priority Data: 60/160,076 20 July 1998 (20.07.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/119,547 (CIP) Filed on 20 July 1998 (20.07.98) (71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). CORLEY, Neil, C. [US/US]; 1240 Dale Avenue #30, Mountain View, CA 94040 (US). GUEGLER, Karl, J. [CH/US]; 1048 Oakland Avenue, Menlo Park, CA 94025 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577	(US). JUNMING, Yang [CN/US]; 7136 Clarendon Street, San Jose, CA 95129 (US). (74) Agents: BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>	
(54) Title: HUMAN CALCIUM REGULATORY PROTEINS (57) Abstract The invention provides human calcium regulatory proteins (CaREG) and polynucleotides which identify and encode CaREG. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating or preventing disorders associated with expression of CaREG.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

HUMAN CALCIUM REGULATORY PROTEINS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of calcium regulatory
5 proteins and to the use of these sequences in the diagnosis, treatment, and prevention of
neurological and cardiovascular disorders and cancer.

BACKGROUND OF THE INVENTION

In nearly all eukaryotic cells, calcium (Ca^{2+}) functions as an intracellular signaling
10 molecule in diverse cellular processes including cell proliferation, neurotransmitter secretion,
glycogen metabolism, and muscle contraction. Within a resting cell, the cytosolic concentration
of Ca^{2+} is less than 10^{-7} M. However, when the cell is stimulated by an external signal, such as a
neural impulse or a growth factor, the cytosolic concentration of Ca^{2+} increases by about 50-fold.
This influx of Ca^{2+} is prompted by the opening of plasma membrane Ca^{2+} channels and by the
15 release of Ca^{2+} from intracellular stores such as the endoplasmic reticulum. Ca^{2+} directly activates
regulatory enzymes, such as protein kinase C, which trigger signal transduction pathways. Ca^{2+}
also binds to specific Ca^{2+} binding proteins which then activate multiple target proteins including
enzymes, membrane transport pumps, and ion channels.

Some Ca^{2+} binding proteins are characterized by the presence of one or more EF-hand
20 Ca^{2+} binding motifs. (Reviewed in Celio, M. R. et al. (1996) Guidebook to the Calcium-Binding
Proteins, Oxford University Press, New York, NY, pp. 15-20.) The EF-hand motif is comprised
of 12 amino acids flanked by α -helices. Acidic residues are generally conserved at positions 1, 3,
5, 7, 9, and 12, and glycine is often conserved at position 6. At least 250 EF-hand proteins
defining 39 families have been identified in various tissues. For example, S100 α is a member of
25 the S100 protein family isolated from human heart. (Celio et al. supra, pp. 135-136; Engelkamp,
D. et al. (1992) *Biochemistry* 31:10258-10264.) S100 α messenger RNA expression is restricted to
the heart, skeletal muscle, and brain. Normally, S100 α is undetectable in serum; however, S100 α
is detectable in serum of patients with renal carcinoma at levels correlated with clinical
progression of the disease. In addition, S100 α levels are increased in serum following myocardial
30 infarction.

Other Ca^{2+} binding proteins include those involved in Ca^{2+} sequestration and protein
folding in the endoplasmic reticulum (ER). For example, Ca^{2+} binding proteins 1 and 2 (CaBP1
and CaBP2) are protein disulfide isomerases that contain two and three thioredoxin-like active

sites, respectively. (Fullekrug, J. et al. (1994) J. Cell Sci. (1994) 107:2719-2727; Van, P. N. et al. (1993) Eur. J. Biochem. 213:789-795.) Each active site contains two invariant cysteines which directly participate in reversible oxidation/reduction reactions. CaBP1 and CaBP2 may facilitate the folding of nascent proteins in the ER by catalyzing the formation and isomerization of disulfide bonds. Like all resident soluble ER proteins, CaBP1 and CaBP2 each contain the C-terminal KDEL/KEEL tetrapeptide required for their retention in the ER.

Other Ca^{2+} binding proteins include the annexins which comprise a ubiquitous family of Ca^{2+} -dependent phospholipid binding proteins. (Celio et al. supra, pp. 181-184.) There are eleven structurally distinct classes of annexins characterized by a variable number of core domain repeats. Although the precise function of the annexins are unknown, evidence suggests that annexins are involved in intracellular signaling, vesicular transport, and cytoskeletal function.

Regulation of intracellular Ca^{2+} levels by Ca^{2+} binding proteins is critical for neural signaling and muscle contraction. In particular, elevated levels of intracellular calcium are a major cause of cardiovascular dysfunction, including myocardial ischemia leading to acute infarction. Current therapies for the treatment of myocardial ischemia block the influx of calcium into myocardial and vascular smooth muscle cells.

The discovery of new Ca^{2+} regulatory proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, treatment, and prevention of neurological and cardiovascular disorders and cancer.

SUMMARY OF THE INVENTION

The invention is based on the discovery of new Ca^{2+} regulatory proteins (CaREG), the polynucleotides encoding CaREG, and the use of these compositions for the diagnosis, treatment, or prevention of neurological and cardiovascular disorders and cancer.

The invention features substantially purified polypeptides, Ca^{2+} regulatory proteins, referred to collectively as "CaREG" and individually as "CaREG-1" and "CaREG-2." In one aspect, the invention provides a substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments thereof.

The invention further provides a substantially purified variant having at least 90% amino acid identity to the amino acid sequence of SEQ ID NO:1, SEQ ID NO:2, or to a fragment of any of these sequences. The invention also provides an isolated and purified polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments thereof. The invention also includes an isolated and purified

polynucleotide variant having at least 70% polynucleotide sequence identity to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments thereof.

Additionally, the invention provides an isolated and purified polynucleotide which
5 hybridizes under stringent conditions to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments thereof, as well as an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide encoding the polypeptide comprising the amino acid
10 sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments thereof.

The invention also provides an isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, and fragments thereof. The invention further provides an isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity to a polynucleotide sequence selected from
15 the group consisting of SEQ ID NO:3, SEQ ID NO:4, and fragments thereof, as well as an isolated and purified polynucleotide having a sequence which is complementary to a polynucleotide sequence selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, and fragments thereof.

The invention further provides an expression vector containing at least a fragment of the
20 polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments thereof. In another aspect, the expression vector is contained within a host cell.

The invention also provides a method for producing a polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments
25 thereof, the method comprising the steps of: (a) culturing the host cell containing an expression vector containing at least a fragment of a polynucleotide encoding the polypeptide under conditions suitable for the expression of the polypeptide; and (b) recovering the polypeptide from the host cell culture.

The invention also provides a pharmaceutical composition comprising a substantially
30 purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

The invention further includes a purified antibody which binds to a polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID

NO:2, and fragments thereof, as well as a purified agonist and a purified antagonist to the polypeptide.

The invention also provides a method for treating or preventing a neurological disorder, the method comprising administering to a subject in need of such treatment an effective amount of
5 a pharmaceutical composition comprising a substantially purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments thereof.

The invention also provides a method for treating or preventing a cardiovascular disorder, the method comprising administering to a subject in need of such treatment an effective amount of
10 a pharmaceutical composition comprising a substantially purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments thereof.

The invention also provides a method for treating or preventing a cancer, the method comprising administering to a subject in need of such treatment an effective amount of an
15 antagonist of the polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments thereof.

The invention also provides a method for detecting a polynucleotide encoding the polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments thereof in a biological sample containing nucleic acids, the
20 method comprising the steps of: (a) hybridizing the complement of the polynucleotide sequence encoding the polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments thereof to at least one of the nucleic acids of the biological sample, thereby forming a hybridization complex; and (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of a
25 polynucleotide encoding the polypeptide in the biological sample. In one aspect, the method further comprises amplifying the polynucleotide prior to hybridization.

BRIEF DESCRIPTION OF THE FIGURES

Figures 1A and 1B show the amino acid sequence (SEQ ID NO:1) and nucleic acid
30 sequence (SEQ ID NO:3) of CaREG-1. The alignment was produced using MACDNASIS PRO software (Hitachi Software Engineering Co. Ltd., Yokohama, Japan).

Figures 2A, 2B, 2C, 2D, 2E, 2F, and 2G show the amino acid sequence (SEQ ID NO:2) and nucleic acid sequence (SEQ ID NO:4) of CaREG-2. The alignment was produced using MACDNASIS PRO software.

Table 1 shows the programs, their descriptions, references, and threshold parameters (where appropriate) used to analyze CaREG.

DESCRIPTION OF THE INVENTION

5 Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors, and reagents described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

10 It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

15 Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods, devices, and materials are now described. All publications mentioned herein are cited for the purpose of
20 describing and disclosing the cell lines, vectors, and methodologies which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

25 DEFINITIONS

"CaREG," as used herein, refers to the amino acid sequences, or variant thereof, of substantially purified CaREG obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and preferably the human species, from any source, whether natural, synthetic, semi-synthetic, or recombinant.

30 The term "agonist," as used herein, refers to a molecule which, when bound to CaREG, increases or prolongs the duration of the effect of CaREG. Agonists may include proteins, nucleic acids, carbohydrates, or any other molecules which bind to and modulate the effect of CaREG.

An "allelic variant," as this term is used herein, is an alternative form of the gene encoding CaREG. Allelic variants may result from at least one mutation in the nucleic acid sequence and

may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. Any given natural or recombinant gene may have none, one, or many allelic forms. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur
5 alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding CaREG, as described herein, include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polynucleotide the same as CaREG or a polypeptide with at least one functional characteristic of CaREG. Included within this definition are polymorphisms which may or may not be readily
10 detectable using a particular oligonucleotide probe of the polynucleotide encoding CaREG, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding CaREG. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent CaREG. Deliberate amino
15 acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of CaREG is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, positively charged amino acids may include lysine and arginine, and amino acids with uncharged polar head groups having similar
20 hydrophilicity values may include leucine, isoleucine, and valine; glycine and alanine; asparagine and glutamine; serine and threonine; and phenylalanine and tyrosine.

The terms "amino acid" or "amino acid sequence," as used herein, refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. In this context, "fragments," "immunogenic
25 fragments," or "antigenic fragments" refer to fragments of CaREG which are preferably at least 5 to about 15 amino acids in length, most preferably at least 14 amino acids, and which retain some biological activity or immunological activity of CaREG. Where "amino acid sequence" is recited herein to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native
30 amino acid sequence associated with the recited protein molecule.

"Amplification," as used herein, relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art. (See, e.g., Dieffenbach, C.W. and G.S. Dveksler (1995) PCR Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainview, NY, pp.1-5.)

The term "antagonist," as it is used herein, refers to a molecule which, when bound to CaREG, decreases the amount or the duration of the effect of the biological or immunological activity of CaREG. Antagonists may include proteins, nucleic acids, carbohydrates, antibodies, or any other molecules which decrease the effect of CaREG.

5 As used herein, the term "antibody" refers to intact molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding the epitopic determinant. Antibodies that bind CaREG polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be
10 derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant," as used herein, refers to that fragment of a molecule
15 (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (given regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

20 The term "antisense," as used herein, refers to any composition containing a nucleic acid sequence which is complementary to the "sense" strand of a specific nucleic acid sequence. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form duplexes and to block either transcription or translation. The designation
25 "negative" can refer to the antisense strand, and the designation "positive" can refer to the sense strand.

As used herein, the term "biologically active," refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic CaREG,
30 or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" or "complementarity," as used herein, refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" binds to the complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules

may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in
5 amplification reactions, which depend upon binding between nucleic acids strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" or a "composition comprising a given amino acid sequence," as these terms are used herein, refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may
10 comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding CaREG or fragments of CaREG may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts, e.g., NaCl, detergents, e.g., sodium dodecyl sulfate (SDS), and other components, e.g.,
15 Denhardt's solution, dry milk, salmon sperm DNA, etc.

"Consensus sequence," as used herein, refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using XL-PCR kit (Perkin-Elmer Corp., Norwalk, CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of more than one Incyte Clone using a computer program for fragment
20 assembly, such as the GELVIEW Fragment Assembly system (GCG, Madison, WI). Some sequences have been both extended and assembled to produce the consensus sequence.

As used herein, the term "correlates with expression of a polynucleotide" indicates that the detection of the presence of nucleic acids, the same or related to a nucleic acid sequence encoding CaREG, by Northern analysis is indicative of the presence of nucleic acids encoding CaREG in a
25 sample, and thereby correlates with expression of the transcript from the polynucleotide encoding CaREG.

A "deletion," as the term is used herein, refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative," as used herein, refers to the chemical modification of a polypeptide
30 sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or

immunological function of the polypeptide from which it was derived.

The term "similarity," as used herein, refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical
5 sequence from hybridizing to a target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or Northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the
10 target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30%
15 similarity or identity). In the absence of non-specific binding, the substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" or "% identity" refer to the percentage of sequence similarity found in a comparison of two or more amino acid or nucleic acid sequences. Percent identity can be determined electronically, e.g., by using the MEGALIGN program (DNASTAR,
20 Inc., Madison WI). The MegAlign™ program can create alignments between two or more sequences according to different methods, e.g., the clustal method. (See, e.g., Higgins, D.G. and P.M. Sharp (1988) *Gene* 73:237-244.) The clustal algorithm groups sequences into clusters by examining the distances between all pairs. The clusters are aligned pairwise and then in groups. The percentage similarity between two amino acid sequences, e.g., sequence A and sequence B, is
25 calculated by dividing the length of sequence A, minus the number of gap residues in sequence A, minus the number of gap residues in sequence B, into the sum of the residue matches between sequence A and sequence B, times one hundred. Gaps of low or of no similarity between the two amino acid sequences are not included in determining percentage similarity. Percent identity between nucleic acid sequences can also be counted or calculated by other methods known in the
30 art, e.g., the Jotun Hein method. (See, e.g., Hein, J. (1990) *Methods Enzymol.* 183:626-645.) Identity between sequences can also be determined by other methods known in the art, e.g., by varying hybridization conditions.

"Human artificial chromosomes" (HACs), as described herein, are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which

contain all of the elements required for stable mitotic chromosome segregation and maintenance.
(See, e.g., Harrington, J.J. et al. (1997) Nat Genet. 15:345-355.)

The term "humanized antibody," as used herein, refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more
5 closely resembles a human antibody, and still retains its original binding ability.

"Hybridization," as the term is used herein, refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing.

As used herein, the term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary
10 bases. A hybridization complex may be formed in solution (e.g., C₀t or R₀t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" or "addition," as used herein, refer to changes in an amino acid or
15 nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively, to the sequence found in the naturally occurring molecule.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which
20 may affect cellular and systemic defense systems.

The term "microarray," as used herein, refers to an arrangement of distinct polynucleotides arrayed on a substrate, e.g., paper, nylon or any other type of membrane, filter, chip, glass slide, or any other suitable solid support.

The terms "element" or "array element" as used herein in a microarray context, refer to
25 hybridizable polynucleotides arranged on the surface of a substrate.

The term "modulate," as it appears herein, refers to a change in the activity of CaREG. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of CaREG.

The phrases "nucleic acid" or "nucleic acid sequence," as used herein, refer to a
30 nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material. In this context, "fragments" refers to those nucleic acid sequences which, comprise a region of unique polynucleotide sequence that specifically identifies

SEQ ID NO:3, SEQ ID NO:4, for example, as distinct from any other sequence in the same genome. For example, a fragment of SEQ ID NO:3, SEQ ID NO:4 is useful in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:3, SEQ ID NO:4 from related polynucleotide sequences. A fragment of SEQ ID NO:3, SEQ ID NO:4 is at least about 15-20 nucleotides in length. The precise length of the fragment of SEQ ID NO:3, SEQ ID NO:4 and the region of SEQ ID NO:3, SEQ ID NO:4 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment. In some cases, a fragment, when translated, would produce polypeptides retaining some functional characteristic, e.g., antigenicity, or structural domain characteristic, e.g., ATP-binding site, of the full-length polypeptide.

The terms "operably associated" or "operably linked," as used herein, refer to functionally related nucleic acid sequences. A promoter is operably associated or operably linked with a coding sequence if the promoter controls the translation of the encoded polypeptide. While operably associated or operably linked nucleic acid sequences can be contiguous and in the same reading frame, certain genetic elements, e.g., repressor genes, are not contiguously linked to the sequence encoding the polypeptide but still bind to operator sequences that control expression of the polypeptide.

The term "oligonucleotide," as used herein, refers to a nucleic acid sequence of at least about 6 nucleotides to 60 nucleotides, preferably about 15 to 30 nucleotides, and most preferably about 20 to 25 nucleotides, which can be used in PCR amplification or in a hybridization assay or microarray. As used herein, the term "oligonucleotide" is substantially equivalent to the terms "amplimer," "primer," "oligomer," and "probe," as these terms are commonly defined in the art.

"Peptide nucleic acid" (PNA), as used herein, refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell. (See, e.g., Nielsen, P.E. et al. (1993) *Anticancer Drug Des.* 8:53-63.)

The term "sample," as used herein, is used in its broadest sense. A biological sample suspected of containing nucleic acids encoding CaREG, or fragments thereof, or CaREG itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a solid support; a tissue; a tissue print; etc.

As used herein, the terms "specific binding" or "specifically binding" refer to that

interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

As used herein, the term "stringent conditions" refers to conditions which permit hybridization between polynucleotides and the claimed polynucleotides. Stringent conditions can be defined by salt concentration, the concentration of organic solvent, e.g., formamide, temperature, and other conditions well known in the art. In particular, stringency can be increased by reducing the concentration of salt, increasing the concentration of formamide, or raising the hybridization temperature.

The term "substantially purified," as used herein, refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably about 75% free, and most preferably about 90% free from other components with which they are naturally associated.

A "substitution," as used herein, refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

"Transformation," as defined herein, describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "variant" of CaREG polypeptides, as used herein, refers to an amino acid sequence that is altered by one or more amino acid residues. The variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties (e.g., replacement of leucine with isoleucine). More rarely, a variant may have "nonconservative" changes (e.g., replacement of glycine with tryptophan). Analogous minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be

substituted, inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, LASERGENE™ software.

The term “variant,” when used in the context of a polynucleotide sequence, may encompass a polynucleotide sequence related to CaREG. This definition may also include, for example, “allelic” (as defined above), “splice,” “species,” or “polymorphic” variants. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or an absence of domains. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass “single nucleotide polymorphisms” (SNPs) in which the polynucleotide sequence varies by one base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

THE INVENTION

The invention is based on the discovery of new human Ca^{2+} regulatory proteins (CaREG), the polynucleotides encoding CaREG, and the use of these compositions for the diagnosis, treatment, or prevention of neurological and cardiovascular disorders and cancer.

Nucleic acids encoding the CaREG-1 of the present invention were first identified in Incyte Clone 1356311 from the fetal lung cDNA library (LUNGNOT09) using a computer search, e.g., BLAST, for amino acid sequence alignments. A consensus sequence, SEQ ID NO:3, was derived from the following overlapping and/or extended nucleic acid sequences: Incyte Clones 1356311H1 and 1356311T6 (LUNGNOT09) and 2220133H1 (LUNGNOT18).

In one embodiment, the invention encompasses a polypeptide comprising the amino acid sequence of SEQ ID NO:1, as shown in Figures 1A and 1B. CaREG-1 is 147 amino acids in length and has one potential N-glycosylation site at N69; two potential cAMP- and cGMP-dependent protein kinase phosphorylation sites at S39 and S54; four potential casein kinase II phosphorylation sites at S7, S54, S71, and S108; and one potential protein kinase C phosphorylation site at S46. BLOCKS and PROFILESCAN analyses show that the region of CaREG-1 from I60 to K144 is similar to S100 Ca^{2+} binding protein signatures. BLOCKS further indicates that the region of CaREG-1 from R104 to F123 is similar to the EF-hand calcium-binding domain. In addition, PRINTS analysis shows that the regions of CaREG-1 from E98 to

D111 and from E115 to L128 are similar to an annexin type I signature. CaREG-1 has chemical and structural similarity with human S100 α (GI 36176). In particular, the region of CaREG-1 from S54 to K147 shares 26% amino acid identity with S100 α . A fragment of SEQ ID NO:3 from about nucleotide 25 to about nucleotide 54 is useful as a hybridization probe. Northern analysis
 5 shows that this sequence is expressed exclusively in cDNA libraries derived from lung tissue, including fetal and cancerous lung tissue.

Nucleic acids encoding the CaREG-2 of the present invention were first identified in Incyte Clone 2176876 from the dermal microvascular endothelial cell cDNA library (ENDCNOT03) using a computer search, e.g., BLAST, for amino acid sequence alignments. A
 10 consensus sequence, SEQ ID NO:4, was derived from the following overlapping and/or extended nucleic acid sequences: Incyte Clones 2176876H1 (ENDCNOT03), 547705X58C1 (BEPINOT01), 1553296F1 (BLADTUT04), 1417868F1 (KIDNNOT09), 1814186F6 (PROSNOT20), 1310941F1 (COLNFET02), 1225462H1 (COLNTUT02), 001150R1 (U937NOT01), and shotgun sequence SADA00307R1.

15 In one embodiment, the invention encompasses a polypeptide comprising the amino acid sequence of SEQ ID NO:2, as shown in Figures 2A, 2B, 2C, 2D, 2E, 2F, and 2G. CaREG-2 is 363 amino acids in length and has one potential cAMP- and cGMP-dependent protein kinase phosphorylation site at S340; ten potential casein kinase II phosphorylation sites at T29, S39, T98, S114, S128, S216, T237, T266, T290, and S295; five potential protein kinase C phosphorylation
 20 sites at T69, T170, S216, S295, and S323; and one potential tyrosine kinase phosphorylation site at Y82. BLOCKS, PRINTS, PFAM, and PROFILESCAN all indicate that the complete amino acid sequence of CaREG-2 shows extensive similarity to thioredoxin protein signatures. In particular, three thioredoxin consensus active sites are conserved in CaREG-2 from F12 to W30, from F140 to W158, and from F273 to W291. Included within each active site region are invariant cysteine
 25 pairs at C20 and C23, C148 and C151, and C281 and C284. A predicted signal peptide from M1 to C23 and an ER retention signal from K360 to L363 suggests that CaREG-2 is a resident ER protein. CaREG-2 has chemical and structural similarity with regions of rat CaBP2 (GI 393203). In particular, CaREG-2 shares 26% overall amino acid identity with CaBP2. A fragment of SEQ ID NO:4 from about nucleotide 163 to about nucleotide 192 is useful as a hybridization probe.
 30 Northern analysis shows the expression of this sequence in various cDNA libraries, at least 63% of which are associated with cancer or proliferating cells and at least 37% of which are associated with the immune response or trauma. Northern analysis further shows that 27% of the libraries expressing CaREG-2 are derived from reproductive tissue, and 17% from cardiovascular tissue. In addition, the highest levels of CaREG-2 expression are observed in libraries derived from

vascular endothelial cells.

The invention also encompasses CaREG variants. A preferred CaREG variant is one which has at least about 80%, more preferably at least about 90%, and most preferably at least about 95% amino acid sequence identity to the CaREG amino acid sequence, and which contains
5 at least one functional or structural characteristic of CaREG.

The invention also encompasses polynucleotides which encode CaREG. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising the sequence of SEQ ID NO:3, which encodes CaREG-1. In a further embodiment, the invention encompasses the polynucleotide sequence comprising the sequence of SEQ ID NO:4, which encodes CaREG-2.

10 The invention also encompasses a variant of a polynucleotide sequence encoding CaREG. In particular, such a variant polynucleotide sequence will have at least about 70%, more preferably at least about 85%, and most preferably at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding CaREG, selected from the group consisting of SEQ ID NO:3 and SEQ ID NO:4. Any one of the polynucleotide variants described above can encode an amino
15 acid sequence which contains at least one functional or structural characteristic of CaREG.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding CaREG, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide
20 sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring CaREG, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode CaREG and its variants are preferably
25 capable of hybridizing to the nucleotide sequence of the naturally occurring CaREG under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding CaREG possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the
30 frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding CaREG and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode CaREG and

CaREG derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding CaREG or any fragment thereof.

5 Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:3, SEQ ID NO:4, and fragments thereof, under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) For example, stringent salt concentration will ordinarily be less
10 than about 750 mM NaCl and 75 mM trisodium citrate, preferably less than about 500 mM NaCl and 50 mM trisodium citrate, and most preferably less than about 250 mM NaCl and 25 mM trisodium citrate. Low stringency hybridization can be obtained in the absence of organic solvent, e.g., formamide, while high stringency hybridization can be obtained in the presence of at least about 35% formamide, and most preferably at least about 50% formamide. Stringent temperature
15 conditions will ordinarily include temperatures of at least about 30°C, more preferably of at least about 37°C, and most preferably of at least about 42°C. Varying additional parameters, such as hybridization time, the concentration of detergent, e.g., sodium dodecyl sulfate (SDS), and the inclusion or exclusion of carrier DNA, are well known to those skilled in the art. Various levels of stringency are accomplished by combining these various conditions as needed. In a preferred
20 embodiment, hybridization will occur at 30°C in 750 mM NaCl, 75 mM trisodium citrate, and 1% SDS. In a more preferred embodiment, hybridization will occur at 37°C in 500 mM NaCl, 50 mM trisodium citrate, 1% SDS, 35% formamide, and 100 µg/ml denatured salmon sperm DNA (ssDNA). In a most preferred embodiment, hybridization will occur at 42°C in 250 mM NaCl, 25 mM trisodium citrate, 1% SDS, 50 % formamide, and 200 µg/ml ssDNA. Useful variations on
25 these conditions will be readily apparent to those skilled in the art.

The washing steps which follow hybridization can also vary in stringency. Wash stringency conditions can be defined by salt concentration and by temperature. As above, wash stringency can be increased by decreasing salt concentration or by increasing temperature. For example, stringent salt concentration for the wash steps will preferably be less than about 30 mM
30 NaCl and 3 mM trisodium citrate, and most preferably less than about 15 mM NaCl and 1.5 mM trisodium citrate. Stringent temperature conditions for the wash steps will ordinarily include temperature of at least about 25°C, more preferably of at least about 42°C, and most preferably of at least about 68°C. In a preferred embodiment, wash steps will occur at 25°C in 30 mM NaCl, 3 mM trisodium citrate, and 0.1% SDS. In a more preferred embodiment, wash steps will occur at

42°C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. In a most preferred embodiment, wash steps will occur at 68°C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS.

Additional variations on these conditions will be readily apparent to those skilled in the art.

Methods for DNA sequencing and analysis are well known in the art. The methods may
5 employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE enzyme (Amersham International PLC, Buckinghamshire England), TAQ polymerase (Perkin-Elmer Corp.), thermostable T7 polymerase (Amersham International PLC), or combinations of polymerases and proofreading exonucleases, such as those found in the ELONGASE amplification system (Life Technologies, Inc., Gaithersburg, MD). Preferably, sequence preparation is
10 automated with machines, e.g., the ABI CATALYST 800 (Perkin-Elmer Corp.) or MICROLAB 2200 systems (Hamilton Co., Reno, NV), in combination with thermal cyclers. Sequencing can also be automated, such as by ABI PRISM 373 or 377 systems (Perkin-Elmer Corp.) or the MEGABACE 1000 capillary electrophoresis system (Molecular Dynamics, Inc., Sunnyvale, CA). Sequences can be analyzed using computer programs and algorithms well known in the art. (See,
15 e.g., Ausubel, et al. (1997) Short Protocols in Molecular Biology, Wiley, New York, NY, unit 7.7; and Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, Inc, New York, NY.)

The nucleic acid sequences encoding CaREG may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect
20 upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived
25 from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an
30 engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-306). Additionally, one may use PCR, nested primers, and PROMOTORFINDER libraries (Clontech, Palo Alto, CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon

junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO™ 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to
5 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of
10 sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for
15 detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR software, (The Perkin-Elmer Corp., Norwalk, CT)), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited
20 amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode CaREG may be cloned in recombinant DNA molecules that direct expression of CaREG, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the
25 same or a functionally equivalent amino acid sequence may be produced and used to express CaREG.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter CaREG-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the
30 gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

In another embodiment, sequences encoding CaREG may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucl. Acids Res. Symp. Ser. 215-223, and Horn, T. et al. (1980) Nucl. Acids Res. Symp. Ser. 225-232.) Alternatively, CaREG itself or a fragment thereof may be synthesized using chemical methods.

- 5 For example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A Peptide Synthesizer (The Perkin-Elmer Corp., Norwalk, CT). Additionally, the amino acid sequence of CaREG, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant
- 10 polypeptide.

- The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH
- 15 Freeman and Co., New York, NY.)

- In order to express a biologically active CaREG, the nucleotide sequences encoding CaREG or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as
- 20 enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding CaREG. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding CaREG. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding CaREG and its initiation
- 25 codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic.
- 30 The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) Results Probl. Cell Differ. 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding CaREG and appropriate transcriptional and

translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview, NY, ch. 4, 8, and 16-17; and Ausubel, F.M. et al. (1995, and periodic supplements) Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding CaREG. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus (CaMV) or tobacco mosaic virus (TMV)) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding CaREG. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding CaREG can be achieved using a multifunctional E. coli vector such as pBLUESCRIPT plasmid (Stratagene) or pSPORT1 plasmid (Life Technologies, Inc.). Ligation of sequences encoding CaREG into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of CaREG are needed, e.g. for the production of antibodies, vectors which direct high level expression of CaREG may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of CaREG. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, supra; and Grant et al. (1987) Methods Enzymol. 153:516-54; Scorer, C. A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of CaREG. Transcription of sequences

encoding CaREG may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV. (Takamatsu, N. (1987) EMBO J. 6:307-311.) Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., Hobbs, S. or Murry, L.E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, NY; pp. 191-196.)

10 In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding CaREG may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses CaREG in host cells. (See, e.g., Logan, J. and T. Shenk
15 (1984) Proc. Natl. Acad. Sci. 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb
20 are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes.

For long term production of recombinant proteins in mammalian systems, stable expression of CaREG in cell lines is preferred. For example, sequences encoding CaREG can be transformed into cell lines using expression vectors which may contain viral origins of replication
25 and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed
30 cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* or *apr*⁻ cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; and Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite,

antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* or *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. 77:3567-3570; Colbere-Garapin, F. et al (1981) J. Mol. Biol. 150:1-14; and Murry, supra.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP) (Clontech, Palo Alto, CA), β glucuronidase and its substrate β -D-glucuronoside, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. et al. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding CaREG is inserted within a marker gene sequence, transformed cells containing sequences encoding CaREG can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding CaREG under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding CaREG and that express CaREG may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of CaREG using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on CaREG is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St Paul, MN, Section IV; Coligan, J. E. et al. (1997 and periodic supplements) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York, NY; and Maddox,

D.E. et al. (1983) J. Exp. Med. 158:1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding CaREG include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding CaREG, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Pharmacia & Upjohn (Kalamazoo, MI), Promega (Madison, WI), and U.S. Biochemical Corp. (Cleveland, OH). Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding CaREG may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode CaREG may be designed to contain signal sequences which direct secretion of CaREG through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to specify protein targeting, folding, and/or activity.

Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC, Bethesda, MD) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding CaREG may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric CaREG protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of CaREG activity.

Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the CaREG encoding sequence and the heterologous protein sequence, so that CaREG may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel, F. M. et al. (1995 and periodic supplements) Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY, ch 10. A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled CaREG may be achieved in vitro using the TNT™ rabbit reticulocyte lysate or wheat germ extract systems (Promega, Madison, WI). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, preferably ³⁵S-methionine.

Fragments of CaREG may be produced not only by recombinant production, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the Applied Biosystems 431A peptide synthesizer (Perkin-Elmer Corp.). Various fragments of CaREG may be synthesized separately and then combined to produce the full length molecule.

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of CaREG-1 and Ca²⁺ binding proteins such as human S100α (GI 36176) and annexin type I. CaREG-1 is expressed in fetal and cancerous lung tissue. In addition, chemical and structural similarity exists among regions of CaREG-2, rat CaBP2 (GI 393203), and thioredoxin. CaREG-2 is expressed in cancerous and proliferating tissue and in cardiovascular tissue. Therefore, CaREG appears to play a role in neurological and cardiovascular disorders and

cancer.

Therefore, in one embodiment, CaREG or a fragment or derivative thereof may be administered to a subject to treat or prevent a neurological disorder. Such disorders can include, but are not limited to, epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including anxiety and schizophrenia, akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, and Tourette's disorder.

In another embodiment, a vector capable of expressing CaREG or a fragment or derivative thereof may be administered to a subject to treat or prevent a neurological disorder including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified CaREG in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a neurological disorder including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of CaREG may be administered to a subject to treat or prevent a neurological disorder including, but not limited to, those listed above.

In another embodiment, CaREG or a fragment or derivative thereof may be administered to a subject to treat or prevent a cardiovascular disorder. Such disorders can include, but are not limited to, cardiomyopathy, myocarditis, Duchenne's muscular dystrophy, myocardial ischemia and infarction, myocardial failure, tachyarrhythmia, long QT syndrome, and myotonic muscular

dystrophy.

In an additional embodiment, a vector capable of expressing CaREG or a fragment or derivative thereof may be administered to a subject to treat or prevent a cardiovascular disorder including, but not limited to, those described above.

5 In a further embodiment, a pharmaceutical composition comprising a substantially purified CaREG in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a cardiovascular disorder including, but not limited to, those provided above.

10 In still another embodiment, an agonist which modulates the activity of CaREG may be administered to a subject to treat or prevent a cardiovascular disorder including, but not limited to, those listed above.

In a further embodiment, an antagonist of CaREG may be administered to a subject to treat or prevent a cancer. Such cancers may include, but are not limited to, adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers
15 of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. In one aspect, an antibody which specifically binds CaREG may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissue which
20 express CaREG.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding CaREG may be administered to a subject to treat or prevent a cancer including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists,
25 complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to
30 achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of CaREG may be produced using methods which are generally known in the art. In particular, purified CaREG may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind CaREG. Antibodies to CaREG

may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use.

5 For the production of polyclonal antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with CaREG or with any fragment or oligopeptide thereof which has immunogenic properties. Rats and mice are preferred hosts for downstream applications involving monoclonal antibody production. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants
10 include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable. (For review of methods for antibody production and analysis, see, e.g., Harlow, E. and Lane, D. (1988) Antibodies: A Laboratory
15 Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.)

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to CaREG have an amino acid sequence consisting of at least about 5 amino acids, and, more preferably, of at least about 14 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and
20 contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of CaREG amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to CaREG may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but
25 are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci.* 80:2026-2030; and Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the
30 splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci.* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce

CaREG-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton D.R. (1991) Proc. Natl. Acad. Sci. 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte
5 population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. 86: 3833-3837; and Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for CaREG may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments
10 produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the
15 desired specificity and minimal cross-reactivity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between CaREG and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering CaREG epitopes is
20 preferred, but a competitive binding assay may also be employed. (Maddox, supra.)

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for CaREG. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of CaREG-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium
25 conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple CaREG epitopes, represents the average affinity, or avidity, of the antibodies for CaREG. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular CaREG epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are
30 preferred for use in immunoassays in which the CaREG-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of CaREG, preferably in active form, from the antibody. (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington, D. C.; and Liddell, J. E. and

Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York, NY.)

The titre and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For
5 example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is preferred for use in procedures requiring precipitation of CaREG-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

10 In another embodiment of the invention, the polynucleotides encoding CaREG, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding CaREG may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding CaREG. Thus, complementary molecules
15 or fragments may be used to modulate CaREG activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding CaREG.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses,
20 or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding CaREG. (See, e.g., Sambrook, supra; and Ausubel, supra.)

Genes encoding CaREG can be turned off by transforming a cell or tissue with expression
25 vectors which express high levels of a polynucleotide, or fragment thereof, encoding CaREG. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector, and may last even longer if appropriate replication
30 elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding CaREG. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, are preferred.

Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al.

- 5 (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing Co., Mt. Kisco, NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

- Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the
10 ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding CaREG.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences:

- 15 GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

- 20 Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding CaREG. Such DNA sequences may
25 be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

- RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3'
30 ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by

endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nature Biotechnology 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of CaREG, antibodies to CaREG, and mimetics, agonists, antagonists, or inhibitors of CaREG. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combining active

compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acid. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of CaREG, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example CaREG or fragments thereof, antibodies of CaREG, and agonists, antagonists or inhibitors of CaREG, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of therapeutic to toxic effects is the therapeutic index, and it can be expressed as the ED_{50}/LD_{50} ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account

include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μg to 100,000 μg , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind CaREG may be used for the diagnosis of disorders characterized by expression of CaREG, or in assays to monitor patients being treated with CaREG or agonists, antagonists, or inhibitors of CaREG. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for CaREG include methods which utilize the antibody and a label to detect CaREG in human body fluids or in extracts of cells or tissues. The antibodies may be used without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring CaREG, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of CaREG expression. Normal or standard values for CaREG expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to CaREG under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, preferably by photometric means. Quantities of CaREG expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding CaREG may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be

used to detect and quantitate gene expression in biopsied tissues in which expression of CaREG may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of CaREG, and to monitor regulation of CaREG levels during therapeutic intervention.

5 In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding CaREG or closely related molecules may be used to identify nucleic acid sequences which encode CaREG. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or
10 amplification (maximal, high, intermediate, or low), will determine whether the probe identifies only naturally occurring sequences encoding CaREG, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and should preferably have at least 50% sequence identity to any of the CaREG encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequences of
15 SEQ ID NO:3 or SEQ ID NO:4 or from genomic sequences including promoters, enhancers, and introns of the CaREG gene.

Means for producing specific hybridization probes for DNAs encoding CaREG include the cloning of polynucleotide sequences encoding CaREG or CaREG derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available,
20 and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

25 Polynucleotide sequences encoding CaREG may be used for the diagnosis of a disorder associated with expression of CaREG. Examples of such a disorder include, but are not limited to, a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron
30 disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic

diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including anxiety and schizophrenia, akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, and Tourette's disorder; a cardiovascular disorder such as cardiomyopathy, myocarditis, Duchenne's muscular dystrophy, myocardial ischemia and infarction, myocardial failure, tachyarrhythmia, long QT syndrome, and myotonic muscular dystrophy; and a cancer such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. The polynucleotide sequences encoding CaREG may be used in Southern or Northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and ELISA assays; and in microarrays utilizing fluids or tissues from patients to detect altered CaREG expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding CaREG may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding CaREG may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding CaREG in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of CaREG, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding CaREG, under conditions suitable for hybridization or

amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from
5 standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period
10 ranging from several days to months.

With respect to cancer, the presence of a high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures
15 or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding CaREG may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a
20 polynucleotide encoding CaREG, or a fragment of a polynucleotide complementary to the polynucleotide encoding CaREG, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantitation of closely related DNA or RNA sequences.

Methods which may also be used to quantitate the expression of CaREG include
25 radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; and Duplaa, C. et al. (1993) Anal. Biochem. 229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or
30 colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic variants, mutations, and polymorphisms. This information may be used to

determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

In another embodiment of the invention, nucleic acid sequences encoding CaREG may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, R.A. (ed.) Molecular Biology and Biotechnology, VCH Publishers New York, NY, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) site. Correlation between the location of the gene encoding CaREG on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal

location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, CaREG, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, 5 affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between CaREG and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test 10 compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with CaREG, or fragments thereof, and washed. Bound CaREG is then detected by methods well known in the art. Purified CaREG can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

15 In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding CaREG specifically compete with a test compound for binding CaREG. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with CaREG.

In additional embodiments, the nucleotide sequences which encode CaREG may be used 20 in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred 25 specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. [Attorney Docket No. PF-0564 P], filed July 20, 1998, are hereby expressly incorporated by reference.

30

EXAMPLES

I. cDNA Library Construction

The LUNGNOT09 cDNA library was constructed using RNA isolated from the lung tissue of a 23-week-old Caucasian male fetus. The pregnancy was terminated following a

diagnosis by ultrasound of infantile polycystic kidney disease.

The frozen tissue was homogenized and lysed in guanidinium isothiocyanate solution using a Brinkmann Homogenizer Polytron PT-3000 (Brinkmann Instruments, Westbury, NY). The lysate was centrifuged over a CsCl cushion to isolate RNA. The RNA was extracted with
5 acid phenol, precipitated with sodium acetate and ethanol, resuspended in RNase free water, and treated with DNase. The RNA was re-extracted with acid phenol and reprecipitated as above. Poly(A⁺) RNA was isolated using the OLIGOTEX kit (QIAGEN Inc, Chatsworth, CA).

The ENDCNOT03 cDNA library was constructed using RNA isolated from dermal microvascular endothelial cells removed from a neonatal Caucasian male. The cells were
10 homogenized in guanidinium isothiocyanate solution, and poly(A⁺) RNA was isolated using oligo (dT)₂₅-streptavidin magnetic particles as recommended by the manufacturer (CPG Inc, Lincoln Park, NJ).

Poly(A⁺) RNA was used for cDNA synthesis and construction of each cDNA library according to the recommended protocols in the SUPERScript plasmid system (Life
15 Technologies). The cDNAs were fractionated on a SEPHAROSE CL4B column (Amersham Pharmacia Biotech), and those cDNAs exceeding 400 bp were ligated into pINCY 1 plasmid (Incyte Pharmaceuticals, Palo Alto, CA). Recombinant plasmids were transformed into DH5 α TM competent cells (Life Technologies).

20 II. Isolation of cDNA Clones

Plasmid DNA was released from the cells and purified using the REAL Prep 96 plasmid kit (QIAGEN Inc). The recommended protocol was employed except for the following changes: 1) the bacteria were cultured in 1 ml of sterile Terrific Broth (Life Technologies) with carbenicillin at 25 mg/l and glycerol at 0.4%; 2) after the cultures were incubated for 19 hours, the cells were
25 lysed with 0.3 ml of lysis buffer; and 3) following isopropanol precipitation, the plasmid DNA pellets were each resuspended in 0.1 ml of distilled water. The DNA samples were stored at 4°C.

III. Sequencing and Analysis

The cDNAs were prepared for sequencing using either an ABI CATALYST 800 (Perkin-
30 Elmer Applied Biosystems, Foster City, CA) or a MICROLAB 2200 (Hamilton Co., Reno, NV) sequencing preparation system in combination with Peltier PTC-200 thermal cyclers (MJ Research, Inc., Watertown, MA). The cDNAs were sequenced using the ABI PRISM 373 or 377 sequencing systems (Perkin-Elmer Corp.) and ABI protocols, base calling software, and kits (Perkin-Elmer Applied Biosystems). Alternatively, solutions and dyes from Amersham Pharmacia

Biotech, Ltd. were used. Reading frames were determined using standard methods (Ausubel, supra). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example V.

The polynucleotide sequences derived from cDNA, extension, and shotgun sequencing
 5 were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 1 summarizes the software programs, their descriptions, references, and threshold parameters where applicable. The references cited in the third column of Table 1 are incorporated by reference herein. Sequences were also analyzed using MACDNASIS PRO software (Hitachi Software Engineering Co., Ltd. San Bruno, CA) and
 10 LASERGENE™ software (DNASTAR).

The polynucleotide sequences were validated by removing vector, linker, and polyA tail sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as GenBank primate, rodent, mammalian,
 15 vertebrate, and eukaryote databases, and BLOCKS to acquire annotation, using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. This was followed by translation of the full length polynucleotide sequences to derive the corresponding full
 20 length amino acid sequences. These full length polynucleotide and amino acid sequences were subsequently analyzed by querying against databases such as the GenBank databases described above and SwissProt, BLOCKS, PRINTS, PFAM, and Prosite.

IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a
 25 gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; and Ausubel, supra, ch. 4 and 16.)

Electronic northern analysis was produced using analogous computer techniques. These techniques apply BLAST to search for identical or related molecules in nucleotide databases such
 30 as GenBank or LIFESEQ database (Incyte Pharmaceuticals). The sensitivity of the computer search was modified to determine the specificity of the match. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

$$100$$

The product score encompasses both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match may have a possibility of a 1% to 2% error, in contrast, a product score of 70 indicates that the match will be exact. Similar molecules were identified by product scores between 15 and 40, although lower
5 scores may identify related molecules.

Electronic northern analysis further involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease categories included cancer, inflammation/trauma, fetal,
10 neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was divided by the total number of libraries across all categories. The results above were reported as a percentage distribution.

V. Extension of CaREG Encoding Polynucleotides

15 The full length nucleic acid sequences, SEQ ID NO:3 and SEQ ID NO:4, were produced by extension of the component fragments disclosed in the invention description using oligonucleotide primers based on those fragments. Primers were used to facilitate the extension of the known sequence "outward" generating amplicons containing new unknown nucleotide sequence for the region of interest. The initial primers were designed from the cDNA using
20 OLIGO 4.06 software (National Biosciences, Plymouth, MN), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries (Life Technologies) were used to extend the sequence. If
25 more than one extension is necessary or desired, additional sets of primers are designed to further extend the known region.

High fidelity amplification was obtained by following the instructions for the XL-PCR kit (The Perkin-Elmer Corp., Norwalk, CT) and thoroughly mixing the enzyme and reaction mix. PCR was performed using the PTC-200 thermal cycler (MJ Research, Inc., Watertown, MA),
30 beginning with 40 pmol of each primer and the recommended concentrations of all other components of the kit, with the following parameters:

Step 1	94° C for 1 min (initial denaturation)
Step 2	65° C for 1 min
Step 3	68° C for 6 min
35 Step 4	94° C for 15 sec

- | | | |
|----|---------|---|
| | Step 5 | 65° C for 1 min |
| | Step 6 | 68° C for 7 min |
| | Step 7 | Repeat steps 4 through 6 for an additional 15 cycles |
| | Step 8 | 94° C for 15 sec |
| 5 | Step 9 | 65° C for 1 min |
| | Step 10 | 68° C for 7:15 min |
| | Step 11 | Repeat steps 8 through 10 for an additional 12 cycles |
| | Step 12 | 72° C for 8 min |
| 10 | Step 13 | 4° C (and holding) |

A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a low concentration (about 0.6% to 0.8%) agarose mini-gel to determine which reactions were successful in extending the sequence. Bands thought to contain the largest products were excised from the gel, purified using QIAQUICK purification kit (Qiagen, Inc.), and trimmed of overhangs using

15 Klenow enzyme to facilitate religation and cloning.

After ethanol precipitation, the products were redissolved in 13 μ l of ligation buffer, 1 μ l T4-DNA ligase (15 units) and 1 μ l T4 polynucleotide kinase were added, and the mixture was incubated at room temperature for 2 to 3 hours, or overnight at 16° C. Competent *E. coli* cells (in 40 μ l of appropriate media) were transformed with 3 μ l of ligation mixture and cultured in 80 μ l

20 of SOC medium. (See, e.g., Sambrook, *supra*, Appendix A, p. 2.) After incubation for one hour at 37° C, the *E. coli* mixture was plated on Luria Bertani (LB) agar (See, e.g., Sambrook, *supra*, Appendix A, p. 1) containing carbenicillin (2x carb). The following day, several colonies were randomly picked from each plate and cultured in 150 μ l of liquid LB/2x carb medium placed in an individual well of an appropriate commercially-available sterile 96-well microtiter plate. The

25 following day, 5 μ l of each overnight culture was transferred into a non-sterile 96-well plate and, after dilution 1:10 with water, 5 μ l from each sample was transferred into a PCR array.

For PCR amplification, 18 μ l of concentrated PCR reaction mix (3.3x) containing 4 units of rTth DNA polymerase, a vector primer, and one or both of the gene specific primers used for the extension reaction were added to each well. Amplification was performed using the following

30 conditions:

- | | | |
|----|--------|--|
| | Step 1 | 94° C for 60 sec |
| | Step 2 | 94° C for 20 sec |
| | Step 3 | 55° C for 30 sec |
| | Step 4 | 72° C for 90 sec |
| 35 | Step 5 | Repeat steps 2 through 4 for an additional 29 cycles |
| | Step 6 | 72° C for 180 sec |
| | Step 7 | 4° C (and holding) |

Aliquots of the PCR reactions were run on agarose gels together with molecular weight

40 markers. The sizes of the PCR products were compared to the original partial cDNAs, and

appropriate clones were selected, ligated into plasmid, and sequenced.

In like manner, the nucleotide sequences of SEQ ID NO:3 and SEQ ID NO:4 are used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for 5' extension, and an appropriate genomic library.

5

VI. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:3 and SEQ ID NO:4 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ - ^{32} P] adenosine triphosphate (Amersham, Chicago, IL), and T4 polynucleotide kinase (DuPont NEN®, Boston, MA). The labeled oligonucleotides are substantially purified using a Sephadex™ G-25 superfine size exclusion dextran bead column (Pharmacia & Upjohn, Kalamazoo, MI). An aliquot containing 10^7 counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN, Boston, MA).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham, NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMAT AR™ film (Kodak, Rochester, NY) is exposed to the blots to film for several hours, hybridization patterns are compared visually.

25

VII. Microarrays

A chemical coupling procedure and an ink jet device can be used to synthesize array elements on the surface of a substrate. (See, e.g., Baldeschweiler, *supra*.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the

scanned images.

Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE software. Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) *Science* 270:467-470; and Shalon, D. et al. (1996) *Genome Res.* 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

VIII. Complementary Polynucleotides

Sequences complementary to the CaREG-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring CaREG. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO™ 4.06 software and the coding sequence of CaREG. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the CaREG-encoding transcript.

IX. Expression of CaREG

Expression and purification of CaREG is achieved using bacterial or virus-based expression systems. For expression of CaREG in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*tac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express CaREG upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of CaREG in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant *Autographica californica* nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding CaREG by either homologous recombination

or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription.

Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic

- 5 modifications to baculovirus. (See Engelhard, E. K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, CaREG is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates.

- 10 GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Pharmacia, Piscataway, NJ). Following purification, the GST moiety can be proteolytically cleaved from CaREG at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and
15 polyclonal anti-FLAG antibodies (Eastman Kodak, Rochester, NY). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN Inc, Chatsworth, CA). Methods for protein expression and purification are discussed in Ausubel, F. M. et al. (1995 and periodic supplements) Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY, ch 10, 16. Purified CaREG obtained by these methods can be used directly
20 in the following activity assay.

X. Demonstration of CaREG Activity

- An assay for CaREG activity measures the binding of CaREG to Ca^{2+} using a Ca^{2+} overlay system. (Weis, K. et al. (1994) J. Biol. Chem. 269:19142-19150.) Purified CaREG is
25 transferred and immobilized onto a nitrocellulose membrane. The membrane is washed three times with buffer (60 mM KCl, 5 mM MgCl_2 , 10 mM imidazole-HCl, pH 6.8) and incubated in this buffer for 10 minutes with 1 μCi [$^{45}\text{Ca}^{2+}$] (NEN-DuPont, Boston, MA). Unbound [$^{45}\text{Ca}^{2+}$] is removed from the membrane by washing with water, and the membrane is dried. Membrane-bound [$^{45}\text{Ca}^{2+}$] is detected by autoradiography and quantified using image analysis systems and
30 software. CaREG activity is proportional to the amount of [$^{45}\text{Ca}^{2+}$] detected on the membrane.

XI. Functional Assays

CaREG function is assessed by expressing the sequences encoding CaREG at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a

mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT (Life Technologies) and PCR 3.1 (Invitrogen, Carlsbad, CA), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, preferably of endothelial or hematopoietic origin, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP) (Clontech, Palo Alto, CA), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP, and to evaluate properties, for example, their apoptotic state. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M. G. (1994) Flow Cytometry, Oxford, New York, NY.

The influence of CaREG on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding CaREG and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success, NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding CaREG and other genes of interest can be analyzed by Northern analysis or microarray techniques.

XII. Production of CaREG Specific Antibodies

CaREG substantially purified using polyacrylamide gel electrophoresis (PAGE)(see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the CaREG amino acid sequence is analyzed using LASERGENE™

software (DNASTAR Inc.) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel supra, ch. 11.)

5 Typically, oligopeptides 15 residues in length are synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry and coupled to KLH (Sigma, St. Louis, MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for
10 antipeptide activity by, for example, binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XIII. Purification of Naturally Occurring CaREG Using Specific Antibodies

Naturally occurring or recombinant CaREG is substantially purified by immunoaffinity
15 chromatography using antibodies specific for CaREG. An immunoaffinity column is constructed by covalently coupling anti-CaREG antibody to an activated chromatographic resin, such as CNBr-activated Sepharose (Pharmacia & Upjohn). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing CaREG are passed over the immunoaffinity column, and the column is
20 washed under conditions that allow the preferential absorbance of CaREG (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/CaREG binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and CaREG is collected.

25 XIV. Identification of Molecules Which Interact with CaREG

CaREG, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton et al. (1973) Biochem. J. 133:529.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled CaREG, washed, and any wells with labeled CaREG complex are assayed. Data obtained using different concentrations of
30 CaREG are used to calculate values for the number, affinity, and association of CaREG with the candidate molecules.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred

embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value= 1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS and PRINTS databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff, Nucl. Acid Res., 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and Probability value= 1.0E-3 or less
PFAM	A Hidden Markov Models-based application useful for protein family search.	Krogh, A. et al. (1994) J. Mol. Biol., 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits, depending on individual protein families

Table 1 cont.

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Score= 4.0 or greater
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score=5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <u>supra</u> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. A substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and fragments thereof.
5
2. A substantially purified variant having at least 90% amino acid identity to the amino acid sequence of claim 1.
3. An isolated and purified polynucleotide encoding the polypeptide of claim 1.
10
4. An isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity to the polynucleotide of claim 3.
5. An isolated and purified polynucleotide which hybridizes under stringent
15 conditions to the polynucleotide of claim 3.
6. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide sequence of claim 3.
- 20 7. An isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:3, SEQ ID NO:4, and fragments thereof.
8. An isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity to the polynucleotide of claim 7.
25
9. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 7.
10. An expression vector comprising at least a fragment of the polynucleotide of
30 claim 3.
11. A host cell comprising the expression vector of claim 10.
12. A method for producing a polypeptide, the method comprising the steps of:

a) culturing the host cell of claim 11 under conditions suitable for the expression of the polypeptide; and

b) recovering the polypeptide from the host cell culture.

5 13. A pharmaceutical composition comprising the polypeptide of claim 1 in conjunction with a suitable pharmaceutical carrier.

14. A purified antibody which specifically binds to the polypeptide of claim 1.

10 15. A purified agonist of the polypeptide of claim 1.

16. A purified antagonist of the polypeptide of claim 1.

17. A method for treating or preventing a neurological disorder, the method
15 comprising administering to a subject in need of such treatment an effective amount of the pharmaceutical composition of claim 13.

18. A method for treating or preventing a cardiovascular disorder, the method
comprising administering to a subject in need of such treatment an effective amount of the
20 pharmaceutical composition of claim 13.

19. A method for treating or preventing a cancer, the method comprising
administering to a subject in need of such treatment an effective amount of the antagonist of claim
16.

25 20. A method for detecting a polynucleotide in a biological sample, the method comprising the steps of:

(a) hybridizing the polynucleotide of claim 6 to at least one of the nucleic acids in the biological sample, thereby forming a hybridization complex; and

30 (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of the polynucleotide encoding the polypeptide in the biological sample.

21. The method of claim 20 further comprising amplifying the polynucleotide prior to hybridization.

5'	GAG	AGA	ATG	GGT	GGC	TGT	ATG	CAC	AGT	ACC	CAG	GAC	AAA	TCT	CTC	CAC	TTG	GAA
	M	G	G	G	C	C	M	H	S	T	Q	D	K	S	L	H	L	E
	9		18		27		36		45		54							
	63		72		81		90		99		108							
	GGA	GAT	CCC	AAT	CCT	TCT	GCA	GCC	CCA	ACA	TCC	ACC	TGC	GCA	CCT	AGG	AAA	ATG
	G	D	P	N	P	S	A	A	P	T	S	T	C	A	P	R	K	M
	117		126		135		144		153		162							
	CCC	AAA	AGG	ATT	TCA	ATA	TCC	AAA	CAA	CTG	GCT	TCA	GTG	AAA	GCT	CTG	AGG	AAG
	P	K	R	I	S	I	S	K	Q	L	A	S	V	K	A	L	R	K
	171		180		189		198		207		216							
	TGC	TCA	GAT	CTG	GAA	AAA	GCT	ATT	GCC	ACC	ACT	GCT	CTG	ATT	TTC	AGA	AAT	TCT
	C	S	D	L	E	K	A	I	A	T	A	L	I	F	R	N	S	
	225		234		243		252		261		270							
	TCT	GAC	TCT	GAT	GGT	AAA	CTT	GAA	AAA	GCT	ATT	GCC	AAA	GAT	CTG	CTG	CAA	ACC
	S	D	S	D	G	K	L	E	K	A	I	A	K	D	L	L	Q	T
	279		288		297		306		315		324							
	CAA	TTT	AGG	AAT	TTC	GCA	GAG	GGA	CAA	GAA	ACC	AAG	CCA	AAA	TAC	AGA	GAG	ATC
	Q	F	R	N	F	A	E	G	Q	E	T	K	P	K	Y	R	E	I
	333		342		351		360		369		378							
	CTT	TCT	GAA	CTT	GAT	GAG	CAC	ACA	GAA	AAT	AAG	CTA	GAT	TTT	GAA	GAC	TTC	ATG
	L	S	E	L	D	E	H	T	E	N	K	L	D	F	E	D	F	M

FIGURE 1A

387	ATC TTG CTC TTA AGC ATC ACT GTC ATG TCA GAT CTG CTA CAA AAT ATA CGG AAT	405	414	423	432
I L L L	S I T V M S D L L Q N I R N				
441	GTA AAA ATT ATG AAA TGA ACA GTT TTA AAT ATG CTG TAT AAA ATA ATG GCA AAA	459	468	477	486
V K I M K					
495	GAC AGT GTT ATT AAA ATG TTT CCA TCT TAT TTT TGA TTA ATT GAA TAT ATC TAT	513	522	531	540
549	CAT GCA TCT GAC ATT GCC TAG GAT GAT GGT TCT GAT TGC TGG TAT TCA GAT CCA ATG	567	576	585	594
603	TAA CTC CAA ATA TTT ACC C 3'	612			

FIGURE 1B

```

5'   9   18   27   36   45   54
    CCC ACG CGT CCG CGG ACG CTG GGC CAT GAC GGC GAC GAG GAC CCG CAC

    63   72   81   90   99   108
    AGC AAG CAC CTG TAC ACG GCC GAC ATG TTC ACG CAC GGG ATC CAG AGC AGC GCC GCG
        M   F   T   H   G   I   Q   S   A   A

    117  126  135  144  153  162
    CAC TTC GTC ATG TTC TTC GCG CCC TGG TGT GGA CAC TGC CAG CGG CTG CAG CCG
    H   F   V   M   F   F   A   P   W   C   G   H   C   Q   R   L   Q   P

    171  180  189  198  207  216
    ACT TGG AAT GAC CTG GGA GAC AAA TAC AAC AGC ATG GAA GAT GCC AAA GTC TAT
    T   W   N   D   L   G   D   K   Y   N   S   M   E   D   A   K   V   Y

    225  234  243  252  261  270
    GTG GCT AAA GTG GAC TGC ACG GCC CAC TCC GAC GTG TGC TCC GCC CAG GGG GTG
    V   A   K   V   D   C   T   A   H   S   D   V   C   S   A   Q   G   V

    279  288  297  306  315  324
    CGA GGA TAC CCC ACC TTA AAG CTT TTC AAG CCA GGC CAA GAA GCT GTG AAG TAC
    R   G   Y   P   T   L   K   L   F   K   P   G   Q   E   A   V   K   Y

    333  342  351  360  369  378
    CAG GGT CCT CGG GAC TTC CAG ACA CTG GAA AAC TGG ATG CTG CAG ACA CTG AAC
    Q   G   P   R   D   F   Q   T   L   E   N   W   M   L   Q   T   L   N

```

FIGURE 2A

387	GAG GAG CCA GTG ACA CCA GAG CCG GAA GTG GAA CCG CCC AGT GCC CCC GAG CTC	405	414	423	432
	E P V T P E P E V E P P S A P E L				
441	AAG CAA GGG CTG TAT GAG CTC TCA GCA AGC AAC TTT GAG CTG CAC GTT GCA CAA	459	468	477	486
	K Q G L Y E L S A S N F E L H V A Q				
495	GGC GAC CAC TTT ATC AAG TTC TTC GCT CCG TGG TGT GGT CAC TGC AAA GCC CTG	513	522	531	540
	G D H F I K F F A P W C G H C K A L				
549	GCT CCA ACC TGG GAG CAG CTG GCT GCT CTG GGC CTT GAA CAT TCC GAA ACT GTC AAG	567	576	585	594
	A P T W E Q L A L G L E H S E T V K				
603	ATT GGC AAG GTT GAT TGT ACA CAG CAC TAT GAA CTC TGC TCC GGA AAC CAG GTT	621	630	639	648
	I G K V D C T Q H Y E L C S G N Q V				
657	CGT GGC TAT CCC ACT CTT CTC TGG TTC CGA GAT GGG AAA AAG GTG GAT CAG TAC	675	684	693	702
	R G Y P T L L L CTC TGG T W F R D G K K V D Q Y				
711	AAG GGA AAG CGG GAT TTG GAG TCA CTG AGG GAG TAC GTG GAG TCG CAG CTG CAG	729	738	747	756
	K G K R D L L E S L R E Y V E S Q L Q				

FIGURE 2B

765	CGC ACA GAG ACT GGA GCG ACC GAG ACC GTC ACG CCC TCA GAG GCC CCG GTG CTG	774	783	792	801	810
	R T E T G A T E T V T P S E A P V L					
819	GCA GCT GAG CCC GAG GCT GAC AAG GGC ACT GTG TTG GCA CTC ACT GAA AAT AAC	828	837	846	855	864
	A A E P E A D K G T V L A L T E N N					
873	TTC GAT GAC ACC ATT GCA GAA GGA ATA ACC TTC ATC AAG TTT TAT GCT CCA TGG	882	891	900	909	918
	F D D T I A E G I T F I K F Y A P W					
927	TGT GGT CAT TGT AAG ACT CTG GCT CCT ACT TGG GAG GAA CTC TCT AAA AAG GAA	936	945	954	963	972
	C G H C K T L A P T T W E L S K K E					
981	TTC CCT GGT CTG GCG GGG GTC AAG ATC GCC GAA GTA GAC TGC ACT GCT GAA CGG	990	999	1008	1017	1026
	F P G L A G V K I A E V D C T A E R					
1035	AAT ATC TGC AGC AAG TAT TCG GTA CGA GGC TAC CCC ACG TTA TTG CTT TTC CGA	1044	1053	1062	1071	1080
	N I C S K Y S V R G Y P T L L F R					
1089	GGA GGG AAG AAA GTC AGT GAG CAC AGT GGA GGC AGA GAC CTT GAC TCG TTA CAC	1098	1107	1116	1125	1134
	G G K K V S E H S G G R D L S L H					

FIGURE 2C

1143	1152	1161	1170	1179	1188
CGC TTT GTC CTG AGC CAA GCG AAA GAC GAA CTT TAG GAA CAC AGT TGG AGG TCA					
R F V L S Q A K D E L					
1197	1206	1215	1224	1233	1242
CCT CTC CTG CCC AGC TCC CGC ACC CTG CGT TTA GGA GTT CAG TCC CAC AGA GGC					
1251	1260	1269	1278	1287	1296
CAC TGG GTT CCC AGT GGT GGC TGT TCA GAA AGC AGA ACA TAC TAA GCG TGA GGT					
1305	1314	1323	1332	1341	1350
ATC TTC TTT GTG TGT GTG TTT TCC AAG CCA ACA CAC TCT ACA GAT TCT TTA TTA					
1359	1368	1377	1386	1395	1404
AGT TAA GTT TCT CTA AGT AAA TGT GTA ACT CAT GGT CAC TGT GTA AAC ATT TTC					
1413	1422	1431	1440	1449	1458
AGT GGC GAT ATA TCC CCT TTG ACC TTC TCT TGA TGA AAT TTA CAT GGT TTC CTT					
1467	1476	1485	1494	1503	1512
TGA GAC TAA AAT AGC GTT GAG GGA AAT GAA ATT GCT GGA CTA TTT GTG GCT CCT					
1521	1530	1539	1548	1557	1566
GAG TTG AGT GAT TTT GGT GAA AGA AAG CAC ATC CAA AGC ATA GTT TAC CTG CCC					
1575	1584	1593	1602	1611	1620
ACG AGT TCT GGA AAG GTG GCC TTG TGG CAG TAT TGA CGT TCC TCT GAT CTT AAG					

FIGURE 2D

1629	1638	1647	1656	1665	1674
GTC ACA GTT GAC TCA ATA CTG TGT TGG TCC GTA GCA TGG AGC AGA TTG AAA TGC					
1683	1692	1701	1710	1719	1728
AAA AAC CCA CAC CTC TGG AAG ATA CCT TCA CGG CCG CTG CTG GAG CTT CTG TTG					
1737	1746	1755	1764	1773	1782
CTG TGA ATA CTT CTC TCA GTG TGA GAG GTT AGC CGT GAT GAA AGC AGC GTT ACT					
1791	1800	1809	1818	1827	1836
TCT GAC CGT GCC TGA GTA AGA GAA TGC TGA TGC CAT AAC TTT ATG TGT CGA TAC					
1845	1854	1863	1872	1881	1890
TTG TCA AAT CAG TTA CTG TTC AGG GGA TCC TTC TGT TTT TCA CGG GGT GAA ACA					
1899	1908	1917	1926	1935	1944
TGT CTT TAG TTC CTC ATG TTA ACA CGA AGC CAG AGC CCA CAT GAA CTG TTG GAT					
1953	1962	1971	1980	1989	1998
GTC TTC CTT AGA AAG GGT AGG CAT GGA AAA TTC CAC GAG GCT CAT TCT CAG TAT					
2007	2016	2025	2034	2043	2052
CTC ATT AAC TCA TTG AAA GAT TCC AGT TGT ATT TGT CAC CTG GGG TGA CAA GAC					
2061	2070	2079	2088	2097	2106
CAG ACA GGC TTT CCC AGG CCT GGG TAT CCA GGG AGG CTC TGC AGC CCT GCT GAA					

FIGURE 2E

2115	2124	2133	2142	2151	2160
GGG CCC TAA CTA GAG TTC TAG AGT TTC TGA TTC TGT TTC TCA GTA GTC CTT TTA					
2169	2178	2187	2196	2205	2214
GAG GCT TGC TAT ACT TGG TCT GCT TCA AGG AGG TCG ACC TTC TAA TGT ATG AAG					
2223	2232	2241	2250	2259	2268
AAT GGG ATG CAT TTG ATC TCA AGA CCA AAG ACA GAT GTC AGT GGG CTG CTC TGG					
2277	2286	2295	2304	2313	2322
CCC TGG TGT GCA CGG CTG TGG CAG CTG TTG ATG CCA GTG TCC TCT AAC TCA TGC					
2331	2340	2349	2358	2367	2376
TGT CCT TGT GAT TAA ACA CCT CTA TCT CCC TTG GGA ATA AGC ACA TAC AGG CTT					
2385	2394	2403	2412	2421	2430
AAG CTC TAA GAT AGA TAG GTG TTT GTC CTT TTA CCA TCG AGC TAC TTC CCA TAA					
2439	2448	2457	2466	2475	2484
TAA CCA CTT TGC ATC CAA CAC TCT TCA CCC ACC TCC CAT ACG CAA GGG GAT GTG					
2493	2502	2511	2520	2529	2538
GAT ACT TGG CCC AAA GTA ACT GGT GGT AGG AAT CTT AGA AAC AAG ACC ACT TAT					
2547	2556	2565	2574	2583	2592
ACT GTC TGT CTG AGG CAG AAG ATA ACA GCA TCT CGA CCA GCC TCT GCC TTA					

FIGURE 2F

2601	2610	2619	2628	2637	2646
AAG GAA ATC TTT ATT AAT CAC GTA TGG TTC ACA GAT AAT TCT TTT TTT AAA AAA					
2655	2664	2673	2682	2691	2700
ACC CAA CCC CCT AGA GAA GCA CAA CTG TCA AGA GTC TTG TAC ACA CAA CTT CAG					
2709	2718	2727	2736	2745	2754
CTT TGC ATC ACG AGT CTT GTA TTC CAA GAA AAT CAA AGT GGT ACA ATT TGT TTG					
2763	2772	2781	2790	2799	
TTT ACA CTA TGA TAC TTT CTA AAT AAA CTC TTT TTT TTT AAA AAA A 3'					

FIGURE 2G

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.
 BANDMAN, Olga
 TANG, Y. Tom
 CORLEY, Neil C.
 GUEGLER, Karl J.
 BAUGHN, Mariah R.
 YANG, Junming

<120> CALCIUM REGULATORY PROTEINS

<130> PF-0564 PCT

<140> To Be Assigned
 <141> Herewith

<150> 09/119,547
 <151> 1998-07-20

<160> 4

<170> PERL program

<210> 1
 <211> 147
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc feature
 <223> Incyte Clone No: 1356311

<300>

<400> 1

Met	Gly	Gly	Cys	Met	His	Ser	Thr	Gln	Asp	Lys	Ser	Leu	His	Leu					
1				5					10					15					
Glu	Gly	Asp	Pro	Asn	Pro	Ser	Ala	Ala	Pro	Thr	Ser	Thr	Cys	Ala					
				20					25					30					
Pro	Arg	Lys	Met	Pro	Lys	Arg	Ile	Ser	Ile	Ser	Lys	Gln	Leu	Ala					
				35					40					45					
Ser	Val	Lys	Ala	Leu	Arg	Lys	Cys	Ser	Asp	Leu	Glu	Lys	Ala	Ile					
				50					55					60					
Ala	Thr	Thr	Ala	Leu	Ile	Phe	Arg	Asn	Ser	Ser	Asp	Ser	Asp	Gly					
				65					70					75					
Lys	Leu	Glu	Lys	Ala	Ile	Ala	Lys	Asp	Leu	Leu	Gln	Thr	Gln	Phe					
				80					85					90					
Arg	Asn	Phe	Ala	Glu	Gly	Gln	Glu	Thr	Lys	Pro	Lys	Tyr	Arg	Glu					
				95					100					105					
Ile	Leu	Ser	Glu	Leu	Asp	Glu	His	Thr	Glu	Asn	Lys	Leu	Asp	Phe					
				110					115					120					
Glu	Asp	Phe	Met	Ile	Leu	Leu	Leu	Ser	Ile	Thr	Val	Met	Ser	Asp					
				125					130					135					
Leu	Leu	Gln	Asn	Ile	Arg	Asn	Val	Lys	Ile	Met	Lys								
				140					145										

<210> 2
 <211> 363
 <212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte Clone No: 2176876

<300>

<400> 2

```

Met Phe Thr His Gly Ile Gln Ser Ala Ala His Phe Val Met Phe
 1      5      10
Phe Ala Pro Trp Cys Gly His Cys Gln Arg Leu Gln Pro Thr Trp
 20      25      30
Asn Asp Leu Gly Asp Lys Tyr Asn Ser Met Glu Asp Ala Lys Val
 35      40      45
Tyr Val Ala Lys Val Asp Cys Thr Ala His Ser Asp Val Cys Ser
 50      55      60
Ala Gln Gly Val Arg Gly Tyr Pro Thr Leu Lys Leu Phe Lys Pro
 65      70      75
Gly Gln Glu Ala Val Lys Tyr Gln Gly Pro Arg Asp Phe Gln Thr
 80      85      90
Leu Glu Asn Trp Met Leu Gln Thr Leu Asn Glu Glu Pro Val Thr
 95     100     105
Pro Glu Pro Glu Val Glu Pro Pro Ser Ala Pro Glu Leu Lys Gln
110     115     120
Gly Leu Tyr Glu Leu Ser Ala Ser Asn Phe Glu Leu His Val Ala
125     130     135
Gln Gly Asp His Phe Ile Lys Phe Phe Ala Pro Trp Cys Gly His
140     145     150
Cys Lys Ala Leu Ala Pro Thr Trp Glu Gln Leu Ala Leu Gly Leu
155     160     165
Glu His Ser Glu Thr Val Lys Ile Gly Lys Val Asp Cys Thr Gln
170     175     180
His Tyr Glu Leu Cys Ser Gly Asn Gln Val Arg Gly Tyr Pro Thr
185     190     195
Leu Leu Trp Phe Arg Asp Gly Lys Lys Val Asp Gln Tyr Lys Gly
200     205     210
Lys Arg Asp Leu Glu Ser Leu Arg Glu Tyr Val Glu Ser Gln Leu
215     220     225
Gln Arg Thr Glu Thr Gly Ala Thr Glu Thr Val Thr Pro Ser Glu
230     235     240
Ala Pro Val Leu Ala Ala Glu Pro Glu Ala Asp Lys Gly Thr Val
245     250     255
Leu Ala Leu Thr Glu Asn Asn Phe Asp Asp Thr Ile Ala Glu Gly
260     265     270
Ile Thr Phe Ile Lys Phe Tyr Ala Pro Trp Cys Gly His Cys Lys
275     280     285
Thr Leu Ala Pro Thr Trp Glu Glu Leu Ser Lys Lys Glu Phe Pro
290     295     300
Gly Leu Ala Gly Val Lys Ile Ala Glu Val Asp Cys Thr Ala Glu
305     310     315
Arg Asn Ile Cys Ser Lys Tyr Ser Val Arg Gly Tyr Pro Thr Leu
320     325     330
Leu Leu Phe Arg Gly Gly Lys Lys Val Ser Glu His Ser Gly Gly
335     340     345
Arg Asp Leu Asp Ser Leu His Arg Phe Val Leu Ser Gln Ala Lys
350     355     360
Asp Glu Leu

```

<210> 3

<211> 613

<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte Clone No: 1356311

<300>

<400> 3

```

gagagaatgg gtggctgtat gcacagtacc caggacaaat ctctccactt ggaaggagat 60
cccaatcctt ctgcagcccc aacatccacc tgcgcaccta ggaaaatgcc caaaaggatt 120
tcaatatcca aacaactggc ttcagtgaat gctctgagga agtgctcaga tctggaaaaa 180
gctattgccca ccaactgctct gattttcaga aattcttctg actctgatgg taaacttgaa 240
aaagctattg ccaaagatct gctgcaaacc caatttagga atttcgcaga gggacaagaa 300
accaagccaa aatacagaga gatcctttct gaacttgatg agcacacaga aaataagcta 360
gattttgaag acttcatgat ctgctotta agcatcactg tcatgtcaga tctgctacaa 420
aatatacgga atgtaaaaat tatgaaatga acagttttta atatgctgta taaaataatg 480
gcaaaagaca gtgttattaa aatgtttcca tcttattttt gattaattga atatatctat 540
catgcatctg acattgccta ggatggttct gattgctggg attcagatcc aatgtaactc 600
caaatattta ccc                                     613

```

<210> 4
<211> 2803
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte Clone No: 2176876

<300>

<400> 4

```

cccacgcgtc cgcggacgct gggccatgac ggcgaggacg gacaggaccc gcacagcaag 60
cacctgtaca cggcgcacat gttcacgcac gggatccaga gcgcccgcga cttcgtcatg 120
ttcttcgcgc cctggtgtgg aactgcccag cggctgcagc cgacttgga tgcctggga 180
gacaaatata acagcatgga agatgccaaa gtctatgtgg cttaaagtga ctgcacggcc 240
cactccgacg tgtgtccgc ccagggggtg cgaggatacc ccacctaata gcttttcaag 300
ccaggccaag aagctgtgaa gtaccagggt cctcgggact tccagacact ggaaaactgg 360
atgctgcaga cactgaacga ggagccagtg acaccagagc cggaagtga accgcccagt 420
gcccccgagc tcaagcaagg gctgtatgag ctctcagcaa gcaactttga gctgcaggtt 480
gcacaaggcg accactttat caagtctctt gctccgtggt gtggtcactg caaagccctg 540
gctccaacct gggagcagct ggctctgggc cttgaacatt ccgaaactgt caagattggc 600
aagggttgatt gtacacagca ctatgaactc tgctccgga accaggttcg tggctatccc 660
actcttctct gggtccgaga tgggaaaaag gtggatcagt acaagggaaa gcgggatttg 720
gagtcactga gggagtacgt ggagtcgcag ctgcagcgca cagagactgg agcgacggag 780
accgtcacgc cctcagaggc cccggtgctg gcagctgagc ccgaggtga caagggcact 840
gtgttggcac tcaactgaaa taacttcgat gacaccattg cagaaggaat aaccttcac 900
aagttttatg ctccatggtg tggctattgt aagactctgg ctctacttg ggaggaactc 960
tctaaaaagg aattccctgg tctggcgggg gtcaagatcg ccgaagtaga ctgcactgct 1020
gaacggaata tctgcagcaa gtattcggta cgaggctacc ccacgttatt gcttttccga 1080
ggaggggaaga aagtcagtga gcacagtgga ggcagagacc ttgactcgtt acaccgcttt 1140
gtcctgagcc aagcgaagaa cgaactttag gaacacagtt ggaggtcacc tctcctgccc 1200
agctcccgca ccctgcgttt aggagttcag tcccacagag gccactgggt tcccagtggt 1260
ggctgttcag aaagcagaac atactaagcg tgaggtatct tctttgtgtg tgtgttttcc 1320
aagccaacac actctacaga ttctttatta agttaagtt ctctaagtaa atgtgtaact 1380
catggtcact gtgtaaacat tttcagtggc gatataatcc ctttgacctt ctcttgatga 1440
aatttacatg gtttcctttg agactaaaat agcgttgagg gaaatgaaat tgctggacta 1500
tttgtggctc ctgagttgag tgattttggt gaaagaaagc acatccaaag catagtttac 1560
ctgcccacga gttctggaaa ggtggccttg tggcagttat gacgttcctc tgatcttaag 1620
gtcacagttg actcaatact gtgttggtcc gtagcatgga gcagattgaa atgcaaaaac 1680
ccacacctct ggaagatacc ttcacggccg ctgctggagc ttctgttgct gtgaataactt 1740

```

ctctcagtgt	gagagggttag	ccgtgatgaa	agcagcgtta	cttctgaccg	tgccctgagta	1800
agagaatgct	gatgccataa	ctttatgtgt	cgatacttgt	caaatcagtt	actgttcagg	1860
ggatccttct	gtttctcacg	gggtgaaaca	tgtcttttagt	tcctcatgtt	aacacgaagc	1920
cagagcccac	atgaactggt	ggatgtcttc	cttagaaagg	gtaggcatgg	aaaattccac	1980
gaggctcatt	ctcagtatct	cattaactca	ttgaaagatt	ccagttgtat	ttgtcacctg	2040
gggtgacaag	accagacagg	ctttcccagg	cctgggtatc	cagggaggct	ctgcagccct	2100
gctgaagggc	cctaactaga	gttctagagt	ttctgattct	gtttctcagt	agtcotttta	2160
gaggcttgct	atacttggtc	tgcttcaagg	aggtcgacct	tctaattgtat	gaagaatggg	2220
atgcatTTga	tctcaagacc	aaagacagat	gtcagtgggc	tgctctggcc	ctgggtgtgca	2280
cggtctgtggc	agctgttgat	gccagtgtcc	tctaactcat	gctgtccttg	tgattaaaca	2340
cctctatctc	ccttggggaat	aagcacatac	aggcttaagc	tctaagatag	atagggtgttt	2400
gtcctttttac	catcgagcta	cttcccataa	taaccacttt	gcaccaaca	ctcttcaccc	2460
acctcccata	cgcaagggga	tgtggatact	tggcccaaag	taactgggtg	taggaatctt	2520
agaaacaaga	ccacttatac	tgtctgtctg	aggcagaaga	taacagcagc	atctcgacca	2580
gcctctgcct	taaaggaaat	ctttattaat	cacgtatggg	tcacagataa	ttcttttttt	2640
aaaaaaaccc	aaccccctag	agaagcacia	ctgtcaagag	tcttgtagac	acaacttcag	2700
ctttgcatca	cgagtcttgt	attccaagaa	aatcaaagt	gtacaatttg	tttgttttaca	2760
ctatgatact	ttctaaataa	actctttttt	tttaaaaaaa	aaa		2803

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/12385

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 C12N5/10 C07K16/18 A61K38/17
C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category ^o	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMST27 'Online! The Institute for Genomic Research Fetal lung II Homo sapiens cDNA, 18 April 1997 (1997-04-18) XP002121205 see Accession AA359824 & ADAMS MD ET AL: "Initial assessment of human gene diversity and expression patterns based upon 83 million nucleotides of cDNA sequence" NATURE, vol. 377, 28 September 1995 (1995-09-28), pages 3-17, XP002914403 --- -/--	1-9

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

^o Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

3 November 1999

Date of mailing of the international search report

16/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

ALCONADA RODRIG..., A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/12385

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL 'Online! National Cancer Institute, Cancer Genome Anatomy Project Homo sapiens cDNA clone, 11 September 1997 (1997-09-11) XP002121206 see Accession AA551623 ---	1-9
X	DATABASE EMBL 'Online! National Cancer Institute, Cancer Genome Anatomy Project Homo sapiens cDNA clone, 5 July 1997 (1997-07-05) XP002121207 see Accession AA507028 ---	1-9
X	DATABASE EMBL 'Online! Wash-Merck EST Project Stratagene endothelial cell H.sapiens cDNA , 15 December 1996 (1996-12-15) XP002121208 see Accession Number AA147831 ---	1-9
A	WO 97 34013 A (HUMAN GENOME SCIENCES INC ;NI JIAN (US); GENTZ REINER L (US); YU G) 18 September 1997 (1997-09-18) page 17 -page 49 claims 1-22 ---	1-16, 19-21
A	WO 98 26068 A (INCYTE PHARMA INC ;GOLI SURYA K (US); HILLMAN JENNIFER L (US)) 18 June 1998 (1998-06-18) page 11 -page 38 claims 1-18 ---	1-16, 18, 20, 21
A	SCHAFER B W ET AL: "The S100 family of EF-hand calcium-binding proteins: functions and pathology" TIBS TRENDS IN BIOCHEMICAL SCIENCES, vol. 21, no. 4, 1 April 1996 (1996-04-01), page 134-140 XP004050923 ISSN: 0968-0004 the whole document -----	17-19

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 12385

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 17-18
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☒ Claims Nos.: 15,16,19
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
See additional sheet
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 15,16,19

Due to the fact that the agonist and antagonist of claims 15 and 16 were not sufficiently disclosed in the description, a reasonable search could not be carried out for the above mentioned claims together with claim 19.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-21 (partially)

Isolation and cloning of a human cDNA coding for a calcium regulatory protein CaREG1, and the recombinant expression of the same.

2. Claims: 1-21 (partially)

Isolation and cloning of a human cDNA coding for a calcium regulatory protein CaREG2, and the recombinant expression of the same.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/12385

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9734013	A	18-09-1997	AU	5420796 A	01-10-1997
WO 9826068	A	18-06-1998	US	5763220 A	09-06-1998
			AU	5520398 A	03-07-1998
			EP	0946722 A	06-10-1999
			US	5935931 A	10-08-1999