



(86) Date de dépôt PCT/PCT Filing Date: 2007/07/05
(87) Date publication PCT/PCT Publication Date: 2008/01/17
(85) Entrée phase nationale/National Entry: 2008/12/19
(86) N° demande PCT/PCT Application No.: EP 2007/005970
(87) N° publication PCT/PCT Publication No.: 2008/006506
(30) Priorité/Priority: 2006/07/10 (EP06014244.5)

(51) Cl.Int./Int.Cl. *A61K 9/50* (2006.01)
(71) Demandeur/Applicant:
DR. R. PFLEGER CHEMISCHE FABRIK GMBH, DE
(72) Inventeurs/Inventors:
JUNG, GERD, DE;
SCHAUPP, ALBERT, DE
(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : PREPARATION PHARMACEUTIQUE POUR ADMINISTRATION PAR VOIE ORALE AVEC LIBERATION
CONTROLEE DE SUBSTANCES ACTIVES DANS L'INTESTIN GRELE ET SON PROCEDE DE FABRICATION
(54) Title: PHARMACEUTICAL PREPARATION FOR ORAL ADMINISTRATION WITH CONTROLLED ACTIVE
INGREDIENT RELEASE IN THE SMALL INTESTINE AND METHOD FOR ITS PRODUCTION

(57) **Abrégé/Abstract:**

A pharmaceutical preparation for oral administration with controlled active ingredient release in the small intestine, on the basis of active ingredient carriers provided with at least one active ingredient which are provided with an inner layer for controlling the active ingredient release and a covering layer, arranged thereon, that is resistant to gastric juices, and is characterized in that the inner layer is constructed from at least two diffusion layers whose permeability for the diffusing active ingredient decreases from the inside to the outside, and a method for its production are described.



WO 2008/006506

PCT/EP2007/005970

Abstract

**Pharmaceutical preparation for oral administration with
controlled active ingredient release in the small
intestine and method for its production**

Any pharmaceutical preparation for oral administration
with controlled release of active ingredient in the
small bowel, on the basis of active ingredient carriers
provided with at least one active ingredient which are
provided with an inner layer to control the release of
active ingredient and with a gastro-resistant coating
layer disposed thereon, which is characterized in that
the inner layer is formed from at least two diffusion
layers whose permeability for the diffusing active
ingredient decreases from the inside to the outside,
and a method for the production thereof, are described.

WO 2008/006506

- 1 -

PCT/EP2007/005970

**Pharmaceutical preparation for oral administration with
controlled active ingredient release in the small
intestine and method for its production**

5

Description

The present invention relates to a pharmaceutical preparation for oral administration with controlled release of active ingredient in the small bowel on the basis of active ingredient carriers which are provided with at least one active ingredient and which are provided with an inner layer to control the release of active ingredient and with a gastro-resistant coating layer disposed thereon, and a method for the manufacture thereof.

Pharmaceutical preparations for oral administration with controlled release of active ingredient in the small bowel have previously been disclosed.

20

Thus, DE 198 23 940 A1 describes enteric fluoxetine pellets which include a neutral core onto which a fluoxetine layer has been applied and which is followed by a separating layer onto which an enteric layer is applied and is optionally provided with a finishing layer. In this case, the enteric layer serves to ensure that the active ingredient passes through the patient's stomach unchanged and dissolves only when the active ingredient leaves the stomach and enters the small bowel.

30

EP 0 941 070 B1 relates to a suckable tablet for modified release of active ingredients in the gastrointestinal tract, which comprises an active ingredient preparation in the form of particles with an at least bilayer coating, where an outer coating layer is saliva-resistant but gastro-soluble, and an inner

35

WO 2008/006506

- 2 -

PCT/EP2007/005970

coating layer is substantially resistant to disintegration in aqueous media but permits sustained release of active ingredient by diffusion. In a further embodiment, the particles with a bilayer coating include an outer coating layer which is saliva-resistant but gastro-soluble, and an inner coating layer which is gastro-resistant but soluble in the small bowel.

10 US patent application US 2004/0142035 A1 discloses a pharmaceutical preparation which has a core which comprises one or more active pharmaceutical ingredients, and a coating layer enveloping the core, where the coating layer includes a combination of two
15 or more enteric coating materials, of which at least two of the enteric coating materials dissolve at different pH values, in order in this way to achieve a controlled release profile of the active ingredient depending on the pH. This design serves to prevent a
20 significant part of the active ingredient still being present in the product after passing through the small bowel, and being lost.

Conventional pharmaceutical preparations for oral
25 administration with controlled release of active ingredient in the small bowel have, however, not proved to be entirely satisfactory because, when the active ingredient is released in the small bowel, there is initially a large rise in the active ingredient release
30 with correspondingly high blood levels, leading in many cases to an increase in the rate and intensity of side effects. With substances mentioned hereinafter, this relates for example in particular to dry mouth, accommodation impairments, gastrointestinal problems or
35 central nervous effects. This is particularly disadvantageous when the entire dose of the active ingredient is to be administered only once a day. It is further observed that oral administration for example

WO 2008/006506

- 3 -

PCT/EP2007/005970

of anticholinergics is followed by a large inter-individual variation in the plasma profiles.

On the other hand, the use of enteric materials with
5 different solubilities at different pH values, as envisaged by the teaching of US 2004/0142035 A1, results in limitations on the choice of materials on the one hand, and on the other hand non-uniform release of active ingredient when there are pH fluctuations.

10

The object of the present invention is therefore to provide a pharmaceutical preparation of the type indicated at the outset for oral administration of active ingredients with controlled release active
15 ingredient in the small bowel, which, in contrast to conventional formulations, exhibits a smaller initial rise in the release of active ingredient, and thus a lower initial plasma peak of the active ingredient, but ensures subsequently a lower but uniform plasma
20 concentration of the active ingredient, and thus leads to a lower rate of side effects and intensity of side effects and to a smaller interindividual variation of the plasma profiles, especially when the entire dose of the active ingredient is to be administered only once a
25 day.

This is achieved with the provided preparation when the release rate of the active ingredient is $\geq 75\%$ in 60 minutes on in vitro testing of the release in
30 accordance with the US Pharmacopeia. It is thus possible without problems to ensure the necessary dose for the whole day with a single administration.

It has surprisingly emerged that this object can be
35 achieved by providing the active ingredient carrier provided with an active ingredient with an inner layer to control the release of active ingredient, which is formed from at least two diffusion layers whose

WO 2008/006506

- 4 -

PCT/EP2007/005970

permeability for the diffusing active ingredient decreases from the inside to the outside and which are further provided with a gastro-resistant coating layer disposed thereon.

5

The invention therefore relates to the pharmaceutical preparation according to the main claim and the use thereof.

10 The invention thus relates to a pharmaceutical preparation for oral administration with controlled release of active ingredient in the small bowel, on the basis of active ingredient carriers provided with at least one active ingredient which are provided with an
15 inner layer to control the release of active ingredient and with a gastro-resistant coating layer disposed thereon, which is characterized in that the inner layer is formed from at least two diffusion layers whose permeability for the diffusing active ingredient
20 decreases from the inside to the outside.

This formation of the inner layer comprising at least two diffusion layers makes it possible to bring about targeted release of the active ingredient in the small
25 bowel, so that, compared with conventional products of this type, a lower initial rate of release of active ingredient can be achieved, leading on administration of the entire daily dose of the active ingredient in a single dose, with appropriately low but sufficient
30 blood levels, to a smaller interindividual variation in the plasma profile and thus to a reduced rate and intensity of side effects.

In a preferred embodiment of the invention, the at
35 least two diffusion layers of the inner layer for controlling the release of active ingredient are formed from a matrix material which is insoluble in small bowel fluid and into which pore formers which control

WO 2008/006506

- 5 -

PCT/EP2007/005970

the permeability for the diffusing active ingredient and are soluble and/or swellable and/or water-absorbing in small bowel fluid are incorporated. It is possible in this connection for the diffusion layers to comprise
5 matrix materials which are identical or of different types, and/or one or more pore formers which are of identical or different types.

Thus, according to the invention, the permeability of
10 the diffusion layers for the diffusing active ingredient is preferably controlled via the nature, the quantity and/or the particle size and/or the solubility and/or the swellability and/or the water-absorbing capacity of the pore formers present in the matrix
15 material.

According to the invention, the inner layer has an inner and an outer diffusion layer for controlling the release of active ingredient, and in the case where the
20 pore former material and matrix material are identical, the particle size and/or the concentration of the pore formers in the inner diffusion layer is greater than in the outer diffusion layer. In a preferred embodiment, when the pore formers are soluble in small bowel fluid,
25 the ratio of the pore former concentration in the inner diffusion layer to the pore former concentration in the outer diffusion layer is in the range from 20:1 to > 1:1, preferably in the range from 10:1 to 1.1:1.

30 The matrix material of the diffusion layers is preferably selected from the group comprising ethylcellulose, cellulose acetate phthalate, cellulose acetate succinate, shellac, hydroxypropyl-methylcellulose acetate succinate, cellulose acetate,
35 cellulose acetate propionate, hydroxypropylmethyl-cellulose phthalate, polyvinyl acetate, polyvinyl acetate phthalate, cellulose acetate butyrate, butyl methacrylate-(2-dimethylaminoethyl) methacrylate-methyl

WO 2008/006506

- 6 -

PCT/EP2007/005970

methacrylate copolymers, ethyl acrylate-methyl
methacrylate copolymers, methacrylic acid-methyl
methacrylate copolymers, methacrylic acid-ethyl
acrylate copolymers, ethyl acrylate-methyl
5 methacrylate-trimethylammoniummethyl methacrylate
chloride copolymers, sodium carboxymethylcellulose,
methylcellulose, hydroxyethylcellulose, methylethyl-
cellulose, methylhydroxypropylcellulose, hydroxypropyl-
methylcellulose, hydroxyethylmethylpropylcellulose,
10 chitosan, chitosan acetate, silicone elastomer latex
suspensions, hydrogenated castor oil, stearic acid,
glycerol monostearate, glycerol distearate, glycerol
dibehenate, stearyl alcohol, white wax, yellow wax,
hydrogenated vegetable oil and microcrystalline wax.

15

The material chosen for the pore formers of the
diffusion layers comprises materials from the group
comprising polyvinyl alcohol-polyethylene glycol
copolymers, polyvinylpyrrolidone, vinylpyrrolidone-
20 vinyl acetate copolymers, crosslinked polyvinyl-
pyrrolidone, microcrystalline cellulose, hydroxypropyl-
cellulose, sodium carboxymethylcellulose, methyl-
cellulose, hydroxyethylcellulose, methylethylcellulose,
methylhydroxypropylcellulose, hydroxypropylmethyl-
25 cellulose, hydroxyethylmethylpropylcellulose, poly-
ethylene glycol, cellulose powder, sucrose, lactose,
mannitol, sorbitol and polysorbate.

The matrix material of the diffusion layers may if
30 necessary additionally comprise plasticizers and
antitack agents in order to adjust the property of the
matrix material of the diffusion layers in the desired
manner. The matrix material of the diffusion layers
preferably provides polyethylene glycol, propylene
35 glycol, triethyl citrate, triacetin, acetyl tributyl
citrate, polysorbates, 2-pyrrolidone, dibutyl sebacate,
stearic acid, castor oil and/or medium-chain
triglycerides as plasticizers, and talc, stearic acid

WO 2008/006506

- 7 -

PCT/EP2007/005970

and salts thereof, fatty alcohols, mono-, di- or triglycerides with straight- and/or branched-chain fatty acids, colloidal silicon dioxide, precipitated silicon dioxide, aluminium oxide, kaolin, maize starch, wheat starch, rice starch, potato starch, titanium dioxide, silicone emulsions and/or Veegum (a magnesium aluminium silicate dispersion) as antitack agent.

The active ingredient carriers present in the pharmaceutical preparation of the invention may comprise as active ingredient any active pharmaceutical ingredients whose effect can be ensured via the small bowel, such as, for example, 4-diethylamino-2-butynyl α -phenylcyclohexaneglyconate hydrochloride, ethyldimethyl(1-methyl-3,3-diphenylpropyl)ammonium carra-
geenate, (+)-[R-2-[α -2-(diisopropylamino)ethyl]benzyl]-p-cresol tartrate, 8-(cyclopropylmethyl)-6 β ,7 β -epoxy-3 α -hydroxy-1 α H,5 α)-(S)-tropate bromide, (1R,3r,5S)-3-[(hydroxydiphenylacetyl)oxy]spiro[8-azonia-
bicyclo[3.2.1]octane-8,1'-pyrrolidinium] chloride, 2-diethylaminoethyl (bicyclohexyl)-1-carboxylate hydrochloride, 1-methyl-4-piperidyl diphenylpropoxyacetate hydrochloride, (3R)-1-azabicyclo[2.2.2]oct-3-yl (1S)-1-phenyl-3,4-dihydroisoquinoline-2(1H)-carboxylate succinate, (S)-1-[2-(2,3-dihydro-5-benzofuranyl)ethyl]- α , α -diphenyl-3-pyrrolidineacetamide hydrobromide, (8R)-3 α -hydroxy-5-isopropyl-1 α H,5 α H-tropanium bromide ($\pm$ tropate), 8-butyl-6 β ,7 β -epoxy-3 α -[(S)-3-hydroxy-2-phenylpropanoyloxy]tropanium bromide, (+)-(S)-N-methyl- γ -(1-naphthyloxy)-2-thiophenepropylamine,
2-[(1R)-3-(diisopropylamino)-1-phenylpropyl]-4-(hydroxymethyl)phenyl isobutyrate, 4-(diethylamino)-but-2-yn-1-yl (2S)-cyclohexyl(hydroxy)phenylacetate, ethyldimethyl(1-methyl-3,3-diphenylpropyl)ammonium bromide,
2-piperidinoethyl 3-methyl-4-oxo-2-phenyl-4H-chromene-8-carboxylate hydrochloride, benzyl(2-chloroethyl)-(1-methyl-2-phenoxyethyl)amine hydrochloride and/or a salt or another salt of these active ingredients.

WO 2008/006506

- 8 -

PCT/EP2007/005970

The active ingredient is preferably present in an amount of from 1 to 250 mg per dose unit on the active ingredient carriers, preferably in an amount which
5 corresponds to the daily dose. It is moreover possible for the active ingredient to be present in the form of a coating on the active ingredient carriers, which coating may, besides the active ingredient, optionally additionally comprise a binder and optionally antitack
10 agent and/or buffer substances in order to control the release of the active ingredient in the desired manner.

The active ingredient carriers are preferably in the form of neutral pellets and/or crystalline substances
15 and/or granulated or extruded carrier substances and advantageously have a particle size of from 0.1 to 3.0 mm, preferably from 0.2 to 2.0 mm.

Such active ingredient carriers are commercially
20 available and include as carrier material for example sugar, starch, microcrystalline cellulose, dicalcium phosphate, sodium chloride, citric acid, tartaric acid, malic acid, sucrose, lactose, sorbitol, mannitol, cellulose, calcium hydrogen phosphate, sodium citrate,
25 tricalcium phosphate, maize starch, wheat starch, potato starch, rice starch and/or mixtures thereof.

In an advantageous embodiment of the invention, the active ingredient carriers provided with the active
30 ingredient comprise per dose unit from 1 to 250 mg, advantageously 5 to 100 mg, more preferably 20 to 60 mg of the active ingredient, from 10 to 500 parts by weight of carrier material, from 1 to 100 parts by weight of binder, from 1 to 100 parts by weight of
35 antitack agent and from 1 to 100 parts by weight of buffer substances.

The binder of the active ingredient carrier provided with the active ingredient can be selected from the group comprising hydroxypropylmethylcellulose, butyl methacrylate-(2-dimethylaminoethyl) methacrylate-methyl
5 methacrylate copolymers, ethyl acrylate-methyl methacrylate copolymers, methacrylic acid-methyl methacrylate copolymers, methacrylic acid-ethyl acrylate copolymers, methyl acrylate-methyl methacrylate-methacrylic acid copolymers, ethyl
10 acrylate-methyl methacrylate-trimethylammoniummethyl methacrylate chloride copolymers, ethylcellulose, sodium carboxymethylcellulose, methylcellulose, hydroxyethylcellulose, methylethylcellulose, hydroxyethylmethylpropylcellulose, polyvinylpyrrolidone, poly-
15 vinyl acetate, vinylpyrrolidone-vinyl acetate copolymers, polyethylene glycol, gelatin, maize starch, wheat starch, rice starch and potato starch and mixtures thereof.

20 The active ingredient carrier provided with the active ingredient may comprise as antitack agent talc, stearic acid and salts thereof, mono-, di-, triglycerides with straight- and/or branched-chain fatty acids, fatty alcohols, colloidal silicon dioxide, precipitated
25 silicon dioxide, aluminium oxide, hydrogenated castor oil and macrogol (polyethylene glycol) and mixtures thereof.

Buffer substances which can be employed to prevent
30 possible pH shifts in the bowel region of the active ingredient carriers provided with the active ingredient are sodium hydroxide, citric acid, tartaric acid, phosphoric acid, ascorbic acid, succinic acid, adipic acid, fumaric acid and their pharmaceutically
35 acceptable salts and mixtures thereof.

The gastro-resistant coating layer disposed on the inner layer to control the release of active ingredient

WO 2008/006506

- 10 -

PCT/EP2007/005970

is preferably formed from the group comprising ethylcellulose, cellulose acetate phthalate, cellulose acetate succinate, shellac, hydroxypropyl-methylcellulose acetate succinate, cellulose acetate, 5 cellulose acetate propionate, hydroxypropylmethyl-cellulose phthalate, polyvinyl acetate phthalate, cellulose acetate butyrate, butyl methacrylate-2-dimethylaminoethyl methacrylate-methyl methacrylate copolymer, ethyl acrylate-methyl methacrylate 10 copolymer, methacrylic acid-methyl methacrylate copolymer, methacrylic acid-ethyl acrylate copolymer, methacrylic acid-methyl methacrylate copolymer, methyl acrylate-methyl methacrylate-methacrylic acid copolymer, ethyl acrylate-methyl methacrylate- 15 trimethylaminoethyl methacrylate chloride copolymer and/or mixtures thereof.

With one type of intake of the active ingredient carriers, in which the latter come directly into 20 contact with the saliva in the mouth, it is possible, because of the pH conditions, for the gastro-resistant layer to dissolve, in which case active ingredient is released there. In order to prevent this, it is preferred according to the invention to provide the 25 active ingredient carriers, which have been provided with the active ingredient, the diffusion layer and, where appropriate, the gastro-resistant coating, with a saliva-resistant outer layer which is preferably formed from ethylcellulose, cellulose acetate phthalate, 30 cellulose acetate succinate, shellac, hydroxypropyl-methylcellulose acetate succinate, cellulose acetate, cellulose acetate propionate, hydroxypropylmethyl-cellulose phthalate, polyvinyl acetate, polyvinyl acetate phthalate, polyvinyl alcohol-polyethylene 35 glycol copolymer, cellulose acetate butyrate, butyl methacrylate-2-dimethylaminoethyl methacrylate-methyl methacrylate copolymer, ethyl acrylate-methyl methacrylate copolymer, methacrylic acid-methyl

WO 2008/006506

- 11 -

PCT/EP2007/005970

methacrylate copolymer, methacrylic acid-ethyl acrylate
copolymer, methacrylic acid-methyl methacrylate
copolymer, methyl acrylate-methyl methacrylate-
methacrylic acid copolymer, ethyl acrylate-methyl
5 methacrylate-trimethylaminoethyl methylacrylate
chloride copolymer and/or mixtures thereof.

The saliva-resistant outer layer can if desired
comprise aromatizing substances, flavourings and/or
10 sweeteners.

In a further preferred embodiment of the invention, the
active ingredient carriers which have been provided
with the active ingredient, the diffusion layers, the
15 gastro-resistant coating layer, and, where appropriate,
the saliva-resistant outer layer, are compressed to
tablets using conventional excipients, or packed into
capsules made of gelatin, cellulose, starch or starch
derivatives, or are in the form of liquids or semisolid
20 or solid preparations for preparing a suspension or a
suspension gel.

The pharmaceutical preparation of the invention can be
produced using procedures known per se, for example by
25 the active ingredient carrier, which is for example in
the form of commercially available pellets, being
coated with a solution or suspension which comprises
the active ingredient, the binder, the antitack agent
and, where appropriate, buffer substances, then
30 providing the intermediate product obtained in this way
successively with at least two diffusion layers whose
permeability for the diffusing active ingredient
decreases from the inside to the outside, applying the
gastro-resistant coating layer and, where appropriate,
35 providing a saliva-resistant outer layer.

It has emerged that the control according to the
invention of the release of active ingredient via the

WO 2008/006506

- 12 -

PCT/EP2007/005970

inner layer composed of at least two diffusion layers whose permeability for the diffusing active ingredient decreases from the inside to the outside has the surprising advantage that, by comparison with other
5 known formulations, there is a smaller increase in release of active ingredient and thus the desired lower blood levels of the active ingredient are obtained, with the active ingredient release characteristics surprisingly being even more favourable by comparison
10 with the teaching of US 2004/0142035 A1 through the fact that two layers differing in diffusion permeability are applied instead of one layer which comprises the two materials with different diffusion characteristics in the form of a mixture.

15

This situation is to be explained in more detail by the following examples and comparative examples.

Two dosage forms are possible and preferred for the
20 pharmaceutical preparation of the invention:

1. A capsule which is filled with the active ingredient carriers provided with the active ingredient, the diffusion layers, the gastro-resistant coating layer and the optional saliva-resistant outer layer;
- 25 2. A tablet which has been obtained by compressing the active ingredient carriers provided with the active ingredient, the diffusion layers, the gastro-resistant coating layer and, where appropriate, the saliva-resistant outer layer with use of conventional
30 excipients.

The tablet may be divisible and can either be swallowed with liquid, taken with suitable food in small pieces, or else sucked or allowed to disintegrate in the mouth,
35 without the desired release of active ingredient in the small bowel being impaired thereby, because with these types of intake the tablet disintegrates into the individual or agglomerated active ingredient carriers,

WO 2008/006506

- 13 -

PCT/EP2007/005970

which active ingredient carriers each comprise the active ingredient, the diffusion layers and the gastro-resistant coating layer, thus ensuring the desired active ingredient release profile.

5

The presence of a saliva-resistant outer layer with the types of intake of the pharmaceutical preparation results in no impairment of the gastro-resistant layer because of the pH conditions in the mouth, and thus the active ingredient is not released in the mouth but, as intended, only in the small bowel.

10

EXAMPLE 1 Batch: 050046/2

15

Pharmaceutical preparation of the invention in the form of pellets with an active ingredient content of 45 mg of (1R,3r,5S)-3-[(hydroxydiphenylacetyl)oxy]spiro[8-azoniabicyclo[3.2.1]octane-8,1'-pyrrolidinium] chloride as active ingredient per dosage unit

20

The ingredients for producing the active ingredient carriers provided with active ingredient are listed in Table 1 below.

25

Table 1		
Formulation, application of the active ingredient		
No.	Starting material	per dose mg
1	Active ingredient*	45.000
2	Neutral pellets	100.000
3	Hydroxypropylmethylcellulose (hypromellose)	4.500
4	Talc	4.500
5	Polyethylene glycol (macrogol 6000)	0.450
	Total amount per dose	154.450

* (1R,3r,5S)-3-[(Hydroxydiphenylacetyl)oxy]spiro-[8-azoniabicyclo[3.2.1]octane-8,1'-pyrrolidinium] chloride

WO 2008/006506

- 14 -

PCT/EP2007/005970

The neutral pellets are sugar-starch pellets with a particle size in the range from 0.1 to 3.0 mm. The active ingredient, the hydroxypropylmethylcellulose and macrogol 6000 are dissolved in water, and talc is suspended in the solution. The suspension is continuously sprayed onto the active ingredient carriers in a fluidized bed granulator, and simultaneously dried at an inlet air temperature of 30°C to 80°C.

10

Subsequently, diffusion layer 1 is applied to the active ingredient carriers provided with the active ingredient in this way, using the ingredients indicated in Table 2 below:

15

Table 2		
Formulation, application of a diffusion layer for optimized release of active ingredient		
Formulation of diffusion layer 1		
No.	Starting material	per dose mg
1	Active ingredient pellets (as in Table 1)	154.450
2	Polyvinyl acetate stabilized with polyvinylpyrrolidone and sodium lauryl sulphate (Kollicoat SR 30 D)	9.000
3	Polyvinyl alcohol-polyethylene glycol copolymer (Kollicoat IR)	1.8000
4	Propylene glycol	0.900
5	Talc	0.360
	Total amount per dose	166.510

Diffusion layer 1 is produced by spraying the active ingredient carriers which have been provided with the active ingredient and obtained using the ingredients indicated in Table 1 with an aqueous suspension of polyvinyl acetate as matrix material and polyvinyl alcohol-polyethylene glycol copolymer as pore former, propylene glycol as plasticizer and talc as antitack agent.

25

Subsequently, diffusion layer 2 is applied using the ingredients indicated in Table 3 below:

Table 3		
Formulation of diffusion layer 2		
No.	Starting material	per dose mg
1	Active ingredient pellets with diffusion layer 1 (as in Table 2)	166.510
2	Polyvinyl acetate stabilized with polyvinylpyrrolidone and sodium lauryl sulphate (Kollicoat SR 30 D)	9.000
3	Polyvinyl alcohol-polyethylene glycol copolymer (Kollicoat IR)	0.405
4	Propylene glycol	0.900
5	Talc	0.360
	Total amount per dose	177.175

5

The active ingredient pellets provided with the active ingredient and diffusion layer 1 and obtained using the ingredients in Table 2 are sprayed with an aqueous suspension of polyvinyl acetate as matrix material and polyvinyl alcohol-polyethylene glycol copolymer as pore former, propylene glycol as plasticizer and talc as antitack agent. As is evident, the material for forming diffusion layer 2 comprises a smaller amount of the pore former polyvinyl alcohol-polyethylene glycol copolymer and accordingly leads to a diffusion layer with a lower permeability for the diffusing active ingredient (1R,3r,5S)-3-[(hydroxydiphenylacetyl)oxy]-spiro[8-azoniabicyclo[3.2.1]octane-8,1'-pyrrolidinium] chloride.

20

The active ingredient carriers provided with the two diffusion layers in this way are subsequently provided with a gastro-resistant coating using the formulation indicated in Table 4 below:

25

Table 4		
Formulation, application of a gastro-resistant coating		
No.	Starting material	per dose mg
1	Active ingredient pellets with diffusion layers 1 and 2 (as in Table 3) one dose comprises 45 mg of active ingredient	177.175
2	Methacrylic acid-ethyl acrylate copolymer (Kollicoat MAE 30 DP)	28.000
3	Talc	12.600
4	Propylene glycol	4.200
5	Sodium carboxymethylcellulose (Tylopur C30 G1)	0.720
	Total amount per dose	222.695

The active ingredient pellets obtained with both diffusion layers 1 and 2 are then provided with the gastro-resistant coating layer using the formulation indicated in Table 4 above, with an aqueous suspension of methacrylic acid-ethyl acrylate copolymer as matrix material, propylene glycol as plasticizer, sodium carboxymethylcellulose as binder and talc as antitack agent being continuously sprayed onto the active ingredient pellets provided with both diffusion layers 1 and 2 and simultaneously dried at an inlet air temperature of 35 to 80°C, resulting in the pharmaceutical preparation of the invention in the form of the active ingredient pellets provided with the active ingredient.

EXAMPLE 2 (comparative) Batch: 050047

For comparison, active ingredient carriers provided with active ingredient using the same ingredients and the same ratios of amounts but comprising the ingredients of diffusion layers 1 and 2 of the invention not in separate layers but in a single diffusion layer are produced.

WO 2008/006506

- 17 -

PCT/EP2007/005970

For this purpose, active ingredient carriers provided with the active ingredient (1R,3r,5S)-3-[(hydroxy-diphenylacetyl)oxy]spiro[8-azoniabicyclo[3.2.1]octane-8,1'-pyrrolidinium] chloride using the ingredients indicated in Table 1 of Example 1 are produced and provided with a diffusion layer using the ingredients listed in Table 5 below:

Table 5		
Formulation of diffusion layer 1 + 2 applied as mixture		
No.	Starting material	per dose mg
1	Active ingredient pellets (as in Table 1)	154.450
2	Polyvinyl acetate stabilized with polyvinylpyrrolidone and sodium lauryl sulphate (Kollicoat SR 30 D)	18.000
3	Polyvinyl alcohol-polyethylene glycol copolymer (Kollicoat IR)	2.205
4	Propylene glycol	1.800
5	Talc	0.720
	Total amount per dose	177.175

10 A gastro-resistant coating is then applied by the procedure of Example 1, and using the ingredients indicated in Table 4 of Example 1, to the active ingredient pellets provided with diffusion layer 1+2, applied as mixture, obtained in this way, and active
 15 ingredient pellets of the composition indicated in Table 6 below are obtained:

Table 6		
No.	Starting material	per dose mg
1	Active ingredient pellets with diffusion layer 1+2 (as in Table 5) one dose comprises 45 mg of active ingredient	177.175
2	Methacrylic acid-ethyl acrylate copolymer (Kollicoat MAE 30 DP)	28.000
3	Talc	12.600
4	Propylene glycol	4.200
5	Sodium carboxymethylcellulose (Tylopur C30 G1)	0.720
	Total amount per dose	222.695

EXAMPLE 3 (comparative) Batch: 050046/1

As further comparative example, active ingredient carriers provided only with the active ingredient (1R,3r,5S)-3-[(hydroxydiphenylacetyl)oxy]spiro[8-azoniabicyclo[3.2.1]octane-8,1'-pyrrolidinium] chloride and not comprising either a diffusion layer or a gastro-resistant coating are prepared, specifically by the procedure of Example 1 and using the ingredients indicated in Table 1, resulting in the active ingredient carriers provided with the active ingredient (1R,3r,5S)-3-[(hydroxydiphenylacetyl)oxy]spiro[8-azoniabicyclo[3.2.1]octane-8,1'-pyrrolidinium] chloride) of the composition indicated in Table 7 below.

Table 7		
No.	Starting material	per dose mg
1	Active ingredient	45.000
2	Neutral pellets	100.000
3	Hydroxypropylmethylcellulose [hypromellose)	4.5000
4	Talc	4.5000
5	Polyethylene glycol 6000 (macrogol 6000)	0.450
	Total amount per dose	154.450

WO 2008/006506

- 19 -

PCT/EP2007/005970

EXAMPLE 4 (comparative) Batch: 050048

This comparative example relates to active ingredient carriers provided with the active ingredient
 5 (1R,3r,5S)-3-[(hydroxydiphenylacetyl)oxy]spiro[8-azoniabicyclo[3.2.1]octane-8,1'-pyrrolidinium] chloride and provided only with one diffusion layer, specifically diffusion layer 1 of Example 1 of the present invention, and the gastro-resistant coating
 10 layer.

The starting material comprises the active ingredient pellets provided with the active ingredient which had been obtained by the procedure of Example 1 using the
 15 ingredients indicated in Table 1.

These active ingredient carriers are provided only with diffusion layer 1 using the ingredients and the procedure indicated for forming diffusion layer 1 in
 20 Example 1 with reference to Table 2.

Application of a gastro-resistant coating layer by the procedure of Example 1 using the ingredients indicated in Table 4 results in active ingredient pellets loaded
 25 with the active ingredient of the composition indicated in Table 8 below:

Table 8		
No.	Starting material	per dose mg
1	Active ingredient pellets with diffusion layer 1 one dose comprises 45 mg of active ingredient	166.510
2	Methacrylic acid-ethyl acrylate copolymer (Kollicoat MAE 30 DP)	28.000
3	Talc	12.600
4	Propylene glycol	4.200
5	Sodium carboxymethylcellulose (Tylopur C30 G1)	0.720
	Total amount per dose	212.030

WO 2008/006506

- 20 -

PCT/EP2007/005970

EXAMPLE 5 (comparative) Batch: 050049

This comparative example relates to active ingredient carriers provided with the active ingredient (1R,3r,5S)-3-[(hydroxydiphenylacetyl)oxy]spiro[8-azoniabicyclo[3.2.1]octane-8,1'-pyrrolidinium] chloride and provided only with diffusion layer 2 of the pharmaceutical preparation of the invention according to Example 1 and with the gastro-resistant coating layer.

The starting material for this purpose comprises the active ingredient carriers provided with the active ingredient which had been produced using the ingredients indicated in Table 1 of Example 1 by the procedure described therein.

Diffusion layer 2 is applied to these active ingredient pellets provided with the active ingredient, using the ingredients indicated in Table 9 below and employing the procedure indicated in connection with Table 3 in Example 1.

Table 9		
No.	Starting material	per dose mg
1	Active ingredient pellets (as in Table 1)	154.450
2	Polyvinyl acetate stabilized with polyvinylpyrrolidone and sodium lauryl sulphate (Kollicoat SR 30D)	9.000
3	Polyvinyl alcohol-polyethylene glycol copolymer (Kollicoat IR)	0.405
4	Propylene glycol	0.900
5	Talc	0.360
	Total amount per dose	165.115

25

Application of a gastro-resistant coating layer by the procedure of Example 1 using the ingredients indicated in Table 4 results in active ingredient pellets with

WO 2008/006506

- 21 -

PCT/EP2007/005970

diffusion layer 2 and the composition indicated in Table 10 below.

Table 10		
Formulation, application of a gastro-resistant coating		
No.	Starting material	per dose mg
1	Active ingredient pellets with diffusion layer 2 one dose comprises 45 mg of active ingredient	165.115
2	Methacrylic acid-ethyl acrylate copolymer (Kollicoat MAE 30 DP)	28.000
3	Talc	12.600
4	Propylene glycol	4.200
5	Sodium carboxymethylcellulose (Tylopur C30 G1)	0.720
	Total amount per dose	210.635

EXAMPLE 6

5

Investigation of the in vitro release of the active ingredient (1R,3r,5S)-3-[(hydroxydiphenylacetyl)oxy]-spiro[8-azoniabicyclo[3.2.1]octane-8,1'-pyrrolidinium] chloride from the active ingredient pellets obtained as in Examples 1 to 5 above.

10

To investigate the release of active ingredient as a function of time, the US Pharmacopeia method for determining the release of products with a gastro-resistant coating No. 724 Method A is employed as follows:

15

Acid stage

20

Apparatus: paddle stirrer
 Speed of rotation: 200 min⁻¹
 Temperature: 37 ± 0.5°C
 Dissolving medium: 750 ml of 0.1 M hydrochloric acid

WO 2008/006506

- 22 -

PCT/EP2007/005970

A dose of active ingredient pellets produced according to the above examples is introduced into the apparatus, the vessel is covered and the paddle stirrer is operated for 2 hours. A sample is then taken to
5 determine the active ingredient (1R,3r,5S)-3-[(hydroxydiphenylacetyl)oxy]spiro[8-azoniabicyclo-[3.2.1]octane-8,1'-pyrrolidinium] chloride.

Buffer stage

10

250 ml of a 0.20 M tribasic sodium phosphate solution equilibrated at $37 \pm 0.5^{\circ}\text{C}$ are added to the acid stage (750 ml of 0.1M hydrochloric acid). The pH is adjusted to a value of 6.8 ± 0.05 if necessary with 2 N
15 hydrochloric acid or 2 N sodium hydroxide solution. The apparatus is operated for 2 hours and then samples are taken after 15, 30, 45, 60, 75, 90, 105 and 120 minutes, and the active ingredient therein is determined by HPLC.

20

Reference solution:

0.045 mg/ml (1R,3r,5S)-3-[(hydroxydiphenylacetyl)oxy]-spiro[8-azoniabicyclo[3.2.1]octane-8,1'-pyrrolidinium]
25 chloride RS in 0.1 N hydrochloric acid.

HPLC chromatography conditions

The chromatography takes place using:

30

- a column of stainless steel with a length of 0.125 m and an internal diameter of 4 mm, which is packed with octadecylsilyl gel for chromatographic purposes (5 μm), specifically Nucleosil 100-5 C18;
- a mixture of 65 volumes of purified water which
35 comprises 2.202 g/l sodium heptane sulphonate and 0.4 ml/l triethylamine and has been adjusted to a pH of 2.5 with concentrated phosphoric acid, and

WO 2008/006506

- 23 -

PCT/EP2007/005970

35 volumes of acetonitrile, is used as mobile phase.

- A spectrophotometer (DAD 210.10, 400.100 nm); set at 210 nm is used as detector;
- 5 - the temperature of the column is kept at 40°C.
- 25 µl of the sample liquid are injected, and the chromatography is performed for about 4.5 minutes.

The concentration of dissolved (1R,3r,5S)-3-[(hydroxy-diphenylacetyl)oxy]spiro[8-azoniabicyclo[3.2.1]octane-
10 8,1'-pyrrolidinium] chloride is calculated in per cent using the following equation:

$$Y = CR_{(w)} \cdot AT_{(w)} \cdot 100 / AR_{(w)} \cdot CT_{(w)}$$

15 in which

Y: dissolved active ingredient (1R,3r,5S)-3-[(hydroxydiphenylacetyl)oxy]spiro[8-
20 azoniabicyclo[3.2.1]octane-8,1'-pyrrolidinium] chloride (% of the stated content)

CT_(w): nominal concentration of active ingredient in the test solution (0.045 mg/ml)

CR_(w): concentration of the active ingredient in the reference solution

25 AT_(w): area of active ingredient peak of the test solution

AR_(w): area of reference substance peak in the reference solution

30 The results of the release of active ingredient obtained in this investigation of the active ingredient pellets of Examples 1 to 5 are compiled in Table 11 below:

WO 2008/006506

- 24 -

PCT/EP2007/005970

PU040014

W 45 tablets MOF

Active ingredient pellets and gastro-resistant pellets
differing in application of the diffusion layers

5

Table 11					
Results of the release of active ingredient*					
	Ex.3 (Ref.)	Ex.1 (invention)	Ex.2 (comp.)	Ex. 4 (comp.)	Ex. 5 (comp.)
Test time/	Active ingredient pellets without diffusion layer	Diffusion layer 1+2 applied successively	Diffusion layer 1+2 applied as mixture	Diffusion layer 1 applied alone	Diffusion layer 2 applied alone
Batch	050046/1	050046/2	050047	050048	050049
0	0.00	0	0	0	0
15	98.02	0.76	73.94	101.02	7.02
30	98.30	8.14	91.12	101.24	34.27
45	98.39	29.89	97.84	101.03	67.08
60	98.47	51.13	98.75	101.20	81.80
75	98.65	65.47	99.59	101.25	88.84
90	98.56	74.35	99.45	101.30	92.39
105	98.49	80.03	99.99	101.35	94.63
120	98.68	84.09	99.89	101.10	95.87

*(1R,3r,5S)-3-[(hydroxydiphenylacetyl)oxy]spiro[8-azo-
niabicyclo[3.2.1]octane-8,1'-pyrrolidinium] chloride

- 10 The results of Table 11 above are depicted as a graph in **Figure 1** below, specifically by plotting the percentage release of active ingredient as a function of time.

WO 2008/006506

- 25 -

PCT/EP2007/005970

As is evident from Figure 1 above, the pharmaceutical preparation (050046/2) of the invention shows the best course as desired according to the invention for the release of active ingredient in vitro with a slow rise and subsequent almost constant release of active ingredient, whereas the comparative products having no diffusion layer (050046/1) or having in each case only one diffusion layer 1 or 2 (050049 and 050048) or the active ingredient pellets of Example 2 in which the ingredients of the two diffusion layers 1 and 2 of the invention are present in a single diffusion layer (050047) show a fast rise in the curves and thus an unwanted rapid release of active ingredient.

It is to be regarded as surprising that the pharmaceutical preparation of the invention in which the inner layer is formed from at least two diffusion layers whose permeability for the diffusing active ingredient decreases from the inside to the outside shows a considerably better course of release of active ingredient than the active ingredient pellets of Example 2, in which the same ingredients for forming the two diffusion layers of the invention are present in a single diffusion layer. This pharmaceutical preparation also shows an unwantedly rapid release of active ingredient at the start, since approximately 95% are released after only 40 minutes, at a time when the pharmaceutical preparation of the invention results in a release of only about 20% active ingredient.

It is thus possible with the pharmaceutical preparation of the invention to achieve, by comparison with a single administration of a pharmaceutical product rapidly releasing the active ingredient (1R,3r,5S)-3-[(hydroxydiphenylacetyl)oxy]spiro[8-azoniabicyclo-[3.2.1]octane-8,1'-pyrrolidinium] chloride, a slower rise in the level of active ingredient in the plasma with an overall lower blood level, and thus to overcome

WO 2008/006506

- 26 -

PCT/EP2007/005970

the disadvantages of conventional products of this type.

WO 2008/006506

- 27 -

PCT/EP2007/005970

AMENDED CLAIMS

**received at the International Bureau on 26 October 2007
(26.10.07)**

1. Pharmaceutical preparation for oral administration
5 with controlled release of active ingredient in the
small bowel, on the basis of active ingredient carriers
provided with at least one active ingredient which are
provided with an inner layer to control the release of
10 active ingredient and with a gastro-resistant coating
layer disposed thereon, characterized in that the inner
layer is formed from at least two diffusion layers
whose permeability for the diffusing active ingredient
decreases from the inside to the outside, the diffusion
15 layers being formed from a matrix material which is
insoluble in small bowel fluid and into which pore
formers which control the permeability for the
diffusing active ingredient and are soluble and/or
swellable and/or water-absorbing in small bowel fluid
are incorporated.

20

2. Pharmaceutical preparation according to Claim 1,
characterized in that the diffusion layers comprise
identical or different types of matrix materials.

25

3. Pharmaceutical preparation according to at least
either of Claims 1 and 2, characterized in that the
diffusion layers comprise one or more pore formers
which are identical or of different types.

30

4. Pharmaceutical preparation according to at least
one of the preceding claims, characterized in that the
permeability of the diffusion layers for the diffusing
active ingredient is controlled via the nature, the
quantity and/or the particle size and/or the solubility
35 and/or the swellability and/or the water-absorbing
capacity of the pore formers.

WO 2008/006506

- 28 -

PCT/EP2007/005970

5. Pharmaceutical preparation according to Claim 4, characterized in that the inner layer for controlling the release of active ingredient includes an inner and an outer diffusion layer.

5

6. Pharmaceutical preparation according to Claim 5, characterized in that in the case of identical pore former material the particle size and/or the concentration of the pore formers in the inner
10 diffusion layer is greater than in the outer diffusion layer.

7. Pharmaceutical preparation according to Claim 6, characterized in that in the case of pore formers which
15 are soluble in small bowel fluid the ratio of the pore former concentration in the inner diffusion layer to the pore former concentration in the outer diffusion layer is in the range from 20:1 to > 1:1, preferably 10:1 to 1.1:1.

20

8. Pharmaceutical preparation according to at least one of the preceding claims, characterized in that the matrix material of the diffusion layers is selected from the group comprising ethylcellulose, cellulose
25 acetate phthalate, cellulose acetate succinate, shellac, hydroxypropylmethylcellulose acetate succinate, cellulose acetate, cellulose acetate propionate, hydroxypropylmethylcellulose phthalate, polyvinyl acetate, polyvinyl acetate phthalate,
30 cellulose acetate butyrate, butyl methacrylate-(2-dimethylaminoethyl) methacrylate-methyl methacrylate copolymers, ethyl acrylate-methyl methacrylate copolymers, methacrylic acid-methyl methacrylate copolymers, methacrylic acid-ethyl acrylate copolymers,
35 ethyl acrylate-methyl methacrylate-trimethyl-ammoniummethyl methacrylate chloride copolymers, sodium carboxymethylcellulose, methylcellulose, hydroxyl-

WO 2008/006506

- 29 -

PCT/EP2007/005970

ethylcellulose, methylethylcellulose, methylhydroxypropylcellulose, hydroxypropylmethylcellulose, hydroxyethylmethylpropylcellulose, chitosan, chitosan acetate, silicone elastomer latex suspensions, hydrogenated castor oil, stearic acid, glycerol monostearate, glycerol distearate, glycerol dibehenate, stearyl alcohol, white wax, yellow wax, hydrogenated vegetable oil and microcrystalline wax.

9. Pharmaceutical preparation according to at least one of the preceding claims, characterized in that the pore former of the diffusion layers is selected from the group comprising polyvinyl alcohol-polyethylene glycol copolymers, polyvinylpyrrolidone, vinylpyrrolidone-vinyl acetate copolymers, crosslinked polyvinylpyrrolidone, microcrystalline cellulose, hydroxypropylcellulose, sodium carboxymethylcellulose, methylcellulose, hydroxyethylcellulose, methylethylcellulose, methylhydroxypropylcellulose, hydroxypropylmethylcellulose, hydroxyethylmethylpropylcellulose, polyethylene glycol, cellulose powder, sucrose, lactose, mannitol, sorbitol and polysorbate.

10. Pharmaceutical preparation according to at least one of the preceding claims, characterized in that the matrix material of the diffusion layers additionally comprises plasticizers and antitack agents.

11. Pharmaceutical preparation according to Claim 10, characterized in that the matrix material of the diffusion layers comprises polyethylene glycol, propylene glycol, triethyl citrate, triacetin, acetyl tributyl citrate, polysorbates, 2-pyrrolidone, dibutyl sebacate, stearic acid, castor oil and/or medium-chain triglycerides as plasticizer, and talc, stearic acid and salts thereof, fatty alcohols, mono-, di- or triglycerides with straight- and/or branched-chain

WO 2008/006506

- 30 -

PCT/EP2007/005970

fatty acids, colloidal silicon dioxide, precipitated silicon dioxide, aluminium oxide, kaolin, maize starch, wheat starch, rice starch, potato starch, titanium dioxide, silicone emulsions and/or Veegum (a magnesium aluminium silicate dispersion) as antitack agent.

12. Pharmaceutical preparation according to at least one of the preceding claims, characterized in that the active ingredient carriers comprise as active ingredient 4-diethylamino-2-butynyl α -phenylcyclohexaneglyconate hydrochloride, ethyldimethyl(1-methyl-3,3-diphenylpropyl)ammonium carrageenate, (+)-[R-2-[α -2-(diisopropylamino)ethyl]benzyl]-p-cresol tartrate, 8-(cyclopropylmethyl)-6 β ,7 β -epoxy-3 α -hydroxy-1 α H,5 α)-(S)-tropate bromide, (1R,3r,5S)-3-[(hydroxyldiphenylacetyl)oxy]spiro[8-azoniabicyclo[3.2.1]octane-8,1'-pyrrolidinium] chloride, 2-diethylaminoethyl(bicyclohexyl)-1-carboxylate hydrochloride, 1-methyl-4-piperidyl diphenylpropoxyacetate hydrochloride, (3R)-1-azabicyclo[2.2.2]oct-3-yl (1S)-1-phenyl-3,4-dihydroisoquinoline-2(1H)-carboxylate succinate, (S)-1-[2-(2,3-dihydro-5-benzofuranyl)ethyl]- α,α -diphenyl-3-pyrrolidineacetamide hydrobromide, (8R)-3 α -hydroxy-5-isopropyl-1 α H,5 α H-tropanium bromide ($\pm$ tropate), 8-butyl-6 β ,7 β -epoxy-3 α -[(S)-3-hydroxy-2-phenylpropanoyloxy]tropanium bromide, (+)-(S)-N-methyl- γ -(1-naphthyloxy)-2-thiophenepropylamine, 2-[(1R)-3-(diisopropylamino)-1-phenylpropyl]-4-(hydroxymethyl)phenyl isobutyrate, 4-(diethylamino)-but-2-yn-1-yl (2S)-cyclohexyl(hydroxy)phenylacetate, ethyldimethyl(1-methyl-3,3-diphenylpropyl)ammonium bromide, 2-piperidinoethyl 3-methyl-4-oxo-2-phenyl-4H-chromene-8-carboxylate hydrochloride, benzyl(2-chloroethyl)(1-methyl-2-phenoxyethyl)amine hydrochloride and/or a salt or another salt of these active ingredients.

WO 2008/006506

- 31 -

PCT/EP2007/005970

13. Pharmaceutical preparation according to Claim 12, characterized in that the active ingredient is present in an amount of from 1 to 250 mg per dose unit on the active ingredient carriers.

5

14. Pharmaceutical preparation according to Claim 12 or 13, characterized in that the active ingredient is present in the form of a coating on the active ingredient carriers.

10

15. Pharmaceutical preparation according to at least one of the preceding claims, characterized in that the active ingredient carriers are in the form of neutral pellets and/or crystalline substances and/or granulated or extruded carrier substances.

15

16. Pharmaceutical preparation according to Claim 15, characterized in that the active ingredient carriers have a particle size of from 0.1 to 3.0 mm.

20

17. Pharmaceutical preparation according to Claim 15 or 16, characterized in that the active ingredient carriers include as carrier material sugar, starch, microcrystalline cellulose, dicalcium phosphate, sodium chloride, citric acid, tartaric acid, malic acid, sucrose, lactose, sorbitol, mannitol, cellulose, calcium hydrogen phosphate, sodium citrate, tricalcium phosphate, maize starch, wheat starch, potato starch, rice starch and/or mixtures thereof.

25

30

18. Pharmaceutical preparation according to at least one of the preceding claims, characterized in that the active ingredient carriers provided with the active ingredient additionally have, besides the active ingredient, on their surface a binder and where appropriate antitack agent and/or buffer substances.

35

WO 2008/006506

- 32 -

PCT/EP2007/005970

19. Pharmaceutical preparation according to Claim 18, characterized in that the active ingredient carriers provided with the active ingredient comprise per dose unit from 1 to 250 mg active ingredient, from 10 to 500 parts by weight of carrier material, from 1 to 100 parts by weight of binder, from 1 to 100 parts by weight of antitack agent and from 1 to 100 parts by weight of buffer substance.
20. Pharmaceutical preparation according to Claims 18 and 19, characterized in that the binder of the active ingredient carriers provided with active ingredient is selected from the group comprising hydroxypropylmethylcellulose, butyl methacrylate-(2-dimethylaminoethyl) methacrylate-methyl methacrylate copolymers, ethyl acrylate-methyl methacrylate copolymers, methacrylic acid-methyl methacrylate copolymers, methacrylic acid-ethyl acrylate copolymers, methyl acrylate-methyl methacrylate-methacrylic acid copolymers, ethyl acrylate-methyl methacrylate-trimethylammoniummethyl methacrylate chloride copolymers, ethylcellulose, sodium carboxymethylcellulose, methylcellulose, hydroxyethylcellulose, methylethylcellulose, hydroxyethylmethylpropylcellulose, polyvinylpyrrolidone, polyvinyl acetate, vinylpyrrolidone-vinyl acetate copolymers, polyethylene glycol, gelatin, maize starch, wheat starch, rice starch and potato starch and mixtures thereof.
21. Pharmaceutical preparation according to Claims 18 to 20, characterized in that the antitack agent of the active ingredient carriers provided with the active ingredient is selected from the group comprising talc, stearic acid and salts thereof, mono-, di-, triglycerides with straight- and/or branched-chain fatty acids, fatty alcohols, colloidal silicon dioxide, precipitated silicon dioxide, aluminium oxide,

WO 2008/006506

- 33 -

PCT/EP2007/005970

hydrogenated castor oil, and macrogol (polyethylene glycol) and mixtures thereof.

22. Pharmaceutical preparation according to Claims 18
5 to 21, characterized in that the buffer substances of
the active ingredient carriers provided with the active
ingredient are selected from the group comprising
sodium hydroxide, citric acid, tartaric acid,
phosphoric acid, ascorbic acid, succinic acid, adipic
10 acid, fumaric acid and their pharmaceutically
acceptable salts and mixtures thereof.

23. Pharmaceutical preparation according to at least
one of the preceding claims, characterized in that the
15 gastro-resistant coating layer disposed on the active
ingredient carrier provided with the active ingredient
and the diffusion layers is formed from the group
comprising ethylcellulose, cellulose acetate phthalate,
cellulose acetate succinate, shellac, hydroxypropyl-
20 methylcellulose acetate succinate, cellulose acetate,
cellulose acetate propionate, hydroxypropylmethyl-
cellulose phthalate, polyvinyl acetate phthalate,
cellulose acetate butyrate, butyl methacrylate-
(2-dimethylaminoethyl methacrylate-methyl methacrylate
25 copolymer, ethyl acrylate-methyl methacrylate
copolymer, methacrylic acid-methyl methacrylate
copolymer, methacrylic acid-ethyl acrylate copolymer,
methacrylic acid-methyl methacrylate copolymer, methyl
acrylate-methyl methacrylate-methacrylic acid copolymer,
30 ethyl acrylate-methyl methacrylate-trimethylaminoethyl
methacrylate chloride copolymer and/or mixtures
thereof.

24. Pharmaceutical preparation according to at least
35 one of the preceding claims, characterized in that the
active ingredient carriers provided with the active
ingredient, the diffusion layers and where appropriate

WO 2008/006506

- 34 -

PCT/EP2007/005970

the gastro-resistant coating are provided with a saliva-resistant outer layer.

25. Pharmaceutical preparation according to claim 24,
5 characterized in that the active ingredient carriers provided with the active ingredient, the diffusion layers and where appropriate the gastro-resistant coating are provided with a saliva-resistant outer layer from the group comprising ethylcellulose,
10 cellulose acetate phthalate, cellulose acetate succinate, shellac, hydroxypropylmethylcellulose acetate succinate, cellulose acetate, cellulose acetate propionate, hydroxypropylmethylcellulose phthalate, polyvinyl acetate, polyvinyl acetate phthalate,
15 polyvinyl alcohol-polyethylene glycol copolymer, cellulose acetate butyrate, butyl methacrylate-2-dimethylaminoethyl methacrylate-methyl methacrylate copolymer, ethyl acrylate-methyl methacrylate copolymer, methacrylic acid-methyl methacrylate
20 copolymer, methacrylic acid-ethyl acrylate copolymer, methacrylic acid-methyl methacrylate copolymer, methyl acrylate-methyl methacrylate-methacrylic acid copolymer, ethyl acrylate-methyl methacrylate-trimethylaminoethyl methacrylate chloride copolymer
25 and/or mixtures thereof.

26. Pharmaceutical preparation according to Claim 25,
characterized in that the saliva-resistant outer layer comprises aromatizing substances, flavourings and/or
30 sweeteners.

27. Pharmaceutical preparation according to at least one of the preceding claims, characterized in that the active ingredient carriers which have been provided
35 with the active ingredient, the diffusion layers, the gastro-resistant coating layer, and, where appropriate, the saliva-resistant outer layer, are compressed to

WO 2008/006506

- 35 -

PCT/EP2007/005970

tablets using conventional excipients, or packed into capsules made of gelatin, cellulose, starch or starch derivatives, or are in the form of liquids or semisolid or solid preparations for preparing a suspension or a
5 suspension gel.

28. Method for producing the pharmaceutical preparation according to Claims 1 to 27, characterized in that, using procedures known per se, the active
10 ingredient carriers are coated with a solution or suspension which comprises the active ingredient, the binder, the antitack agent and, where appropriate, buffer substances, then the at least two diffusion layers whose permeability for the diffusing active
15 ingredient decreases from the inside to the outside are successively applied, the gastro-resistant coating layer is applied and, where appropriate, a saliva-resistant outer layer is provided.




Figure 1