

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 August 2009 (27.08.2009)

PCT

(10) International Publication Number
WO 2009/104199 A1

(51) International Patent Classification:
C07K 14/62 (2006.01)

(21) International Application Number:
PCT/IN2008/000219

(22) International Filing Date:
3 April 2008 (03.04.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
00420/CHE/2008 19 February 2008 (19.02.2008) IN

(71) Applicant (for all designated States except US): **BIO-CON LIMITED** [IN/IN]; 20th KM Hosur Road, Electronic City P.O., Bangalore, Karnataka 560 100 (IN).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HAZRA, Partha** [IN/IN]; Apartment No.S-19, Meenakshi Residency, 41/1 Arekere Bannerghatta Road, Bangalore, Karnataka 560 076 (IN). **DAVE, Nitesh** [IN/IN]; # 424, Mahaveer Woods, 22nd Main, J P Nagar 5th Phase, Near IDBI Bank, Bangalore, Karnataka 560 078 (IN). **KANNAN, Vivekanandan** [IN/IN]; B2-205 L & T South City, Bannerghatta Road, Bangalore, Karnataka 560 076 (IN). **TIWARI, Sanjay** [IN/IN]; 20/A, 11th Cross, 6th A Main, JP Nagar, 3rd Phase, Bangalore, Karnataka 560 078 (IN). **GOEL, Anuj** [IN/IN]; H.No.72, 10th Cross, 29th Main, 1st Phase, JP Nagar, Bangalore, Karnataka 560 078 (IN). **IYER, Harish** [IN/IN]; 5C-221, 3rd Main Road, HRBR 3rd Block, Bangalore, Karnataka 560 043 (IN). **ROY, Nita** [IN/IN]; Flat A301, Wellington Park, Nos. 8 & 9 Wellington Street, Richmond Town, Bangalore, Karnataka 560 025 (IN). **VENKATESAN, Krishnamurthy** [IN/IN]; Plot # 82, Door # 22, Flat C, Grace Enclave, 2nd Main Road, North Jaganathan Nagar, Villivakkam, Chennai, Tamil Nadu 600 049 (IN). **VASUDEVAN, Anoop** [IN/IN]; Naduvathu Mana, Ambala Nada, West Chalakkudy, Trissur District, Kerala 680 037 (IN). **JA-**

GADISH, Anupama [IN/IN]; 602, E Block, 10th Main, 5th Cross, J P Nagar, Mysore, Karnataka 570 008 (IN). **SACHDEV, Goldy** [IN/IN]; No.644, 1st Cross, 1st Main, Indiranagar 1st Stage, Karnataka, Bangalore 560 038 (IN). **PATALE, Mukesh, Babuappa** [IN/IN]; Malli Galli, Ausa, Latur District, Maharashtra 413 520 (IN).

(74) Agents: **BHOLA, Ravi** et al.; K & S Partners, # 134, First Floor, 60 Ft. Domlur Road, Indiranagar, Bangalore, Karnataka 560 008 (IN).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report (Art. 21(3))

(54) Title: A METHOD OF OBTAINING PURIFIED HETEROLOGOUS INSULINS EXPRESSED IN YEAST

(57) Abstract: The invention relates to methods of separation and/or purification of impurities yielding a purified heterologous protein product devoid of related impurities or with substantially minimal quantities of such glycosylated impurities. More specifically, the invention relates to the identification of glycosylated forms of insulin analogues such as glargine impurities characterised post expression in yeast based systems such as *Pichia pastoris*. The invention also relates to methods used to clone gene encoding the protein insulin glargine; inserting the related gene in a suitable yeast host; producing culture of the recombinant strain, stimulating expression of the heterologous polypeptide, its secretion and purification post fermentation and related enzymatic conversions.



WO 2009/104199 A1

A METHOD OF OBTAINING PURIFIED HETEROLOGOUS INSULINS EXPRESSED IN YEAST

FIELD OF THE INVENTION:

The invention relates to methods of separation and/or purification of impurities yielding a purified heterologous protein product devoid of related impurities or with substantially minimal quantities of such glycosylated impurities. More specifically, the invention relates to the identification of glycosylated forms of insulin analogs such as glargine impurities characterized post expression in yeast based systems such as *Pichia pastoris*. The invention also relates to methods used to clone gene encoding the protein insulin glargine; inserting the related gene in a suitable yeast host; producing culture of the recombinant strain, stimulating expression of the heterologous polypeptide, its secretion and purification post fermentation and related enzymatic conversions.

BACKGROUND OF THE INVENTION AND PRIOR ART

Description of related art:

Recombinant forms of insulin, insulin analogues and/or derivatives have been produced in various microbial expression systems. Currently organisms such as *E.coli*, *S.cerevisiae* have been employed for the commercial production of recombinant human insulin and derivatives thereof. Owing to certain disadvantages of these systems such as low expression levels, difficulties in down stream purification etc. the use of methylotrophic yeast *Pichia pastoris* has been favored as a protein expression system. The expression system offers several advantages such as high expression, simple processing, low production cost, high density culture (US6800606).

Yeast expression systems are popular because they are easy to grow, are fast and scalable; however, some yeast expression systems have produced inconsistent results, and it is sometimes difficult to achieve high yields. One yeast expression system that has shown great promise is the methylotrophic *Pichia pastoris*. Compared to other eukaryotic expression systems, *Pichia* offers many advantages because it does not have the endotoxin problem associated with bacteria or the viral contamination problem of proteins produced in animal cell culture (Cino, Am Biotech Lab, May 1999). *Pichia*'s prolific growth rate

makes it easily scalable to large-scale production, although scale-up challenges include pH control, oxygen limitation, nutrient limitation, temperature fluctuation, and other safety considerations (Gottschalk, 2003, BioProcess Intl 1(4):54-61; Cino Am Biotech Lab, May 1999).

Albeit various advantages are attributed to yeast based expression systems such as *Pichia pastoris*, one of the major disadvantages of this system is the post-translational modification of resulting proteins which later exist as impurities in the final product that is difficult to purify. Although there are a number of post translational modifications of proteins known, the most common form of post translational modification is glycosylation. (Hart G.W, Glycosylation, Curr. Opin. Cell.Biol 1992; 4: 1017). Glycosylation can be either N-linked or O-linked depending on the expression system. (Gemmill TR et al., Overview of N- and O- linked oligosaccharide structures found in various yeast species, Biochemica et Biophysica Acta, 1999; 1426:227). Glycosylation affects stability of protein conformation, immunogenicity, clearance rate, protection from proteolysis and improves protein solubility. (Walsh G, Biopharmaceutical benchmarks 2006, Nature Biotechnology, 2006; 24:769).

Despite great advances in improving biotechnological manufacturing, no global solutions exist for every protein. The manufacturing process for a specific therapeutic protein requires novel and innovative solutions to problems that may be specific for that protein or family of proteins. Likewise, successful commercial applications often rely on a combination of specific properties of the protein or family of proteins, and the production processes used for manufacturing that protein or family proteins as pharmaceutical products.

The present invention relates to the identification of various glycoforms of insulin analogues more specifically insulin glargine through chemical methods coupled with mass spectrometry techniques such as electrospray and matrix assisted laser desorption ionisation for identification. The invention shall herewith permit selective purification of the product from the aforesaid impurities through optimized down stream purification methods attributed to the better understanding of the nature of impurities present in the

final product. The end product thus purified shall be substantially free of impurities characterized in the instant invention.

US 4,444,683 and its related applications claim glycosylated insulins, wherein glucose or mannose that is coupled with insulin via a spacer group derived from di-carboxylic acids, acid anhydrides or phenyl amines or a combination thereof.

WO 90/10645 specifically claims glycosylated insulin containing one or more monosaccharide groups or one or more oligosaccharide groups with upto 3 sugar units. Monoglycosylated or triglycosylated insulins at positions A1, B1 or B29.

WO 99/52934 claims a process of separating glycosylated proteins from non-glycosylated proteins by subjecting a solution comprising glycosylated and non-glycosylated proteins to chromatography using a Ca⁺⁺ containing eluent, and obtaining a fraction comprising non-glycosylated proteins, said fraction substantially free from glycosylated proteins.

Glycoforms of Glargine are not disclosed in any of the foregoing prior art. The invention discusses the methods of purifying insulin analogues such as insulin glargine with reduced levels of these characterized glycosylated impurities to obtain a 100% pure glargine product after fermentation through yeast based expression systems.

OBJECTS OF THE INVENTION:

The main object of the present invention is to develop a method of obtaining a purified, biologically active heterologous protein recombinantly expressed in a yeast expression system, characterized in that said purified protein is devoid of or contains substantially minimal amounts of glycosylated byproducts.

Another object of the present invention is to develop a process of obtaining a purified, biologically active heterologous protein recombinantly expressed in a host cell.

Yet another object of the present invention is to develop a process of producing a purified, biologically active heterologous protein.

Still another object of the present invention is to obtain a purified, biologically active heterologous protein product with purity of at least 96%.

Still another object of the present invention is to obtain a purified, biologically active heterologous protein product with purity in the range of 97 - 100%.

Still another object of the present invention is to obtain a purified insulin glargine with purity of at least 96%.

Still another object of the present invention is to obtain a purified insulin glargine containing less than 1% of glycosylated impurities.

Still another object of the present invention is to obtain a purified insulin glargine devoid of glycosylated impurities.

Still another object of the present invention is to obtain a purified insulin glargine devoid of glycosylated impurities, wherein the glycosylated form of the protein may be monoglycosylated, triglycosylated or polyglycosylated.

STATEMENT OF THE INVENTION

Accordingly, the present invention relates to a method of obtaining a purified, biologically active heterologous protein recombinantly expressed in a yeast expression system, characterized in that said purified protein is devoid of or contains substantially minimal amounts of glycosylated byproducts; a process of obtaining a purified, biologically active heterologous protein recombinantly expressed in a host cell which process comprises steps of: a) Culturing host cells transformed by a vector containing a DNA sequence defined by formula I encoding the heterologous protein under conditions suitable for the expression of the protein; b) Recovery of the expressed protein comprising separating the said protein from host cells to produce a recovered protein preparation; c) Subjecting the recovered protein of step (b) to a step of crystallization; d) Performing a step of enzymatic conversion

in the presence of trypsin or trypsin like enzyme; e) Purifying the said protein containing at least one related impurity comprising contacting the said protein mixture with a chromatographic matrix wherein purification is carried out employing a polar organic buffer solvent, in an aqueous phase containing an organic acid buffer; and f) Precipitating the eluted protein; a process of producing a purified, biologically active heterologous protein of claim 9 which comprises: a) Inoculating an aqueous fermentation medium with a transformant yeast strain harboring an expression vector which directs the expression of the said protein; and b) Growing the transformed strain in the fermentation medium under conditions effective for expression of said protein; a purified, biologically active heterologous protein with purity of at least 96%; a purified, biologically active heterologous protein product with purity in the range of 97 - 100%; purified insulin glargine with purity of at least 96%; purified insulin glargine containing less than 1% of glycosylated impurities; purified insulin glargine devoid of glycosylated impurities; and purified insulin glargine wherein the glycosylated form of the protein may be monoglycosylated, triglycosylated or polyglycosylated.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG 1: Graph of % yield versus Time (hrs) of trypsinised Insulin glargine and reaction at different concentrations.

FIG 2: Analytical Chromatogram of the reaction at different time point- before and after enzyme treatment.

FIG 3: HPLC profile of insulin-Glargine

FIG 4: Comparison of retention times of isolated impurities with final product.

FIG 5: RP-HPLC profile of the Insulin-glargine final product.

FIG 6: SEC-HPLC profile of the insulin glargine final product.

FIG 7: Electrospray ionisation mass spectra of glargine, monoglycosylated insulin glargine (mGIG) and triglycosylated (tGIG).


FIG 8: Matrix assisted laser desorption ionisation mass spectrum of monoglycosylated insulin glargine (mGIG)

FIG 9: Matrix assisted laser desorption ionisation mass spectrum of triglycosylated glargine (tGIG)

DESCRIPTION OF PREFERRED EMBODIMENTS:

The present invention relates to a method of obtaining a purified, biologically active heterologous protein recombinantly expressed in a yeast expression system, characterized in that said purified protein is devoid of or contains substantially minimal amounts of glycosylated byproducts.

In another embodiment of the present invention, the DNA encoding a heterologous protein as represented by formula I



wherein,

X is a leader peptide sequence comprising at least one amino acid.

B is the amino acid sequence of the B chain of the insulin molecule, its derivatives or analogs.

Y is a linker peptide comprising at least two amino acids.

A is the amino acid sequence of the A chain of the insulin molecule, its derivatives or analogs and the A and B chain can be modified by amino acid substitution, deletion and/or additions.

In yet another embodiment of the present invention, the linker peptide may be any sequence comprising at least two amino acids with the provision that the first two amino acids represent "RR"

In still another embodiment of the present invention, said protein is defined by SEQ ID: 1 or SEQ ID: 2.

In still another embodiment of the present invention, the host cell of the yeast expression system is selected from *Pichia* sp.

In still another embodiment of the present invention, the host cell of the yeast expression system is selected from the group comprising *Pichia pastoris*, *Pichia methanolica*,

Saccharomyces cerevisiae, *Schisaosaccharomyces pombe*, *Yarrowia lipilitica*, *Hansenula polymorpha*, *Kluyveromyces lactis*.

In still another embodiment of the present invention, the glycosylated form of the protein may be monoglycosylated, triglycosylated or polyglycosylated.

In still another embodiment of the present invention, said purified protein contains less than 1% of glycosylated impurities.

The present invention also relates to a process of obtaining a purified, biologically active heterologous protein recombinantly expressed in a host cell which process comprises steps of:

- a) Culturing host cells transformed by a vector containing a DNA sequence defined by formula I encoding the heterologous protein under conditions suitable for the expression of the protein;
- b) Recovery of the expressed protein comprising separating the said protein from host cells to produce a recovered protein preparation;
- c) Subjecting the recovered protein of step (b) to a step of crystallization;
- d) Performing a step of enzymatic conversion in the presence of trypsin or trypsin like enzyme;
- e) Purifying the said protein containing at least one related impurity comprising contacting the said protein mixture with a chromatographic matrix wherein purification is carried out employing a polar organic buffer solvent, in an aqueous phase containing an organic acid buffer; and
- f) Precipitating the eluted protein.

In yet another embodiment of the present invention, the DNA encoding a heterologous protein is represented by formula I



wherein,

X is a leader peptide sequence comprising at least one amino acid.

B is the amino acid sequence of the B chain of the insulin molecule, its derivatives or analogs

Y is a linker peptide comprising at least two amino acids.

A is the amino acid sequence of the A chain of the insulin molecule, its derivatives or analogs and the A and B chain can be modified by amino acid substitution, deletion and/or addition.

In still another embodiment of the present invention, the linker peptide may be any sequence comprising at least two amino acids with the provision that the first two amino acids represent "RR".

In still another embodiment of the present invention, the gene defined by SEQ ID 1 or 2 is cloned in frame with a signal peptide.

In still another embodiment of the present invention, the gene defined by SEQ ID 1 or 2 is cloned in frame with mat- α -signal peptide.

In still another embodiment of the present invention, the host cell is selected from *Pichia* sp.

In still another embodiment of the present invention, the host cell is selected from the group comprising *Pichia pastoris*, *Pichia methanolica*, *Saccharomyces cerevisiae*, *Schisaosaccharomyces pombe*, *Yarrowia lipilitica*, *Hansenula polymorpha*, *Kluyveromyces lactis*.

In still another embodiment of the present invention, the host cell is *Pichia pastoris*.

In still another embodiment of the present invention, the host strain is GS115.

The present invention also relates to a process of producing a purified, biologically active heterologous protein which comprises:

- a) Inoculating an aqueous fermentation medium with a transformant yeast strain harboring an expression vector which directs the expression of the said protein; and
- b) Growing the transformed strain in the fermentation medium under conditions effective for expression of said protein.

The present invention also relates to a process of producing a purified biologically active heterologous protein wherein the expressed protein is captured from the fermentation broth using ion-exchange chromatography.

In still another embodiment of the present invention, said process comprising a step of crystallizing the protein by addition of zinc chloride and phenol.

The present invention also relates to a process of producing a purified, biologically active heterologous protein, said process comprising a step of enzymatic conversion effected in the presence of trypsin in a water-miscible organic solvent.

In still another embodiment of the present invention, the water-miscible organic solvent is selected from the group comprising DMSO, DMF, ethanol, acetone, acetonitrile, ethyl acetate or mixtures thereof.

In still another embodiment of the present invention, said process comprising a step of RP-HPLC purification of the protein mixture containing at least one related impurity by contacting the said protein mixture with a chromatographic resin matrix wherein purification is carried out employing a polar organic buffer solvent, in an aqueous phase containing an organic acid buffer.

In still another embodiment of the present invention, the polar buffer solvent is acetonitrile.

In still another embodiment of the present invention, the organic acid buffer is selected from the group comprising citric acid, acetic acid, boric acid, formic acid, hydrochloric acid and phosphoric acid.

The present invention also relates to a purified, biologically active heterologous protein product obtained according to any of the preceding claims, with purity of at least 96%.

The present invention also relates to a process a purified, biologically active heterologous protein product with purity in the range of 97 - 100%.

In still another embodiment of the present invention, purified insulin glargine with purity of at least 96%.

In still another embodiment of the present invention, purified insulin glargine containing less than 1% of glycosylated impurities.

In still another embodiment of the present invention, purified insulin glargine devoid of glycosylated impurities.

In still another embodiment of the present invention, purified insulin glargine wherein the glycosylated form of the protein may be monoglycosylated, triglycosylated or polyglycosylated.

Reference will now be made in detail to the presently preferred embodiments of the invention which, together with the following example, serve to explain the principles of the invention.

The Examples which follow are set forth to aid in understanding the invention but are not intended to, and should not be construed to, limit its scope in any way. The Examples do not include detailed descriptions for conventional methods employed in the construction of vectors, the insertion of genes encoding polypeptides into such vectors or the introduction of the resulting plasmids into hosts. The Examples also do not include detailed description for conventional methods employed for assaying the polypeptides produced by such host vector systems. Such methods are well known to those of ordinary skill in the art and are described in numerous publications including by way of examples.

According to one aspect of the invention, a process of obtaining a purified, biologically active heterologous protein recombinantly expressed in a host cell which method comprises steps of :

- a) Culturing host cells transformed by a vector containing a DNA sequence defined by SEQ ID 1 or 2 encoding the heterologous protein under conditions suitable for the expression of the protein.
- b) Recovery of the expressed protein comprising separating the said protein from host cells to produce a recovered protein preparation
- c) Subjecting the recovered protein of step (b) to a step of crystallisation
- d) Performing a step of enzymatic conversion in the presence of trypsin or trypsin like enzyme.
- e) Purifying the said protein containing at least one related impurity comprising contacting the said protein mixture with a chromatographic matrix wherein purification is carried out employing a polar organic buffer solvent, in an aqueous phase containing an organic acid buffer.
- f) Precipitating the eluted protein.


DEFINITIONS:

"Cells" or "cell cultures" or "recombinant host cells" or "host cells" are often used interchangeably as will be clear from the context. These terms include the immediate subject cell which expresses the desired protein of the present invention, and, of course, the progeny thereof. It is understood that not all progeny are exactly identical to the parental cell, due to chance mutations or difference in environment. However, such altered progeny are included in these terms, so long as the progeny retain the characteristics relevant to those conferred on the originally transformed cell.

As used herein, the term "**vector**" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. The term "expression vector" includes plasmids, cosmids or phages capable of synthesizing the subject proteins encoded by their respective recombinant genes carried by the vector. Preferred vectors are those capable of autonomous replication and/expression of nucleic acids to which they are linked. In the present specification, "plasmid" and "vector" are used interchangeably as the plasmid is the most commonly used form of vector. Moreover, the invention is intended to include such other forms of expression vectors which serve equivalent functions and which become known in the art subsequently hereto.

Polypeptides referred to herein as possessing the activity of insulin glargine, e.g. are insulin glargine are understood to have an amino acid sequence with two changes of the human insulin structure: substitution of the amino acid glycine for the native asparagine at position A21 of the A-chain of human insulin and the addition of two arginine molecules to the NH₂-terminal end of the B-chain of human insulin produced by recombinant DNA technology. The primary action of any insulin, including insulin glargine, is regulation of glucose metabolism. Insulin and its analogs lower blood glucose levels by stimulation of peripheral glucose uptake, especially within skeletal muscle and fat, and by inhibition of hepatic glucose production.

The DNA encoding a heterologous protein as represented by formula I



wherein,

X is a leader peptide sequence comprising at least one amino acid.

B is the amino acid sequence of the B chain of the insulin molecule, its derivatives or analogs

Y is a linker peptide comprising at least one amino acid.

A is the amino acid sequence of the A chain of the insulin molecule, its derivatives or analogs and the A and B chain can be modified by amino acid substitution, deletion and/or additions.

The term "C-peptide" or "linker peptide" as used herein includes all forms of insulin C-peptide, including native or synthetic peptides. Such insulin C-peptides may be human peptides, or may be from other animal species and genera, preferably mammals. Thus variants and modifications of native insulin C-peptide are included as long as they retain insulin C-peptide activity. It is known in the art to modify the sequences of proteins or peptides, whilst retaining their useful activity and this may be achieved using techniques which are standard in the art and widely described in the literature e.g. random or site-directed mutagenesis, cleavage and ligation of nucleic acids etc. Thus, functionally equivalent variants or derivatives of native insulin C-peptide sequences may readily be prepared according to techniques well known in the art, and include peptide sequences having a functional, e.g. a biological, activity of a native insulin C-peptide. All such analogues, variants, derivatives or fragments of insulin C-peptide are especially included in the scope of this invention, and are subsumed under the term "an insulin C-peptide".

The linker sequence can be any sequence having at least two amino acids The linker region may comprise from 2 to 25, 2 to 15, 2 to 12 or 2 to 10 amino residues, although the length is not critical and may be selected for convenience or according to choice.

The linker peptide may be any sequence comprising at least two amino acids under the provision that the first two amino acids represent "RR". Moreover, it will be generally

appreciated that, under certain circumstances, it will be advantageous to provide homologs of forms of the subject insulin glargine protein which are either agonists or antagonists of only a subset of that protein's biological activities. Thus, specific biological effects can be elicited by treatment with a homolog of limited function, and with fewer side effects.

In one embodiment, the nucleic acid of the invention encodes a polypeptide which is either an agonist or antagonist the human insulin glargine protein and comprises an amino acid sequence represented by SEQ ID No: 2. Preferred nucleic acids encode a peptide having an insulin glargine protein activity and being at least 90% homologous, more preferably 95% homologous and most preferably 97% homologous with an amino acid sequence shown in SEQ ID No: 2. Nucleic acids which encode agonist or antagonist forms of insulin glargine protein and having at least about 98-99% homology with a sequence shown in SEQ ID No: 2 are also within the scope of the invention. Preferably, the nucleic acid is a cDNA molecule comprising at least a portion of the nucleotide sequence encoding an insulin glargine protein shown in SEQ ID No. 1.

The "**operational elements**," as discussed herein, include at least one promoter, at least one operator, at least one leader sequence, at least one Shine-Dalgarno sequence, at least one terminator codon, and any other DNA sequences necessary or preferred for appropriate transcription and subsequent translation of the vector DNA. In particular, it is contemplated that such vectors will contain at least one origin of replication recognized by the host microorganism along with at least one selectable marker and at least one promoter sequence capable of initiating transcription of the synthetic DNA sequence. It is additionally preferred that the vector, in one embodiment, contains certain DNA sequences capable of functioning as regulators, and other DNA sequences capable of coding for regulator protein. These regulators, in one embodiment, serve to prevent expression of the DNA sequence in the presence of certain environmental conditions and, in the presence of other environmental conditions, allow transcription and subsequent expression of the protein coded for by the DNA sequence.

Additionally, it is preferred that an appropriate secretory leader sequence be present, either in the vector or at the 5' end of the DNA sequence. The leader sequence is in a position which allows the leader sequence to be immediately adjacent to the initial portion of the nucleotide sequence capable of directing expression of the desired protein inhibitor without any intervening translation termination signals. The presence of the leader sequence is desired in part for one or more of the following reasons:

- 1) the presence of the leader sequence may facilitate host processing of the initial product to the mature recombinant protein product ;
- 2) the presence of the leader sequence may facilitate purification of the recombinant protein product, through directing the protease inhibitor out of the cell cytoplasm;
- 3) the presence of the leader sequence may affect the ability of the recombinant protein product to fold to its active structure through directing the protein out of the cell cytoplasm.

"Transcriptional regulatory sequence" is a generic term used throughout the specification to refer to DNA sequences, such as initiation signals, enhancers, and promoters which induce or control transcription of protein coding sequences with which they are operably linked. In preferred embodiments, transcription of a recombinant insulin glargine gene is under the control of a promoter sequence (or other transcriptional regulatory sequence) which controls the expression of the recombinant gene in a cell-type in which expression is intended.

"Control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences which are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, a ribosome binding site, and possibly, other as yet poorly understood sequences. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

"Transformation" means introducing DNA into an organism so that the DNA is replicable, either as an extra chromosomal element or by chromosomal integration.

Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. General aspects of mammalian cell host system transformations have been described by Axel in U.S. Pat. No. 4,399,216 issued Aug. 16, 1983. Transformations into yeast are typically carried out according to the method of Van Solingen, P., et al., J. Bact., 130: 946 (1977) and Hsiao, C. L., et al., Proc. Natl. Acad. Sci. (USA) 76: 3829 (1979).

A recombinant expression system is selected from prokaryotic and eukaryotic hosts. Eucaryotic hosts include yeast cells (e.g., *Saccharomyces cerevisiae* or *Pichia pastoris*), mammalian cells or plant cells. Bacterial and eukaryotic cells are available from a number of different sources including commercial sources to those skilled in the art, e.g., the American Type Culture Collection (ATCC; Rockville, Md.). Commercial sources of cells used for recombinant protein expression also provide instructions for usage of the cells. The choice of the expression system depends on the features desired for the expressed polypeptide.

Consequently, the subject of the present invention is a expression system or cassette which is functional in a cell derived from a yeast selected from the group consisting of strain *Pichia* especially selected from the group consisting of *Pichia pastoris*, *Pichia methanolica* and *Schizosaccharomyces pombe* and allowing the expression of the desired polypeptide thereof encoding the protein fragments thereof, placed under the control of the elements necessary for its expression.

The term "**recombinant**", as used herein to describe a protein or polypeptide means a polypeptide produced by expression of a recombinant polynucleotide. The term "recombinant", as used herein in reference to cells, means cells that can be or have been used as recipients for recombinant vectors or other transfer DNA, and include progeny of the original cell which has been transfected. It shall be understood that progeny of a single parental cell may not be completely identical in morphology or in genomic or total DNA complement to the original parent, due to accidental or deliberate mutation. Progeny of a parental cell which are sufficiently similar to the parent to be characterized by a relevant

property, such as the presence of a nucleotide sequence encoding a desired polypeptide, are also considered progeny.

A "**gene of interest**" (**GOI**) is any nucleic acid sequence for which increased transcriptional expression is desired. The GOI may encode a functional nucleic acid molecule (e.g., an RNA, such as an antisense RNA molecule) or, more typically, encodes a peptide, polypeptide or protein for which increased production is desired. The vectors of the invention can be used to express a "heterologous" protein. As used herein, the term "heterologous" means a nucleic acid sequence or polypeptide that originates from a foreign species, or that is substantially modified from its original form if from the same species. Furthermore, a modified or unmodified nucleic acid sequence or polypeptide that is not normally expressed in a cell is considered heterologous. Vectors of the invention can have one or more GOIs, inserted at the same or different insertion site, where each GOI is operably linked to a regulatory nucleic acid sequence which allows expression of the GOI.

The term "**purifying**" a peptide from a composition comprising the peptide and one or more contaminants thereby increasing the degree of purity of the peptide in the composition by reducing the contents of at least one contaminant from the peptide composition. More specifically the purification process of the present invention results in a biologically active heterologous protein recombinantly expressed in a yeast expression system, characterized in that said purified protein is devoid of or contains substantially minimal amounts of glycosylated byproducts.

Apart from the glycosylated form presence of nonpolar impurity was confirmed by MALDI (m/Z: 6089). The presence of this nonpolar impurity varies from 0-0.5% in the final product which is generated during the processing of Glargine. The difference in mass between glargine (m/z: 6063) and non-polar impurities (m/z: 6089) is 26 units. Chemical identity based on molecular mass indicates the probability of water loss in one of the amino acids and acetylation

The term "**chromatography**" refers to the process by which a solute of interest in a mixture is separated from other solutes in a mixture as a result of differences in rates at

which the individual solutes of the mixture migrate through a stationary medium under the influence of a moving phase, or in bind and elute processes.

The term "**High Performance liquid chromatography**", as used herein, refers to that chromatographic procedure in which the particles (stationary phase) used in the column packing are small (between 3 and 50 microns) and regular with little variation from the selected size. Such chromatography typically employs relatively high (around 500-3500 psi) inlet pressures.

After a host organism has been chosen, the vector is transferred into the host organism using methods generally known by those of ordinary skill in the art. Examples of such methods may be found in *Advanced Bacterial Genetics* by R. W. Davis et. al., Cold Spring Harbor Press, Cold Spring Harbor, N.Y., (1980), which is specifically incorporated herein by reference. It is preferred, in one embodiment, that the transformation occur at low temperatures, temperature regulation is contemplated as a means of regulating gene expression through the use of operational elements as set forth.

The host microorganisms are cultured under conditions appropriate for the expression of the insulin glargine precursor. These conditions are generally specific for the host organism, and are readily determined by one of ordinary skill in the art, in light of the published literature regarding the growth conditions for such organisms, for example *Bergey's Manual of Determinative Bacteriology*, 8th Ed., Williams & Wilkins Company, Baltimore, Md., which is specifically incorporated herein by reference.

Thus, a coding sequence "**operably linked**" to control sequences refers to a configuration wherein the coding sequence can be expressed under the control of these sequences and wherein the DNA sequences being linked are contiguous and, in the case of a secretory leader, contiguous and in reading phase.

According to one aspect of the invention, Insulin glargine precursor of formula X - Glargine B chain [B1-B30] -Y- Glargine A chain [A1-A21], wherein X is the leader sequence, B chain is the Glargine B chain sequence of B1-B30, Y is a linker peptide sequence between B chain and A chain, A chain is the A chain of Glargine. The GIP-A

precursor may be produced by expression in a host expression system not including bacteria other than *E. coli*, yeast such as *Saccharomyces* and *Kluyveromyces* for which methods are known to those skilled in the art (Methods in Enzymology vol. 194); and fungi such as *Aspergillus*, *Neurospora* and *Trichoderma* for which methods are known to those skilled in the art, as described in Applied Molecular Genetics of Fungi, Peberdy, Caten, Ogden, and Bennett, eds. (Cambridge University Press, N.Y. 1991), many of which can be obtained from the American Type Culture Collection, Rockville, Md., which are routinely used by those skilled in the art. Preferably, the host strain is selected from the group comprising yeast such as *Saccharomyces cerevisiae*, *Schisaosaccharomyces pombe*, *Yarrowia lipilitica*, *Hansenula polymorpha*, *Kluyveromyces lactis*, *Pichia methanolica* and *Pichia pastoris*, which are capable of expressing high levels of a recombinant protein. Even more preferably, the host strain is *Pichia pastoris*.

In particular, the present invention provides a process for manufacture of insulin glargine protein including a fermentation process, a recovery process, and a purification process. The process provided herein includes a fermentation process wherein the aforesaid protein is produced in *Pichia pastoris* having at least one sequence encoding the said protein is integrated into the genome, wherein the fermentation process includes a seed fermentation to grow host cells to a desired cell density and a production fermentation process comprising glycerol batch fermentation, methanol induction fermentation and finally production.

The Glargine precursor is then captured from the fermentation broth using cation-exchange chromatography. The recovery process provided herein permits capture of the glargine precursor and removal of cells and cellular debris, where the recovery process provided herein provides for a yield of 95%. Alternately, one skilled in art can test and evaluate alternate methods for capturing the desired protein product and removing cells and cellular debris including but not limited to various other chromatographic methods, centrifugation, filtration, differential precipitation, and other methods to be determined.

The glargine precursor captured from the fermentation broth is further subjected to crystallization for color removal and storage. The precipitation of the precursor in a

crystallized form is suitably performed in an aqueous vehicle at a pH comprised between 3.0 and 8.0, preferably 4.0 and 6.0, most preferably at a pH of 4.0 to 5.0, the concentration of the protein in the aqueous vehicle being comprised between 1 and 80 g/L, preferably between 2 and 50 g/L, most preferably 8 to 14g/L. The crystallisation process can be done at a temperature between 2 and 30°C. The precursor crystals can be separated from the supernatant either by centrifugation, decantation or filtration.

Crystallization of the Glargine precursor GIP-A removes the impurities carried through from the fermentation broth into the cation-exchange elution pool as well as ammonium acetate salt which is used to elute the product from the chromatography. A pure crystalline precursor also helps in reducing cost in the subsequent steps and increase the efficiency of the reaction. The crystalline form can be frozen and stored which would be subsequently stable for the multiple days when stored at -20°C.

Glargine can be prepared from the Glargine precursor through an enzymatic conversion. The conversion is effected by the presence of trypsin or trypsin like enzymes of plant, animal or microbial origin. The reaction is carried out preferably in the presence of water-miscible organic solvents like DMSO (dimethyl sulfoxide) DMF (dimethyl formamide), ethanol, acetone, acetonitrile or ethyl acetate especially DMF and DMSO. The preferable ratio of the organic solvent is about 0-65%, especially about 40-60% of the reaction mixture.

Glargine is produced by treating the precursor GIP-A with Trypsin. The leader (if present) will be cleaved from the precursor by Trypsin. The Trypsin reaction is controlled in such a way that cleavage at the C-terminal of "RR" C-peptide is maximized. But the reaction could not eliminate the unwanted cleavage by trypsin in other trypsin cleavage sites. The probable other trypsin cleavage sites are between "RR" and C-terminus of B-22 (Arg).

The rationale for the use of organic solvents should be determined by the solubility of the starting materials, tendency of enzymes to denature and its hydrolysing activity. As may be appreciated by a skilled artisan, mixtures of organic solvents may also be used. The addition of organic solvents lowers the aqueous concentration of the reaction mixture,

resulting in prevention of hydrolysis of product, and also remarkably increases the solubility of the products.

The concentration of the precursor is generally about 5-50g/L. The reaction is carried out at about pH 5 to 12, preferably about pH 8 to 10. Reaction temperature is about 0-40 °C preferably about 2-25 °C. Trishydroxymethylaminomethane (TRIS) or other buffer systems can be used at different ionic strengths to maintain the required pH. Reaction time is variable and is affected by other reaction conditions. Reaction may be continued until the purity of the product starts decreasing due to hydrolysis of the product, It takes generally about 30 minutes to 24 hours and in most cases about 4-10 hours.

The concentration of enzyme is determined depending on concentration of substrates and enzyme activity. For example, the crystalline trypsin available in the market is used preferably in a concentration about 10-100mg/L of the reaction mixture.

The purification method comprises subjecting the insulin glargine sample to a step of purification through reverse-phase chromatography comprising a polymer based resin matrix under chromatographic conditions sufficient for obtaining said peptide of nearly 98-99% purity, preferably 100% purity.

Thus, unwanted product related impurities formed which is purified in the successive reverse phase chromatography to generate pure Glargine. Glycosylated (mono and Tri) impurities can only be detected after the first purification steps. So, special cares were taken to purify the Glycosylated impurities in the 2nd RP-HPLC purification step. The Glycosylated impurities generated during the fermentation is fully characterised and finally purified from the final product to the maximum extent.

The invention features, in a significant aspect, a method of purifying the protein, the method including the steps of subjecting the instant protein to a reverse phase HPLC column and eluting a sample that includes the peptide with an organic solvent, under conditions that allow the compound to be bound to the resin, and washing the organic solvent from the resin with an aqueous buffer solution. The effective performance of the present invention requires the individuation of right combination of the chromatographic

matrix to be used, the pH value and the ionic strength of the buffer for efficient purifications

Every attribute of the chromatographic procedure has a significant role to play in obtaining the desired protein product. A suitable matrix used in the chromatographic column is C8 Daisogel.

According to one aspect of the invention the yield of the purified insulin glargine product is 75% - 80%, according to another aspect of the invention the yield of the purified insulin glargine product is 80% - 85%, according to yet another aspect of the invention the yield of the purified insulin glargine product is 85% - 90%, according to yet another aspect of the invention the yield of the purified insulin glargine product is 90% - 95%, according to still another aspect of the invention the yield of the purified insulin glargine product is 95% - 100%.

According to one aspect of the invention the purified insulin glargine product is devoid of or contains substantially minimal amounts of glycosylated byproducts. According to one aspect of the invention the purity of the purified insulin glargine product is at least 96%, according to another aspect of the invention the purity of the purified insulin glargine product is at least 97%, according to yet another aspect of the invention the purity of the purified insulin glargine product is at least 98%, according to yet another aspect of the invention the purity of the purified insulin glargine product is at least 99%, according to still another aspect of the invention the yield of the purified insulin glargine product is 100%.

The invention is further elaborated with the help of following examples. However, these examples should not be construed to limit the scope of the invention.

EXAMPLES**EXAMPLE 1:**

GIP-A represent the Glargine precursor of Formula X-[Glargine B chain (B1-B30)]- Y- [Glargine A chain (A1-A21)], where X is a leader peptide sequence, B chain is Glargine B chain sequence of B1-B30, Y is a linker peptide sequence between B chain and A chain, A chain is the A-chain of Glargine. The sequence may be devoid of leader or other leader peptide. The linker peptide Y can be anyone from the example RR, RRDADDR. The GIP-A precursor may be produced by any suitable expression system such as Escherichia coli, Pichia pastoris, Saccharomyces cerevisiae, CHO cells etc.

The GIP-A precursor was cloned in frame with the Mat-alfa signal peptide in Pichia expression vector, pPIC9K. Pichia pastoris host strain GS115 was transformed with the recombinant plasmid to obtain clone expressing Glargine precursor. The secreted precursor was treated with trypsin to prepare Glargine and other product related impurities. Glargine was purified by using reverse phase chromatography.

SEQ ID 1 represents the sequence for the glargine precursor, having a predicted molecular weight of 6045 Da and estimated pI value of 6.88. The sequence has 159 nucleotides and 53 amino acids.

SEQ ID 2 represents the sequence for expression in Pichia pastoris with a predicted molecular weight of 6428.4 da, and estimated pI value of 7.78. The sequence has 171 nucleotides and 57 amino acids.

Glargine precursor GIP-A is secreted by Pichia pastoris into the culture medium. The broth is centrifuged and cells are separated from the supernatant. There are multiple options available for the capture of the precursor GIP-A including Ion-Exchange chromatography and Hydrophobic chromatography. For this invention cation-exchange chromatography and HIC were used to capture the GIP-A.

EXAMPLE 2:**CONSTRUCTION OF THE RECOMBINANT VECTOR CARRYING THE INSULIN GLARGINE PRECURSOR GENE:**

Insulin Glargine Precursor (GIP-A) was cloned in frame with the mat α -signal peptide in the *Pichia pastoris* expression vector, pPIC9K. The recombinant plasmid was used to transform *Pichia pastoris* host strain, GS115, and the secreted GIP-A was Insulin-Glargine Precursor which will be taken to downstream purification process to make the final product.

TRANSFORMATION OF A PICHIA HOST WITH THE RECOMBINANT VECTORS CARRYING THE INSULIN GLARGINE PRECURSOR GENE:

pPIC9K/IGP-A clone harboring recombinant plasmid DNA was digested with Bgl II and used to transform electrocompetent cells of *P.pastoris* GS115 (his4) host. The invitrogen manual provides a detailed protocol of the transformation procedure.

SCREENING FOR MULTI-COPY INTEGRANTS:

Approximately 2000 transformants were inoculated in YPD broth in 384 well micro titer plates along with appropriate controls. The plates were incubated at 30°C for 24 hours and then stamped onto YPD agar plates containing 0.5 mg/mL Geneticin (G418). 49 clones were selected which were stamped further on plates 1, 2 and 3 mg/mL of G418 for second round of screening. Finally, seven clones resistant to 1- 3 mg/ml G418 were selected and used in expression studies.

CONFIRMATION OF GENE INTEGRATION IN THE GENOME BY PCR:

Genomic DNA from selected recombinant *Pichia* clones was subjected to PCR using gene specific primers in order to confirm the integration of GIP-A in the genome.

SMALL SCALE EXPRESSION STUDIES IN P. PASTORIS

Small scale expression studies in *P.pastoris* were carried out as per the protocol given in Invitrogen manual. Briefly, the clones were grown at 30°C in BMGY followed by induction with methanol in BMMY at 24°C. Induction with methanol was carried out for a total of 3 days.

EXAMPLE 3:

FERMENTATION PROCEDURE

Recombinant Insulin-Glargine fermentation process has been optimized at laboratory scale. Presented below is a brief description of the process.

SEED PREPARATION

The seed medium contains following ingredients: Yeast nitrogen base, Ammonium sulfate, Glycerol, potassium dihydrogen phosphate and dipotassium hydrogen phosphate and D-Biotin.

A single vial from freezer was used to inoculate the seed flask. The vials were thawed to ambient temperature before inoculation under sterile conditions. The inoculum was dispensed in the Minimal Glycerol (MGY) medium in an aseptic manner.

The quantity to be added to each flask was ascertained, such that the initial OD600 in the pre-seed flask was between 0.1 - 0.2. The flasks were then incubated for 20-24 h on shaker at 30°C at a speed of 230 RPM.

FERMENTER BATCHING CONDITIONS

Fermentation process has two phases, Growth phase (to build up biomass) followed by induction phase. Product was formed and secreted out to the broth during methanol fed batch phase. Fermentation was performed in an in situ sterilisable fermenter. The fermentation medium comprises of, orthophosphoric acid, potassium sulphate, potassium

hydroxide, magnesium sulphate calcium sulphate and glycerol. These chemicals were dissolved in potable water in the fermenter vessel and sterilized at a set point of 121°C for a specified period. Stock solution of trace salt and biotin were prepared separately and filter sterilized and added aseptically to the fermenter after medium sterilization and pH adjustment to 5.0 with ammonium hydroxide.

Production / Fermentation procedure were adopted. The fermenter was inoculated with 5% seed inoculum from a seed shake flask. Following parameters set at the beginning of fermentation: pH: 5.0, Temperature: 30°C, DO₂: 30%.

EXAMPLE 4:

RELATED CONVERSIONS AND PURIFICATION METHOD:

Insulin glargine precursor that was produced in fermentation of *Pichia pastoris* was subjected to purification the steps which involved in the purification process are as follows.

The column packed with SP Sepharose fast flow matrix (GE Biosciences) was equilibrated with 50mM acetic acid. The cell free supernatant, adjusted to pH 3.8 using ortho phosphoric acid, was loaded on cation exchange column. The loading on the column was < 50g/L. The elution was performed using ammonium acetate. The recovery of the step was 95% at <50 g/L loading.

PRECURSOR CRYSTALLISATION

The Glargine precursor captured from the fermentation broth by using a cation-exchange chromatography step was crystallised for the purpose of colour removal and storage. The crystallisation was done such that the precursor concentration at the start of crystallisation was around 2 to 20 g/L, preferably 8 to 14 g/L. The crystallisation was done by adding ZnCl₂ & phenol and then adjusting the pH to between 3.0 and 8.0, preferably between 3.5 and 5.5. Phenol can be added at 0.1 to 0.5% of the CIEX elution pool volume. A 4% ZnCl₂ solution can be added at 3 to 15% of the CIEX elution pool volume. The pH can be

adjusted by using any alkali, preferably NaOH or TRIS. The crystallisation process can be done at a temperature between 2 and 30 °C and the slurry is stored for some time so that the crystals will be formed completely. The precursor crystals can be separated from the supernatant by either centrifugation or decantation.

Example 4a: 463 ml of elution pool (precursor concentration 13.8 g/L) was taken and 2.315 ml of phenol (0.5% of EP volume) was added after proper thawing. This was followed by addition of 57.875 ml of 4% ZnCl₂ solution (12.5% of EP volume). The pH was 4.08 at this stage and it was adjusted to 4.8 by adding 420 ml of 2.5N NaOH. The mother liquor was kept under slow stirring conditions for 15 minutes and then transferred to cold room (2-8°C), where it was kept overnight. Then the whole mixture was centrifuged at 5000 rpm for 20 minutes in a Beckman Coulter Avanti J-26 XP centrifuge. The loss in supernatant was 1.55%.

Example 4b: 500 ml of elution pool (precursor concentration 2.9 g/L) was taken and 0.625 ml of phenol (0.125% of EP volume) was added after proper thawing. This was followed by addition of 15.625 ml of 4% ZnCl₂ solution (3.125% of EP volume). The pH was 4.08 at this stage and it was adjusted to 4.8 by adding 315 ml of 2.5N NaOH. The mother liquor was kept under slow stirring conditions for 15 minutes and then transferred to cold room (2-8°C), where it was kept for five hours. Then the supernatant was separated by centrifugation. The loss in supernatant was 4%.

OPTIMIZED PRECURSOR CRYSTALLISATION CONDITION WITH YIELD:

The SP-Sepharose elution pool was chilled to 20° + 5 ° C, 5 mL (0.5% V/V) of phenol was added for each litre of elution pool. Required amount of Phenol was added in an aliquot of elution pool and finally added to the main tank for proper redissolution. After addition of phenol the stirring was continued for some time before addition of next reagent for proper redissolution. Now, 12.5 % by volume of 40 g/L Zinc chloride (4%w/v) was added to induce crystallisation. The pH was then adjusted to 4.8 + 0.1 using 2.5N NaOH (approx. 85-90% of SP-EP volume). This entire mixture was stirred for 30 minutes to ensure homogeneity and then held at 2-8 °C for at least 8 hours before centrifugation.

Based on the loss in the supernatant, the yield was approximately 90% at this step.

EXAMPLE 5:

TRYPSINISATION

Glargine was prepared from Glargine precursor through an enzymatic conversion. The conversion was effected by the presence of trypsin or trypsin-like enzymes of plant, animal or microbial origin. The reaction was carried out preferably in presence of water-miscible organic solvents like DMSO, DMF, ethanol, acetone, acetonitrile, ethyl acetate, etc, especially DMF and DMSO. The preferable ratio of organic solvent was about 0 to 65%, especially about 40 to 60 % of the reaction mixture.

The ratio of organic solvent was determined by the solubility of the starting materials, tendency of enzymes to denature and its hydrolyzing activity. Mixtures of organic solvents also can be used. The addition of organic solvents lowers the aqueous concentration of the reaction mixture, resulting in prevention of hydrolysis of product, and also remarkably increases the solubility of the products.

The concentration of the precursor was generally about 5 to 50 g/L. The reaction was carried out at about pH 5 to 12, preferably about pH 8 to 10. Reaction temperature was about 0 – 40 °C, preferably about 2 – 25 °C. Trishydroxymethylaminomethane (TRIS) or other buffer systems was used at different ionic strengths to maintain the required pH. Reaction time was variable and was affected by other reaction conditions. Reaction was continued until the purity of the product starts decreasing due to the hydrolysis of the product. It takes generally about 30 minutes to 24 hours and in most cases about 4 to 10 hours.

The concentration of enzyme was determined depending on concentration of substrates and enzyme activity. For example, the crystalline trypsin available in the market was used preferably in a concentration about 10 to 100 mg/L.

Example 5a: 1g of glargine precursor was dissolved in 1.5ml of 1M TRIS solution. 1 ml each of this solution was taken into two tubes labeled as sample A and sample B. To sample A, 3 ml of water and 166.6mg of TRIS were added. To sample B, 2ml of DMF, 1ml of water and 166.6mg of TRIS were added. Each sample was split into four by adjusting the pH to four different pH. The reaction was carried out in all the samples by adding bovine trypsin at a concentration of 50 μ g/ml. The conditions tried are tabulated as follows.

TABLE 1:

SAMPLE	SOLVENT %	PH	TRIS CONCENTRATION (M)	TRYPSIN CONCENTRATION (μ g/mL)	PRECURSOR CONCENTRATION (mg/mL)
A1	0%	9.2	0.5	50	15.5
A2		8.2			
A3		7.0			
A4		6.2			
B1	50%	8.9			
B2		8.0			
B3		7.0			
B4		6.3			

All the conditions gave product at different proportions. The sample B1 gave best conversion at around 40% whereas conditions A1, A2, A3 and A4 gave less than 15% conversion. Conversions to Glargine in other samples were in the range of 15% and 40%.

Example 5b: 1.6g of Glargine precursor was dissolved in 2ml of 1M TRIS solution. 1ml each of this solution was taken into three tubes labeled as sample A, sample B and sample C. To sample A, 1.8ml of water, 1.2 ml of DMF and 161mg of TRIS were added. To sample B, 0.6ml of water, 2.4ml of DMF and 161mg of TRIS were added. To sample C, 1 ml of water, 2.0ml of DMF and 161mg of TRIS were added. Both sample A and B were split into four and sample C into two by adjusting to different pH. The reaction was carried out in all the samples by adding bovine trypsin at a concentration of 50 μ g/ml. The conditions tried were tabulated as follows.

TABLE 2:

SAMPLE	SOLVENT (DMF)%	PH	TRIS CONCENTRATION (M)	TRYPSIN CONCENTRATION (μ g/mL)	PRECURSOR CONCENTRATION (mg/mL)
A1	30%	10.1	0.5	50	10.5
A2		9.04			
A3		8.05			
A4		6.77			
B1	60%	9.95			
B2		9.13			
B3		8.01			
B4		7.0			
C1	50%	9.0			
C2		8.0			

All the conditions gave product at different proportions. The sample C1 and C2 gave best conversions at around 40% whereas other conditions gave less than 15% conversion.

Example 5c: 1.5g of Glargine precursor was dissolved in 2ml of 1M TRIS solution. 0.75ml each of this solution was taken into three tubes labeled as sample A, sample B and sample C. To sample A, 0.75 ml of water, 1.5ml of ethanol and 106 mg of TRIS were added. To sample B, 0.75ml of water, 1.5 ml of DMF and 106 mg of TRIS were added. To sample C, 0.75ml of water, 1.5ml of DMF and 106 mg of TRIS were added. Both samples A and B were split into two by adjusting to two different pH. The sample C was adjusted to pH 9.0. The reaction was carried out in all the samples by adding bovine trypsin at a concentration of 50 μ g/ml. The conditions tried are tabulated as follows.

TABLE 3:

SAMPLE	SOLVENT %	PH	TRIS CONCENTRATION (M)	TRYPSIN CONCENTRATION (μ G/mL)	PRECURSOR CONCENTRATION (mg/mL)
A1	50%	9.0	0.5	50	17
A2	ETHANOL	7.3			
B1	50%	8.9			
B2	DMSO	6.9			
C1	50% DMF	9.0			

All the conditions gave product at different proportions. The sample C1 gave best conversions at around 40%, followed by B1 (~30%), whereas other conditions gave less than 15% conversion.

Example 5d: 1.5g of Glargine precursor was dissolved in 2ml of 1M TRIS solution. 0.75 ml of this solution was taken and to this 0.75 ml of water, 1.5 ml of ethanol and 106 mg of TRIS were added. The pH was then adjusted to 9.0. Then the solution was split into two samples - A and B. For sample A, the reaction was carried out at room temperature (20 to 25°C) whereas for sample B, the reaction was done at 2 to 8°C. Reaction was started by adding bovine trypsin at a concentration of 50µg/ml.

Both the samples gave conversion of around 40%. The reaction at lower temperature (sample B) took longer duration but showed slightly better yield.

Example 5e: 1g of Glargine precursor was dissolved in 1ml of 1M TRIS solution. 0.75 ml each of this solution was taken into two tubes labeled as sample A, and sample B. To sample A, 0.75 ml of water, 1.5 ml of DMF and 130 mg of TRIS were added. To sample B, 0.75 ml of water, 1.5 ml of DMSO and 130 mg of TRIS were added. Both samples A and B were split into two after adjusting to pH 8.7. The reaction was carried out in all the four samples by adding bovine trypsin at a concentration of 50µg/ml. Reactions in one aliquot each from sample A and B were done at 2-8°C. The conditions tried are tabulated as follows.

TABLE 4:

SAMPLE	SOLVE NT %	PH	TRIS CONCEN TRATIO N (M)	TRYPSIN CONCENTRA TION (µg/mL)	PRECURSOR CONCENTRA TION (mg/mL)	TEMPERAT URE (°C)
A1 (DMF)	50%	8.7	0.5	50	13.5	20 – 25
A2 (DMF)						2- 8
B1 (DMSO)						20 – 25
B2 (DMSO)						2- 8

All the samples gave similar conversion (~40%). The reaction in samples at room temperature was over within one hour whereas those at cold room were over by 5 – 6 hours.

Example 5f: 2g of Glargine precursor was dissolved in a solution containing 0.5g TRIS, 2ml water and 2 ml DMF. From this solution, 6 samples of 3.08 ml were made such that each contained 50% DMF& 0.5M TRIS, at pH 8.7. The precursor concentration was varied among these samples from 10 g/L to 60 g/L. Reaction was started in each sample by adding 5 mg trypsin per gram of precursor. All the samples were at 2-8°C during the reaction. The conditions tried are tabulated as follows.

TABLE 5:

SAMPLE	SOLVENT %	PH	TRIS CONCENTRATION (M)	TRYPSIN CONCENTRATION ($\mu\text{g}/\text{mL}$)	PRECURSOR OR CONCENTRATION (mg/mL)	TEMPERATURE ($^{\circ}\text{C}$)
A	50%	8.7	0.5	50	10	2-8
B				100	20	
C				150	30	
D				200	40	
E				250	50	
F				300	60	

All the samples were showing conversion, but samples with lower concentration gave faster reaction. condition B was considered the best. The maximum yield obtained in each case was around 40 - 45%. The results have been represented in **FIG. 1**

OPTIMIZED PROCESS CONDITION WITH YIELD AND PURITY:**EXAMPLE 5g:**

The wet precursor crystals were stored in -20°C deep freezer which was taken out well before dissolution. These crystals were dissolved by adding that into a TRIS solution (TRIS – 50% w/w of precursor crystals and water – 2L per kg of precursor crystals) under stirring conditions. For example, 1 kg of precursor crystal is added into 2L TRIS solution containing 500 grams of TRIS. This was followed by addition of DMF (2L per kg of precursor crystals). The whole mixture is completely solubilised and this solution was treated as Test solution. The resultant test solution was analyzed for product content using C18 symmetry analytical RP-HPLC method.

According to the product content in the test solution, the control solution for trypsinisation reaction was made, such that the recombinant insulin Glargine precursor concentration was 10 g/L of the reaction mixture. To prepare the control solution, the required amount of TRIS in water was added to keep the final TRIS concentration in the reaction mixture as 500mM. This was followed by the addition of DMF such that the final DMF concentration was 50%. The temperature of this solution was then brought to $6^{\circ}\text{C} \pm 2^{\circ}\text{C}$, and then the pH was adjusted to 8.7 using glacial acetic acid (around 2% of total volume) under mild stirring conditions. This solution was treated as control solution.

The reaction mixture was maintained at $6^{\circ}\text{C} \pm 2^{\circ}\text{C}$ at very slow impeller speed. After complete mixing, 50 mg/L of trypsin was added to initiate the reaction. The enzyme reaction starts once the trypsin is well dissolved by mixing.

The enzyme reaction was carried out at $6^{\circ}\text{C} \pm 2^{\circ}\text{C}$ at very slow impeller speed. The progress of the reaction was monitored by checking every hour, the percentage of recombinant insulin Glargine and the unreacted precursor, using the C18 Symmetry analytical RP-HPLC method. The enzyme reaction was typically completed in less than 10 hours. The reaction was stopped by adjusting the pH of the reaction mixture to 5.0 using glacial acetic acid, when the percentage of unreacted precursor was less than 5%. The

reaction is expected to yield around 40% of Recombinant Insulin Glargine and purity obtained was about 35 - 50 %.

Analytical chromatogram of the reaction at different time points is represented in **FIG 2 & 3**.

EXAMPLE 6: RP-HPLC PURIFICATION:

EXAMPLE 6a:

Insulin glargine precursor produced by fermentation of *Pichia pastoris* was subjected to purification, the steps involved in the process were as follows:

The column packed with SP sepharose fast flow matrix (GE Biosciences) was equilibrated with 50mM acetic acid. The cell free supernatant adjusted to pH 3.8 using orthophosphoric acid was loaded on a cation -exchange column. The loading on the column was < 50g/L. The elution was performed using ammonium acetate. The recovery of the step was 95% at <50 g/L loading. The elution pool was crystallised. The crystal was used for trypsinisation. The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 10% of acetonitrile (Buffer B) in 250mM acetic acid (Buffer A). The loading of trypsinisation end diluted 1:5 using water was performed. A linear gradient of buffer B was used to elute the purified glargine at yield of 75% with purity 94.5% at a loading of <10 g/L. The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 10% of acetonitrile (Buffer B) in 200mM sodium acetate pH 5:0 (Buffer A). The loading of RP eluate from previous step, diluted to contain 10% acetonitrile was performed. A linear gradient of buffer B was used to elute the purified glargine at yield of 80% with purity 98.5% and glycosylated impurity of 0.29%. The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 6% of ethanol (Buffer B) in 50mM acetic acid (Buffer A). The loading of RP eluate from previous step, diluted to 1:3 with water, was performed. A linear gradient of buffer B was used to elute the purified glargine at yield of 85% with purity 99.4% at a loading of 35g/L

EXAMPLE 6b:

The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 10% of acetonitrile (Buffer B) in 250mM acetic acid (Buffer A). The loading of, trypsinisation end diluted 1:5 using water, was performed. A linear gradient of buffer B was used to elute the purified glargine at yield of 75% with purity 94.5% at a loading of <10 g/L. The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 10% of acetonitrile (Buffer B) in 200mM sodium acetate pH 5.0 (Buffer A). The loading of RP eluate from previous step, diluted to contain 10% acetonitrile, was performed. A linear gradient of buffer B was used to elute the purified glargine at yield of 80% with purity 98.5% and glycosylated impurity of 0.29%. The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 6% of ethanol (Buffer B) in 10mM citric acid (Buffer A). The loading of RP eluate from previous step, diluted to 1:3 with water, was performed. A linear gradient of buffer B was used to elute the purified glargine at yield of 79.2% with purity 98.5% at a loading of 35g/L.

EXAMPLE 6c:

The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 10% of acetonitrile (Buffer B) in 250mM acetic acid (Buffer A). The loading of, trypsinisation end diluted 1:5 using water was performed. A linear gradient of buffer B was used to elute the purified glargine at yield of 75% with purity 94.5% at a loading of <10 g/L. The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 10% of acetonitrile (Buffer B) in 100mM Tris+20mM CaCl₂ pH 8.5 (Buffer A). The loading of RP eluate from previous step, diluted to contain 10% acetonitrile, was performed. A linear gradient of buffer B was used to elute the purified glargine at yield of 71% with purity 97.4% and glycosylated impurity of 0.22%. The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 6% of ethanol (Buffer B) in 10mM citric acid (Buffer A). The loading of RP eluate from previous step, diluted to 1:3 with water, was performed.

A linear gradient of buffer B was used to elute the purified glargine at yield of 75.0% with purity 98.0% at a loading of 35g/L.

EXAMPLE 6d:

The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) is equilibrated with 10% of acetonitrile (Buffer B) in 250mM acetic acid (Buffer A). The loading of, trypsinisation end diluted 1:5 using water, is performed. A linear gradient of buffer B is used to elute the purified glargine at yield of 75% with purity 94.5% at a loading of <10 g/L. The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) is equilibrated with 10% of acetonitrile (Buffer B) in 100mM ammonium acetate pH 5.0 (Buffer A). The loading of RP eluate from previous step, diluted to contain 10% acetonitrile, is performed. A linear gradient of buffer B is used to elute the purified glargine at yield of 60.7% with purity 98.0% and glycosylated impurity of 0.91%. The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) is equilibrated with 6% of ethanol (Buffer B) in 10mM citric acid (Buffer A). The loading of RP eluate from previous step, diluted to 1:3 with water, is performed. A linear gradient of buffer B is used to elute the purified glargine at yield of 75.0% with purity 98.5% at a loading of 35g/L.

EXAMPLE 6e:

The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 10% of acetonitrile (Buffer B) in 250mM acetic acid (Buffer A). The loading of, trypsinisation end diluted 1:5 using water, was performed. A linear gradient of buffer B was used to elute the purified glargine at yield of 75% with purity 94.5% at a loading of <10 g/L. The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 10% of acetonitrile (Buffer B) in 20mM perchloric acid pH 3.0 (Buffer A). The loading of RP eluate from previous step, diluted to contain 10% acetonitrile, was performed. A linear gradient of buffer B was used to elute the purified glargine at yield of 37.4% with purity 96.2% and glycosylated impurity of 0.28%. The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 6% of ethanol (Buffer B) in 10mM citric acid (Buffer A). The loading of RP eluate from previous step, diluted to 1:3 with water, was performed. A linear gradient of buffer B was used to elute the purified glargine at yield of 70.0% with purity 97.0% at a loading of 35g/L

EXAMPLE 6f:

The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 10% of acetonitrile (Buffer B) in 250mM acetic acid (Buffer A). The loading of, trypsinisation end diluted 1:5 using water, was performed. A linear gradient of buffer B was used to elute the purified glargine at yield of 75% with purity 94.5% at a loading of <10 g/L. The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 10% of acetonitrile (Buffer B) in 100mM Borate buffer pH 8.5 (Buffer A). The loading of RP eluate from previous step, diluted to contain 10% acetonitrile, was performed. A linear gradient of buffer B was used to elute the purified glargine at yield of 79.4% with purity 98.3% and glycosylated impurity of 0.11%. The column packed with RP-HPLC resin (C8 Daisogel 120 Å pore, 10 µm particle size) was equilibrated with 6% of ethanol (Buffer B) in 10mM citric acid (Buffer A). The loading of RP eluate from previous step, diluted to 1:3 with water, was performed. A linear

gradient of buffer B was used to elute the purified glargine at yield of 75.0% with purity 98.8% at a loading of 35g/L.

EXAMPLE 7:**FINAL PRECIPITATION:**

The glargine purified through different reverse phase chromatography steps was precipitated by adding citric acid buffer and ZnCl₂ solution, and adjusting the pH. (The citric acid buffer comprises 15.4 g/L citric acid (anhydrous), 90 g/L di-sodium orthophosphate (anhydrous) buffer, pH adjusted to 6.3. + 0.1, with o-phosphoric acid). The precipitation was done at a pH range of 4.0 to 10.0, preferably 6.0 to 8.0. The product concentration was preferably 2 – 20 g/L. After the precipitation, the product was separated from the clear supernatant by either centrifuging or decanting the supernatant without disturbing the settled precipitate. The precipitate so obtained was washed with chilled water to remove the unbound ions present.

The yield of the whole process was 85 - 90% and the product purity was around 99%.

EXAMPLE 8:

The final suspension was decanted after settling and the slurry was transferred aseptically to the freeze dryer trays. The slurry was then frozen and lyophilized to get dry powder of Insulin Glargine, which can be stored and formulated later.

EXAMPLE 9:**ISOLATION AND CHARACTERIZATION OF THE GLYCOSYLATED IMPURITY:****ISOLATION OF GLYCOFORMS OF GLARGINE FROM IN-HOUSE ACTIVE PHARMACEUTICAL INGREDIENT:**

EXAMPLE 9a:

To characterise glycoforms of impurities mGIG (monoglycosylated insulin glargine) and tGIG (triglycosylated insulin glargine) were isolated by semi-preparatory reverse phase chromatography. Identification of process impurities were carried out on an Agilent 1100 chromatography system (CA, USA) using a Prochrome C18 (250x 8.0mm) column. The mobile phase solvents were aqueous 0.1% TFA (A) and 0.1% TFA in acetonitrile (B). Impurities of glargine were eluted using a linear gradient program: 0 min 25%B; 0-10 min 27%B; 10-15min 29%B; 15-21min 30%B;21-27min 33%B and 27-35min 25%B at a constant flow rate of 1.5mLmin⁻¹. The injection volume was kept at 20µl and column temperature was maintained at 40°C. The eluants were monitored at 214nm. The chromatographic peaks corresponding to glycosylated impurities were analysed online by Agilent 1100 series LCMSD SL ion trap mass spectrometer (Agilent Technologies, USA). The mass spectrometer was operated in ESI mode. The nebuliser gas was kept at 60psi, drying gas at 12.0 L min⁻¹ and drying temperature was kept at 350°C. Positive ion electrospray mass spectrum was recorded in MS mode in the mass range 600-2200 m/z.

The peaks corresponding to mGIG and tGIG were manually collected. Since mGIG (1.0 %) and tGIG (0.3 %) were observed at trace level in formulations, Glargine active pharmaceutical ingredient (API) of the subject invention was used at a concentration 100mg mL⁻¹ for purifying process impurities.

CHARACTERIZATION OF THE FINAL PURIFIED PRODUCT:**EXAMPLE 9b:****RP-HPLC and SEC-HPLC PROFILES OF THE FINAL PRODUCT:**

The RP-HPLC profile and the SEC-HPLC profile of the final protein product obtained has been represented in **FIG 5** and **FIG 6**.

EXAMPLE 9c:

Electrospray ionization mass spectra of glargine, monoglycosylated insulin glargine (mGIG) and tri-glycosylated insulin glargine (tGIG) are represented in FIG 7, FIG 8 and FIG 9 respectively.

Sample	Time	Area	Height	Width	Area%	Symmetry
Batch No.1	18.192	190417	399.4	0.9616	100	0.881

Introduction of one monosaccharide unit to glargine decrease the retention time due to increase in polarity and increase in mass unit of 162 units; similarly introduction of three monosaccharides further increase in polarity and increase in mass of 486 units. The molecular weights of glycoforms were obtained by deconvoluting multiply charged peaks in ESI-MS.

TABLE 6 : Comparisons of molecular weight observed by electrospray and matrix assisted laser desorption ionization techniques.

	MOLECULAR WEIGHT	
	ESI	MALDI
Insulin Glargine	6063.2	6063.4
mGIG	6225.7	6225.6
tGIG	6549.4	6549.8

MALDI mass spectra obtained for mGIG and tGIG showed molecular weights 6225.6 and 6549.8 respectively. The molecular weight data observed for mGIG and tGIG are consistent with data observed by ESI-MS. The molecular weight difference between IG,

mGIG and tGIG showed multiples of 162 mass units suggesting that glycans are covalently attached to glargine.

Figure 8. Matrix assisted laser desorption ionisation mass spectrum of monoglycosylated insulin glargine (mGIG)

Reduction and alkylation experiments were undertaken to identify the chain at which glycosylation had taken place. The method was standardised using glargine as standard. 1.1mg of glargine was dissolved in 500µl of 8M guanidine HCl, 0.1M TRIS and 1mM EDTA buffer maintained at a pH of 9.0. To this 10µl of 1M dithiothreitol (DTT) was added. The contents were mixed and incubated at 37°C for 2hrs. The incubated samples were brought to room temperature and 10µl of iodoacetamide was added. The samples were covered in aluminium foil to protect from light, incubated at 37°C for two hours and analysed by LCMS. In a similar way mGIG and tGIG was reduced and alkylated for the identification of glycosylation.

TABLE 7. Reduction and Alkylation of Glycosylated Glargines done to identify chains at which glycosylation taken place

	Intact RT* (Min)	Intact Mass (m/z)	Chain A		Chain B	
			Retention Time* (Min)	Mass (m/z)	Retention Time (Min)	Mass (m/z)
Glargine	27.5	6063	5.5	2555	20.5	3857
Mono Glycosylated Glargine	23.4	6225	5.2	2718	21.3	3857

Tri						
Glycosylated Glargine	22.2	6549	5.2	2718	20.1	4180

EXAMPLE 9d:**PEPTIDE MAPPING:**

The glycoforms of glargine were subjected to enzyme digestion to identify the amino acid position at which the glycosylation had taken place. 0.65mg of native glargine dissolved in 200µl of 1M TRIS (pH =9.0). To this freshly prepared 50µl of 1mg mL⁻¹ of V 8 protease (Glu-C) was added and incubated at 37°C for two hours. The tryptic digest of IG, mGIG and tGIG were separated on a C18 250x 4.6mm, 5µ, 300° (Waters; symmetry) at a flow rate of 0.8mL min⁻¹. The column temperature was maintained at 40°C. The following gradient was used 0-60min, 5-80%B, 60-65min, 80-5%B. Solvent A=0.1% aqueous TFA and solvent B = Acetonitrile. 20µl of sample was injected and eluted peaks were monitored at 220nm. The samples were analysed online by Agilent 1100 LCMS for GLU-C fragments analysis. A comparative study was done for IG, mGIG and tGIG to find out the difference in retention time and molecular masses.

TABLE 8. Comparison of intact, fragments retention times and masses of glycosylated glargine:

		Retention time	Molecular Mass
Intact	Glargine	27.9	6063
	mGIG	26.8	6225
	tGIG	26.5	6549
	Glargine	10.3	417

A1-A4	mGIG	11.3	417
	tGIG	12.3	417
B22-B32	Glargine	17.4	1428
	mGIG	17.5	1428
	tGIG	16.9	1752
B14-B21	Glargine	20.3	1320
	mGIG	20.3	1320
A18-A21	tGIG	20.3	1320
B1-B13	Glargine	23.2	2970
	mGIG	22.5	3132
A5-A17	tGIG	22.6	3132

Summary of the invention: The present invention aims to provide a method of producing and purifying recombinant protein which encodes a polypeptide displaying immunological and biological activities of mature insulin glargine. The said purified protein is found to have all physiological, immunological and biochemical characteristics similar to mature insulin glargine.

Another preferred aspect of the present invention is characterisation of the glycosylated impurities of glargine obtained in the final product.

The final and the most significant objective of the present invention is to obtain the insulin glargine protein in a pure form. The invention thus permits simple large scale production and subsequent purification of insulin glargine.

An object of the present invention is therefore constituted by an improved method for the preparation of insulin glargine in the pure form.

According to the first aspect of the invention, the recombinant method of producing insulin glargine comprises:

- (a) Preparation of a DNA sequence capable of directing a host microorganism to produce a protein having insulin glargine activity;
- (b) Cloning the said DNA sequence into a vector capable of being transferred into and replicating in a host microorganism, such vector containing operational elements for the DNA sequence;
- (c) Transferring the vector containing the DNA sequence and operational elements into a host microorganism capable of expressing the insulin glargine protein;
- (d) Culturing the microorganism under conditions appropriate for amplification of the vector and expression of the protein;
- (e) Harvesting the protein.

The second aspect of the invention features a method for constructing the recombinant vector carrying the gene of interest including the steps of cloning the insulin glargine precursor gene in frame with a signal peptide in an suitable expression vector, transforming the host cell with the vector harboring the recombinant gene of interest, screening for multi-copy integrants, selection of the recombinant clones with successful integration of insulin glargine precursor in the genome.

The third aspect of the invention relates to small scale expression of the insulin glargine precursor for the HPLC analysis for detecting the expression level of the desired protein of interest.

The fourth aspect of the invention relates to the fermentation process including steps of growing recombinant cells in a growth medium, the cells being a microorganism or cell

culture transformed with an expression vector containing DNA encoding the desired protein, which in case of the instant invention is insulin glargine.

The fifth aspect of the invention is directed to crystallisation of the insulin glargine product produced post fermentation.

The sixth aspect of the invention is related to the trypsinisation process. Glargine can be prepared from the Glargine precursor through an enzymatic conversion. The conversion is effected by the presence of trypsin or trypsin like enzymes of plant, animal or microbial origin.

A significant aspect of the present invention relates to processes and procedures useful in purifying the protein isolated having biological activities similar to that of the natural insulin glargine protein; and more particularly, to processes which utilize RP-HPLC or other chromatography methods for separating the active peptide substance from other substances which do not have such activity and so may be regarded as impurities.

The most preferred aspect of the invention relates to the identification of various glycoforms of insulin analogues more specifically insulin glargine through chemical methods coupled with mass spectrometry techniques such as electrospray and matrix assisted laser desorption ionisation for identification. The invention shall herewith permit selective purification of the product from the aforesaid impurities through optimised down stream purification methods attributed to the better understanding of the nature of impurities present in the final product.

Additional objects and advantages of the invention will be set forth in part in the description which follows, and in part will be obvious from the description or may be learned from practice of the invention. The objects and advantages may be realised and attained by means of the instrumentalities and combinations particularly pointed out in the appended claims. The present invention relates to a method of obtaining a purified, biologically active heterologous protein.

The accompanying drawings, which are incorporated herein and constitute a part of this application, illustrate various attributes useful in this invention and, together with the description, serve to explain the principles of the invention.

SEQUENCE LISTING

<110> Biocon Ltd

Hazra, Partha

<120> A method of obtaining a purified, biologically active
heterologous Protein

<130> 170108

<160> 2

<170> PatentIn version 3.5

<210> 1

<211> 53

<212> PRT

<213> artificial sequence

<220>

<223> Codon optimised glargine precursor sequence

<221> PEPTIDE

<222> (1)..(53)

<400> 1

Phe Val Asn Gln His Leu Cys Gly Ser His Leu Val Glu Ala Leu Tyr

1 5 10 15

Leu Val Cys Gly Glu Arg Gly Phe Phe Tyr Thr Pro Lys Thr Arg Arg

20 25 30
 Gly Ile Val Glu Gln Cys Cys Thr Ser Ile Cys Ser Leu Tyr Gln Leu
 35 40 45
 Glu Asn Tyr Cys Gly
 50

<210> 2

<211> 57

<212> PRT

<213> artificial sequence

<220>

<223> Codon optimised Glargine Precursor sequence

<221> PEPTIDE

<222> (1)..(57)

<400> 2

Gly Ala Val Arg Phe Val Asn Gln His Leu Cys Gly Ser His Leu Val
 1 5 10 15
 Glu Ala Leu Tyr Leu Val Cys Gly Glu Arg Gly Phe Phe Tyr Thr Pro
 20 25 30
 Lys Thr Arg Arg Gly Ile Val Glu Gln Cys Cys Thr Ser Ile Cys Ser
 35 40 45

Leu Tyr Gln Leu Glu Asn Tyr Cys Gly

50

55

WE CLAIM:

1. A method of obtaining a purified, biologically active heterologous protein recombinantly expressed in a yeast expression system, characterised in that said purified protein is devoid of or contains substantially minimal amounts of glycosylated byproducts.
2. A method according to claim 1, wherein the DNA encoding a heterologous protein as represented by formula I
X – B – Y – A
wherein,
X is a leader peptide sequence comprising at least one amino acid.
B is the amino acid sequence of the B chain of the insulin molecule, its derivatives or analogs
Y is a linker peptide comprising at least two amino acids.
A is the amino acid sequence of the A chain of the insulin molecule, its derivatives or analogs and the A and B chain can be modified by amino acid substitution, deletion and/or additions.
3. A method according to claim 2, wherein the linker peptide may be any sequence comprising at least two amino acids with the provision that the first two amino acids represent "RR".
4. A method according to claim 1, wherein said protein is defined by SEQ ID: 1 or SEQ ID: 2.
5. A method according to claim 1, wherein the host cell of the yeast expression system is selected from *Pichia* sp.
6. A method according to claim 1, wherein the host cell of the yeast expression system is selected from the group comprising *Pichia pastoris*, *Pichia methanolica*,

Saccharomyces cerevisiae, *Schisaosaccharomyces pombe*, *Yarrowia lipilitica*, *Hansenula polymorpha*, *Kluyveromyces lactis*.

7. A method according to claim 1, wherein the glycosylated form of the protein may be monoglycosylated, triglycosylated or polyglycosylated.
8. A method according to claim 1, wherein said purified protein contains less than 1% of glycosylated impurities.
9. A process of obtaining a purified, biologically active heterologous protein recombinantly expressed in a host cell which process comprises steps of:
 - a) Culturing host cells transformed by a vector containing a DNA sequence defined by formula I encoding the heterologous protein under conditions suitable for the expression of the protein;
 - b) Recovery of the expressed protein comprising separating the said protein from host cells to produce a recovered protein preparation;
 - c) Subjecting the recovered protein of step (b) to a step of crystallization;
 - d) Performing a step of enzymatic conversion in the presence of trypsin or trypsin like enzyme;
 - e) Purifying the said protein containing at least one related impurity comprising contacting the said protein mixture with a chromatographic matrix wherein purification is carried out employing a polar organic buffer solvent, in an aqueous phase containing an organic acid buffer; and
 - f) Precipitating the eluted protein.
10. A process according to claim 9, wherein the DNA encoding a heterologous protein is represented by formula I
X – B – Y – A
wherein,
X is a leader peptide sequence comprising at least one amino acid.

B is the amino acid sequence of the B chain of the insulin molecule, its derivatives or analogs

Y is a linker peptide comprising at least two amino acids.

A is the amino acid sequence of the A chain of the insulin molecule, its derivatives or analogs

and the A and B chain can be modified by amino acid substitution, deletion and/or addition.

11. A process according to claim 10, wherein the linker peptide may be any sequence comprising at least two amino acids with the provision that the first two amino acids represent "RR".
12. A process according to claim 9, wherein the gene defined by SEQ ID 1 or 2 is cloned in frame with a signal peptide.
13. A process according to claim 12, wherein the gene defined by SEQ ID 1 or 2 is cloned in frame with mat- α -signal peptide.
14. A process according to claim 9, wherein the host cell is selected from *Pichia* sp.
15. A process according to claim 14, wherein the host cell is selected from the group comprising *Pichia pastoris*, *Pichia methanolica*, *Saccharomyces cerevisiae*, *Schisaosaccharomyces pombe*, *Yarrowia lipilitica*, *Hansenula polymorpha*, *Kluyveromyces lactis*.
16. A process according to claim 15, wherein the host cell is *Pichia pastoris*.
17. A process according to claim 16, wherein the host strain is GS115.
18. A process of producing a purified, biologically active heterologous protein of claim 9 which comprises:

- a) Inoculating an aqueous fermentation medium with a transformant yeast strain harboring an expression vector which directs the expression of the said protein; and
 - b) Growing the transformed strain in the fermentation medium under conditions effective for expression of said protein.
19. A process of producing a purified biologically active heterologous protein of claim 18, wherein the expressed protein is captured from the fermentation broth using ion-exchange chromatography.
20. A process of producing a purified, biologically active heterologous protein according to claim 9, said process comprising a step of crystallizing the protein by addition of zinc chloride and phenol.
21. A process of producing a purified, biologically active heterologous protein according to claim 9, said process comprising a step of enzymatic conversion effected in the presence of trypsin in a water-miscible organic solvent.
22. A process according to claim 9, wherein the water-miscible organic solvent is selected from the group comprising DMSO, DMF, ethanol, acetone, acetonitrile, ethyl acetate or mixtures thereof.
23. A process of producing a purified, biologically active heterologous protein according to claim 9, said process comprising a step of RP-HPLC purification of the protein mixture containing at least one related impurity by contacting the said protein mixture with a chromatographic resin matrix wherein purification is carried out employing a polar organic buffer solvent, in an aqueous phase containing an organic acid buffer.
24. A process according to claim 9, wherein the polar buffer solvent is acetonitrile.

25. A process according to claim 19 ,wherein the organic acid buffer is selected from the group comprising citric acid, acetic acid, boric acid, formic acid, hydrochloric acid and phosphoric acid.
26. A purified, biologically active heterologous protein product obtained according to any of the preceding claims, with purity of at least 96%.
27. A purified, biologically active heterologous protein product obtained according to any of the preceding claims, with purity in the range of 97 - 100%.
28. Purified insulin glargine with purity of at least 96%.
29. Purified insulin glargine containing less than 1% of glycosylated impurities.
30. Purified insulin glargine devoid of glycosylated impurities.
31. Purified insulin glargine according to claims, wherein the glycosylated form of the protein may be monoglycosylated, triglycosylated or polyglycosylated.





FIG 1



Enzyme treatment end (10hr)




FIG 2




FIG 3




FIG 4




FIG 5




FIG 6

7/9




FIG 7




FIG 8




FIG 9

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IN2008/000219

A. CLASSIFICATION OF SUBJECT MATTER		
Int. Cl.		
C07K 14/62 (2006.01)		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
Chemical Abstracts on STN Registry: SEQ. ID. NO: 1 and 2 & (X)BYA		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CN 1 873 006 A (SHANGHAI NEWSUMMIT BIOPHARMA CO., LTD.) 12 June 2006 See abstract inter alia.	1-2, 5-7, 18
<input type="checkbox"/> Further documents are listed in the continuation of Box C <input checked="" type="checkbox"/> See patent family annex		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 24 July 2008		Date of mailing of the international search report 31 JUL 2008
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaustralia.gov.au Facsimile No. +61 2 6283 7999		Authorized officer GAVIN THOMPSON AUSTRALIAN PATENT OFFICE (ISO 9001 Quality Certified Service) Telephone No : (02) 6283 2240

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.:

PCT/IN2008/000219

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report	Patent Family Member
CN 1873006	NO FAMILY
Due to data integration issues this family listing may not include 10 digit Australian applications filed since May 2001.	
END OF ANNEX	