



(86) Date de dépôt PCT/PCT Filing Date: 2001/05/11
(87) Date publication PCT/PCT Publication Date: 2001/11/15
(85) Entrée phase nationale/National Entry: 2002/11/06
(86) N° demande PCT/PCT Application No.: AU 2001/000539
(87) N° publication PCT/PCT Publication No.: 2001/085953
(30) Priorité/Priority: 2000/05/11 (PQ7447) AU

(51) Cl.Int.⁷/Int.Cl.⁷ C12N 15/54, A61K 35/14, A61K 31/13,
A61K 38/00

(71) Demandeur/Applicant:
MEDVET SCIENCE PTY. LTD., AU

(72) Inventeurs/Inventors:
VADAS, MATHEW, AU;
GAMBLE, JENNIFER, AU;
XIA, PU, AU;
WANG, LIJUN, AU

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : SPHINGOSINE KINASE ET SES UTILISATIONS
(54) Title: SPHINGOSINE KINASE AND USES THEREOF

(57) **Abrégé/Abstract:**

The present invention relates generally to a method of modulating the growth of cells and, more particularly, to a method of down-regulating the growth of neoplastic cells. The present invention is useful, inter alia, in the therapeutic and/or prophylactic treatment of cancers such as, but not limited to, solid cancers such as cancers of the colon, stomach, lung, brain, bone, oesophagus, pancreas, breast, ovary or uterus.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 November 2001 (15.11.2001)

PCT

(10) International Publication Number
WO 01/85953 A1

- (51) International Patent Classification⁷: **C12N 15/54**, A61K 31/13, 35/14, 38/00
- (74) Agents: **OBRANOVICH, Tania** et al.; Davies Collison Cave, 1 Little Collins Street, Melbourne, VIC 3000 (AU).
- (21) International Application Number: PCT/AU01/00539
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 11 May 2001 (11.05.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
PQ7447 11 May 2000 (11.05.2000) AU
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): **MED-VET SCIENCE PTY. LTD.** [AU/AU]; 20 Dalglish Street, Thebarton, S.A. 5031 (AU).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **VADAS, Mathew** [AU/AU]; 8 Branch Road, Stirling, S.A. 5162 (AU). **GAMBLE, Jennifer** [AU/AU]; 8 Branch Road, Stirling, S.A. 5152 (AU). **XIA, Pu** [AU/AU]; 97 Shakespeare Avenue, Magill, S.A. 5072 (AU). **WANG, Lijun** [AU/AU]; 97 Shakespeare Avenue, Magill, S.A. 5072 (AU).
- Published:**
— *with international search report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: SPHINGOSINE KINASE AND USES THEREOF

(57) Abstract: The present invention relates generally to a method of modulating the growth of cells and, more particularly, to a method of down-regulating the growth of neoplastic cells. The present invention is useful, *inter alia*, in the therapeutic and/or prophylactic treatment of cancers such as, but not limited to, solid cancers such as cancers of the colon, stomach, lung, brain, bone, oesophagus, pancreas, breast, ovary or uterus.



WO 01/85953 A1

- 1 -

A METHOD OF TREATMENT AND AGENTS USEFUL FOR SAME**FIELD OF THE INVENTION**

5 The present invention relates generally to a method of modulating the growth of cells and, more particularly, to a method of down-regulating the growth of neoplastic cells. The present invention is useful, *inter alia*, in the therapeutic and/or prophylactic treatment of cancers such as, but not limited to, solid cancers such as cancers of the colon, stomach, lung, brain, bone, oesophagus, pancreas, breast, ovary or uterus.

10

BACKGROUND OF THE INVENTION

Bibliographic details of the publications referred to by author in this specification are collected at the end of the description.

15

The reference to any prior art in this specification is not, and should not be taken as, an acknowledgment or any form of suggestion that that prior art forms part of the common general knowledge in Australia.

20 Sphingosine kinase is a key regulatory enzyme in a variety of cellular responses. Sphingosine-1-phosphate is known to be an important second messenger in signal transduction (Meyer *et al.*, 1997). It is mitogenic in various cell types (Alessenko, 1998) and appears to trigger a diverse range of important regulatory pathways including prevention of ceramide-induced apoptosis (Culliver *et al.*, 1996), mobilisation of
25 intracellular calcium by an IP₃-independent pathway, stimulation of DNA synthesis, activation of mitogen-activated protein (MAP) kinase pathway, activation of phospholipase D, and regulation of cell motility (for reviews see (Meyer *et al.*, 1997; Spiegel *et al.*, 1998; Igarashi, 1997)).

30 Recent studies (Xia *et al.*, 1998) have shown that sphingosine-1-phosphate is an obligatory signalling intermediate in the inflammatory response of vascular endothelial cells to

- 2 -

tumour necrosis factor- α (TNF α). In spite of its obvious importance, very little is known of the mechanisms that control cellular sphingosine-1-phosphate levels. It is known that sphingosine-1-phosphate levels in the cell are mediated largely by its formation from sphingosine by sphingosine kinase, and to a lesser extent by its degradation by
5 endoplasmic reticulum-associated sphingosine-1-phosphate lyase and sphingosine-1-phosphate phosphatase (Spiegel *et al.*, 1998). Basal levels of sphingosine-1-phosphate in the cell are generally low, but can increase rapidly and transiently when cells are exposed to mitogenic agents. This response appears correlated with an increase in sphingosine kinase activity in the cytosol and can be prevented by addition of the sphingosine kinase
10 inhibitory molecules *N,N*-dimethylsphingosine and *DL-threo*-dihydrosphingosine. This indicates that sphingosine kinase is an important molecule responsible for regulating cellular sphingosine-1-phosphate levels. This places sphingosine kinase in a central and obligatory role in mediating the effects attributed to sphingosine-1-phosphate in the cell.

15 Sphingosine kinase is speculated to play a role in a number of cellular activities including inflammation, calcium mobilisation, cell motility and adhesion molecule expression. However, the precise nature of the cellular activities which are so regulated and the role and mechanistic actions of sphingosine kinase in this regard are only just beginning to be understood. Many of the signals leading to modulation of various cellular activities have
20 not been precisely defined. Elucidating these cellular signalling mechanisms is necessary for the development of therapeutic and prophylactic strategies to disease conditions involving aberrant or otherwise unwanted cellular activities.

In work leading up to the present invention, the inventors have determined that the
25 signalling cascade stimulated by the lipid kinase, sphingosine kinase, plays a major role in oncogenesis and is, in fact, by itself oncogenic when its level of activity is too high. Even in light of preliminary data indicating a role for sphingosine kinase in various cellular activities, the oncogenic activity of sphingosine kinase is a surprising and completely unexpected function attributable to this molecule. Identification of the link between
30 sphingosine kinase and tumour pathogenesis permits the rational design of therapeutic

- 3 -

and/or prophylactic regimes for modulating cell growth and, further, the identification of a range of molecules for use in the modulation of cell growth.

- 4 -

SUMMARY OF THE INVENTION

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will
5 be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

One aspect of the present invention provides a method of modulating the growth of a cell, said method comprising contacting said cell with an effective amount of an agent for a
10 time and under conditions sufficient to modulate the functional activity of sphingosine kinase.

Another aspect of the present invention provides a method of modulating the growth of a cell, said method comprising contacting said cell with an effective amount of an agent for a
15 time and under conditions sufficient to modulate the level of functional activity of sphingosine kinase.

Still another aspect of the present invention provides a method of modulating the proliferation of a cell, said method comprising contacting said cell with an effective
20 amount of an agent for a time and under conditions sufficient to modulate the functional activity of sphingosine kinase.

Still yet another aspect of the present invention provides a method of down-regulating the proliferation of a cell, said method comprising contacting said cell with an effective
25 amount of an agent for a time and under conditions sufficient to down-regulate the functional activity of sphingosine kinase.

Yet still another aspect of the present invention provides a method of up-regulating the proliferation of a cell, said method comprising contacting said cell with an effective
30 amount of an agent for a time and under conditions sufficient to up-regulate the functional activity of sphingosine kinase.

- 5 -

In a further aspect there is provided a method of down-regulating the proliferation of a neoplastic cell, said method comprising contacting said neoplastic cell with an effective amount of an agent for a time and under conditions sufficient to down-regulate the functional activity of sphingosine kinase.

5

Another further aspect of , the present invention provides a method of up-regulating the proliferation of a cell, said method comprising contacting said cell with an effective amount of an agent, for a time and under conditions sufficient to up-regulate the functional activity of sphingosine kinase.

10

Yet another further aspect of the present invention provides a method for the treatment and/or prophylaxis of a condition characterised by uncontrolled cell proliferation in a mammal, said method comprising administering to said mammal an effective amount of an agent for a time and under conditions sufficient to down regulate the functional activity of sphingosine kinase.

15

Still another further aspect of the present invention relates to the treatment and/or prophylaxis of a neoplastic condition in a mammal, said method comprising administering to said mammal an effective amount of an agent for a time and under conditions sufficient to down-regulate the functional activity of sphingosine kinase.

20

Still yet another further aspect of the present invention relates to the use of an agent capable of modulating the functional activity of sphingosine kinase in the manufacture of a medicament for the modulation of cell growth in a mammal wherein down-regulation of the functional activity of said sphingosine kinase down-regulates the subject cell growth and up-regulation of the functional activity of said sphingosine kinase up-regulates the subject cell growth.

25

Yet another aspect relates to agents for use in modulating the functional activity of sphingosine kinase wherein modulating the functional activity of sphingosine kinase modulates cell growth.

30

- 6 -

In yet another aspect the present invention relates to a pharmaceutical composition comprising a modulatory agent as hereinbefore defined and one or more pharmaceutically acceptable carriers and/or diluents. Said modulatory agents are referred to as the active ingredients.

5

Yet another aspect of the present invention relates to modulatory agents, as hereinbefore defined, when used in the method of the present invention.

10 Still yet another aspect of the present invention relates to diagnostic methodology based on screening individuals for the presence of sphingosine kinase or mRNA or protein or the specific forms of sphingosine kinase which are transcribed and/or translated by a given population of cells. The screening methodology may be directed to qualitative and/or quantitative sphingosine kinase analysis. This is particularly useful, for example, for determining whether a given individual is predisposed to or resistant to diseases/disorders
15 in which aberrant, unwanted or otherwise inappropriate cell growth is a component of the disease state and/or is predisposed or resistant to the development of certain forms of aberrant, unwanted or otherwise inappropriate cell growth.

20

- 7 -

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graphical representation of the overexpression of SphK in NIH 3T3 cells. (A) Cytosolic SphK activity was measured in the stable transfected cells with pcDNA3-SphK-FLAG (SK-3T3) or empty vector alone (N-3T3). (B) Intracellular S1P levels were measured in [³H]sphingosine-labelled SK-3T3 or N-3T3 cell pools. (C) Proteins from soluble cell lysates were probed with anti-FLAG monoclonal antibodies (M2, Kodak).

Figure 2 is a graphical representation of SphK overexpression accelerates proliferation in NIH 3T3 cells. (A) Growth curves reveal that SK-3T3 cells (right panel) proliferate more rapid than N-3T3 cells (left panel). Values are the mean of triplicate determinations and similar results were obtained in three independent experiments. (B) Cell growth at saturation density. Cells were counted at absolute confluence (open bars) and 24 h later of confluence (grey bars), respectively. (C) DMS inhibits SphK overexpression induced proliferation. Equal numbers of N-3T3 and SK-3T3 cells were cultured in DMEM containing 10% serum in the presence or absence of DMS (2.5 μ M) for 5 days. Media was replaced every day. Data in (B) and (C) are the means \pm S.D. from three independent experiments done in triplicate.

Figure 3 is an image of overexpression of SphK induced NIH 3T3 cell transformation. (A) Focus formation of NIH 3T3 cell transfectants. Cultures transfected with SphK (SK-3T3) or vector alone (N-3T3) were photographed 12 days after transfection (40x magnification). (B) Colony formation in soft agar. N-3T3 and SK-3T3 cells were cultured in growth medium containing 0.33% agar and fed with the medium containing various concentrations of DMS every 2 days. Photographs were taken after 2 weeks of cultures followed by staining with MTT. (C) NIH 3T3 cells were transfected with V12-Ras or v-Src, SphK activity was measured after 48 h transfection. (D) Focus formation assays were performed in V12-Ras, v-Src, or SphK transfected NIH 3T3 cells in the absence or presence of DMS (2.5 μ M) over two weeks.

30

- 8 -

Figure 4 is an image of cells overexpressing SphK are tumourigenic. **(A)** Photograph of tumours in the NOD/SCID mice injected with SphK-transfected NIH 3T3 cells. **(B)** Morphology of a tumour (insert) and the paraffin fixed section stained with hematoxylin and eosin (60x magnification). **(C)** Whole cell extracts from three individual tumours
5 (lane 4-6) and their peripheral tissues (lane 1-3) and N-3T3 (lane 7) or SK-3T3 cells (lane 8) were analysed by Western blot. The top blot was probed with anti-FLAG antibody and the bottom with anti-actin antibody (Santa Cruz).

- 9 -

DETAILED DESCRIPTION OF THE INVENTION

- The present invention is predicated, in part, on the determination of a correlation between cell growth, in particular oncogenesis, and modulation in the level of activity of sphingosine kinase. The identification of this correlation permits the identification and rational design of methodology and products for use in therapy, prophylaxis and diagnosis of disease conditions characterised by aberrant, unwanted or otherwise inappropriate cell growth, in particular, uncontrolled oncogene induced proliferation.
- 10 Accordingly, one aspect of the present invention provides a method of modulating the growth of a cell, said method comprising contacting said cell with an effective amount of an agent for a time and under conditions sufficient to modulate the functional activity of sphingosine kinase.
- 15 Reference to "sphingosine kinase" should be understood as including reference to all forms of sphingosine kinase or derivative, homologue, analogue, equivalent or mimetic thereof or other molecules having the function of sphingosine kinase and to nucleic acid molecules encoding sphingosine kinase derivative, homologue, analogue, equivalent or mimetic thereof. This includes, for example, all protein or nucleic acid forms of sphingosine kinase or its functional derivative, homologue, analogue, equivalent or mimetic thereof including, for example, any isoforms which arise from alternative splicing of sphingosine kinase mRNA or mutants or polymorphic variants of sphingosine kinase. It should also be understood that reference to a "nucleic acid form of sphingosine kinase" is a reference to a nucleic acid encoding sphingosine kinase and includes reference to any sphingosine kinase regulatory element (such as promoters or enhancers) which regulate the expression of sphingosine kinase and includes regulatory elements which are located at a position other than between the sphingosine kinase genomic DNA transcription initiation and determination sites. "Sphingosine kinase" should also be understood to include reference to any other molecules which exhibit the functional activity of sphingosine kinase. Such molecules include, for example, endogenously expressed molecules which exhibit sphingosine kinase functional activity or molecules which have been introduced into the
- 20
- 25
- 30

- 10 -

body and which mimic at least one of the sphingosine kinase functions. These molecules may be recombinant, synthetic or naturally occurring. To the extent that it is not specified, any reference to modulating the activity of sphingosine kinase includes modulating the expression of a nucleic acid molecule encoding sphingosine kinase or the functional
5 activity of the sphingosine kinase expression product and should also be understood to include reference to modulating the expression or functional activity or a sphingosine kinase functional equivalent or derivative.

Reference to "modulating the functional activity" of sphingosine kinase should be
10 understood as a reference to up-regulating, down-regulating or otherwise altering any one or more of the functional activities or sphingosine kinase. This includes, for example, modulating the occurrence of one or more of the sphingosine kinase functional activities, modulating the rate at which a given activity is performed, modulating the level at which an activity is performed, modulating the number of activities which are capable of being
15 performed or modulating the role or extent to which any activity is performed or the nature of an activity. Changes in the activity of sphingosine kinase can be effected by any one of a number of means including, but not limited to, post translational modification, associated proteins or other molecules or translation. Modulating said activity should also be understood to encompass increasing or decreasing the concentration levels of sphingosine
20 kinase (for example by modulating expression of sphingosine kinase). In a preferred embodiment, the subject functional activity is the level of activity of sphingosine kinase.

Accordingly, the present invention more preferably provides a method of modulating the growth of a cell, said method comprising contacting said cell with an effective amount of
25 an agent for a time and under conditions sufficient to modulate the level of functional activity of sphingosine kinase.

As detailed above, the present invention is predicated on the identification of a correlation between sphingosine kinase functional activity and cell growth, in particular, oncogenic
30 cell proliferation. Without limiting the present invention to any one theory or mode of action, the inventors have found that the signalling cascade stimulated by the lipid kinase,

- 11 -

sphingosine kinase, has a major role in oncogenesis. Specifically, constitutive activation of sphingosine kinase by overexpression in cells causes cell transformation and tumour formation, thereby indicating that a wild type human lipid kinase is by itself oncogenic. Furthermore, sphingosine kinase is also involved in Ras but not v-Src induced transformation. Finally inhibition of sphingosine kinase activity utilising a sphingosine kinase inhibitor not only reverses transformation in cells overexpressing sphingosine kinase but does so also in Ras transformed cells. In this regard, reference to "modulating" the growth of a cell should be understood as a reference to up-regulating or down-regulating the growth of a cell. More specifically, reference to "down-regulating" should be understood as a reference to preventing, reducing or otherwise inhibiting one or more aspects of the growth of a cell (including inducing the apoptosis of or otherwise killing a cell) while reference to "up-regulating" should be understood to have the converse meaning. Reference to the "growth" of a cell should be understood in its broadest sense to include reference to all aspects of cell division/proliferation and/or differentiation. In a particularly preferred embodiment, the subject growth is proliferation.

According to this preferred embodiment, there is provided a method of modulating the proliferation of a cell, said method comprising contacting said cell with an effective amount of an agent for a time and under conditions sufficient to modulate the functional activity of sphingosine kinase.

Still more preferably, there is provided a method of down-regulating the proliferation of a cell, said method comprising contacting said cell with an effective amount of an agent for a time and under conditions sufficient to down-regulate the functional activity of sphingosine kinase.

In still another preferred embodiment, there is provided a method of up-regulating the proliferation of a cell, said method comprising contacting said cell with an effective amount of an agent for a time and under conditions sufficient to up-regulate the functional activity of sphingosine kinase.

- 12 -

In a most preferred embodiment said proliferation is uncontrolled proliferation such as that caused by the transformation of a cell. Preferably said transformation is caused by the upregulation of an oncogene such as Ras or by sphingosine kinase overexpression oncogenic activity.

5

Preferably said functional activity is the level of sphingosine kinase functional activity.

It should be understood that reference to a "cell" in the context of the present invention is a reference to any form or type of cell, irrespective of its origin. For example, the cell may
10 be a naturally occurring normal or abnormal cell or it may be manipulated, modified or otherwise treated either *in vitro* or *in vivo* such as a cell which has been freeze/thawed or genetically, biochemically or otherwise modified either *in vitro* or *in vivo* (including, for example, cells which are the result of the fusion of two distinct cell types). Preferably, the cell is a neoplastic cell. By "neoplastic cell" is meant a cell exhibiting uncontrolled
15 proliferation. The neoplastic cell may be a benign cell or a malignant cell. Preferably the cell is malignant. In one particular embodiment, the neoplastic cell is a malignant cell the proliferation of which would form a solid tumour such as a malignant cell derived from the colon, stomach, lung, brain, bone, oesophagus or pancreas.

20 According to this most preferred embodiment there is provided a method of down-regulating the proliferation of a neoplastic cell, said method comprising contacting said neoplastic cell with an effective amount of an agent for a time and under conditions sufficient to down-regulate the functional activity of sphingosine kinase.

25 Preferably the neoplastic cell is a malignant cell derived from the colon, stomach, lung, brain, bone, oesophagus, pancreas, breast, ovary or uterus.

Still more preferably said malignant cell has become transformed either due to upregulation of an oncogene such as Ras or due to sphingosine kinase overexpression
30 oncogenic activity.

- 13 -

Even more preferably, said functional activity is the level of sphingosine kinase functional activity.

It should be understood that the cell which is treated according to the method of the present invention may be located *ex vivo* or *in vivo*. By "*ex vivo*" is meant that the cell has been removed from the body of a subject wherein the modulation of its growth will be achieved *in vitro*. For example, the cell may be a non-neoplastic cell which is to be immortalised by up-regulating sphingosine kinase activity. In accordance with the preferred aspects of the present invention, the cell may be a neoplastic cell, such as a malignant cell, located *in vivo* (such as in the colon) and the down-regulation of its growth will be achieved by applying the method of the present invention *in vivo* to down-regulate the level of sphingosine kinase functional activity. It should also be understood that where reference is made to a specific cell type which is located *in vivo*, such as a malignant colorectal cell, this cell may be located in the colorectal area of the patient. If a colorectal primary malignancy has metastasised, the subject colorectal cell may be located in another region of the patient's body. For example, it may form part of a secondary tumour (metastasis) which is located, for example, in the liver, lymph node or bone.

Although the preferred method is to down-regulate the proliferation of a neoplastic cell, for example as a therapeutic treatment for cancer, it may also be desirable to up-regulate cell growth. For example, it may be desirable to immortalise a population of cells *in vitro*, to facilitate their long term *in vitro* use or, for example, to facilitate the *in vitro* growth of tissues such as skin. In another example, it may be useful to adapt cell lines to less fastidious growth conditions such as a capacity to grow in low serum conditions.

25

According to this preferred embodiment, the present invention provides a method of up-regulating the proliferation of a cell, said method comprising contacting said cell with an effective amount of an agent, for a time and under conditions sufficient to up-regulate the functional activity of sphingosine kinase.

30

Preferably, said functional activity is the level of sphingosine kinase functional activity.

- 14 -

Without limiting this aspect of the present invention to any one theory or mode of action, said up-regulation of cellular proliferation is preferably transformation of the cell via the up-regulation of sphingosine kinase oncogenic activity.

5 In this regard, an "effective amount" means an amount necessary to at least partly attain the desired effect, or to delay the onset of, inhibit the progression of, or halt altogether, the onset of progression of a disease condition which is being treated. Such amounts will depend, of course, on the particular conditions being treated, the severity of the condition and individual patient parameters including age, physical conditions, size, weight and
10 concurrent treatment. These factors are well known of those of ordinary skill in the art and can be addressed with no more than routine experimentation. It is preferred generally that a maximum dose be used, that is, the highest safe dose according to sound medical judgement. It will be understood by those of ordinary skill in the art, however, that a lower dose or tolerable dose may be administered for medical reasons, psychological reasons or
15 for virtually any other reasons.

Modulation of the activity of sphingosine kinase by the administration of an agent to a cell can be achieved by one of several techniques including, but in no way limited to, introducing into said cell an agent (said agent being a proteinaceous or non-proteinaceous
20 molecule) which directly or indirectly:

- (i) modulates the expression of sphingosine kinase; or
- (ii) modulates the functional activity of sphingosine kinase expression product.

25 In this regard, modulation of the functional activity of sphingosine kinase can be achieved by any one of several techniques, including, but in no way limited to, introducing into said cell a proteinaceous or non-proteinaceous molecule which directly or indirectly:

- (i) modulates synthesis of said sphingosine kinase;
- 30 (ii) functions as an antagonist to said sphingosine kinase;

- 15 -

(iii) functions as an agonist to said sphingosine kinase (including administration of sphingosine kinase expression product *per se* or functional equivalent, derivative, homologue, analogue or mimetic thereof).

5 Said proteinaceous molecule may be derived from natural, recombinant or synthetic sources including fusion proteins or following, for example, natural product screening. Said non-proteinaceous molecule may be derived from natural sources, such as for example natural product screening or may be chemically synthesised. The present invention contemplates chemical analogs of said sphingosine kinase capable of acting as
10 agonists or antagonists of said sphingosine kinase. Chemical agonists may not necessarily be derived from said sphingosine kinase but may share certain conformational similarities. Alternatively, chemical agonists may be specifically designed to mimic certain physiochemical properties of said sphingosine kinase. Antagonists may be any compound capable of blocking, inhibiting or otherwise preventing said sphingosine
15 kinase from carrying out its normal biological functions (for example N,N-dimethylsphingosine or DL-threo-dihydrosphingosine). Antagonists include monoclonal antibodies specific for said sphingosine kinase, or parts of said sphingosine kinase, and antisense nucleic acids which prevent transcription or translation of genes or mRNA in the subject cells. Modulation of expression may also be achieved utilising antigens,
20 RNA, ribosomes, DNazymes, RNA aptamers, antibodies or molecules suitable for use in co-suppression.

Said proteinaceous or non-proteinaceous molecule may act either directly or indirectly to modulate the expression of sphingosine kinase or the activity of the sphingosine kinase
25 expression product. Said molecule acts directly if it associates with the sphingosine kinase nucleic acid molecule or expression product to modulate expression or activity. Said molecule acts indirectly if it associates with a molecule other than the sphingosine kinase nucleic acid molecule or expression product which other molecule either directly or indirectly modulates the expression or activity of the sphingosine kinase nucleic acid
30 molecule or expression product. Accordingly, the method of the present invention

- 16 -

encompasses regulation of sphingosine kinase nucleic acid molecule expression or expression product functional activity via the induction of a cascade of regulatory steps.

Still without limiting the operation of the present invention to any one theory or mode of
5 action, sphingosine kinase is known to function via a signalling pathway which is commonly referred to as the sphingosine kinase signalling pathway. The "sphingosine kinase signalling pathway" is defined as a signalling pathway which utilises sphingosine kinase. In terms of indirect modulation of sphingosine kinase functional activity, it should be understood that the object of the invention could be achieved by modulating the activity
10 of sphingosine kinase signalling pathway components which function either upstream or downstream of sphingosine kinase, to the extent that it forms part of this pathway. For example, modulation of said "sphingosine kinase activity" may be achieved by:

- 15 (i) modulation of the catalytic activity of sphingosine kinase by competition with substrate (for example, sphingosine or ATP);
- (ii) interference with the catalytic activity of sphingosine kinase by an allosteric mechanisms (binding to sites on the molecule other than the substrate-binding sites); or
- 20 (iii) interfering with enzyme activation, such as by altering:
 - post-translational covalent modification such as phosphorylation, lipid modification
 - 25 - non-covalent coupling to a required co-activator such as a protein, lipid or ion
 - subcellular localisation of the enzyme.

"Derivatives" include fragments, parts, portions, mutants, variants and mimetics from
30 natural, synthetic or recombinant sources including fusion proteins. Parts or fragments include, for example, active regions of sphingosine kinase. Derivatives may be derived from insertion, deletion or substitution of amino acids. Amino acid insertional derivatives

- 17 -

include amino and/or carboxylic terminal fusions as well as intrasequence insertions of single or multiple amino acids. Insertional amino acid sequence variants are those in which one or more amino acid residues are introduced into a predetermined site in the protein although random insertion is also possible with suitable screening of the resulting
5 product. Deletional variants are characterized by the removal of one or more amino acids from the sequence. Substitutional amino acid variants are those in which at least one residue in the sequence has been removed and a different residue inserted in its place. An example of substitutional amino acid variants are conservative amino acid substitutions. Conservative amino acid substitutions typically include substitutions within the following
10 groups: glycine and alanine; valine, isoleucine and leucine; aspartic acid and glutamic acid; asparagine and glutamine; serine and threonine; lysine and arginine; and phenylalanine and tyrosine. Additions to amino acid sequences include fusions with other peptides, polypeptides or proteins.

15 Reference to "homologues" should be understood as a reference to sphingosine kinase nucleic acid molecules or proteins derived from species other than the species being treated.

Chemical and functional equivalents of sphingosine kinase nucleic acid or protein
20 molecules should be understood as molecules exhibiting any one or more of the functional activities of these molecules and may be derived from any source such as being chemically synthesized or identified via screening processes such as natural product screening.

The derivatives include fragments having particular epitopes or parts of the entire protein
25 fused to peptides, polypeptides or other proteinaceous or non-proteinaceous molecules.

Analogues contemplated herein include, but are not limited to, modification to side chains, incorporating of unnatural amino acids and/or their derivatives during peptide, polypeptide or protein synthesis and the use of crosslinkers and other methods which impose
30 conformational constraints on the proteinaceous molecules or their analogues.

Derivatives of nucleic acid sequences may similarly be derived from single or multiple nucleotide substitutions, deletions and/or additions including fusion with other nucleic acid

- 18 -

molecules. The derivatives of the nucleic acid molecules of the present invention include oligonucleotides, PCR primers, antisense molecules, molecules suitable for use in cosuppression and fusion of nucleic acid molecules. Derivatives of nucleic acid sequences also include degenerate variants.

5

Examples of side chain modifications contemplated by the present invention include modifications of amino groups such as by reductive alkylation by reaction with an aldehyde followed by reduction with NaBH₄; amidination with methylacetimidate; acylation with acetic anhydride; carbamoylation of amino groups with cyanate;

10

trinitrobenzylation of amino groups with 2, 4, 6-trinitrobenzene sulphonic acid (TNBS); acylation of amino groups with succinic anhydride and tetrahydrophthalic anhydride; and pyridoxylation of lysine with pyridoxal-5-phosphate followed by reduction with NaBH₄.

15

The guanidine group of arginine residues may be modified by the formation of heterocyclic condensation products with reagents such as 2,3-butanedione, phenylglyoxal and glyoxal.

20

The carboxyl group may be modified by carbodiimide activation *via* O-acylisourea formation followed by subsequent derivitisation, for example, to a corresponding amide.

25

Sulphydryl groups may be modified by methods such as carboxymethylation with iodoacetic acid or iodoacetamide; performic acid oxidation to cysteic acid; formation of a mixed disulphides with other thiol compounds; reaction with maleimide, maleic anhydride or other substituted maleimide; formation of mercurial derivatives using 4-chloromercuribenzoate, 4-chloromercuriphenylsulphonic acid, phenylmercury chloride, 2-chloromercuri-4-nitrophenol and other mercurials; carbamoylation with cyanate at alkaline pH.

30

Tryptophan residues may be modified by, for example, oxidation with N-bromosuccinimide or alkylation of the indole ring with 2-hydroxy-5-nitrobenzyl bromide or sulphenyl halides. Tyrosine residues on the other hand, may be altered by nitration with tetranitromethane to form a 3-nitrotyrosine derivative.

- 19 -

Modification of the imidazole ring of a histidine residue may be accomplished by alkylation with iodoacetic acid derivatives or N-carboethoxylation with diethylpyrocarbonate.

- 5 Examples of incorporating unnatural amino acids and derivatives during protein synthesis include, but are not limited to, use of norleucine, 4-amino butyric acid, 4-amino-3-hydroxy-5-phenylpentanoic acid, 6-aminohexanoic acid, t-butylglycine, norvaline, phenylglycine, ornithine, sarcosine, 4-amino-3-hydroxy-6-methylheptanoic acid, 2-thienyl alanine and/or D-isomers of amino acids. A list of unnatural amino acids contemplated
10 herein is shown in Table 1.

- 20 -

TABLE 1

5	Non-conventional amino acid	Code	Non-conventional amino acid	Code
	α -aminobutyric acid	Abu	L-N-methylalanine	Nmala
	α -amino- α -methylbutyrate	Mgab	L-N-methylarginine	Nmarg
10	aminocyclopropane-carboxylate	Cpro	L-N-methylasparagine	Nmasn
	aminoisobutyric acid	Aib	L-N-methylaspartic acid	Nmasp
	aminonorbornyl-carboxylate	Norb	L-N-methylcysteine	Nmcys
			L-N-methylglutamine	Nmgln
			L-N-methylglutamic acid	Nmglu
15	cyclohexylalanine	Chexa	L-N-methylhistidine	Nmhis
	cyclopentylalanine	Cpen	L-N-methylisoleucine	Nmile
	D-alanine	Dal	L-N-methylleucine	Nmleu
	D-arginine	Darg	L-N-methyllysine	Nmlys
	D-aspartic acid	Dasp	L-N-methylmethionine	Nmmet
20	D-cysteine	Dcys	L-N-methylnorleucine	Nmnle
	D-glutamine	Dgln	L-N-methylnorvaline	Nmnva
	D-glutamic acid	Dglu	L-N-methylornithine	Nmorn
	D-histidine	Dhis	L-N-methylphenylalanine	Nmphe
	D-isoleucine	Dile	L-N-methylproline	Nmpro
25	D-leucine	Dleu	L-N-methylserine	Nmser
	D-lysine	Dlys	L-N-methylthreonine	Nmthr
	D-methionine	Dmet	L-N-methyltryptophan	Nmtrp
	D-ornithine	Dorn	L-N-methyltyrosine	Nmtyr
	D-phenylalanine	Dphe	L-N-methylvaline	Nmval
30	D-proline	Dpro	L-N-methylethylglycine	Nmetg
	D-serine	Dser	L-N-methyl-t-butylglycine	Nmtbug
	D-threonine	Dthr	L-norleucine	Nle

- 21 -

	D-tryptophan	Dtrp	L-norvaline	Nva
	D-tyrosine	Dtyr	α -methyl-aminoisobutyrate	Maib
	D-valine	Dval	α -methyl- -aminobutyrate	Mgab
	D- α -methylalanine	Dmala	α -methylcyclohexylalanine	Mchexa
5	D- α -methylarginine	Dmarg	α -methylcyclopentylalanine	Mcpen
	D- α -methylasparagine	Dmasn	α -methyl- α -naphthylalanine	Manap
	D- α -methylaspartate	Dmasp	α -methylpenicillamine	Mpen
	D- α -methylcysteine	Dmcys	N-(4-aminobutyl)glycine	Nglu
	D- α -methylglutamine	Dmgln	N-(2-aminoethyl)glycine	Naeg
10	D- α -methylhistidine	Dmhis	N-(3-aminopropyl)glycine	Norn
	D- α -methylisoleucine	Dmile	N-amino- α -methylbutyrate	Nmaabu
	D- α -methylleucine	Dmleu	α -naphthylalanine	Anap
	D- α -methyllysine	Dmlys	N-benzylglycine	Nphe
	D- α -methylmethionine	Dmmet	N-(2-carbamylethyl)glycine	Ngln
15	D- α -methylornithine	Dmorn	N-(carbamylmethyl)glycine	Nasn
	D- α -methylphenylalanine	Dmphe	N-(2-carboxyethyl)glycine	Nglu
	D- α -methylproline	Dmpro	N-(carboxymethyl)glycine	Nasp
	D- α -methylserine	Dmser	N-cyclobutylglycine	Ncbut
	D- α -methylthreonine	Dmthr	N-cycloheptylglycine	Nchep
20	D- α -methyltryptophan	Dmtrp	N-cyclohexylglycine	Nchex
	D- α -methyltyrosine	Dmty	N-cyclodecylglycine	Ncdec
	D- α -methylvaline	Dmval	N-cyclododecylglycine	Ncdod
	D-N-methylalanine	Dnmala	N-cyclooctylglycine	Ncoct
	D-N-methylarginine	Dnmarg	N-cyclopropylglycine	Ncpro
25	D-N-methylasparagine	Dnmasn	N-cycloundecylglycine	Ncund
	D-N-methylaspartate	Dnmasp	N-(2,2-diphenylethyl)glycine	Nbhm
	D-N-methylcysteine	Dnmcys	N-(3,3-diphenylpropyl)glycine	Nbhe
	D-N-methylglutamine	Dnmgln	N-(3-guanidinopropyl)glycine	Narg
	D-N-methylglutamate	Dnmglu	N-(1-hydroxyethyl)glycine	Nthr
30	D-N-methylhistidine	Dnmhis	N-(hydroxyethyl)glycine	Nser
	D-N-methylisoleucine	Dnmile	N-(imidazolylethyl)glycine	Nhis
	D-N-methylleucine	Dnmleu	N-(3-indolylyethyl)glycine	Nhtrp
	D-N-methyllysine	Dnmlys	N-methyl- γ -aminobutyrate	Nmgabu

- 22 -

	N-methylcyclohexylalanine	Nmchexa	D-N-methylmethionine	Dnmmtet
	D-N-methylornithine	Dnmorn	N-methylcyclopentylalanine	Nmcpen
	N-methylglycine	Nala	D-N-methylphenylalanine	Dnmphe
	N-methylaminoisobutyrate	Nmaib	D-N-methylproline	Dnmpro
5	N-(1-methylpropyl)glycine	Nile	D-N-methylserine	Dnmser
	N-(2-methylpropyl)glycine	Nleu	D-N-methylthreonine	Dnmthr
	D-N-methyltryptophan	Dnmtrp	N-(1-methylethyl)glycine	Nval
	D-N-methyltyrosine	Dnmtyr	N-methyl- <i>n</i> -naphthylalanine	Nmanap
	D-N-methylvaline	Dnmval	N-methylpenicillamine	Nmpen
10	γ -aminobutyric acid	Gabu	N-(<i>p</i> -hydroxyphenyl)glycine	Nhtyr
	<i>L</i> - <i>t</i> -butylglycine	Tbug	N-(thiomethyl)glycine	Ncys
	<i>L</i> -ethylglycine	Etg	penicillamine	Pen
	<i>L</i> -homophenylalanine	Hphe	<i>L</i> - α -methylalanine	Mala
	<i>L</i> - α -methylarginine	Marg	<i>L</i> - α -methylasparagine	Masn
15	<i>L</i> - α -methylaspartate	Masp	<i>L</i> - α -methyl- <i>t</i> -butylglycine	Mtbug
	<i>L</i> - α -methylcysteine	Mcys	<i>L</i> -methylethylglycine	Metg
	<i>L</i> - α -methylglutamine	Mglu	<i>L</i> - α -methylglutamate	Mglu
	<i>L</i> - α -methylhistidine	Mhis	<i>L</i> - α -methylhomophenylalanine	Mhphe
	<i>L</i> - α -methylisoleucine	Mile	N-(2-methylthioethyl)glycine	Nmet
20	<i>L</i> - α -methylleucine	Mleu	<i>L</i> - α -methyllysine	Mlys
	<i>L</i> - α -methylmethionine	Mmet	<i>L</i> - α -methylnorleucine	Mnle
	<i>L</i> - α -methylnorvaline	Mnva	<i>L</i> - α -methylornithine	Morn
	<i>L</i> - α -methylphenylalanine	Mphe	<i>L</i> - α -methylproline	Mpro
	<i>L</i> - α -methylserine	Mser	<i>L</i> - α -methylthreonine	Mthr
25	<i>L</i> - α -methyltryptophan	Mtrp	<i>L</i> - α -methyltyrosine	Mtyr
	<i>L</i> - α -methylvaline	Mval	<i>L</i> -N-methylhomophenylalanine	Nmhphe
	N-(N-(2,2-diphenylethyl) carbonylmethyl)glycine	Nnbhm	N-(N-(3,3-diphenylpropyl) carbonylmethyl)glycine	Nnbhe
30	1-carboxy-1-(2,2-diphenyl-Nmbc ethylamino)cyclopropane			

- 23 -

Crosslinkers can be used, for example, to stabilise 3D conformations, using homo-bifunctional crosslinkers such as the bifunctional imido esters having $(\text{CH}_2)_n$ spacer groups with $n=1$ to $n=6$, glutaraldehyde, N-hydroxysuccinimide esters and hetero-bifunctional reagents which usually contain an amino-reactive moiety such as N-hydroxysuccinimide
5 and another group specific-reactive moiety.

The molecules which may be administered to a mammal in accordance with the present invention may also be linked to a targeting means such as a monoclonal antibody, which provides specific delivery of these molecules to the target cells.

10

A further aspect of the present invention relates to the use of the invention in relation to the treatment and/or prophylaxis of disease conditions. Without limiting the present invention to any one theory or mode of action, the inventors have determined not only that *constitutive activation of sphingosine kinase causes cell transformation and tumour*
15 *development, thereby indicating that sphingosine kinase is by itself oncogenic, but that sphingosine kinase inhibition is also effective in down-regulating neoplastic cell proliferation where the subject cell has been transformed by certain unrelated oncogenes such as Ras induced transformation. Accordingly, the method of the present invention is particularly useful, but in no way limited to, use in the treatment of primary and secondary*
20 *malignancies such as those associated with solid tumours of the colon, stomach, lung, brain, bone, oesophagus and pancreas and, in particular, tumours which arise from the proliferation of Ras transformed cells. Although the preferred method is to down-regulate uncontrolled cellular proliferation in a subject, up-regulation of cell growth may also be desirable in certain circumstances such as to promote wound healing, angiogenesis or other*
25 *healing process.*

Accordingly, the present invention contemplates a method for the treatment and/or prophylaxis of a condition characterised by aberrant, unwanted or otherwise inappropriate cell growth in a mammal, said method comprising administering to said mammal an
30 effective amount of an agent for a time and under conditions sufficient to modulate the functional activity of sphingosine kinase wherein down-regulation of the functional activity of said sphingosine kinase down-regulates the subject cell growth and up-

- 24 -

regulation of the functional activity of said sphingosine kinase up-regulates the subject cell growth.

Reference to "aberrant, unwanted or otherwise inappropriate" cell growth should be
5 understood as a reference to over active cell growth, to physiologically normal cell growth which is inappropriate in that it is unwanted or to insufficient cell growth. Preferably, said inappropriate cell growth is uncontrolled cell proliferation.

According to this preferred embodiment, there is provided a method for the treatment
10 and/or prophylaxis of a condition characterised by uncontrolled cell proliferation in a mammal, said method comprising administering to said mammal an effective amount of an agent for a time and under conditions sufficient to down regulate the functional activity of sphingosine kinase.

15 Preferably said uncontrolled cell proliferation is caused by the transformation of the cell by oncogene up-regulation or by sphingosine kinase overexpression oncogenic activity.

Still more preferably said cell is a malignant cell which forms a solid tumour of the colon, stomach, lung, brain, bone, oesophagus, pancreas, breast, ovary or uterus.

20

Preferably, said agent is N,N-dimethylsphingosine or DL-threo-dihydrosphingosine.

The method of the present invention preferably facilitates the subject proliferation being reduced, retarded or otherwise inhibited. Reference to "reduced, retarded or otherwise
25 inhibited" should be understood as a reference to inducing or facilitating the partial or complete inhibition of cell proliferation. Said inhibition may occur by either direct or indirect mechanisms and includes the induction of cellular apoptosis or other cellular killing mechanisms.

30 The subject of the treatment or prophylaxis is generally a mammal such as but not limited to human, primate, livestock animal (eg. sheep, cow, horse, donkey, pig), companion

- 25 -

animal (eg. dog, cat), laboratory test animal (eg. mouse, rabbit, rat, guinea pig, hamster),
captive wild animal (eg. fox, deer). Preferably the mammal is a human or primate. Most
preferably the mammal is a human. Although the present invention is exemplified utilising
a murine model, this is not intended as a limitation on the application of the method of the
5 present invention to other species, in particular, humans.

Another aspect of the present invention relates to the treatment and/or prophylaxis of a
neoplastic condition in a mammal, said method comprising administering to said mammal
an effective amount of an agent for a time and under conditions sufficient to down-regulate
10 the functional activity of sphingosine kinase.

Preferably said neoplastic condition is a malignant condition and even more preferably a
solid malignancy such as a tumour of the colon, stomach, lung, brain, bone, oesophagus or
pancreas.

15 More preferably the subject malignancy is caused by transformation of the cell via
oncogene up-regulation or by sphingosine kinase overexpression oncogenic activity.

Still more preferably said agent is *N,N*-dimethylsphingosine or *DL*-threo-
20 dihydrophingosine.

Reference herein to "treatment" and "prophylaxis" is to be considered in its broadest
context. The term "treatment" does not necessarily imply that a mammal is treated until
total recovery. Similarly, "prophylaxis" does not necessarily mean that the subject will not
25 eventually contract a disease condition. Accordingly, treatment and prophylaxis including
amelioration of the symptoms of a particular condition or preventing or otherwise reducing
the risk of developing a particular condition. The term "prophylaxis" may be considered as
reducing the severity or onset of a particular condition. "Treatment" may also reduce the
severity of an existing condition.

30

- 26 -

Administration of the agent (including sphingosine kinase or functional equivalent, derivative, homologue, analogue or mimetic thereof or sphingosine kinase nucleic acid molecule) [herein referred to as "modulatory agent"], in the form of a pharmaceutical composition, may be performed by any convenient means. The modulatory agent of the pharmaceutical composition is contemplated to exhibit therapeutic activity when administered in an amount which depends on the particular case. The variation depends, for example, on the human or animal and the modulatory agent chosen. A broad range of doses may be applicable. Considering a patient, for example, from about 0.1 mg to about 1 mg of modulatory agent may be administered per kilogram of body weight per day.

Dosage regimes may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily, weekly, monthly or other suitable time intervals or the dose may be proportionally reduced as indicated by the exigencies of the situation. The modulatory agent may be administered in a convenient manner such as by the oral, intravenous (where water soluble), intraperitoneal, intramuscular, subcutaneous, intradermal or suppository routes or implanting (e.g. using slow release molecules). The modulatory agent may be administered in the form of pharmaceutically acceptable nontoxic salts, such as acid addition salts or metal complexes, e.g. with zinc, iron or the like (which are considered as salts for purposes of this application). Illustrative of such acid addition salts are hydrochloride, hydrobromide, sulphate, phosphate, maleate, acetate, citrate, benzoate, succinate, malate, ascorbate, tartrate and the like. If the active ingredient is to be administered in tablet form, the tablet may contain a binder such as tragacanth, corn starch or gelatin; a disintegrating agent, such as alginic acid; and a lubricant, such as magnesium stearate.

Routes of administration include, but are not limited to, respiratorally, intratracheally, nasopharyngeally, intravenously, intraperitoneally, subcutaneously, intracranially, intradermally, intramuscularly, intraocularly, intrathecally, intracereberally, intranasally, infusion, orally, rectally, *via* IV drip patch and implant.

In another aspect the present invention relates to the use of an agent capable of modulating the functional activity of sphingosine kinase in the manufacture of a medicament for the

- 27 -

modulation of cell growth in a mammal wherein down-regulation of the functional activity of said sphingosine kinase down-regulates the subject cell growth and up-regulation of the functional activity of said sphingosine kinase up-regulates the subject cell growth.

- 5 Preferably said cell growth is proliferation. Even more preferably said proliferation is uncontrolled cell proliferation which is down-regulated.

Preferably said uncontrolled cell proliferation is caused by the transformation of the cell by oncogene up-regulation or by sphingosine kinase overexpression oncogenic activity.

10

Still more preferably said cell is a malignant cell which forms a solid tumour of the colon, stomach, lung, brain, bone, oesophagus, pancreas, breast, ovary or uterus.

- 15 Yet another aspect relates to agents for use in modulating the functional activity of sphingosine kinase wherein modulating the functional activity of sphingosine kinase modulates cell growth.

Preferably said functional activity is down-regulated thereby down-regulating cell growth.

- 20 In another preferred embodiment said sphingosine kinase functional activity is up-regulated thereby up-regulating cell growth.

Even more preferably said cell growth is cell proliferation.

- 25 Still more preferably said cell is a neoplastic cell and said proliferation is down-regulated.

Preferably said cell proliferation is caused by the transformation of the cell by oncogene up-regulation or by sphingosine kinase overexpression oncogenic activity.

- 30 Still more preferably said cell is a malignant cell which forms a solid tumour of the colon, stomach, lung, brain, bone, oesophagus, pancreas, breast, ovary or uterus.

- 28 -

In yet another aspect the present invention relates to a pharmaceutical composition comprising a modulatory agent as hereinbefore defined and one or more pharmaceutically acceptable carriers and/or diluents. Said modulatory agents are referred to as the active
5 ingredients.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion or may be in the form of a cream or
10 other form suitable for topical application. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol and liquid polyethylene glycol, and the like), suitable mixtures thereof, and
15 vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The preventions of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal and the like. In many cases, it will be
20 preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

25 Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilisation. Generally, dispersions are prepared by incorporating the various sterilised active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those
30 enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze-drying

- 29 -

technique which yield a powder of the active ingredient plus any additional desired ingredient from previously sterile-filtered solution thereof.

When the active ingredients are suitably protected they may be orally administered, for
5 example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed
in hard or soft shell gelatin capsule, or it may be compressed into tablets, or it may be
incorporated directly with the food of the diet. For oral therapeutic administration, the
active compound may be incorporated with excipients and used in the form of ingestible
tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like.
10 Such compositions and preparations should contain at least 1% by weight of active
compound. The percentage of the compositions and preparations may, of course, be varied
and may conveniently be between about 5 to about 80% of the weight of the unit. The
amount of active compound in such therapeutically useful compositions in such that a
suitable dosage will be obtained. Preferred compositions or preparations according to the
15 present invention are prepared so that an oral dosage unit form contains between about 0.1
 μg and 2000 mg of active compound.

The tablets, troches, pills, capsules and the like may also contain the components as listed
hereafter: a binder such as gum, acacia, corn starch or gelatin; excipients such as dicalcium
20 phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the
like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose,
lactose or saccharin may be added or a flavouring agent such as peppermint, oil of
wintergreen, or cherry flavouring. When the dosage unit form is a capsule, it may contain,
in addition to materials of the above type, a liquid carrier. Various other materials may be
25 present as coatings or to otherwise modify the physical form of the dosage unit. For
instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or
elixir may contain the active compound, sucrose as a sweetening agent, methyl and
propylparabens as preservatives, a dye and flavouring such as cherry or orange flavour. Of
course, any material used in preparing any dosage unit form should be pharmaceutically
30 pure and substantially non-toxic in the amounts employed. In addition, the active
compound(s) may be incorporated into sustained-release preparations and formulations.

- 30 -

The pharmaceutical composition may also comprise genetic molecules such as a vector capable of transfecting target cells where the vector carries a nucleic acid molecule encoding a modulatory agent. The vector may, for example, be a viral vector.

5

Screening for the modulatory agents hereinbefore defined can be achieved by any one of several suitable methods known to those of skill in the art including, but in no way limited to, contacting a cell comprising the sphingosine kinase gene or functional equivalent or derivative thereof with an agent and screening for the modulation of sphingosine kinase
10 protein production or functional activity, modulation of the expression of a nucleic acid molecule encoding sphingosine kinase or modulation of the activity or expression of a downstream sphingosine kinase cellular target. Detecting such modulation can be achieved utilising techniques such as Western blotting, electrophoretic mobility shift assays and/or the readout of reporters of sphingosine kinase activity.

15

It should be understood that the sphingosine kinase gene or functional equivalent or derivative thereof may be naturally occurring in the cell which is the subject of testing or it may have been transfected into a host cell for the purpose of testing. Further, the naturally occurring or transfected gene may be constitutively expressed - thereby providing a model
20 useful for, inter alia, screening for agents which down regulate sphingosine kinase activity, at either the nucleic acid or expression product levels, or the gene may require activation - thereby providing a model useful for, inter alia, screening for agents which up regulate sphingosine kinase expression. Further, to the extent that a sphingosine kinase nucleic acid molecule is transfected into a cell, that molecule may comprise the entire sphingosine
25 kinase gene or it may merely comprise a portion of the gene such as the portion which regulates expression of the sphingosine kinase product. For example, the sphingosine kinase promoter region may be transfected into the cell which is the subject of testing. In this regard, where only the promoter is utilised, detecting modulation of the activity of the promoter can be achieved, for example, by ligating the promoter to a reporter gene. For
30 example, the promoter may be ligated to luciferase or a CAT reporter, the modulation of

- 31 -

expression of which gene can be detected via modulation of fluorescence intensity or CAT reporter activity, respectively.

In another example, the subject of detection could be a downstream sphingosine kinase regulatory target, rather than sphingosine kinase itself. Such detection may be achieved
5 utilising technologies such as microarrays (e.g. Chip arrays, nylon arrays), SAGE analysis, RDA or Differential Display.

These methods provide a mechanism for performing high throughput screening of putative
10 modulatory agents such as the proteinaceous or non-proteinaceous agents comprising synthetic, combinatorial, chemical and natural libraries. These methods will also facilitate the detection of agents which bind either the sphingosine kinase nucleic acid molecule or expression product itself or which modulate the expression of an upstream molecule, which upstream molecule subsequently modulates sphingosine kinase expression or
15 expression product activity. Accordingly, these methods provide a mechanism of detecting agents which either directly or indirectly modulate sphingosine kinase expression and/or activity.

As detailed earlier, reference to "sphingosine kinase" should be understood as a reference
20 to either the sphingosine kinase expression product or to a nucleic acid molecule encoding sphingosine kinase. It should also be understood as a reference to a portion or fragment of the sphingosine kinase molecule such as the regulatory region of the sphingosine kinase nucleic acid molecule. Alternatively, the molecule may comprise the binding/active portion of the expression product. In this regard, the sphingosine kinase nucleic acid
25 molecule and/or expression product is expressed in a cell. The cell may be a host cell which has been transfected with the sphingosine kinase nucleic acid molecule or it may be a cell which naturally contains the sphingosine kinase gene.

The screening method herein defined may be based on detecting an "altered expression
30 phenotype associated with said sphingosine kinase". This should be understood as the detection of cellular or cell culture condition changes associated with modulation of the

- 32 -

activity of sphingosine kinase. These may be detectable for example as intracellular changes or changes observable extracellularly. For example, this includes, but is not limited to, detecting changes in expression product levels, cell culture condition changes or, to the extent that the sphingosine kinase regulatory region is ligated to a reporter molecule such as luciferase or CAT, detecting changes in reporter molecule expression. 5 Alternatively, this screening system may be established to detect changes in the expression of downstream molecules which are regulated by the sphingosine kinase expression product.

10 Yet another aspect of the present invention relates to modulatory agents, as hereinbefore defined, when used in the method of the present invention.

Still yet another aspect of the present invention relates to diagnostic methodology based on screening individuals for the presence of sphingosine kinase or mRNA or protein or the 15 specific forms of sphingosine kinase which are transcribed and/or translated by a given population of cells. The screening methodology may be directed to qualitative and/or quantitative sphingosine kinase analysis. This is particularly useful, for example, for determining whether a given individual is predisposed to or resistant to diseases/disorders in which aberrant, unwanted or otherwise inappropriate cell growth is a component of the 20 disease state and/or is predisposed or resistant to the development of certain forms of aberrant, unwanted or otherwise inappropriate cell growth. Such screening can be performed utilising methods which would be known to those skilled in the art including, but not limited to:

25 (i) The use of sphingosine kinase assays to screen for altered sphingosine kinase activity or levels. These parameters can be screened for either in the bodily fluids or tissues of individuals. Although sphingosine kinase is generally not secreted, its down-stream product sphingosine kinase-1-phosphate is secreted. Modulation of the levels of this molecule could therefore be used as an indicator of changes in 30 sphingosine kinase levels or activity. Nevertheless, screening for extracellular

- 33 -

sphingosine kinase should not be excluded.

- (ii) Analysis of sphingosine kinase for possible mutations/polymorphisms (SNPs) and mutations can be performed by melting curve analysis of PCR generated DNA
5 using the LightCycler system (Roche).
- (iii) The screening of SNPs involving sphingosine kinase using Chip Arrays (e.g. Affimetrix GeneChipSNP mapping or using the Incyte technology platform *fSSCP* for SNP discovery.
10

The present invention is further described by the following non-limiting examples.

- 34 -

EXAMPLE 1**TRANSFECTION AND ANALYSIS OF NIH 3T3 FIBROBLASTS**

5 To investigate the oncogenic role of SphK, the nontransformed NIH 3T3 fibroblasts were transfected with human SphK cDNA that was recently cloned in our laboratory (Piston *et al.*, 2000). Pooled stable transfectants (referred as SK-3T3) were used to avoid the phenotypic artifacts that may due to the selection and propagation of individual clones from single transfected cells. In the SK-3T3 cell pools SphK activity was increased by
10 over 600-fold (Fig. 1a) in comparison with the empty vector-transfected NIH 3T3 cells (N-3T3). Immunoblotting analysis showed a specific protein band with an apparent molecular weight consistent with the predicted size of FLAG-tagged human SphK that was detected only in the SK-3T3 cell pools but absent in N-3T3 cells (Fig. 1c). Intracellular levels of S1P, the direct product of SphK, were also increased in SK-3T3 cells by 4- to 5-fold,
15 indicating the stable transfectants with constitutively activation of SphK (Fig. 1b). S1P levels were not directly proportional to the increase in SphK activity assayed *in vitro* perhaps due to rapid degradation of S1P by S1P phosphatase and S1P lyase and the limited availability of sphingosine as a substrate for S1P formation.

20 Growth curves showed a significant difference between the SK-3T3 cell pools and the controls (Fig. 2a). Stable expression of SphK dramatically enhanced cell growth in the media containing either 1% or 10% serum. Even in serum-free medium for up to 7 days, SK-3T3 cells survived and grew whilst the N-3T3 cells underwent death. Furthermore, when the cells reached saturation density SK-3T3 cells continued to proliferate (Fig. 2b)
25 suggesting an escape from contact inhibition. Treatment of SK-3T3 cells with a specific inhibitor of SphK, *N,N*-dimethylsphingosine (DMS), significantly diminished the enhanced proliferation induced by overexpression of SphK, whilst DMS had no effect on proliferation of N-3T3 cells (Fig. 2c). These results indicate that the constitutive activation of SphK in the stably SphK-transfected cells reduces two key growth limiting properties:
30 serum dependence and contact inhibition.

- 35 -

The transforming activity assayed by focus formation in NIH 3T3 cells showed that cells transfected with SphK but not empty vector induced numerous foci (Table 1 and Fig. 3a). Both SphK and control vectors displayed similar efficiency in the generation of G418-resistant colonies indicating that the transforming activity was not due to non-specific effect of transfection. Furthermore, SK-3T3 cells formed vigorous colonies in soft agar (Table 2 and Fig. 3b) revealing the acquisition of anchorage-independent growth. Although N-3T3 cells exhibited a background level of colony formation that may due to spontaneous transformation, overexpression of SphK resulted in a 20~50-fold increase in the number of colonies and an obvious increase in colony size (Table 2). Importantly, DMS inhibited the transforming capacity of SphK in a dose-dependent manner with 2.5 μ M DMS resulting in total reversion to the normal phenotype (Fig. 3b). This suggested that the increased activity of SphK rather than its mere overexpression is responsible for the transforming capacity of this enzyme.

15

EXAMPLE 2

TRANSFECTION AND ANALYSIS OF ONCOGENE TRANSFORMED CELLS

When NIH 3T3 cells were transfected with an activated mutant Ras (V12-Ras), SphK activity was significantly increased by $178 \pm 22\%$ in comparison to the parent cells (Fig. 3c). By contrast the cells transfected with v-Src (Fig. 3c) or dominant-negative Ras (N17-Ras) had no changes in the activity of SphK, suggesting a specific involvement of SphK in some oncogenic transformation. Moreover, when the cells were treated with DMS, the focus formation was reduced by $42 \pm 4\%$ in V12-Ras transformed cells but there were no changes in v-Src transformed cells (Fig. 3d), indicating an important role of SphK in Ras transformation. The likelihood that there are SphK independent pathways operating is suggested by the partial effect of DMS at the dose (2.5 μ M) that was fully effective in SphK transformed cells (Fig 3d). On the other hand, the inability of DMS to inhibit v-Src transformation rules out non-specific effects of DMS or a general toxicity resulting from inhibition of SphK. The increased SphK activity thereby exerts a transforming potential not only in its own right by overexpression but is also involved in oncogenic transformation, eg., induced by Ras.

- 36 -

EXAMPLE 3

ANALYSIS OF TUMORIGENICITY

5 Tumorigenicity was then directly tested in NIH 3T3 cells overexpressing SphK. When the SphK-transfected NIH 3T3 cells from either the stable transfectant pools or selected clones were injected subcutaneously into NOD/SCID mice, tumours became apparent at the site of injection within 3 to 4 weeks (Table 4 and Fig. 4). No mice injected with the vector-transfected 3T3 cells induced tumours during 10-weeks of observation. Histological
10 appearances of tumour sections displayed the morphologies of fibrosarcoma with many mitotic figures (Fig. 4). Western blot analysis of extracts derived from tumours showed high levels of FLAG-tagged protein (Fig. 4), revealing that the neoplastic cells retain and express the SphK transgenes. Thus, the tumours were developed from the injected SphK-transfected cells but not from spontaneously transformed NIH 3T3. This is the first
15 demonstration that a wild type lipid kinase gene, human SphK, acts as an oncogene providing a potential linkage between this enzyme and mammalian tumour pathogenesis. Thus, our finding extends the understanding of phospholipids, particularly sphingolipids, as signal transducers regulating cellular growth, transformation and oncogenesis.

20

EXAMPLE 4

MATERIALS AND METHODS

(a) Sphingosine kinase transfection

25 NIH 3T3 fibroblasts were obtained from the American Type Culture Collection and maintained in DMEM (GIBCO BRL), supplemented with 10% calf serum. The human SphK cDNA tagged with a FLAG tag was subcloned into the pcDNA3 plasmid (Invitrogen Corp.) as previous described (Pitson *et al.*, 2000). For transient transfections, 5 µg of plasmids were transfected to 5×10^5 using Lipofectamine Plus (GIBCO BRL) according to
30 the manufacturer's protocols. For stable expression, the calcium phosphate precipitation method was used and the transfectants were selected in medium containing 500 µg/ml

- 37 -

G418 (GIBCO BRL). The nonclonal pools of G418-resistant transfected cells were collected and used to avoid clonal variability. For some experiments, the selected clones of stable transfectants were used.

5 **(b) Measurement of sphingosine kinase activity**

SphK activity was measured by incubating the cytosolic fraction with 10 μ M sphingosine dissolved in 5% Triton X-100 and [γ - 32 P]ATP (1mM, 0.5mCi/ml) for 15 min at 37°C as described previously (Xia *et al.*, 1999). For assay of intracellular level of S1P, cells were
10 labelled with [3 H]sphingosine (1 μ M, 2 μ Ci/ml) for 30 min and radioactivity incorporated into cellular lipids were extracted. [3 H]S1P was then resolved on TLC with 1-butanol/methanol/acetic acid/water (8:2:1:2, vol/vol), visualized and quantified by Phosphoimager®.

15 **(c) Culture of sphingosine kinase-3T3 cells**

Stably transfected NIH 3T3 cells were plated in 48-well plates (1,000 cells per well) in DMEM containing 10% calf serum. After 8 h, cells were washed twice with DMEM and then grown in DMEM containing 1 or 10% serum and serum-free medium (DMEM
20 containing 0.1% BSA). At the indicated times, cells were incubated with 1 mg/ml MTT for 4 h. The formazan product was solubilized by 10% in SDS in 10 mM HCl and assessed by spectrophotometry at 570 nm and 650 nm absorbance. In some experiments, cells were trypsinized and counted in a hemocytometer.

25 **(d) Focus formation assay**

For focus formation assay, low passage NIH 3T3 cells were transfected with the activated V12 Ras, v-Src (gifts of Dr. Julian Downward), SphK, or empty vector, respectively, using Lipofectamine Plus as described in (a) above. Two days later, the transfected cells were
30 split to 6-well plates. After reaching confluence, they were kept for two weeks in DMEM containing 5% calf serum. The foci were visualized and scored after stained with 0.5%

- 38 -

crystal violet. For soft agar assay, suspensions of 1×10^4 cells from the stable transfected pools in a growth medium containing 0.33% agar were overlaid onto 0.6% agar gel in the absence or presence of DMS at various concentrations. After a 14-day incubation colonies were stained with 0.1 mg/ml MTT and that over 0.1 mm in diameter were scored as
5 positive.

(e) Tumour induction

NOD/SCID mice were bred and maintained under sterile conditions. Four- to six-week-
10 old mice were injected subcutaneously with 5×10^5 cells in 200 μ l sterile PBS from various cell lines (stable SphK- and vector-transfected NIH 3T3 cell pools, two colonies of SphK transfectants). Each cell line was tested in 3 different animals.

Those skilled in the art will appreciate that the invention described herein is susceptible to
15 variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or more of said steps or features.

20

- 39 -

Table 2. Transformation assays in transfected NIH 3T3 cells.

Cell Line	<u>Focus formation</u>	<u>Colonies in soft agar</u>	
	Number	Number	Size (mm)
N-3T3	1.7±1.5	3.3±2.1	<0.1
SK-3T3	65.7±9.6	122.3±17.6	0.1~0.45

5

Focux formation was assayed in NIH 3T3 cells transiently transfected with SphK (SK-3T3) or vector (N-3T3). Transfected cells were plated in 6-well plates and cultured for 2~3 weeks prior to crystal violet staining. Colony formation in soft agar was determined in the stable transfected cells. Results shown are the mean ± SD from 3~5 experiments

10 done in duplicate or triplicate.

- 40 -

Table 3. Tumourigenesis in NOD/SCID mice.

Cell lines	Tumours/ injections	Tumour size	
		cm	cm ³
N-3T3	0/3	0	
SK-3T3 stable pools	3/3	2.8 x 1.8 x 1.1	5.54
		1.9 x 1.5 x 1.0	2.85
		1.1 x 0.9 x 0.8	0.79
Clone KT-2	3/3	3.2 x 1.8. x 1.2	6.91
		1.6 x 1.2 x 0.5	0.96
		2.5 x 1.6 x 1.3	5.20
Clone KT-5	3/3	2.7 x 1.6 x 1.2	5.18
		2.2 x 1.5 x 1.1	3.63
		1.8 x 1.7 x 0.9	2.75

- 5 NOD/SCID mice were injected with cells (5×10^5 cells per mouse) from vector- or SphK-transfected NIH 3T3 cell pools (N-3T3 or SK-3T3), or two individual SphK-transfected clones (KT-2 and KT-5). Tumour size was determined 4 weeks after injection.

- 41 -

CLAIMS:

1. A method of modulating the growth of a cell said method comprising contacting said cell with an effective amount of an agent for a time and under conditions sufficient to modulate the functional activity of sphingosine kinase.
2. A method according to claim 1 wherein said functional activity is the level of functional activity.
3. The method according to claim 1 or 2 wherein said growth is proliferation.
4. The method according to claim 3 wherein said modulation of proliferation is down-regulation of proliferation and said modulation of functional activity is down-regulation of functional activity.
5. The method according to claim 3 wherein said modulation of proliferation is up-regulation of proliferation and said modulation of functional activity is up-regulation of functional activity.
6. The method according to claim 4 wherein said proliferation is uncontrolled proliferation.
7. The method according to claim 6 wherein said cell is a neoplastic cell.
8. The method according to claim 7 wherein said neoplastic cell is a malignant cell.
9. The method according to claim 8 wherein said malignant cell is a cell from the colon, stomach, lung, brain, bone, oesophagus, pancreas, breast, ovary or uterus.
10. The method according to claim 9 wherein said malignant cell has become transfected due to up-regulation of an oncogene.

- 42 -

The method according to claim 10 wherein said oncogene is Ras.

The method according to claim 9 wherein said malignant cell has become transformed by sphingosine kinase overexpression oncogenic activity.

The method according to any one of claims 1-4 or 6-12 wherein said agent is N,N-dimethylsphingosine.

The method according to any one of claims 1-4 or 6-12 wherein said agent is DL-threo-dihydrophingosine.

A method for the treatment and/or prophylaxis of a condition characterized by aberrant, unwanted or otherwise inappropriate cell growth in a mammal, said method comprising administering to said mammal an effective amount of an agent for a time and under conditions sufficient to modulate the functional activity of sphingosine kinase.

A method according to claim 15 wherein said functional activity is the level of functional activity.

The method according to claim 15 or 16 wherein said growth is proliferation.

The method according to claim 17 wherein said modulation of proliferation is down-regulation of proliferation and said modulation of functional activity is down-regulation of functional activity.

The method according to claim 17 wherein said modulation of proliferation is up-regulation of proliferation and said modulation of functional activity is up-regulation of functional activity

- 43 -

20. The method according to claim 18 wherein said proliferation is uncontrolled proliferation.
21. The method according to claim 20 wherein said cell is a neoplastic cell.
22. The method according to claim 21 wherein said neoplastic cell is a malignant cell.
23. The method according to claim 22 wherein said malignant cell forms a solid tumour of the colon, stomach, lung, brain, bone, oesophagus or pancreas.
24. The method according to claim 23 wherein said malignant cell has become transformed due to oncogene up-regulation.
25. The method according to claim 24 wherein said oncogene is Ras.
26. The method according to claim 23 wherein said malignant cell has become transformed by spingosine kinase overexpression oncogenic activity.
27. The method according to any one of claims 15-18 or 20-26 wherein said agent is N,N-dimethylsphingosine.
28. The method according to any one of claims 15-18 or 20-26 wherein said agent is DL-threo-dihydrophingosine.
29. The method according to any one of claims 15-28 wherein said mammal is a human.
30. Use of an agent capable of modulating the functional activity of sphingosine kinase in the manufacture of a medicament for the modulation of cell growth in a mammal.

- 44 -

31. Use according to claim 30 wherein said functional activity is the level of functional activity.
32. Use according to claim 30 or 31 wherein said growth is proliferation.
33. Use according to claim 32 wherein said modulation of proliferation is down-regulation of proliferation and said modulation of functional activity is down-regulation of functional activity.
34. Use according to claim 32 wherein said modulation of proliferation is up-regulation of proliferation and said modulation of functional activity is up-regulation of functional activity.
35. Use according to claim 33 wherein said proliferation is uncontrolled proliferation.
36. Use according to claim 35 wherein said cell is a neoplastic cell.
37. Use according to claim 36 wherein said neoplastic cell is a malignant cell.
38. Use according to claim 37 wherein said malignant cell forms a solid tumour of the colon, stomach, lung, brain, bone, oesophagus, pancreas, breast, ovary or uterus.
39. Use according to claim 38 wherein said malignant cell has become transformed due to oncogene up-regulation.
40. Use according to claim 39 wherein said oncogene is Ras.
41. Use according to claim 38 wherein said malignant cell has become transformed by spingosine kinase overexpression oncogenic activity:

- 45 -

42. Use according to claims 30-33 or 35-41 wherein said agent is N,N-dimethylsphingosine.
43. Use according to claims 30-33 or 35-41 wherein said agent is DL-threo-dihydrophingosine.
44. Use according to claims 30-43 wherein said mammal is a human.
45. A pharmaceutical composition comprising an agent capable of modulating the functional activity of sphingosine kinase together with one or more pharmaceutically acceptable carriers and/or diluents for use in accordance with the method of any one of claims 1-44.
46. The pharmaceutical composition according to claim 45 wherein said agent is N,N-dimethylsphingosine.
47. The pharmaceutical composition according to claim 45 wherein said agent is DL-threo-dihydrophingosine.
48. A method of diagnosing a condition, or a predisposition or resistance to a condition, characterized by aberrant, unwanted or otherwise inappropriate cell growth in a mammal, said method comprising screening a biological sample from said mammal for the presence of sphingosine kinase or nucleic acid molecule encoding sphingosine kinase.

- 46 -

BIBLIOGRAPHY

Allessenko, A.V., (1998) *Biochemistry (Mosc)* 63:62-68.

Culliver, O., Pirianov, G., Kleuser, B., Vanek, P.G., Coso, O.A., Gutkind, J.S. and Spiegel, S. (1996) *Nature* 381:800-803.

Igarashi, Y. (1997) *J. Biochem.* 122:1080-1087.

Meyer zu Heringdorf, D., van Koppen, C.J. and Jakobs, K.H. (1997) *FEBS Lett* 410:34-38.

Pitson, S.M. *et al.* (2000) *Biochem. J.* submitted.

Spiegel, S., Culliver, O., Edsall, L., Kohama, T., Menzeleev, R., Olivera, A., Thomas, D., Tu, Z., Van Brocklyn, J. and Wang, F. (1998) *Biochemistry (Mosc)* 63:69-73.


Xia, P., Gamble, J.R., Rye, K.-A., Wang, L., Hii, C.S.T., Cockerill, P., Khew-Goodall, Y., Bert, A.G., Barter, P.J., and Vadas, M.A. (1998) *Proc. Natl. Acad. Sci. USA* 95:14196-14201.

Xia, P., Vadas, M.A., Rye, K.A., Barter, P.J., Gamble, J.R. (1999) *J. Biol. Chem.* 274:33143-33147.


1/6

Figure 1


A.



B.




C.





2/6

Figure 2

A.




3/6



Figure 2 (CONT'D)**B.****C.**

4/6

Figure 3




5/6

Figure 3 (CONT'D)**C.****D.**


6/6

Figure 4

A.



B.



C.

