

659836


AUSTRALIA
PATENTS ACT 1990
NOTICE OF ENTITLEMENT

We, **Laboratorios Beecham SA**, the applicant/Nominated Person in respect of
Application No. 16498/92 state the following:-

The Nominated Person is entitled to the grant of the patent
because the Nominated Person would, on the grant of a patent for
the invention to the inventors, be entitled to have the patent
assigned to the Nominated Person.

The Nominated Person is entitled to claim priority from the
application listed in the declaration under Article 8 of the PCT
because the Nominated Person made the application listed in the
declaration under Article 8 of the PCT, and because that
application was the first application made in a Convention country
in respect of the invention.

DATED this TWENTY SECOND day of OCTOBER 1993



.....
a member of the firm of
DAVIES COLLISON
CAVE for and on behalf
of the applicant(s)

(DCC ref: 1622275)



AU9216498

(12) PATENT ABRIDGMENT (11) Document No. AU-B-16498/92
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 659836

- (54) Title
PHARMACEUTICAL FORMULATIONS
- International Patent Classification(s)
(51)⁵ A61K 009/20 A61K 031/43
- (21) Application No. : 16498/92 (22) Application Date : 02.05.92
- (87) PCT Publication Number : WO92/19227
- (30) Priority Data
- | | | |
|-------------|-----------|-------------------|
| (31) Number | (32) Date | (33) Country |
| 9109862 | 08.05.91 | GB UNITED KINGDOM |
- (43) Publication Date : 21.12.92
- (44) Publication Date of Accepted Application : 01.06.95
- (71) Applicant(s)
LABORATORIOS BEECHAM SA
- (72) Inventor(s)
LUIS CARVAJAL MARTIN; JUAN DEDIOS ROMERO
- (74) Attorney or Agent
DAVIES COLLISON CAVE , 1 Little Collins Street, MELBOURNE VIC 3000
- (56) Prior Art Documents
GB 2005538
EP 281200
- (57) Claim

1. A tablet formulation having a structure comprising compacted granulates; the granulates comprising at least one compacted medicament optionally together with an intra-granular disintegrant; the medicament being a β -lactam antibiotic optionally in combination with a β -lactamase inhibitor and the granulates being compacted together into a tablet form together with an extra-granular disintegrant and optionally also together with an extra-granular lubricant, provided that if a lubricant is present the amount of lubricant is less than 0.5% by weight of the total tablet.

10. A pharmaceutical granulate formulation comprising a medicament which is a β -lactam antibiotic optionally in combination with a β -lactamase inhibitor and an intra-granular disintegrant which is CLPVP.

PCT

ANNOUNCEMENT OF THE LATER PUBLICATION
OF INTERNATIONAL SEARCH REPORT

16498/92.

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 9/20	A3	(11) International Publication Number: WO 92/19227 (43) International Publication Date: 12 November 1992 (12.11.92)
(21) International Application Number: PCT/EP92/01024 (22) International Filing Date: 2 May 1992 (02.05.92) (30) Priority data: 9109862.4 8 May 1991 (08.05.91) GB (71) Applicant (for MG only): SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). (71) Applicant (for all designated States except US): LABORA- TORIOS BEECHAM SA [ES/ES]; Costa Brava, 14, 28034 Madrid (ES). (72) Inventors; and (75) Inventors/Applicants (for US only) : MARTIN, Luis, Carva- jal [ES/ES]; Ronda de Buenavista, 26, E-45007 Toledo (ES). ROMERO, Juan, Dedios [ES/ES]; SB Pharma- ceuticals, Poligono Industrial, E-45007 Toledo (ES).		(74) Agent: WALKER, Ralph, Francis; SmithKline Beecham, Corporate Patents, Great Burgh, Yew Tree Bottom Road, Epsom, Surrey KT18 5XQ (GB). (81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE, DE (European patent), DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (Eu- ropean patent), GN (OAPI patent), GR (European pa- tent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC (European patent), MG, ML (OAPI patent), MR (OAPI patent), MW, NL, NL (Euro- pean patent), NO, PL, RO, RU, SD, SE, SE (European patent), SN (OAPI patent), TD (OAPI patent), TG (OA- PI patent), US. Published With international search report Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments. (88) Date of publication of the international search report: 23 December 1992 (23.12.92)
(54) Title: PHARMACEUTICAL FORMULATIONS (57) Abstract Tablet formulations having a structure comprising compacted granulates of a mixture of a medicament and an intra-granular disintegrant, the granulates being compacted together into a tablet with an extra-granular disintegrant and optional extra-granular lubricant and excipients.		

PHARMACEUTICAL FORMULATIONS

The present invention relates to pharmaceutical formulations for oral administration in the treatment of bacterial infections, and to processes
5 for the manufacture of such formulations.

It is known to provide formulations for oral administration in the form of water-dispersible granules or tablets which may be swallowed or dispersed in water prior to swallowing.

10

In one known method of tablet manufacture, an intermediate granulate is prepared comprising an intragranular disintegrant and an active material such as an antibiotic. This granulate is then mixed with an intergranular disintegrant (and optional other additives including a lubricant) and
15 compressed into tablets. Such a process, tablets and granulate are for example described in EP 0081200A, CA 1199871 and JP 3240023A.

It is desirable that such solid formulations should rapidly disperse on immersion in water, for example by rapid disintegration of tablets.

20

Novel formulations have now been discovered which assist in achieving some of the above-mentioned desirable features.

The invention therefore provides a tablet formulation having a structure comprising
25 compacted granulates; the granulates comprising at least one compacted medicament optionally together with an intra-granular disintegrant; the medicament being a β -lactam antibiotic optionally in combination with a β -lactamase inhibitor and the granulates being compacted together into a tablet form together with an extra-granular disintegrant and optionally also together with an extra-granular lubricant, provided that if a lubricant is
30 present the amount of lubricant is less than 0.5% by weight of the total tablet.

In the tablets of this invention the granulates may be in a crushed state resulting from the compaction of the tablet, and consequently may not have discrete boundaries, or may be sub-divided or broken up into smaller
35 granulates. The invention is intended to include tablets having such a structure containing crushed granulates. Preferably the size of the granulates is in the range 100 μ m to 2mm, suitably around 1mm \pm 0.25mm, maximum dimension.



The medicament is ~~preferably one which is capable of oral absorption, in particular~~ β -lactam antibiotics optionally in combination with a β -lactamase inhibitor. A preferred antibiotic is amoxycillin, for example
5 present as a hydrate such as the trihydrate. Amoxycillin may be used alone, or may optionally be used in combination with other β -lactam antibiotics and/or β -lactamase inhibitors such as clavulanic acid or salts (especially the potassium salt) thereof, for example in a weight ratio equivalent to amoxycillin: clavulanic acid in the range 12:1 to 1:1 such as
10 around 4:1 or 2:1. Preferably the proportion of the antibiotic in the tablet is 60-98% by weight of the total tablet, in the case of amoxycillin trihydrate calculated as the weight of the trihydrate. Preferably the particles of antibiotic in the granulates are in the size range 1 μ m to 300 μ m, especially 10 μ m to 200 μ m. A typical suitable size distribution of
15 the antibiotic particles is : >200 μ 5% or less, 200-100 μ 5-15%, 100-50 μ 7.5-15%, <50 μ 70% or more.

Suitable intra-granular disintegrants are starches, such as maize starch and rice starch, cross-linked N-vinyl-2-pyrrolidone ("CLPVP"), sodium
20 starch glycollate, croscarmellose sodium and formaldehyde - casein, or combinations thereof. A preferred intra-granular disintegrant is CLPVP, for example as marketed under the trade names Polyplasdone XL and Polyplasdone XL-10.

25 The granulate may consist entirely of antibiotic(s), optionally in the case of a β -lactam antibiotic combined with a β -lactamase inhibitor, and an intra-granular disintegrant. Alternatively, particularly when the granulate contains clavulanic acid or a salt thereof, the granulate may also contain a diluent such as silica gel (eg Syloid-Trade Mark). Suitable
30 intra-granular disintegrants for use with antibiotics are CLPVP and sodium starch glycollate. Typically the proportion of intra-granular disintegrant in the granulate may be 0.1 - 10wt % of the granulate, suitably 1.0 - 8.0wt %, such as 1.25 - 3.5wt %. Typically the proportion of an antibiotic or antibiotic + β -lactamase inhibitor combination in the
35 granulate may be 99.9 - 90wt %, suitably 99 - 92wt %, e.g. 99 - 95wt %, such as 98.75 - 96.5wt % of the weight of the granulate. When the granulate contains a diluent, this may comprise up to 30wt % of the granulate, but is conveniently present in a 1:1 weight ratio with the



- 3 -

amount of clavulanic acid or its salt in the granulate. When the granulate contains a diluent the granulate will contain a correspondingly lower proportion of antibiotic or antibiotic + β -lactamase inhibitor combination, for example 70 - 99.9wt % of the granulate.

5

The intimate contact between the antibiotic and the intra-granular disintegrant in the granulate appears to assist in improved disintegration and dispersion of the granulate in contact with water to release antibiotic particles in the size range referred to above, and to provide finely
10 dispersed suspensions. Problems are associated with preparation of granulates which include clavulanic acid salts, due to their hygroscopicity, and the granulate of the invention facilitates manufacture.

In the tablet formulation the granulate may suitably comprise 70% or
15 more, e.g. 80% or more, 90% or more or 95% or more of the total tablet weight so that a high proportion of medicament is present.

The extra-granular disintegrant may be a conventional disintegrant for example starches such as maize-starch and rice starch, CLPVP, sodium
20 starch glycollate, croscarmellose sodium, microcrystalline or microfine cellulose, low-substituted hydroxypropylcellulose (i.e. cellulose partially substituted with 2-hydroxypropyl groups, e.g. less than 25% substituted, preferably 7-16% substituted), cross-linked sodium
25 carboxymethylcellulose, swellable ion exchange resins, formaldehyde-casein, or alginates. Preferred extra-granular disintegrants are CLPVP, sodium starch glycollate, microfine cellulose and croscarmellose sodium, and combinations thereof. An example of an extra-granular disintegrant combination is a combination of
microcrystalline or microfine cellulose with sodium starch glycollate,
30 croscarmellose sodium, or CLPVP, containing 80-90% by weight cellulose.

The proportion of extra-granular disintegrant to total tablet weight may vary between broad limits, for example 0.1-25 weight %. For example if
35 CLPVP or sodium starch glycollate is used as extra-granular disintegrant it may suitably be used as such in a proportion 0.1-5.0 weight %, suitably 0.1 - 3.0 weight %, preferably 0.1-1.5 weight % of the total tablet weight. If cellulose or a combination containing cellulose is used, e.g. as described above containing around 80-90% by weight of cellulose, the extra-granular

disintegrant may comprise 1-25 weight %, typically around 1-20 weight % of the total tablet.

5 Suitable lubricants are those conventional to the art, such as long-chain fatty acids, such as stearic acid, or salts thereof, in particular Group II metal salts, such as of magnesium or calcium.

10 A preferred lubricant is magnesium stearate. It is preferred to use a lubricant proportion as low as possible e.g. 0.35% by weight or preferably lower, e.g. 0.275% or less, e.g. 0.25% or less, preferably using no lubricant at all.

15 The granulate may also contain an intra-granular lubricant, which may be selected from the same materials as the extra-granular lubricant, such as magnesium stearate. However an advantage of the present tablet formulation is that the granulate and tablet need not contain any lubricant. This can lead to improved wettability and hence improved disintegration of the tablet. Further a reduced lubricant proportion can lead to a lower tablet weight for a given dose of antibiotic and in the case of dispersible formulations can avoid the "smeared" appearance associated with higher lubricant proportions.

25 The tablet may also include conventional excipients, typically present up to about 10% of the total tablet weight. These may include flavouring agents, for example flavourings such as menthol, peppermint, vanilla or fruit flavourings, flavouring agents typically being present up to around 0.5-5% by weight of the whole tablet, and sweeteners, e.g. aspartame, present of up to around 15mg per unit dose. Excipients may also include colouring agents, preservatives, suspending aids and fillers such as silicon dioxide, microcrystalline cellulose, dicalcium phosphate, lactose, sorbitol, calcium carbonate or magnesium carbonate. Such excipients are preferably mixed with the extra-granular disintegrant and lubricant (if present). The materials present in the tablets should have low free moisture content and preferably be pre-dried. In some cases, particularly when the medicament is an antibiotic, and includes clavulanic acid or its salts, it may be necessary to include a desiccant diluent such as silica gel as an excipient, in a proportion of about 1-5% of the weight of the antibiotic, mixed with the antibiotic and intra-granular disintegrant in

35

- 5 -

the granulates. The particle size of the excipients does not appear to be critical but it is desirable to exclude agglomerates.

5 The tablet may also contain an effervescent couple of known type, e.g. a solid acid and an alkali metal carbonate or bicarbonate which generates carbon dioxide on contact with water to assist in disintegration of the tablet.

10 The tablets may be film coated in a conventional manner, e.g. for cosmetic, palatability or production purposes. Suitable coatings include hydroxypropylcellulose, acrylate and/or methacrylate co-polymers, resins etc. Alternatively the coating may be an enteric coating, e.g. which is insoluble in acidic gastric juice but soluble in alkaline digestive juice. Such a coating enables the antibiotic to pass through the stomach into the
15 duodenum, from where it is absorbed. Suitable enteric coatings include cellulose acetate phthalate.

Preferred combinations of components for the tablets of this aspect of the invention therefore comprise:

20

<u>Granulate:</u> <u>Component</u>	<u>wt%</u>	<u>Example</u>
Medicament	70 - 99	Amoxycillin \pm Pot.clavulanate
Disintegrant	0.1 - 4	CLPVP, Microcryst. cellulose, sodium starch glycollate
Diluent	0 - 30	Silica gel

<u>Tablet:</u> <u>Component</u>	<u>wt%</u>	<u>Example</u>
Granulate	70+	above
Disintegrant	0.1 - 25	CLPVP, Microcryst. cellulose, sodium starch glycollate.
Lubricant	0 - 0.35	Magnesium stearate
Excipients	to 100	Aspartame, flavour, colour, silicon dioxide

The invention also provides a process for the manufacture of a tablet in
 which granulates comprising a compacted mixture of at least one
 5 medicament ^{which is} such as a β -lactam antibiotic either alone or in combination
 with a β -lactamase inhibitor, together with an intra-granular disintegrant
 are mixed with an extra-granular disintegrant and optionally with an
 extra-granular lubricant and optionally with any excipients, provided that
 10 if a lubricant is present it amounts to less than 0.5% by weight of the
 mixture, and the mixture is compressed into tablets.

Suitable and preferred antibiotics, intra- and extra-granular
 disintegrants, lubricants, excipients, granulate and particle sizes, and
 relative proportions thereof are as discussed above.

15

The necessary granulate for the process of this aspect of the invention



may be made in a further process by mixing the medicament in a powdered form with the intra-granular disintegrant in a dry state, and compacting the mixture under pressure. Insofar as this further process uses as intra-granular disintegrant CLPVP, sodium starch glycollate, casein-formaldehyde, croscarmellose sodium or combinations thereof, it is believed to be novel, and is a further aspect of this invention.

In this further process, it is desirable to mill and sieve the antibiotic to achieve the desired particle size range. It is also desirable to mill and sieve intra-granular disintegrant to a suitable particle size, for example in the case of CLPVP about 30 μ , but particle size does not appear to be critical.

The compaction of the mixture into granulates may be by conventional dry compaction means, for example pressing, rolling, slugging extrusion etc, and a suitable pressure for the compaction process is 30-200KN, e.g. 35-65KN preferably 40-50 KN. The above-described granulate formulations are particularly suited to formation by roller compaction. It may be necessary to mill and sieve the compacted mixture after compaction so as to achieve a suitable size fraction of the granulate. Compression into tablets may be carried out in a conventional manner, e.g. on a conventional tableting machine. As an optional further step the tablets may be coated as described above.

When the granulates described above contain as a medicament a β -lactam antibiotic such as amoxycillin together in combination with a β -lactamase inhibitor such as clavulanic acid or its salts (especially potassium clavulanate) these granulates are believed to be novel and are a further aspect of this invention. Suitable and preferred features of these granules are as discussed above.

The granulates described above may be suitable for use in the preparation of other pharmaceutical formulations in addition to tablets, for example they may be supplied as a free-flowing granulated formulation in sachets containing a suitable unit dose. This may also for example be dissolved in water together with excipients such as sweetening agents, thickeners, preservatives and buffers such as sodium benzoate, sodium acetate and sodium citrate to form a syrup formulation, for example for administration

to small children.

5 The ability of the granulates to form a loose compact, and their rapid dispersion in contact with water makes them particularly suitable for use in encapsulated formulations. Therefore in a further aspect of this invention there is provided an encapsulated formulation comprising such granulates. The encapsulated formulation may optionally include an extra-granular lubricant, which if present is suitably in an amount of less than 0.5% by weight of the granulates, being contained within a
10 pharmaceutical capsule.

The medicament is ~~preferably one which is capable of oral absorption, in particular~~ a β -lactam antibiotic optionally in combination with a β -lactamase inhibitor. Suitable and preferred antibiotics, β -lactamase
15 inhibitors, intra-granular disintegrant, extra-granular lubricant, granulate and particle sizes, and relative proportions thereof for a capsule formulation are as discussed above, except that a preferred proportion of lubricant is 0.1-0.5%, particularly 0.32-0.35% by weight of the granulate.

20 The pharmaceutical capsule may be an entirely conventional capsule, capable of dissolving in the stomach to release its contents, for example made of gelatine.

The formulations described above preferably contain unit doses of
25 antibiotic, for example 375, 500, 750 or 1000mg of amoxycillin per tablet or capsule. The tablets may be dispersed in water prior to ingestion, or may alternatively be chewed or swallowed whole.

30 The invention further provides a pharmaceutical formulation as described above, for use as an active therapeutic substance.

The invention further provides a pharmaceutical formulation as described above, in which the medicament is a β -lactam antibiotic optionally in combination with a β -lactamase inhibitor, for use in the treatment of
35 bacterial infections.

The invention further provides a method of use of a pharmaceutical formulation as described above in which the medicament is a β -lactam



antibiotic optionally in combination with a β -lactamase inhibitor in the manufacture of a medicament, ~~for use in the treatment of bacterial infections.~~

- 5 The invention further provides a method of treatment of bacterial infections in mammals which comprises the administration to the mammal of an effective amount of a pharmaceutical formulation as described above, in which the medicament is a β -lactam antibiotic, optionally in combination with a β -lactamase inhibitor.

10

The invention will now be described by way of example only.

Example 1: Granulate.

- 15 Amoxycillin trihydrate was milled and sieved using an 0.04 or 0.027 inch (1.0 - 0.7 mm) aperture sieve, and was mixed for 15 minutes in a blender with dried cross-linked polyvinylpyrrolidone having a molecular weight of approximately 1 million and a density of 1.22 mg/cm³ (polyplasdone XL - Trade Mark), the mixture containing 3.4% of CLPVP by weight.

20

The mixture was consolidated using a roller compacter at a controlled pressure of 50KN. The compacted flakes were granulated in a mill, or granulated through a sieve fitted with a 1mm mesh to obtain a suitable size fraction.

25

Example 2: Tablet.

Tablets were prepared having the composition below;

<u>Component</u>	<u>Weight mg.</u>	<u>Weight %</u>
Amoxycillin trihydrate	750 ¹	78.95
CLPVP	26.0	2.73
Sodium Starch	21.6	2.27
Glycollate (Primogel)		
Magnesium Stearate	2.0	0.21
Aspartame	20.0	2.10

as granulate of
example 1

extra granulate



- 10 -

Microcrystalline	130.4	13.74
Cellulose (Avicel PH102)		

(1) Expressed or free acid equivalent:

- To prepare these tablets, the dried sodium starch glycollate, magnesium stearate and microcrystalline cellulose were sieved, then blended with the granulate of example 1. The aspartame was then added, and this mixture was then blended until homogeneous (5 minutes). The mixture was then compressed into tablets on a conventional tableting machine.

Example 3: Granulate.

10

A granulate was prepared using a procedure identical to example 1, comprising 97 weight % of amoxycillin trihydrate and 3 weight % polyplasdone XL, and using a controlled pressure of 40-50 KN.

Example 4: Tablet.

15

Tablets were prepared having the composition below:

Component	wt. mg	wt. mg	wt. mg	wt. mg	wt. %
Amoxycillin	375	500	750	1000	83.00 ¹
CLPVP	17.5	23.33	35	46.65	3.78 ²
Peppermint dry flavour	3	4	6	7.99	0.65
Aspartame	7.5	10	15	19.99	1.62
Magnesium stearate	1	1.34	2	2.67	0.21

- (1) As 95 wt. % of amoxycillin trihydrate.
(2) 3% as intra-granular, and 0.78% as extra-granular disintegrant.

- To prepare these tablets, the dried flavour, aspartame, magnesium stearate and a weight of CLPVP (polyplasdone XL) corresponding to 0.78 wt. % of the total weight of the mixture was mixed for 5 minutes with the granulate of example 3 to give the wt % indicated above. The mixture was then compressed into tablets on a conventional tableting machine.

Typical tablets of this example containing 750mg of amoxycillin as the trihydrate had the following characteristics:

weight	:	925mg \pm 5%
hardness	:	> 16 KP
time of dispersal	:	< 1 minute
in water		
friability	:	<1%
presentation	:	Oval, 17 x 10 x 7mm tablets

5

Example 5: Granulate

A granulate was prepared using a procedure identical to that of example 1, comprising 97.12 weight % amoxycillin trihydrate together with 2.88 weight % sodium starch glycollate (as "Primogel") as intra-granular disintegrant.

10

Example 6: Tablet

15 Tablets were prepared having the composition below:

<u>Component%</u>	<u>Weight mg.</u>	<u>Weight %</u>	
Amoxycillin	750mg ¹	78.95] as granulate of example 5
Sodium starch glycollate	21.6mg	2.27	
Magnesium stearate	2.0mg	0.21] extra granulate
Dried microcrystalline cellulose (Avicel PH102)	to 950mg	18.57	

(1) As free acid equivalent

20 To prepare these tablets, the granulate of example 5 was sieved using a 1mm sieve, and was then blended with appropriate quantities of the magnesium stearate (lubricant) and microcrystalline cellulose, mixing for 15 minutes. The mixture was then compacted to form tablets having the

following characteristics:

weight	:	950mg
hardness	:	12 - 16 KP
time of dispersal	:	10-15 seconds (37°C),
in water		20-25 seconds (20°C)

- 5 These tablets could be provided in the above-described uncoated state for dispersion in water prior to swallowing, or could be film coated for swallowing.

Example 7: Encapsulated Formulation

- 10 The granulate of example 3 was made up into a loose compact under gentle pressure together with an amount of magnesium stearate lubricant to total 0.34% by weight of the total compact. This loose compact was sealed into gelatin capsules containing the following mixture:

<u>Component</u>	<u>Weight mg.</u>	<u>Weight %</u>
Amoxycillin trihydrate:	573.91 ¹	96.8
CLPVP :	17	2.9
magnesium stearate :	2	0.34

(1) corresponds to 500mg
amoxycillin free acid

15

Example 8: Sachet Formulation

<u>Component</u>	<u>Weight mg.</u>	<u>Weight %</u>
Amoxycillin trihydrate		76.12] granulate
Potassium	2711.1	
clavulanate/syloid AL-1		
blend 1:1		
Polyplasdone XL dried		0.38]
Polyplasdone XL dried	13.5	
Lemon dry flavour	408.0	
		11.45] extra granular

- 13 -

Strawberry dry flavour	132.0	3.71
Peach dry flavour	102.0	2.86
Aspartame	45.0	1.26
Xantham Gum	150.0	4.21

Granules were in a manner identical to that of example 1, i.e. by milling and sieving of the granulate components, followed by roller compaction (50KN) and granulation. The granules could be made up into a mixture
5 suitable for a sachet presentation with the extra-granular excipients.

The granulate of this example could be supplied containing appropriate weights of amoxycillin/clavulanate in a sachet, and is also suitable for making up into syrup formulations. For example the weights listed may
10 be made up into 60ml to produce a 156.25mg/5ml syrup or double the listed weights may be made up into 60ml to produce a 312.5mg.5ml syrup. These syrups do not contain any added sugar.

Example 9: Granulate

15

<u>Component</u>	<u>Weight mg.</u>	<u>Weight %</u>
Amoxycillin trihydrate	581.4 ¹	64.0
Potassium clavulanate	152.4 ²	16.8
Syloid AL-1	152.4	16.8
Polyplasdone XL dried	22.0	2.42

as granulate

(1) corresponds to 500mg amoxycillin free acid.

(2) corresponds to 125mg free clavulanic acid.

20 Granules are prepared using this mixture in a manner identical to that of example 8. These granules are suitable for supply in a sachet, together with flavour and sucrose in the proportions listed below for the quantity of granules listed above per sachet:

Lemon dry flavour	136.0mg
Strawberry dry flavour	44.0mg
Peach dry flavour	34.0mg
Sucrose	to 3500mg

25

Sachets containing other weights of amoxycillin, e.g. 250 or 125mg could be made up using proportional amounts of the weights listed and made up to 1750mg total weight with sucrose.

5 Example 10: Tablet

<u>Component</u>	<u>Weight mg.</u>	<u>Weight %</u>
Amoxycillin trihydrate	581.4 ¹	61.2
Potassium clavulanate	152.4 ²	16.0
Syloid AL-1	152.4	16.0
Polyplasdone XL dried	17.4	1.83
Dry flavour (Peppermint or mandarin)	6.0	0.63
Polyplasdone XL dried	25.0	2.63
Aspartame	15.0	1.58
Colouring	5.0	0.53
Magnesium stearate	2.5	0.26

(1) corresponds to 500mg amoxycillin free acid.

(2) corresponds to 125mg free clavulanic acid.

10

Granules are prepared using this mixture in a manner identical to that of example 8. The flavour, polyplasdone XL, colouring and magnesium stearate were sieved then blended with the granulate. The aspartame was then added, and this mixture was then compressed into tablets on a conventional tableting machine. This tablet contains 625.0mg of the amoxycillin: clavulanate combination, and the quantities used may be halved to prepare a tablet containing 312.5mg.

15

Example 11: Tablet

<u>Component</u>	<u>Weight mg.</u>	<u>Weight %</u>
Amoxycillin trihydrate	290.7 ¹	46.3
Potassium clavulanate	152.4 ²	24.3
Syloid AL-1	152.4	24.3
Polyplasdone XL dried	8.7	1.38
}		
Dry flavour (Peppermint or mandarin)	3.0	0.48
Polyplasdone XL dried	12.5	2.00
Aspartame	7.5	1.19
Colouring	2.5	0.39
Magnesium stearate	1.25	0.20
}		

as granulate

extra granulate

- (1) corresponds to 250mg amoxycillin free acid.
 5 (2) corresponds to 125mg free clavulanic acid.

Tablets were made from this mixture using a procedure identical to that of example 10.

10 Example 12 : Sachet or Syrup Formulations

<u>Component</u>	<u>Weight mg</u>	<u>w + %</u>
Amoxycillin : potassium clavulanate 4 : 1 w : w + 3 wt% CLPVP	2255.6	63.3
granulate ¹		
CLPVP	13.5	0.38
Lemon dry flavour	408.0	11.46
Strawberry dry flavour	132.0	3.71
Peach dry flavour	102.0	2.86
Silicon dioxide USNF (Syloid AL-1)	450.0	12.64
Aspartame	45.0	1.26
Xantham gum	150.0	4.21
Total weight	3561.6	100.0

- 16 -

(1) amox : clav expressed as free acid.

The granulate was prepared using the procedure of example 8. This formulation could be supplied in a sachet, or could be made up into a
 5 syrup, for example at concentrations of 3561.6 mg/60ml or 7123.2 mg/60 ml or 7123.2 mg/60 ml (= 156.25 and 312.5 mg amoxycillin : clavulanate / 5 ml respectively). To adjust the syrup to a suitable viscosity and pH, aerosil 200, succinic acid and/or methocel E - 15 (dry) may be used.

10 Example 13 : Sachet Formulation

<u>Component</u>	<u>Weight (mg)</u>				<u>w +%</u>
Granulate (Amox:Kclav 4:1 or 7:1 + 3% PVP)	500	250	125	875	7-25
Lemon dry flavour	136	68	34	136)	
Strawberry dry flavour	44	22	11	44)	3-6.1
Peach dry flavour	34	17	8.5	34)	
Silicon Dioxide U.S.N.F. (Syloid AL-1)	150	75	37.5	150	2.1-4.3
Sucrose	to 3500	1750	1750	3500	to 100

(1) weights and Amox/Kclav expressed as free acid.

15 The granulate was prepared using the procedure of example 8, and was then mixed with the other excipients.

Example 14 : Tablet Formulation

Amox : clav¹ 4 : 1 4 : 1 2 : 1 7 : 1

<u>Component</u>	<u>weight (mg)</u>				<u>w + %</u>
Granulate ²	751.9	376.0	452.1	1201.3	70.90
Dry Flavour ³	6.0	3.0	3.0	8.0	0.48 - 0.63
Poliplasdone XL) dried)	100.0	50.0	66.5	110.0	8.1 - 10.7
Aspartame	15.0	7.5	7.5	15.0	1.1 - 1.6
Colouring	4 - 5	2 - 2.5	2 - 2.5	4 - 5	0.3 - 0.55
Mag. Stearate	2.5	1.25	1.25	3.4	0.19 - 0.26
Silicon Dioxide)					
Syloid AL - 1) to	950	475	628	1350	to 100

- (1) Amox : clav expressed as weight : weight of amoxycillin :
 5 clavulanate free acid.
 (2) Granulate = amox : clav + 3% CLPVP.
 (3) Peppermint or mandarin.

10 The granulate was prepared using the procedure of example 9.

The granulate was prepared using the procedure of example 9. The other excipients except aspartame were sieved and blended then mixed with the granulate. The aspartame was then added, and this mixture was then compressed into tablets in a conventional tableting machine. This tablet
 15 contained 625 mg of the amoxycillin : clavulanate blend. Tablets of different strengths could be formulated correspondingly, eg containing 1000, 375 or 312.5 mg of the amoxycillin : clavulanate combination.

Example 15 : Tablet Formulation

<u>Component</u>	<u>Weight (mg)</u>				<u>w +%</u>
Granulate (Amox.Kclav) 4:1 or 7:1 + 3% PVP	751.9	376.0	188.0	1201.3	71 - 83
Magnesium stearate Ph. Eur	2.6	1.3	0.65	3.9	0.25 - 0.27
Silicon Dioxide USP /NF (Syloid AL-1)	44.0	22.0	11.0	44.0	3 - 4.25
Microcrystalline cellulose Avicel pH 112 dried ...to..	850.0	425.0	212.5	1275.0	1.8-5
Organic film coating	yes	yes	yes	yes	to 100
Actual weight	1050.0	-	-	1450.0	

(1) amox : clav expressed as free acid.

5

The tablet was made up in a manner identical to that of example 14.

The weights and relative proportions of the components of examples 1 to 15 could be varied about the figures listed, but suitably are within $\pm 10\%$ of those listed, desirably within $\pm 5\%$, especially $\pm 2.5\%$.

10

- 18a -

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers.



THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A tablet formulation having a structure comprising compacted granulates; the granulates comprising at least one compacted medicament optionally together with an intra-granular disintegrant; the medicament being a β -lactam antibiotic optionally in combination with a β -lactamase inhibitor and the granulates being compacted together into a tablet form together with an extra-granular disintegrant and optionally also together with an extra-granular lubricant, provided that if a lubricant is present the amount of lubricant is less than 0.5% by weight of the total tablet.
2. A tablet formulation according to claim 1 wherein the antibiotic is amoxycillin, optionally in combination with a salt of clavulanic acid in a weight ratio equivalent to amoxycillin : clavulanic acid in the range 12:1 to 1:1.
3. A tablet formulation according to claim 1 or 2 wherein the intra-granular disintegrant is selected from maize starch, rice starch, cross linked N-vinyl-2-pyrrolidone ("CLPVP"), sodium starch glycollate, croscarmellose sodium, formaldehyde-casein or combinations thereof.
4. A tablet formulation according to any one of the preceding claims wherein the proportion of intra-granular disintegrant is 0.1 to 10 wt% of the weight of the granulate.
5. A tablet formulation according to any one of the preceding claims in which the granulate comprises a medicament which is amoxycillin or amoxycillin plus potassium clavulanate in combination, an intra-granular disintegrant which is CLPVP or sodium starch glycollate, and optionally one or more diluent(s), in a proportion 70-99 wt% medicament, 1-5 wt% disintegrant and up to 30 wt% diluent.
6. A tablet formulation according to any one of the preceding claims wherein the granulate comprises 70 wt% or more of the tablet weight.



7. A tablet formulation according to any one of the preceding claims in which the extra-granular disintegrant is selected from maize starch, rice starch, CLPVP, sodium starch glycollate, croscarmellose sodium, microcrystalline or microfine cellulose, low-substituted hydroxypropylcellulose, cross-linked sodium carboxymethylcellulose, swellable ion exchange resins, formaldehyde-casein, or alginates.

8. A tablet formulation according to any one of the preceding claims wherein the proportion of extra-granular disintegrant in the tablet is between 0.1 - 25 wt% of the total tablet weight.

10

9. A tablet formulation according to any one of the preceding claims which contains 0 - 0.35 wt% lubricant.

10. A pharmaceutical granulate formulation comprising a medicament which is a β -lactam antibiotic optionally in combination with a β -lactamase inhibitor and an intra-granular disintegrant which is CLPVP.

11. A formulation according to claim 10 wherein the medicament is amoxycillin in combination with a salt of clavulanic acid in a weight ratio equivalent to amoxycillin : clavulanic acid in the range 12:1 to 1:1.

12. A formulation according to claim 10 or 11 wherein the proportion of intra-granular disintegrant is 0.1 to 10 wt% of the formulation.

13. A formulation according to claim 10 in which the granulate comprises a medicament which is amoxycillin plus potassium clavulanate in combination, an intra-granular disintegrant which is CLPVP or sodium starch glycollate, and optionally one or more diluent(s) in a proportion 70-99 wt% medicament, 1-5 wt% disintegrant and up to 30 wt% diluent.

30

14. A formulation according to any one of claims 10 to 13 or a granulate as defined in claim 1 when encapsulated in a pharmaceutical capsule.



15. A process for the manufacture of a pharmaceutical tablet, in which granulates comprising at least one compacted medicament which is a β -lactam antibiotic optionally in combination with a β -lactamase inhibitor, optionally together with an intra-granular disintegrant are mixed with an extra-granular disintegrant and optionally with an extra granular lubricant and excipients, provided that if a lubricant is present it amounts to less than 0.5 wt% of the mixture, and the mixture is compressed into tablets.

16. A process for the manufacture of a pharmaceutical granulate, in which a medicament which is a β -lactam antibiotic optionally in combination with a β -lactamase inhibitor is compacted under pressure, together with an intra-granular disintegrant which is CLPVP.

17. A process according to claim 16 wherein the compaction is carried out using roller compaction.

15

18. A method for the treatment of bacterial infections in mammals which comprises administering to the mammal an effective amount of a pharmaceutical formulation according to any one of claims 1 to 14 in which the medicament is a β -lactam antibiotic optionally in combination with a β -lactamase inhibitor.

20

19. Use of a pharmaceutical formulation according to any one of claims 1 to 14 in which the medicament is a β -lactam antibiotic optionally in combination with a β -lactamase inhibitor, in the manufacture of a medicament.

20. Tablet or pharmaceutical granulate formulations, processes for their manufacture or methods of treatment involving them, substantially as hereinbefore described with reference to the Examples.

30

DATED this 27th day of March, 1995

Laboratorios Beecham SA

By Its Patent Attorneys

DAVIES COLLISON CAVE

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 92/01024

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all)*		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 A61K9/20		
II. FIELDS SEARCHED		
Minimum Documentation Searched?		
Classification System	Classification Symbols	
Int.Cl. 5	A61K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ¹		
III. DOCUMENTS CONSIDERED TO BE RELEVANT²		
Category ³	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	GB,A,2 005 538 (BEECHAM GROUP LIMITED) 25 April 1979 see claims 1,4-6 see page 1, line 47 - line 94 see examples 1,2	1-23
X	EP,A,0 281 200 (GIST BROCADES N.V.) 7 September 1988 cited in the application see the whole document	1-3,5, 7-10,18, 21-23
A	FR,A,2 320 731 (BEECHAM GROUP LIMITED) 11 March 1977 see claims 1,3,5 see page 1, line 10 - line 14 see page 1, line 21 - page 2, line 18 see page 2, line 33 - line 40 see page 3, line 16 - line 29	6,13-15, 17,19-20
<p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
21 OCTOBER 1992		09.11.92
International Searching Authority		Signature of Authorized Officer
EUROPEAN PATENT OFFICE		Dagmar Frank

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

EP 9201024
SA 63757

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 21/10/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A-2005538	25-04-79	AU-B- 525089	21-10-82
		AU-A- 4050678	17-04-80
		BE-A- 870988	03-04-79
		CA-A- 1105385	21-07-81
		CH-A- 642258	13-04-84
		DE-A- 2043318	12-04-79
		FR-A, B 2405711	11-05-79
		JP-C- 1494273	20-04-89
		JP-A- 54076831	19-06-79
		JP-B- 63041886	19-08-88
		NL-A- 7810176	17-04-79
		SE-B- 435899	29-10-84
		SE-A- 7810591	11-04-79
		SE-B- 451668	26-10-87
		SE-A- 8107592	17-12-81
		US-A- 4441609	10-04-84
		US-A- 4301149	17-11-81
EP-A-0281200	07-09-88	AU-B- 603870	29-11-90
		AU-A- 1252088	01-09-88
		JP-A- 63301820	08-12-88
		US-A- 4950484	21-08-90
FR-A-2320731	11-03-77	GB-A- 1552416	12-09-79