

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
14 September 2006 (14.09.2006)

PCT

(10) International Publication Number
WO 2006/094385 A1

(51) International Patent Classification:

C12N 7/02 (2006.01) A61K 39/275 (2006.01)
C12Q 1/04 (2006.01) A61K 35/76 (2006.01)
A61K 31/436 (2006.01)

N6G 2K5 (CA). **BARRETT, John** [CA/CA]; C/o Robarts Research Institute, 100 Perth Drive, P.o. Box 5015, London, Ontario N6A 5K8 (CA). **STANFORD, Marianne** [CA/CA]; C/o Robarts Research Institute, 100 Perth Drive, P.o. Box 5015, London, Ontario N6A 5K8 (CA).

(21) International Application Number:

PCT/CA2006/000315

(74) **Agent: SMART & BIGGAR**; ATTN: KANG, Yoon (Ms.), 438 University Avenue, Suite 1500, Box 111, Toronto, Ontario M5G 2K8 (CA).

(22) International Filing Date: 6 March 2006 (06.03.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/658,816 7 March 2005 (07.03.2005) US

(71) **Applicant (for all designated States except US): RO-BARTS RESEARCH INSTITUTE** [CA/CA]; 100 Perth Drive, P.o. Box 5015, London, Ontario N6A 5K8 (CA).

(72) Inventors; and

(75) **Inventors/Applicants (for US only): MCFADDEN, Grant** [CA/CA]; 1435 Corley Drive, S., London, Ontario


(81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,

[Continued on next page]


(54) **Title:** USE OF A COMBINATION OF MYXOMA VIRUS AND RAPAMYCIN FOR THERAPEUTIC TREATMENT

Multistep Growth Curve: BGMK (vMyxLac)



(57) **Abstract:** The present invention relates to therapeutic use of a combination of Myxoma virus, including in combination with rapamycin. Treatment with rapamycin enhances the ability of Myxoma virus to selectively infect cells that have a deficient innate anti-viral response, including cells that are not responsive to interferon. The combination of rapamycin and Myxoma virus can be used to treat diseases characterized by the presence of such cells, including cancer. The invention also relates to therapeutic use of Myxoma virus that does not express functional M135R.

Multistep Growth Curve: MCF7 (vMyxLac)



WO 2006/094385 A1



ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

- *with international search report*
- *with amended claims*

USE OF A COMBINATION OF MYXOMA VIRUS AND RAPAMYCIN FOR THERAPEUTIC TREATMENT

CROSS-REFERENCE TO RELATED APPLICATION

[0001] This application claims benefit and priority from U.S. provisional patent application No. 60/658,816, filed on March 7, 2005, the contents of which are incorporated herein by reference.

FIELD OF THE INVENTION

[0002] The present invention relates generally to therapeutic use of Myxoma virus and rapamycin.

BACKGROUND OF THE INVENTION

[0003] Current treatments used to treat various types of cancer tend to work by poisoning or killing the cancerous cell. Unfortunately, treatments that are toxic to cancer cells typically tend to be toxic to healthy cells as well. Moreover, the heterogenous nature of tumours is one of the primary reasons that effective treatments for cancer remain elusive. Current mainstream therapies such as chemotherapy and radiotherapy tend to be used within a narrow therapeutic window of toxicity. These types of therapies are considered blunt tools that have limited applicability due to the varying types of tumour cells and the limited window in which these treatments can be administered.

[0004] Modern anticancer therapies currently being developed attempt to selectively target tumour cells while being less toxic to healthy cells, thereby being more likely to leave healthy cells unaffected.

[0005] Oncolytic viral therapy is one approach that aims to exploit cellular differences between tumour cells and normal cells. This therapy uses replication-competent, tumour-selective viral vectors as anti-cancer agents. The oncolytic virus either specifically targets cancer cells for infection, or is more suited for efficient replication in cancer cells versus healthy cells. These replication-competent, oncolytic viruses are either naturally occurring or genetically engineered to be a highly selective and highly potent means of targeting the heterogeneous tumour population. Since the replication selective oncolytic virus does not replicate efficiently in normal cells, toxicity to the patient should be low, particularly in

comparison to traditional therapies such as radiation or chemotherapy.

[0006] Numerous studies have reported oncolytic activity for various virus strains, with the most promising oncolytic viruses being a naturally occurring or genetically modified version of adenovirus, herpes simplex virus 1 ("HSV1"), Reovirus, Vaccinia Virus, Vesicular Stomatitis Virus ("VSV") or Poliovirus. Modified oncolytic viruses currently under investigation as anticancer agents include HSV, adenovirus, Newcastle disease virus ("NDV"), Reovirus and Vaccinia virus, measles, VSV and poliovirus. Various oncolytic viruses are in Phase I and Phase II clinical trials with some showing sustained efficacy. However, it is unknown which viruses will best fulfill the oncolytic goals of sustained replication, specificity and potent lytic activity. A completely efficient candidate for an oncolytic viral vector would be one that has a short lifecycle, forms mature virions quickly, spreads efficiently from cell to cell and has a large genome ready for insertions. As well, evidence suggests that inhibiting the early innate immune response and slowing the development of Th1 responses are important for the efficacy of oncolytic therapy. It is clear that human viruses are highly immunogenic, as measured by the high level of antibody and T cell responses that are observed in the normal population for many of the viruses being considered for the development of oncolytic viruses.

[0007] Clinical work has shown that current oncolytic viruses are indeed safe, but are not potent enough as monotherapies to be completely clinically effective. As insufficient or inefficient infection of tumour cells is usually observed, the current movement is to arm candidate viruses by genetically engineering them to express therapeutic transgenes to increase their efficiency. Most of the above-mentioned oncolytic viruses are also being tested in combination with other common oncolytic therapies.

[0008] Adenovirus can be easily genetically manipulated and has well-known associated viral protein function. In addition, it is associated with a fairly mild disease. The ONYX-015 human adenovirus (Onyx Pharmaceuticals Inc.) is one of the most extensively tested oncolytic viruses that has been optimized for clinical use. It is believed to replicate preferentially in p53-negative tumours and shows potential in clinical trials with head and neck cancer patients. However, reports show that ONYX-015 has only produced an objective clinical response in 14% of treated patients (Nemunaitis J, Khuri F, Ganly I, Arseneau J, Posner M, Vokes E, Kuhn J, McCarty T, Landers S, Blackburn A,

Romel L, Randlev B, Kaye S, Kirn D. *J. Clin. Oncol.* 2001 Jan 15;19(2):289-98).

[0009] WO96/03997 and W097/26904 describe a mutant oncolytic HSV that inhibits tumour cell growth and is specific to neuronal cells. Further advantages are that the HSV can be genetically modified with ease, and drugs exist to shut off any unwanted viral replication. However, the application of such a common human pathogen is limited, as it is likely that the general population has been exposed and acquired an immune response to this virus, which would attenuate the lytic effect of the virus. HSV can also cause serious side effects or a potentially fatal disease.

[0010] Reovirus type III is associated with relatively mild diseases and its viral gene function is fairly well understood. Reovirus type III is currently being developed by Oncolytic Biotech as a cancer therapeutic which exhibits enhanced replication properties in cells expressing mutant ras oncogen and preferentially grows in PKR $-/-$ cells (Strong J.E. and P.W. Lee, *J. Virology*, 1996. 70:612-616). However, Reovirus is difficult to genetically manipulate and its viral replication cannot be easily shut off.

[0011] VSV is associated with relatively mild diseases and also has well-known viral gene function. WO99/04026 discloses the use of VSV as a vector in gene therapy for the expression of wide treatment of a variety of disorders. However, VSV suffers from the same problems as the Reovirus in that it is difficult to genetically manipulate and its viral replication cannot be easily shut off.

[0012] Vaccina virus and Poliovirus are other candidate oncolytic viruses described in the art but have been associated with a serious or potentially fatal disease.

[0013] US 4,806,347 discloses the use of gamma interferon and a fragment of IFN γ against human tumour cells. WO99/18799 discloses a method of treating disease in a mammal in which the diseased cells have defects in an interferon-mediated antiviral response, comprising administering to the mammal a therapeutically effective amount of an interferon-sensitive, replication competent clonal virus. It specifically discloses that VSV particles have toxic activity against tumour cells but that alleviation of cytotoxicity in normal cells by VSV occurs in the presence of interferon. WO99/18799 also discloses that NDV-induced sensitivity was observed with the interferon-treated tumour cells but that adding interferon to normal cells makes these cells resistant to NDV. This method

aims to make cells sensitive to interferon by infecting them with interferon sensitive viruses.

SUMMARY OF THE INVENTION

[0014] The present invention is based on the unexpected discovery that rabbit Myxoma virus, including a novel Myxoma virus that does not express functional M135R protein, can selectively infect cells, including human tumour cells, that have a deficient innate anti-viral response, including those that are non-responsive to interferon, and that such infection is enhanced by treating such cells with the drug rapamycin. The term “innate” as used in this context describes non-antigen specific immune response. Since Myxoma virus does not replicate efficiently in normal human cells, the virus can therefore be used as a treatment for various disorders and conditions characterized by cells that have a deficient innate anti-viral response, including cells that are non-responsive to interferon, for example, as an oncolytic treatment for cancer. The virus can also be used to identify cells that have a deficient innate anti-viral response and to image these cells *in vivo*.

[0015] In one aspect, the present invention provides a method for inhibiting a cell that has a deficient innate anti-viral response comprising administering to the cell an effective amount of a combination of Myxoma virus and rapamycin.

[0016] In one aspect, the invention provides a method for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response, comprising administering to a patient in need thereof an effective amount of a combination of Myxoma virus and rapamycin.

[0017] The present invention further provides use of an effective amount of a combination of Myxoma virus and rapamycin for inhibiting a cell that has a deficient innate anti-viral response and for the manufacture of a medicament for inhibiting a cell that has a deficient innate anti-viral response.

[0018] The present invention further provides use of an effective amount of a combination of Myxoma virus and rapamycin for treating a disease state in a patient, wherein the disease state is characterized by the presence of cells that have a deficient innate anti-viral response and for the manufacture of a medicament for treating such a

disease state in a patient.

[0019] In another aspect, the present invention provides a pharmaceutical composition comprising Myxoma virus and rapamycin. The pharmaceutical composition may be useful for inhibiting a cell that has a deficient innate anti-viral response or for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response.

[0020] In another aspect, the present invention provides a kit comprising Myxoma virus, rapamycin and instructions for inhibiting a cell that has a deficient innate anti-viral response or for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response. The disease states include cancer and a chronic viral infection.

[0021] The present invention further provides a method of detection a cell that has a deficient innate anti-viral response, comprising exposing a population of cells to a combination of Myxoma virus and rapamycin; allowing the virus to infect a cell that has a deficient innate anti-viral response; and determining the infection of any cells of the population of cells by the Myxoma virus.

[0022] The present invention is further based on the unexpected discovery that rabbit Myxoma virus protein M135R is involved in eliciting an immune response in rabbits and that a Myxoma virus strain that does not express functional M135R can kill cells *in vitro*, but does not cause myxomatosis disease in animals. Such a viral strain can be used to treat cells having a deficient innate anti-viral response, including those that are non-responsive to interferon, and including treatments given in combination with the drug rapamycin, without the need for increased containment of the virus, leading to improved safety.

[0023] In one aspect, the present invention provides a method for inhibiting a cell that has a deficient innate anti-viral response comprising administering to the cell an effective amount of Myxoma virus that does not express functional M135R, optionally in combination with an effective amount of rapamycin.

[0024] In one aspect, the invention provides a method for treating a disease state

characterized by the presence of cells that have a deficient innate anti-viral response, comprising administering to a patient in need thereof an effective amount of Myxoma virus that does not express functional M135R, optionally in combination with an effective amount of rapamycin.

[0025] The present invention further provides use of an effective amount of Myxoma virus that does not express functional M135R, optionally in combination with an effective amount of rapamycin, for inhibiting a cell that has a deficient innate anti-viral response and in the manufacture of a medicament for inhibiting a cell that has a deficient innate anti-viral response.

[0026] The present invention further provides use of an effective amount of Myxoma virus that does not express functional M135R, optionally in combination with an effective amount of rapamycin, for treating a disease state in a patient, wherein the disease state is characterized by the presence of cells that have a deficient innate anti-viral response and in the manufacture of a medicament for treating such a disease state in a patient.

[0027] In a further aspect, the present invention provides a Myxoma virus that does not express functional M135R.

[0028] In another aspect, the present invention provides a pharmaceutical composition comprising Myxoma virus that does not express functional M135R. The pharmaceutical composition may be useful for inhibiting a cell that has a deficient innate anti-viral response or for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response. The pharmaceutical composition may further comprise rapamycin.

[0029] In another aspect, the present invention provides a kit comprising Myxoma virus that does not express functional M135R and instructions for inhibiting a cell that has a deficient innate anti-viral response or for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response. The kit may further comprise rapamycin. The disease state includes cancer and a chronic viral infection.

[0030] The present invention further provides a method for detecting a cell that has a deficient innate anti-viral response, comprising exposing a population of cells to a

Myxoma virus that does not express functional M135R, optionally in combination with rapamycin; allowing the virus to infect a cell that has a deficient innate anti-viral response; and determining the infection of any cells of the population of cells by the Myxoma virus.

[0031] Other aspects and features of the present invention will become apparent to those of ordinary skill in the art upon review of the following description of specific embodiments of the invention in conjunction with the accompanying figures. It should be understood, however, that the detailed description and the specific examples while indicating preferred embodiments of the invention are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

[0032] In the figures, which illustrate embodiments of the present invention, by way of example only,

[0033] Figure 1 is a schematic diagram of an interferon mediated anti-viral signalling scheme induced upon viral infection of a cell;

[0034] Figure 2 is a phase contrast micrograph of nonpermissive WT murine embryonic fibroblasts ("MEFs") after exposure to Myxoma virus, demonstrating that the MEFs become permissive after inhibition of interferon α/β with neutralizing antibody;

[0035] Figure 3 is a Western blot showing phosphorylation states (activation) of STAT1 and STAT2 after Myxoma virus infection, demonstrating that nonpermissive infections of MEF cells is associated with activation of STAT 1 and STAT 2;

[0036] Figure 4 is a Western blot showing phosphorylation states (inactivation) of STAT3, STAT4, STAT5 and STAT6 after Myxoma virus infection, demonstrating that nonpermissive infections of MEF cells does not activate any of these species;

[0037] Figure 5 is a phase contrast micrograph of IFN α/β R $^{-/-}$ MEFs and STAT1 $^{-/-}$ MEFs, IFN α/β R $^{-/-}$ MEFs and STAT1 $^{-/-}$ MEFs after infection with Myxoma virus, showing that inactivation of IFN/STAT/JAK signalling renders cells permissive for Myxoma infection;

[0038] Figure 6 is a Western blot showing phosphorylation states of PKR in nonpermissive wildtype MEFs after Myxoma virus infection, demonstrating that PKR is not activated by Myxoma virus infection;

[0039] Figure 7 is a Western blot showing phosphorylation states of PKR in wildtype MEFs either mock infected or pre-infected with Myxoma virus, showing that Myxoma virus blocks PKR activation in MEF cells;

[0040] Figure 8 is a Western blot showing phosphorylation states of PERK in wildtype MEFs after Myxoma virus infection, demonstrating that Myxoma virus blocks PERK activation in MEF cells;

[0041] Figure 9 is a phase contrast micrograph of PKR^{-/-}, RNase L^{-/-} and Mx1^{-/-} triple knockout after exposure to Myxoma virus, showing that the antiviral state in MEF cells is mediated by a distinct pathway;

[0042] Figure 10 is a phase contrast micrograph of PKR^{-/-}, RNase L^{-/-} and Mx1^{-/-} triple knockout after exposure to Myxoma virus;

[0043] Figure 11 is a phase contrast micrograph of PKR^{-/-}, RNase L^{-/-} and Mx1^{-/-} triple knockout after treatment with neutralizing antibody to IFN α/β and after exposure to Myxoma virus;

[0044] Figure 12 is a Western blot showing phosphorylation levels of eIF2 α and PKR in nonpermissive MEFs after treatment with neutralizing antibody to IFN α/β and after exposure to Myxoma virus, showing that eIF2 α phosphorylation in nonresponsive cells is catalysed by a PKR-independent pathway;

[0045] Figure 13 is a Western blot showing STAT1 phosphorylation states in PKR^{-/-}, RNase L^{-/-} and Mx1^{-/-} triple knockout after Myxoma virus infection, indicating normal IFN-induced signalling responses

[0046] Figure 14 is a phase contrast micrograph illustrating subcellular localization of tyrosine-phosphorylated STAT1 in nonpermissive PKR^{-/-} + RNaseL^{-/-} + Mx1^{-/-} cells at 12 hours post-infection, indicating that the activated STAT localizes to the nucleus, as predicted for normal IFN/STAT signalling responses;

[0047] Figure 15 is a fluorescent image of brains from nude mice having intracranial gliomas mock-infected or infected with dead or live Myxoma virus expressing GFP, showing targeting of Myxoma to the glioma cells;

[0048] Figure 16 is a fluorescent image and a photograph of a thin-sectioned mouse glioma infected with Myxoma virus expressing GFP showing that the Myxoma virus replicated only in tumour cells;

[0049] Figure 17 is a phase contrast micrograph of HT29 human tumour cells, stained with either X-Gal or Crystal violet after infection with Myxoma virus, showing an example of a non-permissive infection in human cells;

[0050] Figure 18 is a phase contrast micrograph of HOP92 human tumour cells, stained with X-Gal or Crystal Violet after infection with Myxoma virus, showing an example of a permissive infection of human cells;

[0051] Figure 19 is phase contrast micrograph of OVCAR4 human tumour cells, stained with either X-Gal or Crystal Violet after infection with Myxoma virus, showing an example of a permissive infection of human cells;

[0052] Figure 20 is a phase contrast micrograph of SK-MEL3 human tumour cells, stained with either X-Gal or Crystal Violet after infection with Myxoma virus, showing an example of a permissive infection of human cells;

[0053] Figure 21 is a phase contrast micrograph of SK-MEL28 human tumour cells, stained with either X-Gal or Crystal Violet after infection with Myxoma virus, showing an example of a semi-permissive infection of human tumour cells;

[0054] Figure 22 is a phase contrast micrograph of BGMK cells, stained with either X-Gal or Crystal Violet after infection with Myxoma virus, showing a typical permissive control infection;

[0055] Figure 23 is a phase contrast micrograph of positive control BGMK cells and human tumour lines U87, A172 and U373 infected with increasing concentrations of Myxoma virus expressing the LacZ protein, stained with X-Gal, showing that these human glioma cells were all permissive for Myxoma virus replication;

[0056] Figure 24 is a graph depicting survival rate of BGMK, U87, A172 and U373 cells infected with Myxoma virus, 72 hours post-infection, at increasing concentrations of the virus, demonstrating the ability of Myxoma to kill all of these cells;

[0057] Figure 25 is a phase contrast micrograph and fluorescence micrograph of SF04 1585 astrocytoma cells infected with MV GFP, showing the infection in primary human glioma cells;

[0058] Figure 26 is a phase contrast micrograph of U373 glioma cells infected with Myxoma virus expressing the LacZ protein and stained with X-Gal, showing infection of these human tumour cells;

[0059] Figure 27 is a graph depicting the survival rate of SF04 1585 cells infected with MV GFP 48 hours post-infection, showing killing of these infected human tumour cells;

[0060] Figure 28 is a fluorescence micrograph of Daoy and D384 medulloblastoma lines infected with Myxoma virus expressing GFP, showing infection of these human tumour cells.

[0061] Figure 29 is graphical representations of the rate of virus production in various cell lines with or without pre-treatment with rapamycin: BGMK (primate control cell line); RK-13 and RL5 (rabbit control cell lines); 4T1 and B16F10 (mouse cancer cell lines); HOS, PC3, 786-0, HCT116, ACHN, MCF-7, M14 and COLO205 (human cancer cell lines); using wildtype virus vMyxLac and the M-T5 knock out virus vMyxT5KO as indicated;

[0062] Figure 30 is photographs of virally infected cell lines, infected with either vMyxLac or vMyxLacT5-;

[0063] Figure 31 is graphical representations of the rate of virus production in various cell lines (BGMK; A9; MCF-7; MDA-MB-435; M14; and COLO205) with or without pre-treatment with rapamycin;

[0064] Figure 32 is (A) a schematic alignment of Myxoma virus protein M135R and Vaccinia virus protein B18R and (B) an amino acid sequence alignment between M135R

and the first 179 amino acids of B18R;

[0065] Figure 33 is (A) a Western blot of M135R expressed in BGMK cells infected with Myxoma virus Lausanne (vMyxLau) and (B) a Western blot of M135R expressed in BGMK cells infected with vMyxLau and treated with araC, tunicamycin or monensin;

[0066] Figure 34 is (A) a fluorescence micrograph of BGMK cells mock infected or infected with Myxoma virus and stained for M135R and (B) a Western blot against immunoprecipitations or cell lysates of cells infected with wildtype Myxoma virus (vMyxgfp) or an M135R knockout strain (vMyx135KO) using anti-M135R antibody;

[0067] Figure 35 is (A) is a schematic diagram of the cloning strategy to produce vMyx135KO, (B) an agarose gel of the PCR insert product and (C) a Western blot of cells infected with wildtype and M135R knockout Myxoma virus;

[0068] Figure 36 is a growth curve of viral foci in BGMK cells infected with vMyxgfp or vMyx135KO;

[0069] Figure 37 is light and fluorescent micrographs of rabbit embryo fibroblasts infected with vMyxgfp or vMyx135KO;

[0070] Figure 38 is light and fluorescent micrographs of rabbit HIG82 fibroblasts infected with vMyxgfp or vMyx135KO;

[0071] Figure 39 is light and fluorescent micrographs of human primary fibroblasts infected with vMyxgfp or vMyx135KO;

[0072] Figure 40 is a graph of body temperature in rabbits infected with vMyxLau or vMyx135KO;

[0073] Figure 41 is a graph of ^{125}I emissions of cells mock infected or infected with vMyxgfp or vMyx135KO and treated with ^{125}I -labelled rabbit interferon α/β ;

[0074] Figure 42 is a graph of foci formed by infecting RK13 or BGMK cells with vMyxgfp or vMyx135KO, in which cells were untreated or treated with rabbit interferon α/β 24 hours prior to infection; and

[0075] Figure 43 is photographs of Western blots using cell lysates from 786-0 human cancer cells that were pre-treated with either 20nM rapamycin (R) or with the vehicle control (D), probed using antibodies directed against the indicated proteins.

DETAILED DESCRIPTION

[0076] Previously, the inventors have discovered that wildtype Myxoma virus, a virus that normally infects rabbits, can selectively infect and kill cells, including human cells, that have a deficient innate anti-viral response, for example, cells that are non-responsive to interferon, as described in the application PCT/CA2004/000341, which is herein fully incorporated by reference. Myxoma virus does not replicate efficiently in normal human cells. Since many diseases or conditions are characterized by the presence of cells that have a deficient innate anti-viral response, including cells that are not responsive to interferon, for example, cancer, Myxoma virus can be used to treat such diseases and conditions, including cancer, with low toxicity for normal healthy cells. Myxoma virus can also be used to treat chronically infected cells as such cells have a deficient innate anti-viral response. For example, many viruses encode gene products that function to inhibit the antiviral, interferon response of cells; Myxoma virus can selectively infect such cells.

[0077] Myxoma virus ("MV") is the causative agent of myxomatosis in rabbits. MV belongs to the *Leporipoxvirus* genus of the *Poxviridae* family, the largest of the DNA viruses. MV induces a benign disease in its natural host, the *Sylvilagus* rabbit in the Americas. However, it is a virulent and host-specific poxvirus that causes a fatal disease in European rabbits, characterized by lesions found systemically and especially around the mucosal areas. (Cameron C, Hota-Mitchell S, Chen L, Barrett J, Cao JX, Macaulay C, Willer D, Evans D, McFadden G. *Virology* 1999, 264(2): 298-318; Kerr P & McFadden G. *Viral Immunology* 2002, 15(2): 229-246).

[0078] MV is a large virus with a double-stranded DNA genome of 163 kb which replicates in the cytoplasm of infected cells (B. N. Fields, D. M. Knipe, P. M. Howley, Eds., *Virology* Lippincott Raven Press, New York, 2nd ed., 1996). MV is known to encode a variety of cell-associated and secreted proteins that have been implicated in down-regulation of the host's immune and inflammatory responses and inhibition of

apoptosis of virus-infected cells. MV can be taken up by all human somatic cells. However, other than in normal somatic rabbit cells, if the cells have a normal innate anti-viral response, the virus will not be able to productively infect the cell, meaning the virus will not be able to replicate and cause cell death.

[0079] Interferons (“IFNs”) are a family of cytokines that are secreted in response to a variety of stimuli. Interferons bind to cell surface receptors, activating a signaling cascade that leads to numerous cellular responses, including an anti-viral response and induction of growth inhibition and/or apoptotic signals. Interferons are classified as either type I or type II. Type I IFNs include IFN- α , - β , - τ , and - ω , which are all monomeric; the only type II IFN is IFN- γ , a dimer. Twelve different subtypes of IFN- α are produced by 14 genes, but all other IFNs are monogenic (Arduini et al., 1999). IFNs exert direct anti-tumour activity via the modulation of oncogene expression. Overexpression of growth-stimulating oncogenes or loss of tumour suppressor oncogenes can lead to malignant transformation. Some oncogenes implicated in the genesis of cancer are p53, Rb, PC, NF1, WT1, DCC.

[0080] Myxoma virus, as well as other oncolytic viruses such as Reovirus and VSV, needs to bypass the anti-viral defenses that exist in normal healthy cells in order to be able to replicate within cells. MV and other oncolytic viruses induce interferon production, and are generally sensitive to the anti-viral effect of the IFN pathway. Relevant proteins induced by the IFN anti-viral response, and which principally affect virus multiplication include PKR, OAS synthetase and Rnase L nuclease. PKR activates eIF2 α , leading to inhibition of translation and induction of apoptosis. A schematic representation of the IFN response pathway is depicted in Figure 1. In normal cells, MV is directly affected by PKR and eIF2 α .

[0081] Anti-viral response pathways are often disrupted in cancerous cells. For example, reduced or defective response to IFN is a genetic defect that often arises during the process of transformation and tumour evolution. Over 80% of tumour cell lines do not respond to, or exhibit impaired responses to, interferon. (Stojdl et al., *Cancer Cell* (2003) 4: 263-275 and references cited therein; Wong et al. *J Biol Chem.* (1997) 272(45):28779-85; Sun et al. *Blood.* (1998) 91(2):570-6; Matin et al. *Cancer Res.* (2001) 61(5):2261-6; Balachandran et al *Cancer Cell* (2004) 5(1):51-65). As previously disclosed in PCT/CA2004/000341, MV can infect and kill cancer cells, including human tumour cells,

and without being limited by any particular theory, it is believed that MV can infect these cells because they have a deficient innate anti-viral response.

[0082] Evidence suggests that inhibiting the early innate immune response and slowing the development of Th1 responses are important for the efficacy of oncolytic therapy. Although Myxoma virus is a virulent virus, it is host-specific and has a very narrow host range; it does not infect humans or mice. Without being limited by any specific theory, it is believed that since Myxoma virus is a non-human virus, it should encounter no pre-existing immune recognition in humans. Therefore, its potential as an oncolytic virus will be less compromised and Myxoma virus should provide more potent infection of permissive tumour cells than native human viruses, and thereby can provide an effective oncolytic treatment for cancer.

[0083] The Myxoma virus host range gene M-T5 appears to play a critical role during Myxoma virus infection of many human tumour cell lines (Sypula et al, (2004) *Gene Ther. Mol. Biol.* 8:103). The MT-5 gene encodes an ankyrin repeat protein that is required for Myxoma replication in rabbit lymphocytes, and Myxoma virus with the MT-5 gene deleted cannot cause myxomatosis in susceptible rabbits (Mossman et al, (1996) *J. Virol.* 70: 4394). Available evidence suggests that differences in the intracellular signalling within an infected human tumour cell are critical for distinguishing human tumour cells that are permissive to Myxoma virus infection and productive replication (Johnston et al, (2003) *J. Virol.* 77: 5877).

[0084] Furthermore, Myxoma virus possesses a protein, M135R, which displays homology to the amino terminus portion of interferon α/β receptor ("IFN α/β -R"). It has been suggested that M135R mimics the host IFN α/β -R in order to prevent IFN α/β from triggering a host anti-viral response (Barrett et al., *Seminars in Immunology* (2001)13:73-84). The prediction is based on sequence homology to the viral IFN α/β -R from vaccinia virus, B18R, and it has been demonstrated that Vaccinia virus ("VV") employs such an immune evasion strategy. However, M135R is only half the size of VV B18R and all other IFN α/β -R homologs from sequenced poxviruses, and in all cases aligns only to the amino terminus half of the homolog.

[0085] The inventors have discovered that even though immunofluorescence results

suggest that M135R localizes to the cell surface, attempts to demonstrate the ability of M135R to interact with IFN α/β have been negative. Despite these results, the inventors have discovered that deletion of M135R severely attenuates the ability of Myxoma virus to cause disease in host animals although Myxoma virus having such a deletion is equally effective at infecting and killing cells *in vitro* compared to wildtype MV. Thus, in one aspect, the present invention relates to the discovery that Myxoma virus that does not express functional M135R is useful for treatment of cells having a deficient innate anti-viral response, including for oncolytic studies, since this virus provides a safer alternative for oncolytic viral therapy as no unusual containment strategies should be needed for patients undergoing treatment.

[0086] In another aspect, the present invention relates to the discovery that the anti-cancer agent rapamycin acts to enhance the levels of infectivity of Myxoma virus in human tumour cells which are permissive for Myxoma virus infection, and that rapamycin allows replication of certain strains of Myxoma virus in human tumour cells which, without rapamycin, are restrictive for the replication of those strains of Myxoma virus. A cell that is permissive for Myxoma virus infection is a cell that the virus can enter and in which the virus can productively reproduce. Permissive cells may have defects or mutations in one or more of the pathways that involve the proteins PTEN, PDK, AKT, GSK, Raf, mTOR or P70S6K. A restrictive cell is a cell which is permissive to Myxoma virus only under certain conditions, but does not allow productive infection under other conditions. For example, a restrictive cell may be permissive to wildtype strains of the virus, but does not allow certain mutant Myxoma strains, for example a strain having the MT-5 gene knocked out, to productively reproduce. In another example, a cell restrictive for Myxoma virus may not permit productive infection of Myxoma virus alone, but when treated with rapamycin, the same Myxoma virus is able to productively infect the cell. Abortive cell lines are non-permissive for Myxoma virus infection, meaning that the virus may be able to enter the cell, but does not productively infect the cell.

[0087] Thus, rapamycin, when used in combination with Myxoma virus, enhances the infectivity of Myxoma virus for cells having a deficient innate anti-viral response. The present invention relates to the use of rapamycin in combination with Myxoma virus to treat cells having a deficient innate anti-viral response.

[0088] Rapamycin is a macrocyclic lactone that has been shown to be the active antifungal compound purified from the soil bacterium *Streptomyces hygroscopicus*. Rapamycin as used herein refers to rapamycin (also referred to as sirolimus) and analogs or derivatives thereof capable of complexing with FKBP12 and inhibiting mTOR, including the analogs CCI-779 (also referred to as cell cycle inhibitor-779 or rapamycin-42,2,2-bis(hydroxymethyl)-propionic acid) and RAD001 (also referred to as everolimus or 40-O-(2-hydroxyethyl)-rapamycin). Rapamycin, CCI-779 and RAD001 are commercially available, and rapamycin is available under the name RapamuneTM, from Wyeth-Ayerst. The term rapamycin further includes pharmaceutically acceptable salts and esters of rapamycin, its hydrates, solvates, polymorphs, analogs or derivatives, as well as pro-drugs or precursors which are metabolized or converted to rapamycin or its analogs or derivatives during use, for example when administered to a patient.

[0089] Rapamycin as an inhibitor of cellular signaling is highly specific: it enters the cell and binds to a cellular protein known as FKBP12. The rapamycin/FKBP12 complex then binds to the specific cellular target mTOR (mammalian Target of Rapamycin). Many cancers have been shown to develop from an over activity of signaling molecules such as P13K, or a loss of the tumor suppressor gene PTEN. Both of these molecules lie upstream of mTOR. mTOR has been shown to be a central regulator of cell proliferation, growth, differentiation, migration and survival, and is therefore an ideal target in stemming the uncontrolled growth of cancer cells. Cancer cell lines that are sensitive to rapamycin are generally those that have resulted from an activation of the pathway through mTOR.

[0090] Rapamycins are used primarily in transplant patients as an alternative or complementary treatment to cyclosporine treatment. In transplant patients, rapamycin treatment generally has fewer side effects than cyclosporine A or FK506. In addition, retrospective studies have indicated that patients on rapamycin treatment generally develop fewer cancers and have a lower incidence of CMV (cytomegalovirus; a herpes virus) infection. It is therefore surprising that rapamycin treatment enhances Myxoma virus infection of cancer cells, particularly in light of research postulating that CMV replication should be reduced by rapamycin (reviewed by Ponticelli: "The pleiotropic effects of mTOR inhibitors" in *J Nephrology* 2004; 17: 762). Without being limited to a particular theory, it is possible that Myxoma virus takes advantage of aberrant signaling

through the mTOR pathway that may be associated with the neoplastic phenotype of these cells. Manipulation of this pathway by mTOR inhibitors could then be a selective advantage to the virus.

[0091] Thus, there is provided a method for inhibiting a cell that has a deficient innate anti-viral response comprising administering to the cell an effective amount of Myxoma virus. In a further embodiment, the virus is administered in combination with an effective amount of rapamycin.

[0092] The Myxoma virus may be any virus that belongs to the Leporipoxvirus species of pox viruses that is replication-competent. The Myxoma virus may be a wild-type strain of Myxoma virus or it may be a genetically modified strain of Myxoma virus, including an MT-5 knockout strain of Myxoma. The Myxoma virus may be a strain that has an attenuated affect in rabbits, thereby causing lower risk of disease, including a strain that does not express functional M135 protein, as described below.

[0093] In a particular embodiment, the Myxoma virus is a Myxoma virus that does not express functional M135R.

[0094] A Myxoma virus that does not express functional M135R includes a Myxoma virus that has part, or all, of the open reading frame that encodes M135R deleted, replaced or interrupted such that no gene product, no stable gene product, or no functional gene product is expressed. Such a virus also includes a Myxoma virus that has part, or all, of the M135R gene regulatory region deleted, replaced or interrupted such that no protein can be expressed from the gene encoding M135R. Functional M135R protein is M135R that is transcribed, translated, folded, post-translationally modified and localized within the cell, and which allows Myxoma virus to cause myxomatosis in an infected host. If the M135R protein is not, or not properly or not sufficiently, transcribed, translated, folded, post-translationally modified or localized within the cell such that an infected host does not develop myxomatosis, then no functional M135R protein is expressed in the cell.

[0095] In a further embodiment, the cell is non-responsive to interferon.

[0096] In specific embodiments, the cell is a mammalian cancer cell. In one embodiment the cell is a human cancer cell including a human solid tumour cell.

[0097] In another embodiment, the cell is chronically infected with a virus.

[0098] A “combination” of rapamycin and Myxoma virus for administration may be formulated together in the same dosage form or may be formulated in separate dosage forms, and the separate dosage forms may be the same form or different forms, for administration by the same mode or by different modes of administration. Furthermore, administration of a combination of rapamycin and Myxoma virus, when not together in the same dosage form, means that the rapamycin and Myxoma virus are administered concurrently to the mammal being treated, and may be administered at the same time or sequentially in any order or at different points in time. Thus, rapamycin and Myxoma virus may be administered separately but sufficiently closely in time so as to provide the desired therapeutic effect.

[0099] The term “effective amount” as used herein means an amount effective, at dosages and for periods of time necessary to achieve the desired result.

[00100] The term “a cell that has a deficient innate anti-viral response” as used herein refers to a cell that, when exposed to a virus or when invaded by a virus, does not induce anti-viral defence mechanisms, which include inhibition of viral replication, production of interferon, induction of the interferon response pathway, and apoptosis, which may or may not be mediated by interferon, and is thereby infectable by MV, alone or in combination with rapamycin treatment. The term includes a cell that has a reduced or defective innate anti-viral response upon exposure to or infection by a virus as compared to a normal cell, for example, a non-infected, or non-cancer cell. This includes a cell that is non-responsive to interferon and a cell that has a reduced or defective apoptotic response or induction of the apoptotic pathway. The deficiency may be caused by various causes, including infection, genetic defect, or environmental stress. It will however be understood that when the deficiency is caused by a pre-existing infection, superinfection by MV may be excluded and a skilled person can readily identify such instances. A skilled person can readily determine without undue experimentation whether any given cell type has a deficient innate anti-viral response and therefore infectable by Myxoma virus, either alone or in combination with rapamycin treatment. For example, VSV is commonly used to measure an anti-viral response of a cell.

[00101] To assess whether a given cell type, for example a given cancer cell type, has a deficient innate anti-viral response, a skilled person can take an explant, grow some of the cells *in vitro* and determine infectability by VSV or alternatively, by Myxoma virus, including Myxoma virus in combination with rapamycin.

[00102] The term “a cell that is non-responsive to interferon” as used throughout the specification means a cell that does not respond to the activity of interferon, for example anti-viral or anti-tumour activity of interferon or that has an abnormal interferon response, for example, a reduced or ineffective response to interferon, or abnormal interferon signalling as measured by, for example, phosphorylation or activation of signalling molecules such as transcription factors, for example STAT1. For example, without limitation, the cell may not undergo inhibition of proliferation or it may not be killed when exposed to interferon levels sufficient to induce such a response in a cell that is responsive to interferon. The cell that is non-responsive to interferon may have a defect in the intracellular signalling pathway or pathways that are normally activated in the responsive cells. Typically, susceptibility to infection by VSV is indicative of non-responsiveness to interferon, and a skilled person can readily determine whether a particular cell is non-responsive to interferon by its ability, or lack thereof, to inhibit VSV infection in the presence of interferon or using other markers of interferon activity known in the art, for example, the level of expression of IFN stimulated genes such as PKR, STAT, OAS, MX.

[00103] The term “replication-competent” as used throughout the specification refers to a virus that is capable of infecting and replicating within a particular host cell. This includes a virus which alone is restricted for replication in a particular host cell, but when the host cell is treated with rapamycin, the virus can then productively infect that cell.

[00104] The term “a cell” as used herein includes a single cell as well as a plurality or population of cells. Administering an agent to a cell includes both *in vitro* and *in vivo* administrations.

[00105] The term “animal” as used herein includes all members of the animal kingdom, including particularly mammals, especially humans.

[00106] The term “inhibiting” a cell that has a deficient innate anti-viral response includes cell death by lysis or apoptosis or other mechanisms of cell death, in addition to rendering the cell incapable of growing or dividing or reducing or retarding cell growth or division.

[00107] The Myxoma virus genome may be readily modified to express one or more therapeutic transgenes using standard molecular biology techniques known to a skilled person, and described for example in Sambrook et al. ((2001) *Molecular Cloning: a Laboratory Manual*, 3rd ed., Cold Spring Harbour Laboratory Press). A skilled person will be able to readily determine which portions of the Myxoma viral genome can be deleted such that the virus is still capable of productive infection. For example, non-essential regions of the viral genome that can be deleted can be deduced from comparing the published viral genome sequence with the genomes of other well-characterized viruses (see for example C. Cameron, S. Hota-Mitchell, L. Chen, J. Barrett, J.-X. Cao, C. Macaulay, D. Willer, D. Evans, and G. McFadden, *Virology* (1999) 264: 298-318)).

[00108] The term “therapeutic gene” or “therapeutic transgenes” as used herein is intended to describe broadly any gene the expression of which effects a desired result, for example, anti-cancer effect. For example, the virus may be modified to carry a gene that will enhance the anti-cancer effect of the viral treatment. Such a gene may be a gene that is involved in triggering apoptosis, or is involved in targeting the infected cell for immune destruction, such as a gene that repairs a lack of response to interferon, or which results in the expression of a cell surface marker that stimulates an antibody response, such as a bacterial cell surface antigen. The virus may also be modified to express genes involved in shutting off the neoplastic or cancer cell’s proliferation and growth, thereby preventing the cells from dividing. As well, the virus may be modified to include therapeutic genes, such as genes involved in the synthesis of chemotherapeutic agents, or it may be modified to have increased replication levels in cells of the particular species from which the cells to be inhibited or killed are derived, for example, human cells. Specific examples of genes that may be inserted into the Myxoma virus to increase its anti-cancer effect include the human gene for the TRAIL protein or the adenoviral gene that encodes the E4 orf4 polypeptide, both of which proteins are involved in killing human tumour cells.

[00109] It will be understood that therapeutic effect of the Myxoma virus, including

when used in combination with rapamycin, may be achieved by cell lysis by the virus or by delivery of therapeutic products by the virus. The inclusion of rapamycin in combination with the Myxoma virus should allow for enhancement of the effect of Myxoma virus alone. That is, the Myxoma virus, when administered in combination with rapamycin should be able to productively infect a greater number of target cells than Myxoma virus alone, or should be able to productively infect target cells having a deficient innate anti-viral response which are restrictive for productive infection by Myxoma virus in the absence of rapamycin.

[00110] The virus may be prepared using standard techniques known in the art. For example, the virus may be prepared by infecting cultured rabbit cells with the Myxoma virus strain that is to be used, allowing the infection to progress such that the virus replicates in the cultured cells and can be released by standard methods known in the art for disrupting the cell surface and thereby releasing the virus particles for harvesting. Once harvested, the virus titre may be determined by infecting a confluent lawn of rabbit cells and performing a plaque assay (see Mossman et al. (1996) *Virology* 215:17-30).

[00111] There is also provided a method for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response in a patient in need of such treatment comprising administering to the patient an effective amount of Myxoma virus, optionally in combination with rapamycin. The patient may be any animal, including a mammal, including a human.

[00112] “A disease state characterized by the presence of cells that have a deficient innate anti-viral response” as used herein refers to any disease, disorder or condition which is associated with, related to, or a characteristic of which is, the presence of cells that have a deficient innate anti-viral response and which disease, disorder, condition or symptoms thereof may be treated by killing these cells. For example, the disease state may be cancer. The disease state may also include chronic infection with a virus.

[00113] “Treating” a disease state refers to an approach for obtaining beneficial or desired results, including clinical results. Beneficial or desired clinical results can include, but are not limited to, alleviation or amelioration of one or more symptoms or conditions, diminishment of extent of disease, stabilization of the state of disease, prevention of

development of disease, prevention of spread of disease, delay or slowing of disease progression, delay or slowing of disease onset, amelioration or palliation of the disease state, and remission (whether partial or total). "Treating" can also mean prolonging survival of a patient beyond that expected in the absence of treatment. "Treating" can also mean inhibiting the progression of disease, slowing the progression of disease temporarily, although more preferably, it involves halting the progression of the disease permanently.

[00114] In one embodiment, the disease state is cancer. The cancer may be any type of cancer wherein at least some of the cells, although not necessarily all of the cells have a deficient innate anti-viral response. In one embodiment, the cancer may be a cancer wherein at least some of the cells are non-responsive to interferon. As used herein, the terms "tumour", "tumour cells", "cancer" and "cancer cells", (used interchangeably) refer to cells that exhibit abnormal growth, characterized by a significant loss of control of cell proliferation or cells that have been immortalized. The term "cancer" or "tumour" includes metastatic as well as non-metastatic cancer or tumours. As used herein, "neoplastic" or "neoplasm" broadly refers to a cell or cells that proliferate without normal growth inhibition mechanisms, and therefore includes benign tumours, in addition to cancer as well as dysplastic or hyperplastic cells.

[00115] A cancer may be diagnosed using criteria generally accepted in the art, including the presence of a malignant tumor.

[00116] Types of cancer that may be treated according to the present invention include, but are not limited to, hematopoietic cell cancers including leukemias and lymphomas, colon cancer, lung cancer, kidney cancer, pancreas cancer, endometrial cancer, thyroid cancer, oral cancer, ovarian cancer, laryngeal cancer, hepatocellular cancer, bile duct cancer, squamous cell carcinoma, prostate cancer, breast cancer, cervical cancer, colorectal cancer, melanomas and any other tumours. Solid tumours such as sarcomas and carcinomas include but are not limited to fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, and other sarcomas, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, lymphoid malignancy, pancreatic cancer, breast cancer, lung cancers, ovarian cancer, prostate cancer, hepatocellular carcinoma, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma,

papillary adenocarcinomas, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, Wilms' tumor, cervical cancer, testicular tumor, bladder carcinoma, and CNS tumors (such as a glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma and retinoblastoma).

[00117] In another embodiment, the disease state is a chronic viral infection.

[00118] The chronically infecting virus may be any virus that infects and replicates in cells of an animal in a persistent manner over a prolonged period so as to cause a pathological condition. The chronically infecting virus may be a virus that is associated or correlated with the development of cancer.

[00119] A chronic infection with a virus may be diagnosed using standard methods known in the art. For example, a chronic viral infection may be detected by the presence of anti-viral antibodies in the patient or a positive test for the presence of viral RNA or DNA in cells of the patient.

[00120] When administered to a patient, an effective amount of the Myxoma virus, and optionally the combination of Myxoma virus with rapamycin, is the amount required, at the dosages and for sufficient time period, for the virus to alleviate, improve, mitigate, ameliorate, stabilize, prevent the spread of, slow or delay the progression of or cure the disease. For example, it may be an amount sufficient to achieve the effect of reducing the number of or destroying cancerous cells or neoplastic cells, or reducing the number of or destroying cells chronically infected with a virus, or inhibiting the growth and/or proliferation of such cells.

[00121] The effective amount to be administered to a patient can vary depending on many factors such as the pharmacodynamic properties of the Myxoma virus and the optionally rapamycin, the modes of administration, the age, health and weight of the patient, the nature and extent of the disease state, the frequency of the treatment and the type of concurrent treatment, if any, and the virulence and titre of the virus.

[00122] One of skill in the art can determine the appropriate amount of Myxoma

virus for administration based on the above factors. The virus may be administered initially in a suitable amount that may be adjusted as required, depending on the clinical response of the patient. The effective amount of virus can be determined empirically and depends on the maximal amount of the virus that can be administered safely, and the minimal amount of the virus that produces the desired result.

[00123] Myxoma virus may be administered to the patient using standard methods of administration. In one embodiment, the virus is administered systemically. In another embodiment, the virus is administered by injection at the disease site. In a particular embodiment, the disease state is a solid tumour and the virus is administered by injection at the tumour site. In various embodiments, the virus may be administered orally or parenterally, or by any standard method known in the art.

[00124] To produce the same clinical effect when administering the virus systemically as that achieved through injection of the virus at the disease site, administration of significantly higher amounts of virus may be required. However, the appropriate dose level should be the minimum amount that would achieve the desired result.

[00125] The concentration of virus to be administered will vary depending on the virulence of the particular strain of Myxoma that is to be administered and on the nature of the cells that are being targeted. In one embodiment, a dose of less than about 10^9 plaque forming units ("pfu") is administered to a human patient. In various embodiments, between about 10^2 to about 10^9 pfu, between about 10^2 to about 10^7 pfu, between about 10^3 to about 10^6 pfu, or between about 10^4 to about 10^5 pfu may be administered in a single dose.

[00126] One of skill in the art can also determine, using the above factors, the appropriate amount of rapamycin to administer to a patient. The effective amount of rapamycin can be determined empirically and will depend on the amount and strain of virus being administered, the maximum amount of rapamycin that can be safely administered and the minimal amount of rapamycin that can be administered in order to achieve an enhancement of the infectivity of Myxoma virus.

[00127] Rapamycin may be administered to the patient using standard methods of

administration. In one embodiment, the rapamycin is administered systemically. In another embodiment, the rapamycin is administered by injection at the disease site. In a particular embodiment, the disease state is a solid tumour and the rapamycin is administered by injection at the tumour site. In various embodiments, the rapamycin may be administered orally or parenterally, or by any standard method known in the art.

[00128] The total amount of rapamycin may be administered in a single dose or in multiple doses spread out over 1 day or several days. The frequency and duration of administration of doses can be readily determined. The schedule of dosing will depend on the length of time that the Myxoma virus is to be administered. For example, rapamycin may be administered once to a patient, or may be administered 2 to 4 times per day.

[00129] In various embodiments, the dose of rapamycin may be from about 0.01 to about 250 mg per kg of body weight per day, from about 0.01 to 50 mg per kg of body weight per day, from about 0.05 to 10 mg per kg of body weight per day, or from about 0.1 to 7.5 mg per kg of body weight per day.

[00130] Effective amounts of a combination of Myxoma virus and rapamycin can be given repeatedly, depending upon the effect of the initial treatment regimen. Administrations are typically given periodically, while monitoring any response. It will be recognized by a skilled person that lower or higher dosages than those indicated above may be given, according to the administration schedules and routes selected.

[00131] The Myxoma virus, optionally in combination with rapamycin, may be administered as a sole therapy or may be administered in combination with other therapies, including chemotherapy, radiation therapy or other anti-viral therapies. For example, the Myxoma virus, optionally in combination with rapamycin, may be administered either prior to or following surgical removal of a primary tumour or prior to, concurrently with or following treatment such as administration of radiotherapy or conventional chemotherapeutic drugs. In one embodiment, the Myxoma virus, optionally in combination with rapamycin can be administered in combination with, or in a sequential fashion with, other oncolytic viruses, which may demonstrate specificity for varying tumour cell types.

[00132] To aid in administration, the Myxoma virus, optionally in combination

together with rapamycin, may be formulated as an ingredient in a pharmaceutical composition. Therefore, in a further embodiment, there is provided a pharmaceutical composition comprising Myxoma virus, and optionally rapamycin, and a pharmaceutically acceptable diluent. The invention in one aspect therefore also includes such pharmaceutical compositions for use in inhibiting a cell that has a deficient innate anti-viral response or treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response. The compositions may routinely contain pharmaceutically acceptable concentrations of salt, buffering agents, preservatives and various compatible carriers. For all forms of delivery, the recombinant Myxoma virus may be formulated in a physiological salt solution.

[00133] The pharmaceutical compositions may additionally contain additional therapeutic agents, such as additional anti-cancer agents. In one embodiment, the compositions include a chemotherapeutic agent. The chemotherapeutic agent, for example, may be substantially any agent which exhibits an oncolytic effect against cancer cells or neoplastic cells of the patient and that does not inhibit or diminish the tumour killing effect of the Myxoma virus. For example, the chemotherapeutic agent may be, without limitation, an anthracycline, an alkylating agent, an alkyl sulfonate, an aziridine, an ethylenimine, a methylmelamine, a nitrogen mustard, a nitrosourea, an antibiotic, an antimetabolite, a folic acid analogue, a purine analogue, a pyrimidine analogue, an enzyme, a podophyllotoxin, a platinum-containing agent or a cytokine. Preferably, the chemotherapeutic agent is one that is known to be effective against the particular cell type that is cancerous or neoplastic.

[00134] The proportion and identity of the pharmaceutically acceptable diluent is determined by chosen route of administration, compatibility with a live virus, and where applicable compatibility with the chemical stability of rapamycin, and standard pharmaceutical practice. Generally, the pharmaceutical composition will be formulated with components that will not significantly impair the biological properties of the live Myxoma virus, or cause degradation of or reduce the stability or efficacy of the rapamycin where included.

[00135] The pharmaceutical composition can be prepared by known methods for the preparation of pharmaceutically acceptable compositions suitable for administration to

patients, such that an effective quantity of the active substance or substances is combined in a mixture with a pharmaceutically acceptable vehicle. Suitable vehicles are described, for example, in Remington's Pharmaceutical Sciences (Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., USA 1985). On this basis, the compositions include, albeit not exclusively, solutions of the Myxoma virus, optionally with rapamycin, in association with one or more pharmaceutically acceptable vehicles or diluents, and contained in buffer solutions with a suitable pH and iso-osmotic with physiological fluids.

[00136] The pharmaceutical composition may be administered to a patient in a variety of forms depending on the selected route of administration, as will be understood by those skilled in the art. The composition of the invention may be administered orally or parenterally. Parenteral administration includes intravenous, intraperitoneal, subcutaneous, intramuscular, transepithelial, nasal, intrapulmonary, intrathecal, rectal and topical modes of administration. Parenteral administration may be by continuous infusion over a selected period of time.

[00137] The pharmaceutical composition may be administered orally, for example, with an inert diluent or with an assimilable carrier, or it may be enclosed in hard or soft shell gelatin capsules, or it may be compressed into tablets. For oral therapeutic administration, the Myxoma virus may be incorporated, optionally together with rapamycin, with an excipient and be used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers and the like.

[00138] Solutions of Myxoma virus, optionally together with rapamycin, may be prepared in a physiologically suitable buffer. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms, but that will not inactivate the live virus. A person skilled in the art would know how to prepare suitable formulations. Conventional procedures and ingredients for the selection and preparation of suitable formulations are described, for example, in Remington's Pharmaceutical Sciences and in The United States Pharmacopeia: The National Formulary (USP 24 NF19) published in 1999.

[00139] In different embodiments, the composition is administered by injection

(subcutaneously, intravenously, intramuscularly, etc.) directly at the disease site, such as a tumour site, or by oral administration, alternatively by transdermal administration.

[00140] The forms of the pharmaceutical composition suitable for injectable use include sterile aqueous solutions or dispersion and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions, wherein the term sterile does not extend to the live Myxoma virus itself that is to be administered. In all cases the form must be sterile and must be fluid to the extent that easy syringability exists.

[00141] The dose of the pharmaceutical composition that is to be used depends on the particular condition being treated, the severity of the condition, the individual patient parameters including age, physical condition, size and weight, the duration of the treatment, the nature of concurrent therapy (if any), the specific route of administration and other similar factors that are within the knowledge and expertise of the health practitioner. These factors are known to those of skill in the art and can be addressed with minimal routine experimentation.

[00142] The Myxoma virus, optionally in combination with rapamycin, or pharmaceutical compositions comprising the Myxoma virus and rapamycin, either together in the same formulation or different formulations, may also be packaged as a kit, containing instructions for use of Myxoma virus and rapamycin, including the use of Myxoma virus, or use of Myxoma virus in combination with rapamycin, to inhibit a cell that has a deficient innate anti-viral response, or use of Myxoma virus, or use of Myxoma virus in combination with rapamycin, to treat a disease state characterized by the presence of cells that have a deficient innate anti-viral response, in a patient in need thereof. The disease state may be cancer, or it may be a chronic viral infection.

[00143] The present invention also contemplates the use of Myxoma virus, optionally in combination with rapamycin, for inhibiting a cell that has a deficient innate anti-viral response. In one embodiment, the cell is non-responsive to interferon. There is further provided use of Myxoma virus, optionally in combination with rapamycin, for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response, in a patient in need thereof. In one embodiment the disease state is cancer. There is also provided use of Myxoma virus, optionally in combination with

rapamycin, in the manufacture of a medicament, for inhibiting a cell that has a deficient innate anti-viral response, or for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response in a patient in need thereof.

[00144] MV can selectively infect cells in or derived from animals other than the natural host of MV, from a population of cells, which have a deficient innate anti-viral response. This ability of MV provides for the use of MV in detecting cells from a population of cells, either in culture or in an animal, that have a deficient innate anti-viral response, including cells that are non-responsive to interferon. Such cells may otherwise not be easily detectable, for example certain cancer cells that have not yet advanced to palpable tumour, or have not yet induced noticeable symptoms in the animal.

[00145] Thus, in one embodiment, there is provided a method for detecting cells that have a deficient innate anti-viral response in a patient, comprising administering to the patient Myxoma virus modified to express a detectable marker, optionally in combination with rapamycin; allowing the virus to infect a cell that has a deficient innate anti-viral response in the patient; and detecting the cell expressing the detectable marker in the patient.

[00146] The infected cells may be detected using any conventional method for visualizing diagnostic images. The method of detection will depend on the particular detectable marker that is used. For example, cells infected with Myxoma virus genetically modified to express a fluorescent protein may be detected using fluorescence digital imaging microscopy. Other methods include computed tomography (CT), whole body scan such as position emission tomography (PET), magnetic resonance imaging (MRI), and sonography. Skilled artisans will be able to determine the appropriate method for detecting a particular detectable marker.

[00147] The detectable marker includes, but is not limited to, any marker for which genes for its expression or synthesis can be inserted into the Myxoma genome so as to result in expression or synthesis of the marker within cells that are infected by the modified virus. For example, in one embodiment, the detectable marker may be a fluorescent protein. The infected cells may be detected at a suitable time interval after administration of the modified virus to the patient, so as to allow for the virus to infect any

cells that have a deficient innate anti-viral response, and to express the detectable marker in such cells at levels that would allow for detection. For example, detection may occur anywhere between 2 and 20 days following administration to the patient of the virus genetically modified to express a fluorescent protein.

[00148] The detecting method may be carried out repeatedly at intervals in a patient in order to monitor the presence of cells that have a deficient innate anti-viral response in that patient. For example, the method for detecting such cells using Myxoma virus may be carried out on a patient at 6 month, 1 year or 2 year intervals, as is necessary, depending on the nature of the cells that has a deficient innate anti-viral response and the nature of any disease state caused as a result of the presence of such cells in a patient. Repeating the method over a time period allows for monitoring of the progression or remission of disease state, or the spread of disease within the body of the patient.

[00149] Myxoma virus is capable of selectively infecting cells that have a deficient innate anti-viral response, and can be used as an indicator of such a deficiency in cells. Thus, cells removed from a patient may be assayed for deficiency in innate anti-viral response using the methods of the present invention. Such determination may indicate, when combined with other indicators, that the patient may be suffering from a particular disease state, for example, cancer.

[00150] In one embodiment therefore, there is provided a method for detecting in a sample a cell that has a deficient innate anti-viral response comprising culturing the cell, exposing cultured cells to Myxoma virus, optionally in combination with rapamycin; and determining infectivity of cells by Myxoma virus.

[00151] The cells may be removed from a subject, including a human subject, using known biopsy methods. The biopsy method will depend on the location and type of cell that is to be tested.

[00152] Cells are cultured according to known culturing techniques, and are exposed to MV, and optionally rapamycin, by adding live Myxoma virus, and optionally rapamycin, to the culture medium. Where Myxoma virus is added in combination with rapamycin, the virus and rapamycin may be added either simultaneously or sequentially. The multiplicity of infection ("MOI"), including in the presence of rapamycin, may be

varied to determine an optimum MOI for a given cell type, density and culture technique, and a particular rapamycin concentration, using a positive control cell culture that is known to be infected upon exposure to MV.

[00153] The amount of rapamycin, and the timing of addition of rapamycin and Myxoma virus to the cultured cells may be varied depending on cell type, method of culturing and strain of virus. Such parameters can be readily tested and adjusted with minimal testing using routine methods.

[00154] Infectivity of the cultured cells by MV, including in the presence of rapamycin, may be determined by various methods known to a skilled person, including the ability of the MV to cause cell death. It may also involve the addition of reagents to the cell culture to complete an enzymatic or chemical reaction with a viral expression product. The viral expression product may be expressed from a reporter gene that has been inserted into the MV genome.

[00155] In one embodiment the MV may be modified to enhance the ease of detection of infection state. For example, the MV may be genetically modified to express a marker that can be readily detected by phase contrast microscopy, fluorescence microscopy or by radioimaging. The marker may be an expressed fluorescent protein or an expressed enzyme that may be involved in a colorimetric or radiolabelling reaction. In another embodiment the marker may be a gene product that interrupts or inhibits a particular function of the cells being tested.

[00156] The invention is further illustrated by the following non-limiting examples.

EXAMPLES

Example 1: Infection of Mouse and Human Cell Lines with Myxoma Virus

Virus Strains

[00157] Viral strains used include wildtype MV, MV modified to express either green fluorescence protein ("GFP") or β -galactosidase ("LacZ"), and killed ("dead") MV. Viruses were prepped and titred using standard techniques.

Cell Strains

[00158] Mouse experiments were performed using mouse embryo fibroblasts (“MEFs”) derived from a wild-type mouse, and from the following mouse knockouts: IFN α / β receptor homozygous knockout; STAT1 homozygous knockout; PKR heterozygous; RNaseL heterozygous knockout; Mx1 heterozygous knockout; triple PKR/RNaseL/Mx1 homozygous knockout.

[00159] Human experiments were performed on BGMK control cells and human tumour cell lines HT29, HOP92, OVCAR4, OVCAR5, SK-MEL3, SK-MEL28, M14, SKOV3, PC3, DU145, CAKI-1, 786-0, T47D, MDAMB 435, SF04, U87, A172, U373, Daoy and D384 as described in Stojdl et al., *Cancer Cell* (2003) 4: 263-275.

Methods

[00160] Generally, assays and experiments were performed as described in Lalani et al. *Virology* (1999) 256: 233-245; Johnston et al. *J Virology* (2003) 77(13): 7682-7688; and Sypula et al. *Gen Ther Mol Biol* (2004) 8: 103.

[00161] For the *in vivo* mouse studies, nude mice were implanted with intracranial human gliomas U87. 15 days after implantation, mice were intratumourally injected with live or dead MV GFP, at a titre of 5×10^6 , or mock-infected. 72 hours post-infection, animals were sacrificed, the brains removed, embedded in OCT (Optimal Cutting Temperature compound), and frozen sections were cut. Myxoma-GFP was visualized in whole brain sections by fluorescence microscopy. Sections were then fixed and stained with H&E (hemotoxylin and eosin) to visualize the tumor.

[00162] For human tumour cell assays, the tumours were trypsonized and plated immediately after surgery and infected with virus the next day at an MOI of 0.1, 1.0 or 10. Data was gathered regarding cytotoxicity and viral expression using phase microscopy and fluorescent microscopy, respectively, at 24 and 48 hours post-infection. Assays using the yellow tetrazolium salt MTT were performed to quantify the % cell survival (as a percentage of cells surviving mock infection) at 48, 72 or 96 hours post-infection.

[00163] Human pediatric medulloblastoma cell lines, Daoy and D384, were infected with 10 M.O.I. of Myxoma-GFP. 72 hours after infection, cell viability was measured using MTT.

Results: Infections of Mouse Cell Lines

[00164] Previous research showed that some clones of mouse 3T3 cells transfected with chemokine receptors were infectable by Myxoma virus while other clones were not. To investigate whether Myxoma virus tropism in other mouse cells was dependent on any particular receptors, we exploited primary mouse embryo fibroblasts (MEFs) from wild-type (WT) mice and various gene knock-outs.

[00165] Since IFNs play a key role in mounting anti-viral responses, we hypothesized that the restrictive phenotype was related to the “antiviral state” mediated by IFN. Disruption of the chain of events of the IFN system, neutralizing circulating IFN with antibodies or generating IFN receptor negative mice, or mice with deleted genes in the intracellular pathway of signal transmission, would severely compromise the host’s resistance to the Myxoma virus which typically does not infect normal mouse cells.

[00166] In order to test this hypothesis we needed to demonstrate if the non-infectivity of Myxoma virus in the nonpermissive cells was due to the antiviral action of IFNs. Various MEF cell types having knock-outs of one or more proteins involved in intracellular IFN signaling response were tested for the effect of MV infection on the IFN pathway.

[00167] Experiments performed on primary MEFs demonstrated that wildtype (“WT”) MEFs are not infectable by Myxoma virus. The MEFs are fully infectable by Myxoma virus when the IFN pathway is blocked by neutralizing antibody to IFN α/β (Figure 2). However, MEFs exposed to neutralizing antibodies to IFN γ remained nonpermissive. This outlined the importance of IFN α/β but not IFN γ in creating a permissive environment for Myxoma virus to infect MEFs *in vitro*. Different intracellular signaling pathways for IFN α/β and IFN γ have been identified in the literature. However, both IFN α/β and IFN γ likely play an important role in infected hosts, unlike cultured fibroblasts. We predict that human tumors deficient in either IFN α/β and/or IFN γ pathway will be susceptible to Myxoma virus infection *in vivo*.

[00168] We examined the activity of STAT1 and STAT2 in nonpermissive WT MEFs that were infected with MV. The results shown in Figure 3 indicated that STAT1

and STAT2 were activated. Further study showed that STAT3, STAT4 and STAT5 are not activated (Figure 4).

[00169] In order to confirm the importance of the IFN α/β intracellular pathway in maintaining a nonpermissive state in MEFs, genetic deletion studies were performed to provide disruptions in the IFN α/β receptors and in the intracellular cascade. Genetic deletion of IFN receptors or JAK1 or STAT1 was performed. MV was used to infect WT MEFs, IFN α/β R $^{-/-}$ MEFs and STAT1 $^{-/-}$ MEFs. IFN α/β R $^{-/-}$ MEFs and STAT1 $^{-/-}$ MEFs were permissive to MV demonstrating the IFN α/β and STAT1 signalling cascades are critical for MV infection (Figure 5).

[00170] Protein Kinase R (PKR) is an enzyme induced in a wide variety of cells by IFN α/β . This kinase, in the presence of dsRNA, undergoes autophosphorylation and then phosphorylates several cellular proteins including eukaryotic protein synthesis initiation factor (eIF-2 α) whose phosphorylation can induce an inhibition of protein translation and apoptosis. PKR is also indicated in the activation of RNaseL. We examined the activation of PKR in nonpermissive MEFs following MV infection. PKR is not phosphorylated in nonpermissive MEFs in which the antiviral state is well established (Figure 6). Furthermore MV infection inhibits PKR phosphorylation (Figure 7). In addition, PERK (PKR-like, ER kinase) is not phosphorylated in the primary WT MEFs following Myxoma virus infection (Figure 8).

[00171] MV was used to infect MEFs with single gene knockouts of PKR, RNaseL or Mx1 (Figure 9). It was discovered that PKR, RNaseL and Mx1 are nonessential for maintaining nonpermissiveness for Myxoma virus infection. To further confirm the nonessential role of PKR, RNaseL and Mx1 a Triple knockout of PKR $^{-/-}$, RNase L $^{-/-}$ and Mx1 $^{-/-}$ in MEFs was performed. A PKR $^{-/-}$, RNase L $^{-/-}$ and Mx1 $^{-/-}$ triple knockout does not support Myxoma virus infection (Figure 10), however MEFs with a triple KO of PKR, RNaseL and Mx1 treated with a neutralizing antibody to Interferon α/β becomes permissive to Myxoma virus infection (compare Figures 10 and 11). These experiments demonstrate that PKR, RNaseL and Mx1 are not essential in mediating the nonpermissiveness of MEFs to MV.

[00172] Further studies were performed to examine the activation of eIF-2 α and

PKR in nonpermissive wildtype MEFs and permissive IFN α/β R $^{-/-}$ MEFs and STAT1 $^{-/-}$ MEFs after MV infection. After MV infection, eIF-2 α is phosphorylated in nonpermissive and permissive MEFs although PKR is not phosphorylated in either case (Figure 12). This demonstrates that without the involvement of PKR and PERK, the antiviral state is mediated by another pathway that causes eIF2 α phosphorylation.

[00173] STAT1 is both serine- and tyrosine-phosphorylated following Myxoma infections in nonpermissive PKR, RNaseL and Mx1 Triple KO MEFs (Figure 13). Subcellular localization of tyrosine-phosphorylated STAT1 in nonpermissive PKR $^{-/-}$ + RNaseL $^{-/-}$ + Mx1 $^{-/-}$ MEFs following Myxoma virus infection is also shown (Figure 14).

[00174] In summary, these results indicate that a parallel PKR/PERK-independent antiviral pathway involving IFN/STAT1 is critical for poxvirus tropism. Furthermore, eIF2 α phosphorylation is the best marker for the antiviral action by INF.

Results: Human Tumour Studies

[00175] We studied the ability of MV to infect human tumour cells in an *in vivo* system. Nude mice were injected with human glioma cells, and subsequently developed intracranial gliomas. Live virus was able to infect these human tumours cells but did not infect surrounding cells (Figure 15). The localization of fluorescent signal from GFP to the tumour is depicted in Figure 16.

[00176] Given that many human tumours are non-responsive to interferon, and that the tumour cells do not have normal IFN signaling cascades compared to those found in normal human cells, studies were performed to investigate the effect of Myxoma virus on human tumours. The results are summarized below.

[00177] Initially, Myxoma virus was used to study the infectivity and cytolytic effects on various control and human tumour cell lines: BGМК, HT29, HOP92, OVCAR4, SK-MEL3, and SK-MEL28. MV demonstrated various infectivity and cytolytic results: HT29 (Figure 17) HOP92 (Figure 18), OVCAR4 (Figure 19) SK-MEL3 (Figure 20), SK-MEL28 (Figure 21) and BGМК (Figure 22).

[00178] Additional tumour cells were tested and Table 1 below classifies the various tumour types tested as permissive or non-permissive.

Table 1 Myxoma Virus Trophism for Human Tumour Cells

Cell Line	Cell Origin	Species	Permissive	Non-Permissive
BGMK	Kidney	Monkey	X	
RK-13	Kidney	Rabbit	X	
RL5	T-Lymphocyte	Rabbit	X	
HOS	Osteosarcoma	Human	X	
PC3	Prostate cancer	Human	X	
Caki-1	Renal cancer	Human	X	
HCT116	Colon cancer	Human	X	
786-0	Renal cancer	Human	X	
SK-OV-3	Ovarian cancer	Human	X	
ACHN	Renal cancer	Human	X	
HOP92	Lung cancer	Human	X	
SK-MEL3	Melanoma	Human	X	
SK-MEL28	Melanoma	Human	X	
OVCAR4	Ovarian cancer	Human	X	
OVCAR5	Ovarian cancer	Human	X	
DU145	Prostate cancer	Human	X	
A498	Renal cancer	Human	X	
T47D	Breast cancer	Human	X	
Colo205	Colon cancer	Human		X
HT29	Colon cancer	Human		X
MDAMB435	Breast cancer	Human		X
M14	Melanoma	Human		X
MCF7	Breast cancer	Human		X
SK-MEL5	Melanoma	Human		X

[00179] Various human tumour lines demonstrated varying responsiveness to infection with increasing concentrations of MV-LacZ. For example, U373 cells required higher virus titres to achieve the levels of cell killing achieved with lower virus titres in U87 (Figure 23 and Figure 24). Myxoma efficiently infected astrocytoma cells (Figure 25), and glioma cells (Figure 26). Myxoma was effective at 48 hours post-infection at killing human astrocytoma and pediatric medulloblastoma cells (Figure 27 and 28).

Example 2: Effect of Rapamycin on the Kinetics of Myxoma Virus Replication in Restrictive Cell Lines

Virus Strains

[00180] Viral strains used include wildtype MV ("vMyxLac"), and MV modified to have the MT-5 gene knocked out ("vMyxLacT5-"). Viruses were prepped and titred using standard techniques.

Cell Strains

[00181] Human experiments were performed on BGMK primate control cells, RK-13 rabbit control cells and normal human fibroblasts A9, restrictive human tumour cell lines 786-0 (renal), ACHN (renal), HCT116 (colon), MCF-7 (breast), MDA-MB-435 (breast), M14 (melanoma) and COLO205 (colon).

Methods

[00182] Generally, assays and experiments were performed as described in Lalani et al. *Virology* (1999) 256: 233-245; Johnston et al. *J Virology* (2003) 77(13): 7682-7688; and Sypula et al. *Gen Ther Mol Biol* (2004) 8: 103.

[00183] For viral growth curves, cells were grown *in vitro* in a monolayer, and pretreated with 20 nM rapamycin or a control (1:5000 dilution of DMSO) prior to infection with virus.

[00184] Samples of indicated cell lines infected with the indicated viral strain were collected at 72 hours post infection and lysed. The virus contained within the cell lysates was titrated and used to infect BGMK monolayers. At 48 hours post infection, cells were fixed and stained using X-gal.

Results

[00185] Myxoma virus has been previously demonstrated by the inventors to be able to infect and replicate in many types of human tumor cells (Sypula et al. (2004) *Gene Ther. Mol. Biol.* 8:103). This rabbit specific virus can preferentially infect a majority (approximately 70%) of human cancer cell lines from the NCI reference collection. In addition, the host range gene M-T5 was found to play a critical role during Myxoma virus infection of many of these cell lines.

[00186] In the present investigation of potential intracellular molecules that may

be affecting the ability of Myxoma to selectively replicate within human tumour cells, the effect of rapamycin was tested.

[00187] As seen in Figure 29, the ability of Myxoma virus to replicate and spread following a low multiplicity of infection (MOI) was performed using a multistep growth curve, using BGМК (control primate cell line); RK-13 and RL5 (control rabbit cell lines); 4T1 and B16F10 (mouse cancer cell lines); HOS and PC3 (permissive human cancer cell lines); 786-0, HCT116 and ACHN (restrictive human cancer cell lines); MCF-7, M14 and COLO205 (abortive human cancer cell lines). Both wild type vMyxLac and the M-T5 knock out virus vMyxT5KO were tested to investigate the ability of both viruses to infect and spread throughout the monolayer in the presence and absence of pre-treatment with rapamycin. Virus titre was assessed by foci formation on BGМК cells. Cells were pretreated with 20 nM rapamycin or appropriate vehicle control (1:5000 dilution of DMSO) for 6 hours before infection.

[00188] As demonstrated, rapamycin has no effect on control BGМК cells, nor on either of the rabbit cell lines tested, including the RL-5 cells, which are non permissive for the MT-5 knock out virus. However, rapamycin does enhance the replication of myxoma virus in mouse tumour cell lines, and marginally in permissive (Type I) cell lines, such as PC-3. Rapamycin has less of an effect on highly permissive cells such as HOS cells, likely due to the fact that such cell lines are already maximally permissive for the Myxoma virus. The greatest effect with rapamycin was observed in the restrictive (Type II) cell lines (786-0, HCT116 and ACHN), which are permissive for wildtype virus but non-permissive for the vMyxT5KO strain. Some effect was seen even in abortive (Type III) cell lines MCF-7 and COLO205, although not in abortive cell line M14.

[00189] Samples of the BGМК and 786-0 infected cells were then collected and lysed, and the isolated virus was used to infect monolayers of BGМК cells (Figure 30). Virally infected cells were visualized using X-Gal staining.

[00190] Pretreatment of tumour cells that are “restrictive” for Myxoma infection, i.e. those cells that permit the replication of the wild type Myxoma virus but not the MT-5 knock-out virus, with rapamycin resulted in a restoration of the ability of Myxoma virus to replicate in these cancer cell lines, which include renal, colon and ovarian cancer cell lines

(Figures 29 and 30).

[00191] In addition, the treatment with rapamycin enhanced the ability of the wild type virus to replicate in these same cells, but not control rabbit or primate cells. These results indicate that rapamycin acts to enhance Myxoma virus infection. In addition, rapamycin appears to influence the ability of cancer cells that are poorly infectable by this virus to permit virus replication.

[00192] Subsequent experiments examined the effect of rapamycin treatment on human tumour cells that could not support wild type Myxoma virus infection (Figure 31). The pretreatment had little effect on control primate cells or normal human fibroblasts, yet could enhance virus infectivity in several cell lines, including the breast cancer cell line MCF-7. As several of the human tumour cell lines remained resistant to rapamycin treatment, as well as the control cell lines, it is unlikely that rapamycin treatment could permit Myxoma virus to productively infect non-transformed tissue.

Example 3: Myxoma virus M135KO variant as an improved oncolytic virus candidate

M135R is expressed from Myxoma virus as an early gene

[00193] Myxoma virus encodes a protein (M135R) identified from the sequencing of the MV genome (Cameron et al. *Virology* (1999) 264: 298-318) predicted to mimic the host IFN α/β receptor and prevent IFN α/β from triggering a host anti-viral response (Barrett et al. *Seminars in Immunology* (2001) 13:73-84). This prediction is based on sequence homology to the viral IFN α/β receptor homolog from vaccinia virus (B18R), which virus has been demonstrated to employ such an immune evasion strategy (Symons et al. *Cell* (1995) 81:551-560). However M135R is only half the size of VV B18R and all other IFN α/β -R homologs sequenced from poxviruses, and in all cases aligns only to the amino terminus half of poxviral IFN α/β -R homologs. Figure 32 indicates the predicted structure and sequence similarity between M135R from MV and B18R from VV. Only the first 179 amino acid residues of B18R are shown in the sequence alignment. Table 2 indicates the % identity between M135R and the indicated poxviral IFN α/β -R homologs. Numbers above the diagonal represent % identity and numbers below the diagonal represent % similarity between any two species. The numbers in brackets across the top represent the number of amino acids in the putative proteins. Comparison was done

between the predicted full length copy of M135R (178 amino acids) and the first 178 residues of each homolog only.

Table 2 Comparison of M135R to Other Poxviral Homologs

<i>species</i>	% Identity									
	<i>Myxoma</i> (178)	<i>Vaccinia</i> (351)	<i>Variola</i> (354)	<i>Monkeypox</i> (352)	<i>Cowpox</i> (351)	<i>Ectromelia</i> (358)	<i>Camelpox</i> (355)	<i>YLDV</i> (351)	<i>Swinepox</i> (344)	<i>LSDV</i> (360)
Myxoma	-----	24	21	24	23	21	22	20	18	17
Vaccinia	39	-----	80	93	90	84	79	20	23	25
Variola	36	88	-----	79	87	88	79	19	24	24
Monkeypox	38	95	87	-----	86	83	78	20	20	26
Cowpox	38	93	93	91	-----	92	88	21	23	25
Ectromelia	35	89	93	87	94	-----	87	17	21	25
Camelpox	34	86	94	85	92	93	-----	17	23	24
YLDV	38	37	37	37	38	34	34	-----	23	28
Swinepox	32	39	39	36	39	35	37	38	-----	25
LSDV	32	39	39	41	38	39	36	43	38	-----

[00194] Peptides against predicted immunogenic regions of M135R were synthesized and used to generate polyclonal antibodies in rabbits that were used in western blot analysis, immuno-precipitations and immuno-fluorescence. Immunoblotting confirmed that M135R is synthesized as an early gene whose expression can be detected as early as three hours post infection (Figure 33A; lane 1: mock infected BGMK cells; lanes 2-6: BGMK cells infected with vMyxLau 0, 3, 6, 18 and 36 hours post infection, respectively). Treatment of infected cells with AraC indicates that synthesis of M135R was not altered by inhibition of late protein expression and is therefore an early gene (Figure 33B). However treatment with tunicamycin indicates that M135R is N-linked glycosylated, likely at the single site predicted from the sequence (Figure 33B). Monensin treatment suggests that there is no O-linked glycosylation. For the results shown in Figure 33, BGMKs were infected at an moi of 10 with Myxoma virus. Cells were treated with AraC at a concentration of 40 µg/ml, tunicamycin at 1 µg/ml and monensin at 1 µg/ml, or were untreated, at the times indicated. M135R was detected with a peptide antibody.

M135R encodes a signal sequence but is not secreted

[00195] Sequence analysis of M135R indicates the presence of a predicted signal sequence (Figure 32B). However there is also a predicted transmembrane domain at the carboxy terminus (Figure 32B). Immunoblots of supernatants from infected BGMK cells

indicate that M135R is not secreted. However, M135R is easily detected in whole cell lysates (Figure 33). To test whether the signal sequence functioned to drive M135R to the cell surface, we deleted the transmembrane domain and cloned the mutant into a baculovirus expression system. Comparison of AcM135R and Ac135 Δ TM infected supernatants indicated that full length M135R is found in the cell lysate there is no evidence of secretion. In contrast Ac135 Δ TM is secreted and confirms that the signal sequence functions to drive M135R into the extracellular environment (data not shown).

M135R protein localizes to the surface of infected cells

[00196] The observation that M135R has a functional signal sequence as well as a transmembrane domain prompted us to test the localization of M135R. Two pieces of evidence indicate that M135R localizes to the cell surface. First, when BGMKs were seeded onto glass coverslips and infected with vMyxLau (moi of 10) for 24 hours then M135R was detected by immunostaining with affinity purified anti-M135R followed by FITC-conjugated secondary antibody (Figure 34A). M135R staining pattern indicates localization to the cell surface of infected cells. vMyxLau is a true wildtype strain of Myxoma virus which has not been altered by insertion of the β -gal or EGFP gene.

[00197] The second piece of evidence for cell surface localization M135R follows biotinylation of cell surface proteins of GHOST cells infected with either vMyxgfp or vMyx135KO. Twenty-four hours post infection cell lysates were prepared. Streptavidin agarose beads were mixed with 500 μ g of total cellular protein from cell lysates for 45 minutes. The beads were washed and separated on a 15% PAGE-SDS gel and then probed with anti-M135R. 50 μ g of total protein from the infected cell lysates were run as controls. Immunoprecipitation of biotinylated surface proteins indicates that m135R is at the surface of infected cells (Figure 34B).

M135R is non-essential for Myxoma virus replication in vitro

[00198] To test the ability of M135R to act as a virulence factor we constructed a recombinant virus in which M135R was deleted and replaced by a cassette encoding EGFP and gpt under VV early/late promoters (460 nucleotides, or 86% of the orf was deleted). The cloning strategy and cassette is shown in Figure 35A. The recombinant was plaque-purified by selecting virus clones expressing EGFP. The purity of the recombinant

was confirmed by PCR (Figure 35B; Lane 1 is the 1Kb plus DNA ladder, Lane 2 and 3 are PCR products from two purified vMyx135KO clones. The PCR product represents the region into which the M135R coding region has been deleted and the EGFP/gpt marker has been inserted. Lane 2 is plaque 1 and Lane 3 is plaque 2. Lane 4 represents the same region and covers the native, uninterrupted M135R locus.). Immunoblotting of BGMK cells infected with either vMyxLau or vMyx135KO confirmed that vMyx135KO had lost M135R expression (Figure 35C; time course of expression of M135R: Lane 1 is uninfected BGMK cells. Lanes 2-6 represent BGMK cells infected with vMyxLau at times 0 (lane 2), 3 (lane 3), 6 (lane 4), 18 (lane 5), and 36 hours post infection (lane 6). Lanes 7 and 8 represent BGMK cells infected with vMyx135KO at 6 (lane 7) and 18 (lane 8) hours post infection. Lane 9 is a positive control with M135R expressed in AcNPV.).

[00199] Single step growth curves were used to test the ability of vMyx135KO to replicate in BGMK cells. BGMK cells were infected with vMyxgfp or vMyx135KO at an moi of 5 and cells were collected at the times indicated. Virus titres were determined on BGMK cells. There was no difference in the replication pattern between vMyxgfp and vMyx135KO (Figure 36). These results indicate that M135R is not required for replication *in vitro*.

[00200] During our studies of the ability of another gene of Myxoma to influence Myxoma replication in rabbit primary embryo fibroblasts (REFs), we used vMyx135KO as a knockout control and observed a curious phenomenon. Infection of the REFs with vMyxgfp resulted in a normal focus of infection however vMyx135KO produced a plaque-like zone of infection (Figure 37). When we tested other cells to confirm this phenotype we were able to replicate the plaque formation in other rabbit fibroblasts (HIG-82, Figure 38) and human primary fibroblasts (ccd922-sk, Figure 39).

M135R is a critical virulence factor for pathogenesis in rabbits

[00201] We next tested the ability of vMyx135KO to produce myxomatosis in lab rabbits. In contrast to the animals injected with vMyxLau or vMyxgfp which developed normal myxomatosis and had to be euthanized between days 9 and 10 post injection, the rabbits injected with vMyx135KO recovered completely (Table 3). To confirm that loss of M135R caused the attenuation of vMyx135KO we generated a revertant virus in which M135R was restored and we tested the ability of this revertant (vMyx135REV) to restore

the ability to produce myxomatosis. All four treated groups of rabbits responded in a similar manner for the first six days following injection of the respective viruses (Table 3). We observed a large, red, raised lesion at the site of injection in all treatment groups by 4 days post infection. However beginning at day 6 and continuing over the next 3-4 days the differences between the different viruses became evident. Those animals injected with the wildtype or revertant virus had numerous secondary lesions in the ears, eyes and nose which were not observed in the animals injected with vMyx135KO (Table 3). We conclude that loss of M135R drastically attenuated MV in animal models and indicates that M135R is a critical virulence factor.

Table 3 Pathogenesis of vMyx135KO Compared to Wildtype Controls

Clinical Signs	Observations and Time of onset (number + days indicates first appearance in days post injection)		
	Lausanne (4 animals)	vMyx135KO (6 animals)	vMyx135REV (3 animals)
inoculation site	2 days: red, visible slightly raised 4 days: red, dark centre	4 days: 11-16 mm red, raised, dark centre	3 days: small red lump, slightly raised
satellites	4 days	6 days: just beginning, over course of infection very few observed	6 days: 5-10 visible increasing to 30-40 satellites visible by day 8
conjunctival inflammation	none observed	9 days: single rabbit discharge from eye	none observed
anogenital edema	7 days: swelling	7 days: redness, swelling	
secondary lesions	6-7 days: first around eyes then ears	7 days: few small red spots not yet lesions, ears eyes	6 days: first observed as red areas on eyelids, clearly lesion by day 7
respiratory difficulty	little or none	little or none	little or none
lesion regression		11 days: 25mm, black, scabby - satellites losing colour and becoming scabby 13 days: scab beginning to separate from healthy tissue	
	- two animals euthanized day 9 - two animals euthanized day 10	all animals recovered	three animals euthanized day 10

[00202] The temperature of rabbits was taken daily for the three days preceeding the study. This was considered the baseline body temperatures of the animals. We

continued to take the temperatures daily of each animal for the duration of the study. However there was no difference in body temperature between the treatment groups (Figure 40). This suggests that M135R does not play a role in the febrile response of infected animals.

M135R does not bind or inhibit rabbit IFN α/β

[00203] The sequence of M135R is similar to the vaccinia B18R, an IFN α/β receptor mimic. We tested the ability of M135R to bind rabbit type 1 IFN. We first iodinated rabbit IFN (5 μ g, using Iodobeads) and tested the ability of vMyx135KO infected cells to bind 125 I-rabbit IFN in comparison to cells infected with vMyxgfp (moi of 10). Cells were collected, washed and counted in a gamma counter. Deletion of M135R did not affect IFN α/β binding to infected cells and we did not observe any difference in the amount of IFN bound to the cell surface of either RK13 or BGMK cells (Figure 41). As well, treatment of RK13 or BGMK cells with exogenous rabbit type1 IFN did not affect infection of cells by vMyx135KO (Figure 42; cells were seeded in 12 well dishes and infected with the indicated virus at an moi of 0.01; fluorescent foci were counted 72-96 hours post infection; 200 units of rabbit IFN α/β was either added 24 hours prior to infection or cells were untreated). This same result was observed when cells were pretreated 24 h before infection to induce an anti-viral state in the cell. We did not notice any significant difference in the foci formed following infection in either RK13 or BGMK cells (data not shown). This phenomenon was also true if cells were treated with human IFN α/β (data not shown). As well, we were unable to observe any binding when Ac135 Δ TM supernatants were applied to rabbit IFN α/β adhered to a BIAcore chip (data not shown).

Example 3: Molecular consequences of inhibiting mTOR in the context of myxoma virus infection

[00204] Western Blot analysis (Figure 43) was performed using cell lysates from 786-0 cells, a Type II cancer cell line where rapamycin enhances myxoma virus infection. Lysates were collected 16 hours post infection with either vMyxLac or vMyxT5KO at an MOI of 3, or without virus infection. Indicated lanes contain protein from cells that were pretreated with 20 nM rapamycin (designated R) or appropriate vehicle control (1:5000 dilution of DMSO, designated D) for 6 hours before infection. The blots were probed

using primary antibodies directed against the indicated proteins.

[00205] As demonstrated, myxoma virus infection affects many of the signaling pathways that converge on mTOR, the physiologic target of rapamycin. In the context of infection with either wild type (vMyxLac) or MT-5 deficient (vMyxT5KO) virus, where rapamycin has a beneficial effect on virus replication, global effects are observed in many of these signaling molecules that would not be predictable based on treatment with rapamycin alone (see mock infected lanes). These effects include an increase in the kinase activity of AKT-1, Raf-1, GSK-3 β and mTOR itself, as well as a decrease in the kinase activity of PTEN and p70S6K. This data indicate that these pathways are likely to play a role in myxoma virus permissiveness in human cancer cells lines.

[00206] As can be understood by one skilled in the art, many modifications to the exemplary embodiments described herein are possible. The invention, rather, is intended to encompass all such modification within its scope, as defined by the claims.

[00207] Although various embodiments of the invention are disclosed herein, many adaptations and modifications may be made within the scope of the invention in accordance with the common general knowledge of those skilled in this art. Such modifications include the substitution of known equivalents for any aspect of the invention in order to achieve the same result in substantially the same way. All technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art of this invention, unless defined otherwise.

[00208] All reference cited herein are fully incorporated by reference.

WHAT IS CLAIMED IS:

1. A method for inhibiting a cell that has a deficient innate anti-viral response comprising administering to the cell an effective amount of a combination of Myxoma virus and rapamycin.
2. The method of claim 1 wherein the cell is non-responsive to interferon.
3. The method of claim 2 wherein the cell exhibits abnormal interferon signalling.
4. The method of claim 3 wherein the cell is a human cancer cell.
5. The method of claim 4 wherein the Myxoma virus is a wild-type virus.
6. The method of claim 4 wherein the Myxoma virus is genetically modified.
7. The method of claim 6 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
8. The method of claim 4 wherein the cell is lung cancer cell, melanoma cell, ovarian cancer cell, prostate cancer cell, renal cancer cell, glioma cell or astrocytoma cell.
9. The method of claim 1 wherein the cell is a human cell chronically infected with a virus.
10. A method for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response, comprising administering to a patient in need thereof an effective amount of a combination of Myxoma virus and rapamycin.
11. The method of claim 10 wherein the disease state is cancer.
12. The method of claim 11 wherein the cancer is a solid tumour, hematopoietic cell cancer, colon cancer, lung cancer, kidney cancer, pancreas cancer, endometrial cancer, thyroid cancer, oral cancer, ovarian cancer, laryngeal cancer, hepatocellular cancer, bile duct cancer, squamous cell carcinoma, prostate cancer, breast cancer, cervical cancer, colorectal cancer or melanoma.

13. The method of claim 11 wherein the cancer is lung cancer, melanoma, ovarian cancer, prostate cancer, renal cancer, glioma or astrocytoma.
14. The method of claim 13 wherein the patient is a human.
15. The method of claim 14 wherein the Myxoma virus is a wild-type virus.
16. The method of claim 14 wherein the Myxoma virus is genetically modified.
17. The method of claim 16 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
18. The method of claim 14 wherein the virus and the rapamycin are administered to the site of the cancer by injection.
19. The method of claim 14 wherein the virus and the rapamycin are administered systemically.
20. The method of claim 10 wherein the disease state is a chronic viral infection.
21. Use of an effective amount of a combination of Myxoma virus and rapamycin for inhibiting a cell that has a deficient innate anti-viral response.
22. Use of an effective amount of a combination of Myxoma virus and rapamycin in the manufacture of a medicament for inhibiting a cell that has a deficient innate anti-viral response.
23. The use of claim 21 or 22 wherein the cell is non-responsive to interferon.
24. The use of claim 23 wherein the cell exhibits abnormal interferon signalling.
25. The use of claim 24 wherein the cell is chronically infected with a virus.
26. The use of claim 24 wherein the cell is a human cancer cell.
27. The use of claim 26 wherein the cell is lung cancer cell, melanoma cell, renal cancer cell, glioma cell or astrocytoma cell.
28. The use of any one of claims 20 to 27 wherein the Myxoma virus is a wild-type

virus.

29. The use of any one of claims 20 to 27 wherein the Myxoma virus is genetically modified.

30. The use of claim 29 wherein the Myxoma virus is genetically modified to express a therapeutic gene.

31. Use of an effective amount of a combination of Myxoma virus and rapamycin for treating a disease state in a patient, wherein the disease state is characterized by the presence of cells that have a deficient innate anti-viral response.

32. Use of an effective amount of a combination of Myxoma virus and rapamycin in the manufacture of a medicament for treating a disease state in a patient, wherein the disease state is characterized by the presence of cells that have a deficient innate anti-viral response.

33. The use of claim 31 or 32 wherein the disease state is a cancer.

34. The use of claim 33 wherein the cancer is a solid tumour, hematopoietic cell cancer, colon cancer, lung cancer, kidney cancer, pancreas cancer, endometrial cancer, thyroid cancer, oral cancer, ovarian cancer, laryngeal cancer, hepatocellular cancer, bile duct cancer, squamous cell carcinoma, prostate cancer, breast cancer, cervical cancer, colorectal cancer or melanoma.

35. The use of claim 33 wherein the cancer is lung cancer, melanoma, ovarian cancer, prostate cancer, renal cancer, glioma or astrocytoma.

36. The use of claim 31 or 32 wherein the disease state is a chronic viral infection.

37. The use of any one of claims 31 to 36 wherein the patient is a human.

38. The use of any one of claims 31 to 37 wherein the Myxoma virus is a wild-type virus.

39. The use of any one of claims 31 to 37 wherein the Myxoma virus is genetically modified.

40. The use of claim 39 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
41. A pharmaceutical composition comprising Myxoma virus and rapamycin.
42. The pharmaceutical composition of claim 41 further comprising an additional therapeutic agent.
43. The pharmaceutical composition of claim 42 wherein the additional therapeutic agent is a chemotherapeutic agent.
44. A kit comprising Myxoma virus, rapamycin and instructions for inhibiting a cell that has a deficient innate anti-viral response or for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response.
45. A method for detecting a cell that has a deficient innate anti-viral response, comprising exposing a population of cells to a combination of Myxoma virus and rapamycin; allowing the virus to infect a cell that has a deficient innate anti-viral response; and determining the infection of any cells of the population of cells by the Myxoma virus.
46. The method of claim 45 wherein the population of cells is in a patient; said exposing comprises administering to the patient the combination of Myxoma virus and rapamycin, the Myxoma virus being modified to express a detectable marker; and said determining comprises detecting a cell expressing the detectable marker in the patient.
47. The method of claim 45 wherein the population of cells is in culture.
48. A method for inhibiting a cell that has a deficient innate anti-viral response comprising administering to the cell an effective amount of Myxoma virus that does not express functional M135R, optionally in combination with an effective amount of rapamycin.
49. The method of claim 48 wherein the cell is non-responsive to interferon.
50. The method of claim 49 wherein the cell exhibits abnormal interferon signalling.
51. The method of claim 50 wherein the cell is a human cancer cell.

52. The method of claim 51 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
53. The method of claim 51 wherein the cell is lung cancer cell, melanoma cell, ovarian cancer cell, prostate cancer cell, renal cancer cell, glioma cell or astrocytoma cell.
54. The method of claim 48 wherein the cell is a human cell chronically infected with a virus.
55. A method for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response, comprising administering to a patient in need thereof an effective amount of Myxoma virus that does not express functional M135R, optionally in combination with an effective amount of rapamycin.
56. The method of claim 55 wherein the disease state is cancer.
57. The method of claim 56 wherein the cancer is a solid tumour, hematopoietic cell cancer, colon cancer, lung cancer, kidney cancer, pancreas cancer, endometrial cancer, thyroid cancer, oral cancer, ovarian cancer, laryngeal cancer, hepatocellular cancer, bile duct cancer, squamous cell carcinoma, prostate cancer, breast cancer, cervical cancer, colorectal cancer or melanoma.
58. The method of claim 56 wherein the cancer is lung cancer, melanoma, ovarian cancer, prostate cancer, renal cancer, glioma or astrocytoma.
59. The method of claim 58 wherein the patient is a human.
60. The method of claim 59 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
61. The method of claim 56 wherein the virus, and optionally the rapamycin, is administered to the site of the cancer by injection.
62. The method of claim 56 wherein the virus, and optionally the rapamycin, is administered systemically.
63. The method of claim 55 wherein the disease state is a chronic viral infection.

64. Use of an effective amount of Myxoma virus that does not express functional M135R, optionally in combination with an effective amount of rapamycin.
65. Use of an effective amount of Myxoma virus that does not express functional M135R, optionally in combination with an effective amount of rapamycin, for the manufacture of a medicament for inhibiting a cell that has a deficient innate anti-viral response.
66. The use of claim 64 or 65 wherein the cell is non-responsive to interferon.
67. The use of claim 66 wherein the cell exhibits abnormal interferon signalling.
68. The use of claim 67 wherein the cell is chronically infected with a virus.
69. The use of claim 67 wherein the cell is a human cancer cell.
70. The use of claim 69 wherein the cell is lung cancer cell, melanoma cell, renal cancer cell, glioma cell or astrocytoma cell.
71. The use of any one of claims 64 to 70 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
72. Use of an effective amount of Myxoma virus that does not express functional M135R, optionally in combination with an effective amount of rapamycin, for treating a disease state in a patient, wherein the disease state is characterized by the presence of cells that have a deficient innate anti-viral response.
73. Use of an effective amount of Myxoma virus that does not express functional M135R, optionally in combination with an effective amount of rapamycin, for the manufacture of a medicament for treating a disease state in a patient, wherein the disease state is characterized by the presence of cells that have a deficient innate anti-viral response.
74. The use of claim 72 or 73 wherein the disease state is a cancer.
75. The use of claim 74 wherein the cancer is a solid tumour, hematopoietic cell cancer, colon cancer, lung cancer, kidney cancer, pancreas cancer, endometrial cancer,

thyroid cancer, oral cancer, ovarian cancer, laryngeal cancer, hepatocellular cancer, bile duct cancer, squamous cell carcinoma, prostate cancer, breast cancer, cervical cancer, colorectal cancer or melanoma.

76. The use of claim 74 wherein the cancer is lung cancer, melanoma, ovarian cancer, prostate cancer, renal cancer, glioma or astrocytoma.
77. The use of claim 72 or 73 wherein the disease state is a chronic viral infection.
78. The use of any one of claims 72 to 77 wherein the patient is a human.
79. The use of any one of claims 72 to 78 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
80. A Myxoma virus that does not express functional M135R.
81. A pharmaceutical composition comprising Myxoma virus that does not express functional M135R and optionally rapamycin.
82. The pharmaceutical composition of claim 80 further comprising an additional therapeutic agent.
83. The pharmaceutical composition of claim 81 wherein the additional therapeutic agent is a chemotherapeutic agent.
84. A kit comprising Myxoma virus that does not express functional M135R, and optionally rapamycin, and instructions for inhibiting a cell that has a deficient innate anti-viral response or for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response.
85. A method for detecting a cell that has a deficient innate anti-viral response, comprising exposing a population of cells to a Myxoma virus that does not express functional M135R, optionally in combination with rapamycin; allowing the virus to infect a cell that has a deficient innate anti-viral response; and determining the infection of any cells of the population of cells by the Myxoma virus.
86. The method of claim 84 wherein the population of cells is in a patient; said

exposing comprises administering to the patient the Myxoma virus that does not express functional M135R and optionally the rapamycin, the Myxoma virus that does not express functional M135R being modified to express a detectable marker; and said determining comprises detecting a cell expressing the detectable marker in the patient.

87. The method of claim 85 wherein the population of cells is in culture.

**ARTICLE 19 RECEIVED BY THE INTERNATIONAL BUREAU ON 28
AUGUST 2006 (28.08.2006)**

54

WHAT IS CLAIMED IS:

1. A method for inhibiting a cell that has a deficient innate anti-viral response comprising administering to the cell an effective amount of a combination of Myxoma virus and rapamycin.
2. The method of claim 1 wherein the cell is non-responsive to interferon.
3. The method of claim 2 wherein the cell exhibits abnormal interferon signalling.
4. The method of claim 3 wherein the cell is a human cancer cell.
5. The method of claim 4 wherein the Myxoma virus is a wild-type virus.
6. The method of claim 4 wherein the Myxoma virus is genetically modified.
7. The method of claim 6 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
8. The method of claim 4 wherein the cell is lung cancer cell, melanoma cell, ovarian cancer cell, prostate cancer cell, renal cancer cell, glioma cell or astrocytoma cell.
9. The method of claim 1 wherein the cell is a human cell chronically infected with a virus.
10. A method for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response, comprising administering to a patient in need thereof an effective amount of a combination of Myxoma virus and rapamycin.
11. The method of claim 10 wherein the disease state is cancer.
12. The method of claim 11 wherein the cancer is a solid tumour, hematopoietic cell cancer, colon cancer, lung cancer, kidney cancer, pancreas cancer, endometrial cancer, thyroid cancer, oral cancer, ovarian cancer, laryngeal cancer, hepatocellular cancer, bile duct cancer, squamous cell carcinoma, prostate cancer, breast cancer, cervical cancer, colorectal cancer or melanoma.

13. The method of claim 11 wherein the cancer is lung cancer, melanoma, ovarian cancer, prostate cancer, renal cancer, glioma or astrocytoma.
14. The method of claim 13 wherein the patient is a human.
15. The method of claim 14 wherein the Myxoma virus is a wild-type virus.
16. The method of claim 14 wherein the Myxoma virus is genetically modified.
17. The method of claim 16 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
18. The method of claim 14 wherein the virus and the rapamycin are administered to the site of the cancer by injection.
19. The method of claim 14 wherein the virus and the rapamycin are administered systemically.
20. The method of claim 10 wherein the disease state is a chronic viral infection.
21. Use of an effective amount of a combination of Myxoma virus and rapamycin for inhibiting a cell that has a deficient innate anti-viral response.
22. Use of an effective amount of a combination of Myxoma virus and rapamycin in the manufacture of a medicament for inhibiting a cell that has a deficient innate anti-viral response.
23. The use of claim 21 or 22 wherein the cell is non-responsive to interferon.
24. The use of claim 23 wherein the cell exhibits abnormal interferon signalling.
25. The use of claim 24 wherein the cell is chronically infected with a virus.
26. The use of claim 24 wherein the cell is a human cancer cell.
27. The use of claim 26 wherein the cell is lung cancer cell, melanoma cell, renal cancer cell, glioma cell or astrocytoma cell.
28. The use of any one of claims 20 to 27 wherein the Myxoma virus is a wild-type

virus.

29. The use of any one of claims 20 to 27 wherein the Myxoma virus is genetically modified.

30. The use of claim 29 wherein the Myxoma virus is genetically modified to express a therapeutic gene.

31. Use of an effective amount of a combination of Myxoma virus and rapamycin for treating a disease state in a patient, wherein the disease state is characterized by the presence of cells that have a deficient innate anti-viral response.

32. Use of an effective amount of a combination of Myxoma virus and rapamycin in the manufacture of a medicament for treating a disease state in a patient, wherein the disease state is characterized by the presence of cells that have a deficient innate anti-viral response.

33. The use of claim 31 or 32 wherein the disease state is a cancer.

34. The use of claim 33 wherein the cancer is a solid tumour, hematopoietic cell cancer, colon cancer, lung cancer, kidney cancer, pancreas cancer, endometrial cancer, thyroid cancer, oral cancer, ovarian cancer, laryngeal cancer, hepatocellular cancer, bile duct cancer, squamous cell carcinoma, prostate cancer, breast cancer, cervical cancer, colorectal cancer or melanoma.

35. The use of claim 33 wherein the cancer is lung cancer, melanoma, ovarian cancer, prostate cancer, renal cancer, glioma or astrocytoma.

36. The use of claim 31 or 32 wherein the disease state is a chronic viral infection.

37. The use of any one of claims 31 to 36 wherein the patient is a human.

38. The use of any one of claims 31 to 37 wherein the Myxoma virus is a wild-type virus.

39. The use of any one of claims 31 to 37 wherein the Myxoma virus is genetically modified.

40. The use of claim 39 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
41. A pharmaceutical composition comprising Myxoma virus and rapamycin.
42. The pharmaceutical composition of claim 41 further comprising an additional therapeutic agent.
43. The pharmaceutical composition of claim 42 wherein the additional therapeutic agent is a chemotherapeutic agent.
44. A kit comprising Myxoma virus, rapamycin and instructions for inhibiting a cell that has a deficient innate anti-viral response or for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response.
45. A method for detecting a cell that has a deficient innate anti-viral response, comprising exposing a population of cells to a combination of Myxoma virus and rapamycin; allowing the virus to infect a cell that has a deficient innate anti-viral response; and determining the infection of any cells of the population of cells by the Myxoma virus.
46. The method of claim 45 wherein the population of cells is in a patient; said exposing comprises administering to the patient the combination of Myxoma virus and rapamycin, the Myxoma virus being modified to express a detectable marker; and said determining comprises detecting a cell expressing the detectable marker in the patient.
47. The method of claim 45 wherein the population of cells is in culture.
48. A method for inhibiting a cell that has a deficient innate anti-viral response comprising administering to the cell an effective amount of Myxoma virus that does not express functional M135R, optionally in combination with an effective amount of rapamycin.
49. The method of claim 48 wherein the cell is non-responsive to interferon.
50. The method of claim 49 wherein the cell exhibits abnormal interferon signalling.
51. The method of claim 50 wherein the cell is a human cancer cell.

52. The method of claim 51 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
53. The method of claim 51 wherein the cell is lung cancer cell, melanoma cell, ovarian cancer cell, prostate cancer cell, renal cancer cell, glioma cell or astrocytoma cell.
54. The method of claim 48 wherein the cell is a human cell chronically infected with a virus.
55. A method for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response, comprising administering to a patient in need thereof an effective amount of Myxoma virus that does not express functional M135R.
56. The method of claim 55, further comprising administering an effective amount of rapamycin to the patient.
57. The method of claim 55 or 56 wherein the disease state is cancer.
58. The method of claim 57 wherein the cancer is a solid tumour, hematopoietic cell cancer, colon cancer, lung cancer, kidney cancer, pancreas cancer, endometrial cancer, thyroid cancer, oral cancer, ovarian cancer, laryngeal cancer, hepatocellular cancer, bile duct cancer, squamous cell carcinoma, prostate cancer, breast cancer, cervical cancer, colorectal cancer or melanoma.
59. The method of claim 57 wherein the cancer is lung cancer, melanoma, ovarian cancer, prostate cancer, renal cancer, glioma or astrocytoma.
60. The method of claim 59 wherein the patient is a human.
61. The method of claim 59 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
62. The method of claim 55 or 56 wherein the administration is by injection at the site of the cancer.
63. The method of claim 55 or 56 wherein the administration is systemic.
64. The method of claim 55 wherein the disease state is a chronic viral infection.

65. Use of an effective amount of Myxoma virus that does not express functional M135R.
66. The use of claim 65, further in combination with an effective amount of rapamycin.
67. Use of an effective amount of Myxoma virus that does not express functional M135R for the manufacture of a medicament for inhibiting a cell that has a deficient innate anti-viral response.
68. The use of claim 67, wherein the medicament is formulated for administration in combination with an effective of rapamycin.
69. The use of any one of claims 65 to 68, wherein the cell is non-responsive to interferon.
70. The use of claim 69 wherein the cell exhibits abnormal interferon signalling.
71. The use of claim 70 wherein the cell is chronically infected with a virus.
72. The use of claim 70 wherein the cell is a human cancer cell.
73. The use of claim 72 wherein the cell is lung cancer cell, melanoma cell, renal cancer cell, glioma cell or astrocytoma cell.
74. The use of any one of claims 65 to 73 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
75. Use of an effective amount of Myxoma virus that does not express functional M135R for treating a disease state in a patient, wherein the disease state is characterized by the presence of cells that have a deficient innate anti-viral response.
76. The use of claim 75, wherein the use further comprises an effective amount of rapamycin.
77. Use of an effective amount of Myxoma virus that does not express functional M135R for the manufacture of a medicament for treating a disease state in a patient, wherein the disease state is characterized by the presence of cells that have a deficient

innate anti-viral response.

78. The use of claim 77, wherein the medicament is formulated for administration in combination with an effective amount of rapamycin.
79. The use of any one of claims 75 to 78 wherein the disease state is a cancer.
80. The use of claim 79 wherein the cancer is a solid tumour, hematopoietic cell cancer, colon cancer, lung cancer, kidney cancer, pancreas cancer, endometrial cancer, thyroid cancer, oral cancer, ovarian cancer, laryngeal cancer, hepatocellular cancer, bile duct cancer, squamous cell carcinoma, prostate cancer, breast cancer, cervical cancer, colorectal cancer or melanoma.
81. The use of claim 79 wherein the cancer is lung cancer, melanoma, ovarian cancer, prostate cancer, renal cancer, glioma or astrocytoma.
82. The use of any one of claims 75 to 78 wherein the disease state is a chronic viral infection.
83. The use of any one of claims 75 to 82 wherein the patient is a human.
84. The use of any one of claims 75 to 83 wherein the Myxoma virus is genetically modified to express a therapeutic gene.
85. A Myxoma virus that does not express functional M135R.
86. A pharmaceutical composition comprising Myxoma virus that does not express functional M135R.
87. The pharmaceutical composition of claim 86, further comprising rapamycin.
88. The pharmaceutical composition of claim 86 further comprising an additional therapeutic agent.
89. The pharmaceutical composition of claim 86 wherein the additional therapeutic agent is a chemotherapeutic agent.
90. A kit comprising Myxoma virus that does not express functional M135R and

instructions for inhibiting a cell that has a deficient innate anti-viral response or for treating a disease state characterized by the presence of cells that have a deficient innate anti-viral response.

91. The kit of claim 90, further comprising rapamycin.

92. A method for detecting a cell that has a deficient innate anti-viral response, comprising exposing a population of cells to a Myxoma virus that does not express functional M135R; allowing the virus to infect a cell that has a deficient innate anti-viral response; and determining the infection of any cells of the population of cells by the Myxoma virus.

93. The method of claim 92 wherein the population of cells is in a patient; said exposing comprises administering to the patient the Myxoma virus that does not express functional M135R, the Myxoma virus that does not express functional M135R being modified to express a detectable marker; and said determining comprises detecting a cell expressing the detectable marker in the patient.

94. The method of claim 92 or 93, wherein the exposure is performed in combination with rapamycin.

95. The method of claim 92 wherein the population of cells is in culture.

1/41




FIGURE 1

2/41




FIGURE 2




FIGURE 4

3/41




FIGURE 3




FIGURE 5

4/41




FIGURE 6

5/41







FIGURE 7

6/41

**FIGURE 8**


7/41

**FIGURE 9****FIGURE 10**

8/41



Mock Neutralization



Anti-IFN α/β
(2000 u/ml)

FIGURE 11

9/41




FIGURE 12

10/41




FIGURE 13




FIGURE 14

12/41




FIGURE 15

13/41




FIGURE 16

14/41




FIGURE 17

15/41




FIGURE 18

16/41






FIGURE 19

17/41

**FIGURE 20**

18/41

**FIGURE 21**

19/41




FIGURE 22

20/41




FIGURE 23

21/41





FIGURE 24

22/41

**FIGURE 25**

23/41




FIGURE 26




FIGURE 27

24/41

Daoy

D384




FIGURE 28

25/41




FIGURE 29

26/41




FIGURE 29 (continued)

27/41




FIGURE 29 (continued)

28/41




FIGURE 29 (continued)

29/41


vMyxLac

FIGURE 29 (continued)

30/41







FIGURE 30

31/41


**FIGURE 31**

32/41

**FIGURE 31 (continued)**

33/41

Multistep Growth Curve: MDAMB435 (vMyxLac)



Multistep Growth Curve: COLO205 (vMyxLac)




FIGURE 31 (continued)

34/41




FIGURE 32

35/41




FIGURE 33






FIGURE 34

36/41



B



C



FIGURE 35

37/41




FIGURE 36




FIGURE 37

38/41




FIGURE 38




FIGURE 39

39/41




FIGURE 40




FIGURE 41

40/41




FIGURE 42

41/41




FIGURE 43

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2006/000315

A. CLASSIFICATION OF SUBJECT MATTER IPC: <i>C12N 7/02</i> (2006.01) , <i>C12Q 1/04</i> (2006.01) , <i>A61K 31/436</i> (2006.01) , <i>A61K 39/275</i> (2006.01) , <i>A61K 35/76</i> (2006.01) According to International Patent Classification (IPC) or to both national classification and IPC																							
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC: <i>C12N 7/02</i> (2006.01) , <i>C12Q 1/04</i> (2006.01) , <i>A61K 31/436</i> (2006.01) , <i>A61K 39/275</i> (2006.01) , <i>A61K 35/76</i> (2006.01) Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic database(s) consulted during the international search (name of database(s) and, where practicable, search terms used) Canadian Patent Database, PUBMED, Esp@cenet, USPTO, CAPLUS, BIOSIS, GOOGLE Terms: Myxoma, Rapamycin, Cancer, Macrolide, Mutant, M135R																							
C. DOCUMENTS CONSIDERED TO BE RELEVANT <table border="1"> <thead> <tr> <th>Category*</th> <th>Citation of document, with indication, where appropriate, of the relevant passages</th> <th>Relevant to claim No.</th> </tr> </thead> <tbody> <tr> <td>P, X, O</td> <td>STANFORD, M. "Rapamycin enhances myxoma virus replication in human tumor cells", 2005 Margaret P. Moffat Graduate Research Day, University of Western Ontario, May 10, 2005. See abstract no. 33 submitted for a poster competition.</td> <td>ALL</td> </tr> <tr> <td>A</td> <td>WO 2005/113018 A2 (WELLSTAT BIOLOGICS CORPORATION [US/US]) 01-12-2005</td> <td>ALL</td> </tr> <tr> <td>A</td> <td>US 2004/0147541 A1 (NOVARTIS [US/US]) 29-07-2004</td> <td>1 to 79 and 81 to 87</td> </tr> <tr> <td>A</td> <td>CA 2,517,147 A1 (ROBARTS RESEARCH INSTITUTE [CA/CA]) 16-09-2004</td> <td>ALL</td> </tr> <tr> <td>A</td> <td>CAMERON, C., et. al. "The Complete DNA Sequence of Myxoma Virus", VIROLOGY, 1999, Vol. 264, pages 298-318. ISSN 0042-6822</td> <td>ALL</td> </tr> <tr> <td>A</td> <td>LUN, X., et. al. "Myxoma Virus is a Novel Oncolytic Virus with Significant Antitumor Activity against Experimental Human Gliomas", CANCER RESEARCH, 2005, Vol. 65, No. 21, pages 9982-9990. ISSN 0008-5472</td> <td>ALL</td> </tr> </tbody> </table>			Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	P, X, O	STANFORD, M. "Rapamycin enhances myxoma virus replication in human tumor cells", 2005 Margaret P. Moffat Graduate Research Day, University of Western Ontario, May 10, 2005. See abstract no. 33 submitted for a poster competition.	ALL	A	WO 2005/113018 A2 (WELLSTAT BIOLOGICS CORPORATION [US/US]) 01-12-2005	ALL	A	US 2004/0147541 A1 (NOVARTIS [US/US]) 29-07-2004	1 to 79 and 81 to 87	A	CA 2,517,147 A1 (ROBARTS RESEARCH INSTITUTE [CA/CA]) 16-09-2004	ALL	A	CAMERON, C., et. al. "The Complete DNA Sequence of Myxoma Virus", VIROLOGY, 1999, Vol. 264, pages 298-318. ISSN 0042-6822	ALL	A	LUN, X., et. al. "Myxoma Virus is a Novel Oncolytic Virus with Significant Antitumor Activity against Experimental Human Gliomas", CANCER RESEARCH, 2005, Vol. 65, No. 21, pages 9982-9990. ISSN 0008-5472	ALL
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.																					
P, X, O	STANFORD, M. "Rapamycin enhances myxoma virus replication in human tumor cells", 2005 Margaret P. Moffat Graduate Research Day, University of Western Ontario, May 10, 2005. See abstract no. 33 submitted for a poster competition.	ALL																					
A	WO 2005/113018 A2 (WELLSTAT BIOLOGICS CORPORATION [US/US]) 01-12-2005	ALL																					
A	US 2004/0147541 A1 (NOVARTIS [US/US]) 29-07-2004	1 to 79 and 81 to 87																					
A	CA 2,517,147 A1 (ROBARTS RESEARCH INSTITUTE [CA/CA]) 16-09-2004	ALL																					
A	CAMERON, C., et. al. "The Complete DNA Sequence of Myxoma Virus", VIROLOGY, 1999, Vol. 264, pages 298-318. ISSN 0042-6822	ALL																					
A	LUN, X., et. al. "Myxoma Virus is a Novel Oncolytic Virus with Significant Antitumor Activity against Experimental Human Gliomas", CANCER RESEARCH, 2005, Vol. 65, No. 21, pages 9982-9990. ISSN 0008-5472	ALL																					
☐ Further documents are listed in the continuation of Box C. ☒ See patent family annex. <table border="1"> <tbody> <tr> <td>* Special categories of cited documents :</td> <td>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>"A" document defining the general state of the art which is not considered to be of particular relevance</td> <td>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>"E" earlier application or patent but published on or after the international filing date</td> <td>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>"&" document member of the same patent family</td> </tr> <tr> <td>"O" document referring to an oral disclosure, use, exhibition or other means</td> <td></td> </tr> <tr> <td>"P" document published prior to the international filing date but later than the priority date claimed</td> <td></td> </tr> </tbody> </table>			* Special categories of cited documents :	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"E" earlier application or patent but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family	"O" document referring to an oral disclosure, use, exhibition or other means		"P" document published prior to the international filing date but later than the priority date claimed										
* Special categories of cited documents :	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention																						
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone																						
"E" earlier application or patent but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art																						
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family																						
"O" document referring to an oral disclosure, use, exhibition or other means																							
"P" document published prior to the international filing date but later than the priority date claimed																							
Date of the actual completion of the international search 2 June 2006 (02-06-2006)		Date of mailing of the international search report 29 June 2006 (29-06-2006)																					
Name and mailing address of the ISA/CA Canadian Intellectual Property Office Place du Portage I, C114 - 1st Floor, Box PCT 50 Victoria Street Gatineau, Quebec K1A 0C9 Facsimile No.: 001(819)953-2476		Authorized officer Nancy Trus (819) 953-3355																					

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.
PCT/CA2006/000315

Patent Document Cited in Search Report	Publication Date	Patent Family Member(s)	Publication Date
CA2517147 A1	16-09-2004	AU2004216928 A1 EP1601376 A1 RU2005131020 A WO2004078206 A1	16-09-2004 07-12-2005 10-03-2006 16-09-2004
WO2005113018 A2	01-12-2005	NONE	
US2004147541 A1	29-07-2004	BR0207378 A CA2438504 A1 CN1551767 A CN1679559 A CZ20032209 A3 EP1363627 A2 GB0104072D D0 GB0124957D D0 HU0303271 A2 IL157425D D0 JP2004525899T T MXPA03007418 A NO20033651 A NZ527692 A PL363918 A1 RU2003127391 A SK10382003 A3 WO02066019 A2 ZA200306403 A	15-06-2004 29-08-2002 01-12-2004 12-10-2005 14-01-2004 26-11-2003 04-04-2001 05-12-2001 28-01-2004 28-03-2004 26-08-2004 18-11-2003 17-10-2003 27-05-2005 29-11-2004 27-03-2005 02-03-2004 29-08-2002 29-04-2004

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2006/000315

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of the first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons :

1. ☒ Claim Nos. :
because they relate to subject matter not required to be searched by this Authority, namely :
Claims 1 to 20, 45, 46, 48 to 63, 85 and 86 are directed to, or may encompass, a method for treatment of the human or animal body by surgery or therapy which the International Search Authority is not required to search under Rule 39.1(iv) of the PCT. Regardless, this Authority has carried out a search based on the alleged effects or use of the product defined in claims 1 to 20, 45, 46, 48 to 63, 85 and 86.
2. ☐ Claim Nos. :
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically :
3. ☐ Claim Nos. :
because they are dependant claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows :

1. Claims: 1 to 44 and 48 to 84
directed to the combination of Myxoma virus (including MVΔM135R) and rapamycin and their use together to treat cancer, as well as claims to MVΔM135R per se.
2. Claims: 45 to 47 and 85 to 87
directed to methods of detecting cells with deficient innate anti-viral responses.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claim Nos. :
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim Nos. :

- Remark on Protest** ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.