



Office de la Propriété

Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2684577 C 2013/04/02

(11)(21) **2 684 577**

(12) **BREVET CANADIEN**
CANADIAN PATENT

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2008/04/18
(87) Date publication PCT/PCT Publication Date: 2008/10/30
(45) Date de délivrance/Issue Date: 2013/04/02
(85) Entrée phase nationale/National Entry: 2009/10/19
(86) N° demande PCT/PCT Application No.: IB 2008/051504
(87) N° publication PCT/PCT Publication No.: 2008/129493
(30) Priorité/Priority: 2007/04/19 (US60/925,352)

(51) Cl.Int./Int.Cl. *A61K 8/02*(2006.01),
A61K 8/73(2006.01), *A61K 8/81*(2006.01),
A61Q 19/00(2006.01), *A61Q 5/02*(2006.01)

(72) Inventeurs/Inventors:
WELLS, ROBERT LEE, US;
BROWN, MARK ANTHONY, US;
ROYCE, DOUGLAS ALLAN, US;
LANE, BRANDON SCOTT, US

(73) Propriétaire/Owner:
THE PROCTER & GAMBLE COMPANY, US

(74) Agent: MBM INTELLECTUAL PROPERTY LAW LLP

(54) Titre : COMPOSITIONS DE SOINS D'HYGIENE PERSONNELLE CONTENANT AU MOINS DEUX POLYMERES CATIONIQUES ET UN AGENT TENSIOACTIF ANIONIQUE
(54) Title: PERSONAL CARE COMPOSITIONS CONTAINING AT LEAST TWO CATIONIC POLYMERS AND AN ANIONIC SURFACTANT

(57) Abrégé/Abstract:

A personal cleansing composition comprising: a. from about 5% to about 50% by weight of an anionic detergents surfactant; b. from about 0.025% to about 5% by weight of a first cationic polymer having a cationic charge density of less than about 4 meq/gm, wherein said first cationic polymer forms an isotropic coacervate; c. from about 0.025% to about 5% by weight of a second cationic polymer having a cationic charge density of greater than or equal to about 4 meq/gm, wherein said second cationic polymer forms a lyotropic liquid crystal coacervate; and d. from about 20% to about 94% by weight of water.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
30 October 2008 (30.10.2008)

PCT

(10) International Publication Number
WO 2008/129493 A3

(51) International Patent Classification:

A61K 8/02 (2006.01) *A61Q 5/02* (2006.01)
A61K 8/81 (2006.01) *A61Q 19/00* (2006.01)
A61K 8/73 (2006.01)

(21) International Application Number:

PCT/IB2008/051504

(22) International Filing Date: 18 April 2008 (18.04.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/925,352 19 April 2007 (19.04.2007) US

(71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, Ohio 45202 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **WELLS, Robert**, Lee [US/US]; 4245 Rose Hill Avenue, Cincinnati, Ohio 45229 (US). **BROWN, Mark, Anthony** [US/US]; 1807 Sumpter Court, Union, Kentucky 41091-7663 (US). **ROYCE, Douglas, Allan** [US/US]; 5555N - 400E, Sunman, Indiana 47041 (US). **LANE, Brandon, Scott** [US/US]; 550 Leo Drive, Hamilton, Ohio 45013 (US).

(74) Common Representative: THE PROCTER & GAMBLE COMPANY; c/o Eileen L. Huggett, Winton Hill Business Center, 6250 Center Hill Avenue, Cincinnati, OH 45224 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available):

AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available):

ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:

18 December 2008

WO 2008/129493 A3

(54) Title: PERSONAL CARE COMPOSITIONS CONTAINING AT LEAST TWO CATIONIC POLYMERS AND AN ANIONIC SURFACTANT

(57) Abstract: A personal cleansing composition comprising: a. from about 5% to about 50% by weight of an anionic detergents surfactant; b. from about 0.025% to about 5% by weight of a first cationic polymer having a cationic charge density of less than about 4 meq/gm, wherein said first cationic polymer forms an isotropic coacervate; c. from about 0.025% to about 5% by weight of a second cationic polymer having a cationic charge density of greater than or equal to about 4 meq/gm, wherein said second cationic polymer forms a lyotropic liquid crystal coacervate; and d. from about 20% to about 94% by weight of water.

PERSONAL CARE COMPOSITIONS CONTAINING AT LEAST TWO CATIONIC POLYMERS AND AN ANIONIC SURFACTANT

FIELD OF THE INVENTION

The present invention relates to conditioning personal care compositions with improved conditioning performance which comprise select cationic polymers.

BACKGROUND OF THE INVENTION

Conditioning personal care compositions comprising various combinations of detergents, surfactant and hair conditioning agents are known. These personal care products typically comprise an anionic detergents, surfactants in combination with a conditioning agent such as silicone, hydrocarbon oil, fatty esters, or combinations thereof. These personal care compositions have become more popular among consumers as a means of conveniently obtaining hair conditioning and hair cleansing performance all from a single hair care product.

Many personal care compositions, however, do not provide sufficient deposition of conditioning agents onto hair or skin during application. Obtaining good deposition of a conditioning agent onto hair is complicated by the action of detergents, surfactants in the personal care composition. Detergents, surfactants are designed to carry away or remove, oil, grease, dirt, and particulate matter from the hair and scalp. In doing so, the detergents, surfactants can also interfere with deposition of the conditioning agent, and carry away both deposited and non-deposited conditioning agent during rinsing. This reduces deposition of the conditioning agent onto the hair after rinsing, thus reducing hair conditioning performance. Without sufficient deposition of the conditioning agent on the hair, relatively high levels of conditioning agents may be needed in the personal care composition to provide adequate hair conditioning performance. Such high levels of a conditioning agent, however, can increase raw material costs, reduce lathering, and present product stability concerns.

One known method for improving deposition of a hair conditioning agent onto hair involves the use of certain cationic polymers. Such cationic polymers may form coacervate phases which aid deposition of conditioning agents and provide good wet hair conditioning benefits such as wet hair feel, detangling, and wet combing ease.

One class of cationic polymers comprises high charge density polymers which form a lyotropic liquid crystal coacervate in the presence of an anionic surfactant. Such polymers are characterized by excellent wet combing benefits when used in shampoo compositions, even in

the absence of secondary conditioning agents. When conditioning agents are used, liquid crystal coacervates aid in deposition of large particle size silicone to skin and hair. However, liquid crystal forming polymers do not provide an extended soft wet conditioning and feel throughout the rinse process—as would be expected by a consumer from a desirable conditioning shampoo.

Another class of cationic polymers comprises low charge density cationic polymers which form isotropic coacervates upon dilution of the personal care composition with water. They are often characterized by their favorable wet conditioning and feel, even in the absence of conditioning agents. When conditioning agents are used, isotropic coacervates have been found to aid deposition of small particle size silicone, to skin and hair. However, such low charge density cationic polymers lack the level of wet combing benefit delivered by lyotropic liquid crystal coacervate forming cationic polymers.

Furthermore, formulation of compositions comprising both lyotropic liquid crystal coacervate and isotropic coacervate has been, thus far, unsuccessful. Previous attempts to formulate such compositions have resulted in the destruction of one, or both, of the coacervate phases upon combination.

Based on the foregoing, there still exists a need for a personal care composition with both excellent wet conditioning and wet combing benefits.

SUMMARY OF THE INVENTION

An object of the present invention is to provide personal care compositions containing at least two cationic polymers and an anionic surfactant.

The present invention relates to a personal care composition comprising:

- a. from about 5% to about 50% by weight of an anionic detergents surfactant;
- b. from about 0.025% to about 5% by weight of a first cationic polymer having a cationic charge density of less than about 4 meq/gm, wherein said first cationic polymer forms an isotropic coacervate;
- c. from about 0.025% to about 5% by weight of a second cationic polymer having a cationic charge density of greater than or equal to about 4 meq/gm, wherein said second cationic polymer forms a lyotropic liquid crystal coacervate; and
- d. from about 20% to about 94% by weight of water.

In accordance with another aspect of the present invention, there is provided a hair care composition comprising:

- a) from about 5% to about 50% by weight of an anionic surfactant;
- b) from about 0.025% to about 5% by weight of a first cationic polymer having a cationic charge density of less than 4 meq/g, wherein said first cationic polymer forms an isotropic coacervate in the neat composition or upon dilution with water;
- c) from about 0.025% to about 5% by weight of a second cationic polymer having a cationic charge density of greater than or equal to 4 meq/g, wherein said second cationic polymer forms a lyotropic liquid crystal coacervate upon combination with said anionic surfactant;
- d) an anionic polymeric thickener; and
- e) from about 20% to about 94% by weight of water;

wherein the composition is formed upon mixing a pre-mixture of the second cationic polymer and the anionic surfactant with the first cationic polymer.

In accordance with another aspect of the present invention, there is provided a hair care composition comprising:

- a) an anionic surfactant;
- b) a naturally derived cationic polymer having a cationic charge density of less than 4 meq/g which forms an isotropic coacervate in said hair care composition;
- c) a synthetic cationic polymer having a cationic charge density of greater than or equal to 4 meq/g which forms a lyotropic liquid crystal coacervate in said hair care composition;
- d) an anionic polymeric thickener; and
- e) water;

wherein the composition is formed upon mixing a pre-mixture of the synthetic cationic polymer and the anionic surfactant with the naturally derived cationic polymer.

In accordance with another aspect of the present invention, there is provided a hair care composition comprising:

- a) an anionic surfactant;

- b) a lyotropic liquid crystal coacervate formed in a premix of a cationic polymer having a charge density of greater than or equal to 4 meq/g with said anionic surfactant;
- c) a cationic polymer having a charge density of less than 4 meq/g, which forms an isotropic coacervate in the neat composition or upon dilution with water;
- d) an anionic polymeric thickener; and
- e) water.

In accordance with another aspect of the present invention, there is provided a method of making a hair care composition comprising:

- a) an anionic surfactant;
- b) a first cationic polymer having a cationic charge density of less than 4 meq/g, wherein said first cationic polymer forms an isotropic coacervate in the neat composition or upon dilution with water;
- c) a second cationic polymer having a cationic charge density of greater than or equal to 4 meq/g, wherein said second cationic polymer forms a lyotropic liquid crystal coacervate upon combination with said anionic surfactant; and
- d) water;

wherein prior to combining said first cationic polymer and said second cationic polymer, said second cationic polymer is mixed with said anionic surfactant to form a premix.

DETAILED DESCRIPTION OF THE INVENTION

The personal care compositions of the present invention can comprise, consist of, or consist essentially of the essential elements and limitations of the invention described herein, as well any of the additional or optional ingredients, components, or limitations described herein.

It has been surprisingly discovered that by combining low charge density cationic polymers which form isotropic coacervates with higher charge density cationic polymers, which combine with an anionic surfactant to form lyotropic liquid crystals in a personal care composition, provides both excellent wet hair feel and excellent wet combing benefits.

Furthermore, it has been discovered that in order for the lyotropic liquid crystal coacervate and isotropic coacervates to coexist in a personal care composition the high charge density liquid crystal forming polymer must be premixed with an anionic surfactant prior to the addition of the low charge density isotropic coacervate forming polymer. A detailed description of the formulation process is described herein.

Moreover, without being limited to a particular theory, it appears that when dispersed conditioning agent particles are added to the matrix, the concentrated polymer lyotropic liquid crystal phase and isotropic coacervate phase provide an improved mechanism for conditioning agent deposition, resulting in an overall conditioning benefit.

The liquid crystalline state exists structurally between the solid crystalline phase and the liquid phase (i.e. an intermediate between the three dimensionally ordered crystalline state and the completely disordered liquid state).

In one embodiment, the personal cleansing compositions include an anionic surfactant, a cationic polymer having a cationic charge density of less than about 4 meq/gm, a cationic polymer having a cationic charge density of greater than or equal to about 4 meq/gm, and water. Each of these components, as well as other preferred or optional components are described in detail hereinafter.

All percentages, parts and ratios are based upon the total weight of the compositions of the present invention, unless otherwise specified. All such weights as they pertain to listed ingredients are based on the active level and, therefore, do not include solvents or by-products that may be included in commercially available materials, unless otherwise specified.

All molecular weights as used herein are weight average molecular weights expressed as grams/mole, unless otherwise specified.

The term "amphiphilic" as used herein, refers to complexes having both hydrophilic and hydrophobic properties.

The term "isotropic" as used herein, means a particular phase structure of coacervate wherein the structure is "[random or disordered] along any three orthogonal directions in space, and is therefore dark or 'nonbirefringent' when viewed between crossed polarized light. (One

direction is ‘orthogonal’ to another if the vector component of the first, in the direction of the second, is zero.)’(Laughlin, R. G. (1994) *The Aqueous Phase Behavior of Surfactants*, 182, 8.2).

The term ‘liquid crystal’ as used herein, means a material having phases that are ordered and/or crystalline in only one or two of their three possible orthogonal directions and are disordered (random and/or liquid-like) in the other dimensions.

The term ‘lyotropic’ as used herein, means that the ordering effects of a material are induced by changing both its concentration and temperature. Lyotropic liquid crystals are in the class of amphiphilic materials.

The term ‘neat’ as used herein, means the unadulterated form of the present composition (i.e. prior to altering of the composition through dilution with water).

The term, “nonvolatile” refers to any material having little or no significant vapor pressure under ambient conditions, and a boiling point under one atmosphere (atm) preferably at least about 250°C. The vapor pressure under such conditions is preferably less than about 0.2 mm.

The term “water soluble” as used herein, means that the polymer is soluble in water in the present composition. In general, the polymer should be soluble at 25°C at a concentration of 0.1% by weight of the water solvent, preferably at 1%, more preferably at 5%, more preferably at 15%.

Detensive Surfactant Component

The personal care compositions comprise an anionic detensive surfactant component to provide cleaning performance to the composition. The anionic detensive surfactant component in turn comprises anionic detensive surfactant, zwitterionic or amphoteric detensive surfactant which has an attached group that is anionic at the pH of the composition, or a combination thereof, preferably anionic detensive surfactant. Such surfactants should be physically and chemically compatible with the essential components described herein, or should not otherwise unduly impair product stability, aesthetics or performance.

Suitable anionic detensive surfactant components include those which are known for use in hair care or other personal care cleansing compositions. The concentration of the anionic surfactant component in the personal care composition should be sufficient to provide the desired cleaning and lather performance, and generally range from about 5% to about 50%, preferably from about 8% to about 30%, more preferably from about 10% to about 25%, even more preferably from about 12% to about 20%, by weight of the composition.

Preferred anionic detergents for use in the compositions include ammonium lauryl sulfate, ammonium laureth sulfate, triethylamine lauryl sulfate, triethylamine laureth sulfate, triethanolamine lauryl sulfate, triethanolamine laureth sulfate, monoethanolamine lauryl sulfate, monoethanolamine laureth sulfate, diethanolamine lauryl sulfate, diethanolamine laureth sulfate, lauric monoglyceride sodium sulfate, sodium lauryl sulfate, sodium laureth sulfate, potassium lauryl sulfate, potassium laureth sulfate, sodium lauryl sarcosinate, sodium lauroyl sarcosinate, lauryl sarcosine, cocoyl sarcosine, ammonium cocoyl sulfate, ammonium lauroyl sulfate, sodium cocoyl sulfate, sodium lauroyl sulfate, potassium cocoyl sulfate, potassium lauryl sulfate, triethanolamine lauryl sulfate, triethanolamine lauryl sulfate, monoethanolamine cocoyl sulfate, monoethanolamine lauryl sulfate, sodium tridecyl benzene sulfonate, sodium dodecyl benzene sulfonate, sodium cocoyl isethionate and combinations thereof.

Suitable amphoteric or zwitterionic detergents for use in the personal care composition herein include those which are known for use in hair care or other personal care cleansing composition, and which contain a group that is anionic at the pH of the personal care composition. Concentration of such amphoteric detergents preferably ranges from about 0.5% to about 20%, preferably from about 1% to about 10%, by weight of the composition. Non limiting examples of suitable zwitterionic or amphoteric detergents are described in U.S. Patent Nos. 5,104,646 and 5,106,609.

The personal care compositions may further comprise additional detergents for use in combination with the anionic detergent component described hereinbefore. Suitable optional detergents include nonionic detergents, cationic detergents, and combinations thereof. Any such detergent known in the art for use in hair or personal care products may be used, provided that the optional additional detergent is also chemically and physically compatible with the essential components of the personal care composition, or does not otherwise unduly impair product performance, aesthetics or stability. The concentration of the optional additional detergents in the personal care composition may vary with the cleansing or lather performance desired, the optional detergent selected, the desired product concentration, the presence of other components in the composition, and other factors well known in the art.

Non limiting examples of other anionic, zwitterionic, amphoteric or optional additional detergents suitable for use in the personal care compositions are described in McCutcheon's, Emulsifiers and Detergents, 1989 Annual, published by M. C. Publishing Co., and U.S. Patent Nos. 3,929,678; 2,658,072; 2,438,091; and 2,528,378.

Naturally Derived Cationic Polymer

The personal care compositions of the present invention comprise a first cationic polymer which is a low charge density cationic polymer. In a preferred embodiment, the first cationic polymer is a naturally derived cationic polymer. The term, "naturally derived cationic polymer" as used herein, refers to cationic polymers which are obtained from natural sources. The natural sources may be polysaccharide polymers. Therefore, the naturally derived cationic polymer may be selected from celluloses, starches, guars, non-guar-galactomannans, and other sources found in nature.

The first cationic polymer has a molecular weight of from about 1,000 to about 10,000,000, and a cationic charge density of at least about 0.2 meq/g, more preferably at least about 0.5 meq/g. The first cationic polymer also has a charge density of less than about 4.0 meq/gm, and more preferably less than or equal to about 2 meq/gm. The polymers are typically present in a concentration of from about 0.025% to about 5%, and more preferably from about 0.10% to about 2% by weight of the personal care composition. The first cationic polymers form an isotropic coacervate in the neat composition or upon dilution with water. The isotropic coacervate aids in deposition of optional small particle size conditioning agents, and provides excellent wet conditioning performance. Such deposition and wet conditioning enhancement result in hair feel, shine, and other appreciable benefits.

The cationic polymers herein are either soluble in the personal care composition, or are soluble in a complex coacervate phase in the personal care composition formed by the cationic polymer and the anionic detergents surfactant component described hereinbefore. Complex coacervates of the cationic polymer can also be formed with other charged materials, such as anionic polymers, in the personal care composition.

Isotropic coacervate formation is dependent upon a variety of criteria such as molecular weight, component concentration, and ratio of interacting ionic components, ionic strength (including modification of ionic strength, for example, by addition of salts), charge density of the cationic and anionic components, pH, temperature, and the aforementioned surfactant system. Isotropic coacervate systems and the effect of these parameters have been described, for example, by J. Caelles, et al., "Anionic and Cationic Compounds in Mixed Systems", Cosmetics & Toiletries, Vol. 106, April 1991, pp 49-54, C. J. van Oss, "Coacervation, Complex Coacervation and Flocculation", J. Dispersion Science and Technology, Vol. 9 (5,6), 1988-89, pp 561-573, and

D. J. Burgess, "Practical Analysis of Complex Coacervate Systems", *J. of Colloid and Interface Science*, Vol. 140, No. 1, November 1990, pp 227-238.


It is believed to be particularly advantageous for the first cationic polymer to be present in the personal care composition in an isotropic coacervate phase, or to form an isotropic coacervate phase upon application or rinsing of the composition to or from the hair. Complex isotropic coacervates are believed to more readily deposit on the hair than a dissolved polymer. Thus, in general, it is preferred that the first cationic polymer exist in the personal care composition as an isotropic coacervate phase or form an isotropic coacervate phase upon dilution.

Techniques for analysis of formation of complex isotropic coacervates are known in the art. For example, microscopic analyses of the personal care compositions, at any chosen stage of dilution, can be utilized to identify whether an isotropic coacervate phase has formed. Such isotropic coacervate phases will be identifiable as an additional emulsified phase in the composition. The use of dyes can aid in distinguishing the isotropic coacervate phase from other insoluble phases dispersed in the personal care composition.

Cationic Polysaccharide Polymers

The personal care compositions of the present invention may include a naturally derived cationic polymer which is a cationic polysaccharide polymer. Cationic polysaccharide polymers encompass cellulose polymers, starch polymers, and polymers made up of multiple monosaccharides joined together by glycosidic linkages.

Suitable polysaccharide cationic polymers include those which conform to the following formula:



wherein A is an anhydroglucose residual group, such as a cellulose anhydroglucose residual; R is an alkylene oxyalkylene, polyoxyalkylene, or hydroxyalkylene group, or combination thereof; R₁, R₂, and R₃ independently are alkyl, aryl, alkyl aryl, aryl alkyl, alkoxyalkyl, or alkoxyaryl groups, each group containing up to about 18 carbon atoms, and the total No of carbon atoms for each cationic moiety (i.e., the sum of carbon atoms in R₁, R₂ and R₃) preferably being about 20 or less; and X is an anionic counterion. Non-limiting examples of such counterions include halides (e.g., chlorine, fluorine, bromine, iodine), sulfate and methylsulfate. The degree of cationic substitution in these polysaccharide polymers is typically from about 0.01 to about 1 cationic groups per anhydroglucose unit.

In one embodiment of the invention, the cellulose cationic polymers are salts of hydroxyethyl cellulose reacted with trimethyl ammonium substituted epoxide, referred to in the industry (CTFA) as Polyquaternium 10 and available from Amerchol Corp. (Edison, N.J., USA) under the trade name, UcareTM Polymer KG-30M, having a cationic charge density of about 1.9 meq/gm.

Cationically Modified Starch Polymer

The personal care compositions may comprise a naturally derived cationic polymer which is a water-soluble cationically modified starch polymer. As used herein, the term, "cationically modified starch", refers to a starch to which a cationic group is added prior to degradation of the starch to a smaller molecular weight, or wherein a cationic group is added after modification of the starch to achieve a desired molecular weight. The definition of the term "cationically modified starch" also includes amphotERICALLY modified starches having a net positive charge. The term "amphotERICALLY modified starch" refers to a starch hydrolysate to which a cationic group and an anionic group are added.

A method of chemically modifying the charge densities of the cationically modified starch polymers includes, but is not limited to, the addition of amino and/or ammonium groups into the starch molecules. Non-limiting examples of these ammonium groups may include substituents such as hydroxypropyl trimmonium chloride, trimethylhydroxypropyl ammonium chloride, dimethylstearylhydroxypropyl ammonium chloride, and dimethyldodecylhydroxypropyl ammonium chloride. See Solarek, D. B., *Cationic Starches in Modified Starches: Properties and Uses*, Wurzburg, O.B., Ed., CRC Press, Inc., Boca Raton, Florida 1986, pp 113-125. The cationic groups may be added to the starch prior to degradation to a smaller molecular weight or the cationic groups may be added after such modification.

The cationically modified starch polymers may comprise maltodextrin. Thus, in one embodiment of the present invention, the cationically modified starch polymers may be further characterized by a Dextrose Equivalence (DE) value of less than about 35, and more preferably from about 1 to about 20. The DE value is a measure of the reducing equivalence of the hydrolyzed starch referenced to dextrose and expressed as a percent (on dry basis). Starch completely hydrolyzed to dextrose has a DE value of 100, and unhydrolyzed starch has a DE value of 0. A suitable assay for DE value includes one described in "Dextrose Equivalent", Standard Analytical Methods of the Member Companies of the Corn Industries Research Foundation, 1st ed., Method E-26. Additionally, the cationically modified starch polymers may

comprise a dextrin. Dextrin is typically a pyrolysis product of starch with a wide range of molecular weights.

The source of starch before chemical modification can be chosen from a variety of sources such as tubers, legumes, cereal, and grains. Non-limiting examples of this source starch may include corn starch, wheat starch, rice starch, waxy corn starch, oat starch, cassava starch, waxy barley, waxy rice starch, glutenous rice starch, sweet rice starch, amioica, potato starch, tapioca starch, oat starch, sago starch, sweet rice, or mixtures thereof. Waxy corn starch is preferred.

In one embodiment of the present invention, cationically modified starch polymers are selected from degraded cationic maize starch, cationic tapioca, cationic potato starch, and mixtures thereof. In another embodiment, cationically modified starch polymers are cationic corn starch.

The starch, prior to degradation or after modification to a smaller molecular weight, may comprise one or more additional modifications. For example, these modifications may include cross-linking, stabilization reactions, phosphorylations, and hydrolyzations. Stabilization reactions may include alkylation and esterification.

The cationically modified starch polymers in the present invention may be incorporated into the composition in the form of hydrolyzed starch (e.g., acid, enzyme, or alkaline degradation), oxidized starch (e.g., peroxide, peracid, hypochlorite, alkaline, or any other oxidizing agent), physically/mechanically degraded starch (e.g., via the thermo-mechanical energy input of the processing equipment), or combinations thereof.

Suitable cationically modified starches are available from known starch suppliers, such as National Starch. Also suitable for use in the present invention is nonionic modified starch that could be further derivatized to a cationically modified starch as is known in the art. Other suitable modified starch starting materials may be quaternized, as is known in the art, to produce a cationically modified starch polymer suitable for use in the present invention.

One method of conducting starch degradation involves preparing a starch slurry by mixing granular starch in water. The temperature is raised to about 35°C. An aqueous solution of potassium permanganate is then added at a concentration of about 50 ppm based on starch. The pH is raised to about 11.5 with sodium hydroxide and the slurry is stirred sufficiently to prevent settling of the starch. Then, about a 30% solution of hydrogen peroxide diluted in water is added to a level of about 1% of peroxide based on starch. The pH of about 11.5 is then restored by adding additional sodium hydroxide. The reaction is completed over about a 1 to

about 20 hour period. The mixture is then neutralized with dilute hydrochloric acid. The degraded starch is recovered by filtration followed by washing and drying.

Cationic Galactomannan Polymer

The personal care compositions may comprise a naturally derived cationic polymer which may be a guar or non-guar galactomannan polymer. In one embodiment, the galactomannan polymer is a polymer derivative having a mannose to galactose ratio of 2 : 1 or greater, on a monomer to monomer basis, and the galactomannan polymer derivative is selected from the group consisting of a cationic galactomannan polymer derivative and an amphoteric galactomannan polymer derivative having a net positive charge. The term "galactomannan polymer derivative", means a compound obtained from a galactomannan polymer (ie. a galactomannan gum) which is chemically modified. As used herein, the term "cationic galactomannan" refers to a galactomannan polymer to which a cationic group is added. The term "amphoteric galactomannan" refers to a galactomannan polymer to which a cationic group and an anionic group are added such that the polymer has a net positive charge.

Galactomannan polymers are present in the endosperm of seeds of the leguminosae family. Galactomannan polymers are made up of a combination of mannose monomers and galactose monomers. The galactomannan molecule is a straight chain mannan branched at regular intervals with single membered galactose units on specific mannose units. The mannose units are linked to each other by means of μ (1-4) glycosidic linkages. The galactose branching arises by way of an μ (1-6) linkage. The ratio of mannose monomers to galactose monomers varies according to the species of the plant and also is affected by climate. Guar is an example of one type of a galactomannan polymer, specifically having a mannose to galactose ratio of 2 monomers of mannose to 1 monomer of galactose. In one embodiment, the galactomannan polymer derivatives have a ratio of mannose to galactose of greater than 2 : 1 on a monomer to monomer basis (i.e., non-guar galactomannan polymers). Preferably, the ratio of mannose to galactose is greater than about 3 : 1, and more preferably the ratio of mannose to galactose is greater than about 4 : 1. Analysis of mannose to galactose ratios is well known in the art and is typically based on the measurement of the galactose content.

The gum for use in preparing the non-guar galactomannan polymer derivatives is typically obtained as naturally occurring material such as seeds or beans from plants. Examples of various non-guar galactomannan polymers include but are not limited to tara gum (3 parts mannose / 1 part galactose), locust bean or carob (4 parts mannose / 1 part galactose), and cassia

gum (5 parts mannose / 1 part galactose). Herein, the term “non-guar galactomannan polymer derivatives” refers to cationic polymers which are chemically modified from a non-guar galactomannan polymer. A preferred non-guar galactomannan polymer derivative is cationic cassia and is sold under the trade name Cassia EX-906, which is commercially available from Noveon Inc.

Suitable galactomannan polymer derivatives are described in U.S. Patent Publication No. 2006/0099167A1 to Staudigel et al.

Synthetic Cationic Polymer

The personal care compositions of the present invention comprise a second cationic polymer having a high charge density. In a preferred embodiment, the second cationic polymer is a synthetic cationic polymer. The second cationic polymer, in combination with the anionic surfactant component, forms lyotropic liquid crystals. The polymers can be formulated in a stable personal care composition that provides improved conditioning performance, and also provides improved deposition of the optional large particle size conditioning agents (described herein) onto hair. In one embodiment, the synthetic cationic polymer may be formed from


- i) one or more cationic monomer units, and optionally
- ii) one or more momomer units bearing a negative charge, and/or
- iii) a nonionic momomer,

wherein the subsequent charge of the copolymer is positive. The ratio of the three types of monomers is given by m, p and q where m is the number of cationic monomers, p the No of momomers bearing a negative charge and q is the number of nonionic momomers.


The concentration of the second cationic polymer in the personal care composition ranges about 0.025% to about 5%, preferably from about 0.1% to about 3%, more preferably from about 0.2% to about 1%, by weight of the composition.

The second cationic polymers have a cationic charge density of at least about 4 meq/gm or at least about 4.5 meq/gm, and they have a molecular weight of at least about 500,000 to about 5,000,000, more preferably from about 500,000 to about 2,000,000.

In one embodiment, the second cationic polymers are water soluble or dispersible, non-crosslinked, synthetic cationic polymers having the following structure:



Where A, may be one or more of the following cationic moieties:



Where @ = amido, alkylamido, ester, ether, alkyl or alkylaryl.

Where Y = C1-C22 alkyl, alkoxy, alkylidene, alkyl or aryloxy

Where ψ = C1-C22 alkyl, alkyloxy, alkyl aryl or alkyl aryloxy

Where Z = C1-C22 alkyl, alkyloxy, aryl or aryloxy

Where R1 = H, C1-C4 linear or branched alkyl

Where s = 0 or 1, n = 0 or ≥ 1

Where T and R7 = C1-C22 alkyl

Where X- = halogen, hydroxide, alkoxide, sulfate or alkylsulfate

Examples of cationic monomers include aminoalkyl (meth)acrylates, (meth)aminoalkyl (meth)acrylamides; monomers comprising at least one secondary, tertiary or quaternary amine function, or a heterocyclic group containing a nitrogen atom, vinylamine or ethylenimine;

diallyldialkyl ammonium salts; their mixtures, their salts, and macromonomers deriving from therefrom.


Further examples of cationic monomers include dimethylaminoethyl (meth)acrylate, dimethylaminopropyl (meth)acrylate, ditertiobutylaminoethyl (meth)acrylate, dimethylaminomethyl (meth)acrylamide, dimethylaminopropyl (meth)acrylamide, ethylenimine, vinylamine, 2-vinylpyridine, 4-vinylpyridine, trimethylammonium ethyl (meth)acrylate chloride, trimethylammonium ethyl (meth)acrylate methyl sulphate, dimethylammonium ethyl (meth)acrylate benzyl chloride, 4-benzoylbenzyl dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl (meth)acrylamido chloride, trimethyl ammonium propyl (meth)acrylamido chloride, vinylbenzyl trimethyl ammonium chloride, diallyldimethyl ammonium chloride.

Preferred cationic monomers comprise quaternary ammonium group of formula -NR₃+, wherein R, which is identical or different, represents a hydrogen atom, an alkyl group comprising 1 to 10 carbon atoms, or a benzyl group, optionally carrying a hydroxyl group, and comprise an anion (counter-ion). Examples of anions are halides such as chlorides, bromides, sulphates, hydrosulphates, alkylsulphates (for example comprising 1 to 6 carbon atoms), phosphates, citrates, formates, and acetates.

Preferred cationic monomers include trimethylammonium ethyl (meth)acrylate chloride, trimethylammonium ethyl (meth)acrylate methyl sulphate, dimethylammonium ethyl (meth)acrylate benzyl chloride, 4-benzoylbenzyl dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl (meth)acrylamido chloride, trimethyl ammonium propyl (meth)acrylamido chloride, vinylbenzyl trimethyl ammonium chloride.

More preferred cationic monomers include trimethyl ammonium propyl (meth)acrylamido chloride.

Where the monomer bearing a negative charge is defined by R₂ = H, C₁-C₄ linear or branched alkyl and R₃ as:



Where D = electronegative element chosen between oxygen, nitrogen, sulfur

Where Q = NH₂ or O

Where u = 1-6


Where t = 0-1

Where J = oxygenated functional group containing the following elements P, S, C

Examples of monomers bearing a negative charge include alpha ethylenically unsaturated monomers comprising a phosphate or phosphonate group, alpha ethylenically unsaturated monocarboxylic acids, monoalkylesters of alpha ethylenically unsaturated dicarboxylic acids, monoalkylamides of alpha ethylenically unsaturated dicarboxylic acids, alpha ethylenically unsaturated compounds comprising a sulphonic acid group, and salts of alpha ethylenically unsaturated compounds comprising a sulphonic acid group.

Preferred monomers with a negative charge include acrylic acid, methacrylic acid, vinyl sulphonic acid, salts of vinyl sulfonic acid, vinylbenzene sulphonic acid, salts of vinylbenzene sulphonic acid, alpha-acrylamidomethylpropanesulphonic acid, salts of alpha-acrylamidomethylpropanesulphonic acid, 2-sulphoethyl methacrylate, salts of 2-sulphoethyl methacrylate, acrylamido-2-methylpropanesulphonic acid (AMPS), salts of acrylamido-2-methylpropanesulphonic acid, and styrenesulphonate (SS).

Where the nonionic monomer is defined by R₂ = H, C₁-C₄ linear or branched alkyl, R₆ = linear or branched alkyl, alkyl aryl, aryl oxy, alkyloxy, alkylaryl oxy and β is defined as



Where G' and G'' are, independently of one another, O, S or N-H and L = 0 or 1.

Examples of such nonionic monomers include vinyl acetate, amides of alpha ethylenically unsaturated carboxylic acids, esters of an alpha ethylenically unsaturated monocarboxylic acids with an hydrogenated or fluorinated alcohol, polyethylene oxide (meth)acrylate (i.e. polyethoxylated (meth)acrylic acid), monoalkylesters of alpha ethylenically unsaturated dicarboxylic acids, monoalkylamides of alpha ethylenically unsaturated dicarboxylic acids, vinyl nitriles, vinylamine amides, vinyl alcohol, vinyl pyrrolidone, and vinyl aromatic compounds.

Preferred nonionic monomers include styrene, acrylamide, methacrylamide, acrylonitrile, methylacrylate, ethylacrylate, n-propylacrylate, n-butylacrylate, methylmethacrylate, ethylmethacrylate, n-propylmethacrylate, n-butylmethacrylate, 2-ethyl-hexyl acrylate, 2-ethyl-hexyl methacrylate, 2-hydroxyethylacrylate and 2-hydroxyethylmethacrylate.

The anionic counterion (X⁻) in association with the synthetic cationic polymers may be any known counterion so long as the polymers remain soluble or dispersible in water, in the personal care composition, or in a coacervate phase of the personal care composition, and so long as the counterions are physically and chemically compatible with the essential components of the personal care composition or do not otherwise unduly impair product performance, stability or aesthetics. Non limiting examples of such counterions include halides (e.g., chlorine, fluorine, bromine, iodine), sulfate and methylsulfate.

Optional Ingredients

The present personal care composition may further comprise optional ingredients selected from the group consisting of oily conditioning agents, hydrocarbon oils, polyolefins, fatty esters, fluorinated conditioning compounds, fatty alcohols, quaternary ammonium compounds, polyethylene glycols, anti-dandruff actives, anti-microbial actives, inorganic or synthetic particles, opacifying agents, suspending agents, propellants, paraffinic hydrocarbons, mono or divalent salts, fragrances, vitamins, chelating agents, colorants, pigments, dyes and mixtures thereof. These optional components are described in detail in U.S. Patent Publication No. 2003/0223951A1.

Silicone Conditioning Agent

If oily conditioning agents are included, they are preferably in the form of a water-insoluble silicone conditioning agent. The silicone conditioning agent may comprise volatile silicone, non-volatile silicone, or combinations thereof. Preferred are non-volatile silicone conditioning agents. If volatile silicones are present, it will typically be incidental to their use as a solvent or carrier for commercially available forms of non-volatile silicone materials, such as

silicone gums and resins. The silicone conditioning agent particles may comprise a silicone fluid conditioning agent and may also comprise other ingredients, such as a silicone resin to improve silicone fluid deposition efficiency or enhance glossiness of the hair.

Non-limiting examples of suitable silicone conditioning agents, and optional suspending agents for the silicone, are described in U.S. Reissue Patent No. 34,584, U.S. Patent Nos. 5,104,646, and 5,106,609. The silicone conditioning agents for use in the compositions of the present invention preferably have a viscosity, as measured at 25°C, from about 20 to about 2,000,000 centistokes (“csk”), more preferably from about 1,000 to about 1,800,000 csk, even more preferably from about 5,000 to about 1,500,000 csk, more preferably from about 10,000 to about 1,000,000 csk.

In one embodiment of the present invention, the personal care composition is opaque. The personal care composition comprises a non-volatile silicone oil having a particle size as measured in the personal care composition from about 1 μm to about 50 μm . The lyotropic liquid crystal coacervate, described herein, enhances deposition of these large particle size silicones (larger than about 1 μm).

In another embodiment of the present invention, the personal care composition comprises a non-volatile silicone oil having a particle size as measured in the personal care composition from about 100 nm to about 1 μm . A substantially clear composition embodiment of the present invention comprises a non-volatile silicone oil having a particle size as measured in the personal care composition of less than about 100 nm. The isotropic coacervate, described herein, enhances deposition of small particle size silicones having a particle size of less than or equal to about 1 μm .

Non-volatile silicone oils suitable for use in compositions of the present invention may be selected from organo-modified silicones and fluoro-modified silicones. In one embodiment of the present invention, the non-volatile silicone oil is an organo-modified silicone which comprises an organo group selected from the group consisting of alkyl groups, alkenyl groups, hydroxyl groups, amine groups, quaternary groups, carboxyl groups, fatty acid groups, ether groups, ester groups, mercapto groups, sulfate groups, sulfonate groups, phosphate groups, propylene oxide groups, and ethylene oxide groups.

In a preferred embodiment of the present invention, the non-volatile silicone oil is polydimethylsiloxane.

Silicone fluids suitable for use in the compositions of the present invention are disclosed in U.S. Patent Nos. 2,826,551; 3,964,500; and 4,364,837, British Patent No 849,433, and Silicon Compounds, Petrarch Systems, Inc. (1984).

Method of Making

All exemplified amounts are listed as weight percents and exclude minor materials such as diluents, preservatives, color solutions, imagery or conceptual ingredients, botanicals, and so forth, unless otherwise specified.

One method of making the present compositions involves mixing water and cocamide, which are heated to above 55°C in the main mix tank. A polymeric anionic thickener, such as acrylates copolymer may optionally be added. Acrylates copolymer, which is commercially sold under the trade name, Aqua SF1, is available from National Starch and other commercial suppliers. Surfactants are then added and the pH is raised to >6.5 in order to activate thickening of the acrylates copolymer. Next, the synthetic cationic polymer is mixed, in a concentrated salt solution, with an anionic surfactant to form a liquid crystal coacervate in a premix. This premix is then added to the main mix tank. Then naturally derived cationic polymer is added to the main mix tank. Alternately, the naturally derived cationic polymer may be added to the main mix tank, followed by the aforementioned premix. It has been discovered that if the naturally derived cationic polymer is added to the main mix tank before the synthetic cationic polymer, then liquid crystal coacervate formation is inhibited, unless the synthetic cationic polymer is subsequently added to the main mix tank in the form of the aforementioned premix. Glycol distearate can be crystallized in a separate surfactant premix and then added to the main mix. A silicone/surfactant premix can be made to achieve the desired particle size and then added to the main mix tank. The pH is finally adjusted to the desired level and preservatives and minors added.

If acrylates copolymer is not used then the step of increasing the pH to > 6.5 is not needed.

If a multiple phase shampoo is formulated, the multiple phases are made separately and then added separately to the package to achieve the desired appearance. The multiple phases may differ in color, composition, texture, transparency or mixtures of these attributes. Multiple phase personal care compositions are described in detail in U.S. Patent No. 3,932,610.

The following examples are representative of shampoo compositions of the invention. The present compositions may comprise one, two, or more phases in accordance with the examples below:

Ingredient (Phase 1)	Chem Conc. %	% Active in Final			% Active (Phase 1)		
Sodium Laureth Sulfate (28% active in water)	28.0	5.0000		5.0000	5.0000	5.0000	4.0000
Sodium Lauryl Sulfate (29% active in water)	29.0	9.0000		15.0000	15.0000	9.0000	8.0000
Ammonium Laureth Sulfate (25% active in water)	25.0		10.0000				
Ammonium Lauryl Sulfate (25% active in water)	25.0		6.0000				
Polydimethyl siloxane	100.0	1.0000	2.0000	1.5000	3.0000	4.0000	3.0000
Glycol distearate	100.0	1.5000	1.5000	1.5000	1.5000	1.5000	1.5000
Carbopo TM Aqua SF-1 (Acrylates copolymer) (Available from National Starch)	30.0				1.5000	1.2500	1.2500
Polyquaternium 10 (LR400) (Available from Americhol)		0.5000					
Polyquaternium 10 (LR30M) (Available from Americhol)			0.2500	0.2500	0.2500		
Polyquaternium 10 (KG30M) (Available from Americhol)	100.0					0.5000	0.2500
Mirapol TM 100 (Polyquaternium 6) (Available from Rhodia)	31.5	0.0500				0.2500	0.2500
Polycare TM 133 (Polymethacrylamidopropyl trimonium CL) (Available from Rhodia)			0.1000	0.1000	0.1000		
cocodimethyl amide	85.0	0.8000	0.8000	0.8000	0.8000	0.8000	0.8000
Brij TM 30 (Laureth-4)	100.0	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
NaOH (50%)	50.0	as needed	as needed	as needed	as needed	as needed	as needed
Sodium Benzoate	100.0	0.2500	0.2500	0.2500	0.2500	0.2500	0.2500
Disodium EDTA	100.0	0.1274	0.1274	0.1274	0.1274	0.1274	0.1274
Citric Acid	100.0	0.5000	0.5000	0.5000	0.5000	0.5000	0.5000
NaCl	100.0	as needed	as needed	as needed	as needed	as needed	as needed
Sodium Xylene Sulfonate	41.5	as needed	as needed	as needed	as needed	as needed	as needed
Kathon TM CG (Methylchloroisothiazolinone and Methylisothiazolinone)	100.0	0.0005	0.0005	0.0005	0.0005	0.0005	0.0005
Perfume/colors/other minors	100.0	as needed	as needed	as needed	as needed	as needed	as needed
Q.S. Water- USP Purified	100.0						
Ingredient (Phase 2)	Chem Conc. %	No 2nd Phase			% Active (Phase 2)		
Sodium Laureth Sulfate (28% active in water)	28.0			5.0000	5.0000	4.0000	
Sodium Lauryl Sulfate (29% active in water)	29.0			15.0000	9.0000	8.0000	
Polydimethyl siloxane	100.0			1.0000	2.0000	1.0000	
Carbopol Aqua SF-1 (Acrylates copolymer) (Available from National Starch)	30.0			1.5000	1.2500	1.2500	
Polyquaternium 10 (LR30M) (Available from Americhol)				0.2500			
Polyquaternium 10 (KG30M) (Available from Americhol)	100.0				0.5000	0.2500	
Mirapol 100 (Polyquaternium 6)	31.5				0.2500	0.2500	
Polycare 133 (Polymethacrylamidopropyl trimonium CL)				0.1000			
cocodimethyl amide	85.0			0.8000	0.8000	0.8000	
Brij 30 (Laureth-4)	100.0			1.0000	1.0000	1.0000	
NaOH (50%)	50.0			as needed	as needed	as needed	
Sodium Benzoate	100.0			0.2500	0.2500	0.2500	
Disodium EDTA	100.0			0.1274	0.1274	0.1274	
Citric Acid	100.0			0.5000	0.5000	0.5000	
NaCl	100.0			as needed	as needed	as needed	
Sodium Xylene Sulfonate	41.5			as needed	as needed	as needed	
Kathon CG (Methylchloroisothiazolinone and Methylisothiazolinone)	100.0			0.0005	0.0005	0.0005	

The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and a functionally equivalent range

surrounding that value. For example, a dimension disclosed as "40 mm" is intended to mean "about 40 mm."

The citation of any document is not to be construed as an admission that it is prior art with respect to the present invention. To the extent that any meaning or definition of a term in this document conflicts with any meaning or definition of the same term in a document cited herein, the meaning or definition assigned to that term in this document shall govern.

THE EMBODIMENTS OF THE INVENTION FOR WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A hair care composition comprising:
 - a) from about 5% to about 50% by weight of an anionic surfactant;
 - b) from about 0.025% to about 5% by weight of a first cationic polymer having a cationic charge density of less than 4 meq/g, wherein said first cationic polymer forms an isotropic coacervate in the neat composition or upon dilution with water;
 - c) from about 0.025% to about 5% by weight of a second cationic polymer having a cationic charge density of greater than or equal to 4 meq/g, wherein said second cationic polymer forms a lyotropic liquid crystal coacervate upon combination with said anionic surfactant;
 - d) an anionic polymeric thickener; and
 - e) from about 20% to about 94% by weight of water;wherein the composition is formed upon mixing a pre-mixture of the second cationic polymer and the anionic surfactant with the first cationic polymer.
2. A hair care composition according to claim 1, wherein said first cationic polymer is a naturally derived cationic polymer.
3. A hair care composition according to claim 2, wherein said naturally derived cationic polymer is selected from the group consisting of celluloses, starches, guar, and non-guar-galactomannans.
4. A hair care composition according to claim 2 or 3, wherein said naturally derived cationic polymer has a cationic charge density of less than or equal to 2 meq/g.
5. A hair care composition according to claim 2, wherein said naturally derived cationic polymer is polyquaternium-10.

6. A hair care composition according to claim 5, wherein said polyquaternium-10 has a cationic charge density of about 1.9 meq/g.
7. A hair care composition according to claim 2, wherein said naturally derived cationic polymer is cationic cassia.
8. A hair care composition according to any one of claims 1 to 7, wherein said second cationic polymer is a synthetic cationic polymer.
9. A hair care composition according to claim 8, wherein said synthetic cationic polymer comprises monomers selected from the group consisting of dimethylaminoethyl methacrylate, dimethylaminoethyl acrylate, dimethylaminopropyl methacrylate, dimethylaminopropyl acrylate, ditertiobutylaminoethyl methacrylate, ditertiobutylaminoethyl acrylate, dimethylaminomethyl methacrylamide, dimethylaminomethyl acrylamide, dimethylaminopropyl methacrylamide, dimethylaminopropyl acrylamide, ethylenimine, vinylamine, 2-vinylpyridine, 4-vinylpyridine, trimethylammonium ethyl methacrylate chloride, trimethylammonium ethyl acrylate chloride, trimethylammonium ethyl methacrylate methyl sulphate, trimethylammonium ethyl acrylate methyl sulphate, dimethylammonium ethyl methacrylate benzyl chloride, dimethylammonium ethyl acrylate benzyl chloride, 4-benzoylbenzyl dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl methacrylamido chloride, trimethyl ammonium ethyl acrylamido chloride, trimethyl ammonium propyl methacrylamido chloride, trimethyl ammonium propyl acrylamido chloride, vinylbenzyl trimethyl ammonium chloride, diallyldimethyl ammonium chloride, trimethylammonium ethyl methacrylate chloride, trimethylammonium ethyl acrylate chloride, trimethylammonium ethyl methacrylate methyl sulphate, trimethylammonium ethyl acrylate methyl sulphate, dimethylammonium ethyl methacrylate benzyl chloride, dimethylammonium ethyl acrylate benzyl chloride, 4-benzoylbenzyl

dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl methacrylamido chloride, trimethyl ammonium ethyl acrylamido chloride, trimethyl ammonium propyl methacrylamido chloride, trimethyl ammonium propyl acrylamido chloride, vinylbenzyl trimethyl ammonium chloride, trimethyl ammonium propyl methacrylamido chloride, and trimethyl ammonium propylacrylamido chloride.

10. A hair care composition according to claim 8, wherein said synthetic cationic polymer comprises monomers selected from the group consisting of dimethylaminoethyl methacrylate, dimethylaminopropyl methacrylate, ditertiobutylaminoethyl methacrylate, dimethylaminomethyl methacrylamide, dimethylaminopropyl methacrylamide, ethylenimine, vinylamine, 2-vinylpyridine, 4-vinylpyridine, trimethylammonium ethyl methacrylate chloride, trimethylammonium ethyl methacrylate methyl sulphate, dimethylammonium ethyl methacrylate benzyl chloride, 4-benzoylbenzyl dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl methacrylamido chloride, trimethyl ammonium propyl methacrylamido chloride, vinylbenzyl trimethyl ammonium chloride, diallyldimethyl ammonium chloride, trimethylammonium ethyl methacrylate chloride, trimethylammonium ethyl methacrylate methyl sulphate, dimethylammonium ethyl methacrylate benzyl chloride, 4-benzoylbenzyl dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl methacrylamido chloride, trimethyl ammonium propyl methacrylamido chloride, vinylbenzyl trimethyl ammonium chloride and trimethyl ammonium propyl methacrylamido chloride.
11. A hair care composition according to any one of claims 1 to 10, further comprising an ingredient selected from the group consisting of oily conditioning agents, hydrocarbon oils, polyolefins, fatty esters, fluorinated conditioning compounds, fatty alcohols, quaternary ammonium compounds, polyethylene glycols, anti-dandruff actives, anti-microbial actives, inorganic particles, synthetic particles, opacifying agents, suspending agents, propellants, paraffinic

hydrocarbons, monovalent salts, divalent salts, fragrances, vitamins, chelating agents, colorants, pigments, dyes and mixtures thereof.

12. A hair care composition according to claim 11, wherein said oily conditioning agent is a silicone conditioning agent having a particle size of from about 1 μm to about 50 μm .
13. A hair care composition according to claim 11, wherein said oily conditioning agent is a silicone conditioning agent having a particle size of less than 1 μm .
14. A hair care composition according to claim 11, wherein said oily conditioning agent is polydimethylsiloxane.
15. A hair care composition according to any one of claims 1 to 14, wherein said composition is a multiple phase composition.
16. A hair care composition comprising:
 - a) an anionic surfactant;
 - b) a naturally derived cationic polymer having a cationic charge density of less than 4 meq/g which forms an isotropic coacervate in said hair care composition;
 - c) a synthetic cationic polymer having a cationic charge density of greater than or equal to 4 meq/g which forms a lyotropic liquid crystal coacervate in said hair care composition;
 - d) an anionic polymeric thickener; and
 - e) water;wherein the composition is formed upon mixing a pre-mixture of the synthetic cationic polymer and the anionic surfactant with the naturally derived cationic polymer.
17. A hair care composition comprising:

- a) an anionic surfactant;
- b) a lyotropic liquid crystal coacervate formed in a premix of a cationic polymer having a charge density of greater than or equal to 4 meq/g with said anionic surfactant;
- c) a cationic polymer having a charge density of less than 4 meq/g, which forms an isotropic coacervate in the neat composition or upon dilution with water;
- d) an anionic polymeric thickener; and
- e) water.

18. A hair care composition of any one of claims 1 to 17, wherein said anionic polymeric thickener comprises acrylates copolymer.

19. A method of making a hair care composition comprising:

- a) an anionic surfactant;
- b) a first cationic polymer having a cationic charge density of less than 4 meq/g, wherein said first cationic polymer forms an isotropic coacervate in the neat composition or upon dilution with water;
- c) a second cationic polymer having a cationic charge density of greater than or equal to 4 meq/g, wherein said second cationic polymer forms a lyotropic liquid crystal coacervate upon combination with said anionic surfactant; and
- d) water;

wherein prior to combining said first cationic polymer and said second cationic polymer, said second cationic polymer is mixed with said anionic surfactant to form a premix.

20. A method according to claim 19, wherein said premix is added to additional ingredients comprising an anionic polymeric thickener prior to combining the premix with the first cationic polymer.

21. A method according to claim 20, wherein said anionic polymeric thickener comprises acrylates copolymer.
22. A method according to any one of claims 19 to 21, wherein said first cationic polymer is a naturally derived cationic polymer.
23. A method according to claim 22, wherein said naturally derived cationic polymer is selected from the group consisting of celluloses, starches, guars, and non-guar-galactomannans.
24. A method according to claim 22 or 23, wherein said naturally derived cationic polymer has a cationic charge density of less than or equal to 2 meq/g.
25. A method according to claim 22, wherein said naturally derived cationic polymer is polyquaternium-10.
26. A method according to claim 25, wherein said polyquaternium-10 has a cationic charge density of about 1.9 meq/g.
27. A method according to claim 22, wherein said naturally derived cationic polymer is cationic cassia.
28. A method according to any one of claims 19 to 27 wherein said second cationic polymer is a synthetic cationic polymer.
29. A method according to claim 28, wherein said synthetic cationic polymer comprises monomers selected from the group consisting of dimethylaminoethyl methacrylate, dimethylaminoethyl acrylate, dimethylaminopropyl methacrylate, dimethylaminopropyl acrylate, ditertiobutylaminoethyl methacrylate, ditertiobutylaminoethyl acrylate, dimethylaminomethyl methacrylamide, dimethylaminomethyl acrylamide, dimethylaminopropyl methacrylamide;

dimethylaminopropyl acrylamide, ethylenimine, vinylamine, 2-vinylpyridine, 4-vinylpyridine, trimethylammonium ethyl methacrylate chloride, trimethylammonium ethyl acrylate chloride, trimethylammonium ethyl methacrylate methyl sulphate, trimethylammonium ethyl acrylate methyl sulphate, dimethylammonium ethyl methacrylate benzyl chloride, dimethylammonium ethyl acrylate benzyl chloride, 4-benzoylbenzyl dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl methacrylamido chloride, trimethyl ammonium ethyl acrylamido chloride, trimethyl ammonium propyl methacrylamido chloride, trimethyl ammonium propyl acrylamido chloride, vinylbenzyl trimethyl ammonium chloride, diallyldimethyl ammonium chloride, trimethylammonium ethyl methacrylate chloride, trimethylammonium ethyl acrylate chloride, trimethylammonium ethyl methacrylate methyl sulphate, trimethylammonium ethyl acrylate methyl sulphate, dimethylammonium ethyl methacrylate benzyl chloride, dimethylammonium ethyl acrylate benzyl chloride, 4-benzoylbenzyl dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl methacrylamido chloride, trimethyl ammonium ethyl acrylamido chloride, trimethyl ammonium propyl methacrylamido chloride, trimethyl ammonium propyl acrylamido chloride, vinylbenzyl trimethyl ammonium chloride, trimethyl ammonium propyl methacrylamido chloride, and trimethyl ammonium propylacrylamido chloride.

30. A method according to claim 28, wherein said synthetic cationic polymer comprises monomers selected from the group consisting of dimethylaminoethyl methacrylate, dimethylaminopropyl methacrylate, ditertiobutylaminoethyl methacrylate, dimethylaminomethyl methacrylamide, dimethylaminopropyl methacrylamide, ethylenimine, vinylamine, 2-vinylpyridine, 4-vinylpyridine, trimethylammonium ethyl methacrylate chloride, trimethylammonium ethyl methacrylate methyl sulphate, dimethylammonium ethyl methacrylate benzyl chloride, 4-benzoylbenzyl dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl methacrylamido chloride, trimethyl ammonium propyl methacrylamido chloride, trimethyl ammonium propyl acrylamido chloride, vinylbenzyl trimethyl ammonium chloride, trimethyl ammonium propyl methacrylamido chloride, and trimethyl ammonium propylacrylamido chloride, trimethyl ammonium propyl

methacrylamido chloride, vinylbenzyl trimethyl ammonium chloride, diallyldimethyl ammonium chloride, trimethylammonium ethyl methacrylate chloride, trimethylammonium ethyl methacrylate methyl sulphate, dimethylammonium ethyl methacrylate benzyl chloride, 4-benzoylbenzyl dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl methacrylamido chloride, trimethyl ammonium propyl methacrylamido chloride, vinylbenzyl trimethyl ammonium chloride and trimethyl ammonium propyl methacrylamido chloride.

31. A method according to any one of claims 19 to 30, further comprising an ingredient selected from the group consisting of oily conditioning agents, hydrocarbon oils, polyolefins, fatty esters, fluorinated conditioning compounds, fatty alcohols, quaternary ammonium compounds, polyethylene glycols, anti-dandruff actives, anti-microbial actives, inorganic particles, synthetic particles, opacifying agents, suspending agents, propellants, paraffinic hydrocarbons, monovalent salts, divalent salts, fragrances, vitamins, chelating agents, colorants, pigments, dyes and mixtures thereof.
32. A method according to claim 31, wherein said oily conditioning agent is a silicone conditioning agent having a particle size of from about 1 μm to about 50 μm .
33. A method according to claim 31, wherein said oily conditioning agent is a silicone conditioning agent having a particle size of less than 1 μm .
34. A method according to claim 31, wherein said oily conditioning agent is polydimethylsiloxane.
35. A method according to any one of claims 19 to 34, wherein said composition is a multiple phase composition.

36. A hair care composition made by the method according to any one of claims 19 to 35.