



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) **Date de dépôt PCT/PCT Filing Date:** 2022/06/03
(87) **Date publication PCT/PCT Publication Date:** 2022/12/08
(85) **Entrée phase nationale/National Entry:** 2023/11/08
(86) **N° demande PCT/PCT Application No.:** EP 2022/065165
(87) **N° publication PCT/PCT Publication No.:** 2022/253997
(30) **Priorité/Priority:** 2021/06/03 (IN202111024711)

(51) **Cl.Int./Int.Cl. C07D 281/10** (2006.01),
A61P 1/16 (2006.01), **C07D 285/36** (2006.01)
(71) **Demandeur/Applicant:**
ALBIREO AB, SE
(72) **Inventeurs/Inventors:**
STARKE, INGEMAR, SE;
KULKARNI, SANTOSH S., IN;
GILLBERG, PER-GORAN, SE
(74) **Agent:** C6 PATENT GROUP INCORPORATED,
OPERATING AS THE "CARBON PATENT GROUP"

(54) **Titre : COMPOSES DE BENZOTHIA(DI)AZEPINE ET LEUR UTILISATION EN TANT QUE MODULATEURS DE L'ACIDE BILIAIRE**
(54) **Title: BENZOTHIA(DI)AZEPINE COMPOUNDS AND THEIR USE AS BILE ACID MODULATORS**

(57) **Abrégé/Abstract:**

The invention relates to 1,5-benzothiazepine and 1,2,5-benzothiadiazepine derivatives of formula (I). These compounds are bile acid modulators having apical sodium-dependent bile acid transporter(ASBT) and/or liver bile acid transport (LBAT) inhibitory activity. The invention also relates to pharmaceutical compositions comprising these compounds and to the use of these compounds in the treatment of cardiovascular diseases, fatty acid metabolism and glucose utilization disorders, gastrointestinal diseases and liver diseases.

Date Submitted: 2023/11/08

CA App. No.: 3218474

Abstract:

The invention relates to 1,5-benzothiazepine and 1,2,5-benzothiadiazepine derivatives of formula (I). These compounds are bile acid modulators having apical sodium-dependent bile acid transporter (ASBT) and/or liver bile acid transport (LBAT) inhibitory activity. The invention also relates to pharmaceutical compositions comprising these compounds and to the use of these compounds in the treatment of cardiovascular diseases, fatty acid metabolism and glucose utilization disorders, gastrointestinal diseases and liver diseases.

BENZOTHIA(DI)AZEPINE COMPOUNDS AND THEIR USE AS BILE ACID MODULATORS

CROSS-REFERENCE TO RELATED APPLICATIONS

- 5 This application claims priority to Indian Patent Application No. 202111024711, filed June 3, 2021, the disclosure of which is incorporated herein by reference in its entirety.

TECHNICAL FIELD

- 10 The invention relates to 1,5-benzothiazepine and 1,2,5-benzothiadiazepine derivatives of formula (I). These compounds are bile acid modulators having apical sodium-dependent bile acid transporter (ASBT) and/or liver bile acid transport (LBAT) inhibitory activity. The invention also relates to pharmaceutical compositions comprising these compounds and to the use of these compounds in the treatment of cardiovascular diseases, fatty acid metabolism and glucose utilization disorders,
15 gastrointestinal diseases and liver diseases.

BACKGROUND

- Bile acids are physiological detergents that play an important role in the intestinal absorption and
20 transport of lipids, nutrients and vitamins. They are also signaling molecules that activate nuclear receptors and cell signaling pathways that regulate lipid, glucose and energy metabolism. Bile acids are steroid acids that are synthesized from cholesterol in the liver and stored in the gallbladder as mixed micelles. During digestion, the duodenum triggers the release of hormones that cause the gallbladder to contract, thereby releasing bile acids in the small intestine where they enable
25 absorption of fat-soluble vitamins and cholesterol. When they reach the ileum, bile acids are reabsorbed from the intestine and secreted into portal blood to return to the liver via the portal venous circulation. Over 90% of the bile acids are thus recycled and returned to the liver. These bile acids are then transported across the sinusoidal membrane of hepatocytes and re-secreted across the canalicular membrane into bile. In this first pass, 75-90% of bile acids are taken up by
30 hepatocytes, completing one round of enterohepatic circulation. The fraction of bile acids that escapes being cleared in the liver enters the systemic circulation where the free bile acids are filtered by the renal glomerulus, efficiently reclaimed in the proximal tubules and exported back into the systemic circulation. Interestingly, most of the bile acids secreted across the canalicular membrane into bile are derived from the recirculating pool with less than 10% coming from new *de novo* hepatic
35 synthesis. The small fraction of bile acids that is not reabsorbed in the ileum reaches the colon.

Within the intestinal lumen, the primary bile acids are transformed into secondary bile acids under the action of intestinal bacteria, mainly by single or dual dehydroxylation reactions of the steroid nucleus. The bile acids that escape intestinal absorption are thereafter excreted into the faeces.

5 Overall, the efficient transport system helps maintain a constant bile acid pool, ensuring sufficiently high levels of conjugated bile acids in the intestine to promote lipid absorption as well as reduce the small intestinal bacterial load. The system also minimizes fecal and urinary bile acid loss and protects the intestinal and hepatobiliary compartments by eliminating potentially cytotoxic detergents (as reviewed by Kusters and Karpen (*Xenobiotica* 2008, vol. 38, p. 1043-1071); by Chiang (*J. Lipid Res.*
10 2009, vol. 50, p. 1955-1966); and by Dawson (*Handb. Exp. Pharmacol.* 2011, vol. 201, p. 169-203)).

The regulation of the bile acid pool size has been found to play a key role in cholesterol homeostasis by hepatic conversion of cholesterol to bile acid, which represents a major route for elimination of cholesterol from the body. The liver plays an essential role in removing endogenous and xenobiotic
15 compounds from the body. The normal hepatobiliary secretion and enterohepatic circulation are required for the elimination of endogenous compounds such as cholesterol and bilirubin and their metabolites from the body, thereby maintaining lipid and bile acid homeostasis. (Kusters and Karpen, *Xenobiotica* 2008, vol. 38, p. 1043-1071).

20 The reabsorption of bile acids in the ileum may be inhibited by apical sodium-dependent bile acid transporter (ASBT) inhibitor compounds. Inhibition of bile acid reabsorption has been reported useful in the treatment of several diseases, including dyslipidemia, diabetes, obesity, constipation, cholestatic liver diseases, non-alcoholic steatohepatitis and other hepatic diseases. A number of ASBT inhibitor compounds has been disclosed over the past decades, see e.g. WO 93/16055,
25 WO 94/18183, WO 94/18184, WO 96/05188, WO 96/08484, WO 96/16051, WO 97/33882, WO 98/03818, WO 98/07449, WO 98/40375, WO 99/35135, WO 99/64409, WO 99/64410, WO 00/47568, WO 00/61568, WO 00/38725, WO 00/38726, WO 00/38727, WO 00/38728, WO 00/38729, WO 01/66533, WO 01/68096, WO 02/32428, WO 02/50051, WO 03/020710, WO 03/022286, WO 03/022825, WO 03/022830, WO 03/061663, WO 03/091232, WO 03/106482,
30 WO 2004/006899, WO 2004/076430, WO 2007/009655, WO 2007/009656, WO 2011/137135, WO 2019/234077, WO 2020/161216, WO 2020/161217, WO 2021/110883, WO 2021/110884, WO 2021/110885, WO 2021/110886, WO 2021/110887, WO 2022/029101, DE 19825804, EP 864582, EP 489423, EP 549967, EP 573848, EP 624593, EP 624594, EP 624595, EP 624596, EP 0864582, EP 1173205, EP 1535913 and EP 3210977.


35

Despite the number of ASBT inhibitor compounds that have been previously reported, there is a need for additional bile acid modulating compounds that have an optimized profile with respect to potency, selectivity and bioavailability.

5 DETAILED DESCRIPTION OF THE INVENTION

It has been discovered that certain 1,5-benzothiazepine and 1,2,5-benzothiadiazepine derivatives are potent inhibitors of apical sodium-dependent bile acid transporter (ASBT) and/or liver bile acid transporter (LBAT), and may be useful for treating diseases wherein inhibition of bile acid circulation is desirable.

In a first aspect, the invention relates to a compound of formula (I)



15 wherein

M is selected from -CH₂- and -NR⁵-;

R¹ is C₁₋₄ alkyl;

R² is independently selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄ alkyl, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkoxy, cyano, nitro, amino, *N*-(C₁₋₄ alkyl)amino, *N,N*-di(C₁₋₄ alkyl)amino, C₁₋₆ alkylcarbonylamino, C₃₋₆ cycloalkylcarbonylamino, *N*-(C₁₋₄ alkyl)amino-carbonyl, *N,N*-di(C₁₋₄ alkyl)aminocarbonyl, C₁₋₄ alkyloxycarbonylamino, C₃₋₆ cycloalkyloxycarbonyl-amino, C₁₋₄ alkylsulfonamido and C₃₋₆ cycloalkylsulfonamido;

n is an integer 1, 2 or 3;

R³ is selected from the group consisting of hydrogen, halogen, cyano, C₁₋₄ alkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, C₃₋₆ cycloalkyloxy, C₁₋₄ alkylthio, C₃₋₆ cycloalkylthio, amino, *N*-(C₁₋₄ alkyl)amino and *N,N*-di(C₁₋₄ alkyl)amino;

R^{4A} and R^{4B} are each independently selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄ alkyl and C₁₋₄ alkoxy; or R^{4A} and R^{4B}, together with the carbon atom to which they are attached, form a 3- to 5-membered saturated carbocyclic ring;

R^{4C} and R^{4D} are each independently selected from the group consisting of hydrogen and C₁₋₄ alkyl; and

R⁵ is selected from the group consisting of hydrogen and C₁₋₄ alkyl; or a pharmaceutically acceptable salt thereof.

In some embodiments, R¹ is C₂₋₄ alkyl. In a preferred embodiment, R¹ is *n*-propyl. In another preferred embodiment, R¹ is *n*-butyl.

In some embodiments, R² is selected from the group consisting of hydrogen, fluoro, chloro, bromo, hydroxy, methoxy, amino, methylamino and dimethylamino. In a preferred embodiment, *n* is 1, i.e. the phenyl-ring is substituted with only one substituent R². In another preferred embodiment, R² is in the para-position.


In some embodiments, R³ is selected from the group consisting of fluoro, chloro, bromo, methyl, cyclopropyl, methoxy, ethoxy, methylthio, ethylthio, amino, methylamino and dimethylamino.

In some embodiments, R^{4A} and R^{4B} are each independently selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄ alkyl and C₁₋₄ alkoxy, or R^{4A} and R^{4B}, together with the carbon atom to which they are attached, form a cyclopropyl ring. In some embodiments, R^{4A} and R^{4B} are each independently fluoro, methyl or methoxy, or together with the carbon atom to which they are attached form a cyclopropyl ring.

In some embodiments, R^{4C} and R^{4D} are each independently hydrogen or methyl. In some embodiments, R^{4C} and R^{4D} are each hydrogen.

In some embodiments, R⁵ is hydrogen. In some embodiments, R⁵ is methyl.

In a preferred embodiment, the compound of formula (I) is a compound of formula (I-a):



wherein

M is selected from the group consisting of $-CH_2-$, $-NH-$ and $-NCH_3-$;

5 R^1 is C_{2-4} alkyl;


R^2 is independently selected from the group consisting of hydrogen, halogen, hydroxy, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkoxy, amino, N -(C_{1-4} alkyl)amino, N,N -di(C_{1-4} alkyl)amino; n is an integer 1 or 2;

10 R^3 is selected from the group consisting of halogen, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{1-4} alkylthio, amino, N -(C_{1-4} alkyl)amino and N,N -di(C_{1-4} alkyl)amino;

R^{4A} and R^{4B} are each independently selected from the group consisting of hydrogen, halogen, hydroxy, C_{1-4} alkyl and C_{1-4} alkoxy, or R^{4A} and R^{4B} , together with the carbon atom to which they are attached, form a cyclopropyl ring; or a pharmaceutically acceptable salt thereof.

15

In a preferred embodiment, the compound of formula (I) is a compound of formula (I-b):



wherein

20

M is selected from the group consisting of $-CH_2-$, $-NH-$ and $-N(CH_3)-$;

R^1 is C_{2-4} alkyl, more preferably n -propyl or n -butyl;

R^2 is independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, hydroxy, methoxy, amino, methylamino, dimethylamino;

R³ is selected from the group consisting of fluoro, chloro, bromo, methyl, cyclopropyl, methoxy, ethoxy, methylthio, ethylthio, amino, methylamino and dimethylamino;

R^{4A} and R^{4B} are each independently hydrogen, fluoro, methyl, methoxy or ethoxy, or together with the carbon atom to which they are attached form a cyclopropyl ring;

5 or a pharmaceutically acceptable salt thereof.

Preferred compounds of the invention are compounds of formula (I-b), as defined above, wherein M, R¹, R², R³, R^{4A} and R^{4B} are as indicated in Table 1 below, or a pharmaceutically acceptable salt thereof:

10 **Table 1**

M	R ¹	R ²	R ³	R ^{4A}	R ^{4B}
CH ₂	CH ₂ CH ₂ CH ₃	H	SCH ₃	CH ₃	CH ₃
CH ₂	CH ₂ CH ₂ CH ₃	H	SCH ₃	-CH ₂ CH ₂ -	
CH ₂	CH ₂ CH ₂ CH ₃	H	SCH ₃	F	F
CH ₂	CH ₂ CH ₂ CH ₃	F	SCH ₃	CH ₃	CH ₃
CH ₂	CH ₂ CH ₂ CH ₃	F	SCH ₃	-CH ₂ CH ₂ -	
CH ₂	CH ₂ CH ₂ CH ₃	F	SCH ₃	F	F
NH	CH ₂ CH ₂ CH ₃	H	SCH ₃	CH ₃	CH ₃
NH	CH ₂ CH ₂ CH ₃	H	SCH ₃	-CH ₂ CH ₂ -	
NH	CH ₂ CH ₂ CH ₃	H	SCH ₃	F	F
NH	CH ₂ CH ₂ CH ₃	F	SCH ₃	CH ₃	CH ₃
NH	CH ₂ CH ₂ CH ₃	F	SCH ₃	-CH ₂ CH ₂ -	
NH	CH ₂ CH ₂ CH ₃	F	SCH ₃	F	F
NCH ₃	CH ₂ CH ₂ CH ₃	H	SCH ₃	CH ₃	CH ₃
NCH ₃	CH ₂ CH ₂ CH ₃	H	SCH ₃	-CH ₂ CH ₂ -	
NCH ₃	CH ₂ CH ₂ CH ₃	H	SCH ₃	F	F
NCH ₃	CH ₂ CH ₂ CH ₃	F	SCH ₃	CH ₃	CH ₃
NCH ₃	CH ₂ CH ₂ CH ₃	F	SCH ₃	-CH ₂ CH ₂ -	
NCH ₃	CH ₂ CH ₂ CH ₃	F	SCH ₃	F	F
CH ₂	CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	CH ₃	CH ₃
CH ₂	CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	-CH ₂ CH ₂ -	
CH ₂	CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	F	F
CH ₂	CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	CH ₃	CH ₃
CH ₂	CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	-CH ₂ CH ₂ -	
CH ₂	CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	F	F
NH	CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	CH ₃	CH ₃
NH	CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	-CH ₂ CH ₂ -	
NH	CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	F	F
NH	CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	CH ₃	CH ₃
NH	CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	-CH ₂ CH ₂ -	

M	R ¹	R ²	R ³	R ^{4A}	R ^{4B}
NH	CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	F	F
NCH ₃	CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	CH ₃	CH ₃
NCH ₃	CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	-CH ₂ CH ₂ -	
NCH ₃	CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	F	F
NCH ₃	CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	CH ₃	CH ₃
NCH ₃	CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	-CH ₂ CH ₂ -	
NCH ₃	CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	F	F
CH ₂	CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	CH ₃	CH ₃
CH ₂	CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	-CH ₂ CH ₂ -	
CH ₂	CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	F	F
CH ₂	CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	CH ₃	CH ₃
CH ₂	CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	-CH ₂ CH ₂ -	
CH ₂	CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	F	F
NH	CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	CH ₃	CH ₃
NH	CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	-CH ₂ CH ₂ -	
NH	CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	F	F
NH	CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	CH ₃	CH ₃
NH	CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	-CH ₂ CH ₂ -	
NH	CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	F	F
NCH ₃	CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	CH ₃	CH ₃
NCH ₃	CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	-CH ₂ CH ₂ -	
NCH ₃	CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	F	F
NCH ₃	CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	CH ₃	CH ₃
NCH ₃	CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	-CH ₂ CH ₂ -	
NCH ₃	CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	F	F
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₃	CH ₃	CH ₃
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₃	-CH ₂ CH ₂ -	
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₃	F	F
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₃	CH ₃	CH ₃
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₃	-CH ₂ CH ₂ -	
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₃	F	F
NH	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₃	CH ₃	CH ₃
NH	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₃	-CH ₂ CH ₂ -	
NH	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₃	F	F
NH	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₃	CH ₃	CH ₃
NH	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₃	-CH ₂ CH ₂ -	
NH	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₃	F	F
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₃	CH ₃	CH ₃
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₃	-CH ₂ CH ₂ -	
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₃	F	F
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₃	CH ₃	CH ₃
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₃	-CH ₂ CH ₂ -	

M	R ¹	R ²	R ³	R ^{4A}	R ^{4B}
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₃	F	F
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	CH ₃	CH ₃
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	-CH ₂ CH ₂ -	
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	F	F
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	CH ₃	CH ₃
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	-CH ₂ CH ₂ -	
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	F	F
NH	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	CH ₃	CH ₃
NH	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	-CH ₂ CH ₂ -	
NH	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	F	F
NH	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	CH ₃	CH ₃
NH	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	-CH ₂ CH ₂ -	
NH	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	F	F
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	CH ₃	CH ₃
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	-CH ₂ CH ₂ -	
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	H	SCH ₂ CH ₃	F	F
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	CH ₃	CH ₃
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	-CH ₂ CH ₂ -	
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	F	SCH ₂ CH ₃	F	F
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	CH ₃	CH ₃
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	-CH ₂ CH ₂ -	
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	F	F
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	CH ₃	CH ₃
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	-CH ₂ CH ₂ -	
CH ₂	CH ₂ CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	F	F
NH	CH ₂ CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	CH ₃	CH ₃
NH	CH ₂ CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	-CH ₂ CH ₂ -	
NH	CH ₂ CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	F	F
NH	CH ₂ CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	CH ₃	CH ₃
NH	CH ₂ CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	-CH ₂ CH ₂ -	
NH	CH ₂ CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	F	F
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	CH ₃	CH ₃
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	-CH ₂ CH ₂ -	
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	H	N(CH ₃) ₂	F	F
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	CH ₃	CH ₃
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	-CH ₂ CH ₂ -	
NCH ₃	CH ₂ CH ₂ CH ₂ CH ₃	F	N(CH ₃) ₂	F	F

In a particular embodiment, the compound of formula (I) is selected from the group consisting of:

3-((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

(S)-3-((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

(R)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

5 1-(((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

(S)-1-(((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

10 (R)-1-(((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

3-((3-Butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

(S)-3-((3-Butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

15 (R)-3-((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

1-(((3-Butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

20 (S)-1-(((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

(R)-1-(((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

3-((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

25 1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

1-(((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

30 (S)-1-(((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

(R)-1-(((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

3-((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

(S)-3-((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

(R)-3-((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

5 1-(((3-butyl-7-(ethylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

10 (S)-1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

(R)-1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

1-(((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

15 (S)-1-(((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

(R)-1-(((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

20 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-difluoropropanoic acid;

(S)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-difluoropropanoic acid;

(R)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-difluoropropanoic acid;

25 3-((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid;

(S)-3-(((R)-3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid;

30 (S)-3-(((S)-3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid;

(R)-3-(((R)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid;

(R)-3-(((S)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid;

3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)propanoic acid;

(S)-3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)propanoic acid;

5 (R)-3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)propanoic acid;

3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-ethoxypropanoic acid;

10 3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)-2-hydroxypropanoic acid;

3-((3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)propanoic acid;

3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid;

15 3-(((S)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid;

3-(((R)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid;

20 (S)-3-(((R)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid;

(R)-3-(((R)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid;

(S)-3-(((S)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid;

25 (R)-3-(((S)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid; and

3-((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-2-methyl-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid;

or a pharmaceutically acceptable salt thereof.

30

As used herein, the term "halo" refers to fluoro, chloro, bromo and iodo.

As used herein, the term "C₁₋₆ alkyl" refers to a straight or branched alkyl group having from 1 to 6 carbon atoms, and the term "C₁₋₄ alkyl" refers to a straight or branched alkyl group having from 1 to 4

carbon atoms. Examples of C₁₋₄ alkyl include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl and tert-butyl.

As used herein, the term "C₁₋₄ haloalkyl" refers to a straight or branched C₁₋₄ alkyl group, as defined herein, wherein one or more hydrogen atoms have been replaced with halogen. Examples of C₁₋₄ haloalkyl include chloromethyl, fluoroethyl and trifluoromethyl.

As used herein, the terms "C₁₋₄ alkoxy" and "C₁₋₄ alkylthio" refer to a straight or branched C₁₋₄ alkyl group attached to the remainder of the molecule through an oxygen or sulphur atom, respectively.

As used herein, the term "C₃₋₆ cycloalkyl" refers to a monocyclic saturated hydrocarbon ring having from 3 to 6 carbon atoms. Examples of C₃₋₆ cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

The term "aryl" denotes an aromatic monocyclic ring composed of 6 carbon atoms or an aromatic bicyclic ring system composed of 10 carbon atoms. Examples of aryl include phenyl, naphthyl and azulenyl.

The term "amino" refers to an -NH₂ group. As used herein, the terms "*N*-(C₁₋₄ alkyl)amino" and "*N,N*-di(C₁₋₄ alkyl)amino" refer to an amino group wherein one or both hydrogen atom(s), respectively, are replaced with a straight or branched C₁₋₄ alkyl group. Examples of *N*-(C₁₋₄ alkyl)amino include methylamino, ethylamino and tert-butylamino, and examples of *N,N*-di-(C₁₋₄ alkyl)amino include dimethylamino and diethylamino.

As used herein, the term "*N*-(aryl-C₁₋₄ alkyl)amino" refers to an amino group wherein a hydrogen atom is replaced with an aryl-C₁₋₄ alkyl group. Examples of *N*-(aryl-C₁₋₄ alkyl)amino include benzylamino and phenylethylamino. The term "C₁₋₆ alkylcarbonylamino" refers to an amino group wherein a hydrogen atom is replaced with a C₁₋₆ alkylcarbonyl group. Examples of C₁₋₆ alkanoylamino include acetylamino and tert-butylcarbonylamino. The term "C₁₋₄ alkyloxycarbonylamino" refers to an amino group wherein a hydrogen atom is replaced with a C₁₋₄ alkyloxycarbonyl group. An example of C₁₋₄ alkyloxycarbonylamino is tert-butoxycarbonylamino. The terms "C₁₋₄ alkylsulfonamido" and "C₃₋₆ cycloalkylsulfonamido" refer to an amino group wherein a hydrogen atom is replaced with a C₁₋₄ alkylsulfonyl or a C₃₋₆ cycloalkylsulfonyl group, respectively.

As used herein, the term "pharmaceutically acceptable" refers to those compounds, materials, compositions and/or dosage forms that are suitable for human pharmaceutical use and that are generally safe, non-toxic and neither biologically nor otherwise undesirable.

5 As used herein, the term "about" refers to a value or parameter herein that includes (and describes) embodiments that are directed to that value or parameter per se. For example, description referring to "about 20" includes description of "20." Numeric ranges are inclusive of the numbers defining the range. Generally speaking, the term "about" refers to the indicated value of the variable and to all values of the variable that are within the experimental error of the indicated value (e.g., within the
10 95% confidence interval for the mean) or within 10 percent of the indicated value, whichever is greater.

The 1,5-benzothiazepine and 1,2,5-benzothiadiazepine compounds of formula (I), or pharmaceutically acceptable salts thereof, are inhibitors of the apical sodium-dependent bile acid
15 transporter (ASBT inhibitors), of the liver bile acid transporter (LBAT inhibitors), or of both the apical sodium-dependent bile acid and liver bile acid transporters (dual ASBT/LBAT inhibitors). They are therefore useful in the treatment or prevention of conditions, disorders and diseases wherein inhibition of bile acid circulation is desirable, such as cardiovascular diseases, fatty acid metabolism and glucose utilization disorders, gastrointestinal diseases and liver diseases.

20 Cardiovascular diseases and disorders of fatty acid metabolism and glucose utilization include, but are not limited to, hypercholesterolemia; disorders of fatty acid metabolism; type 1 and type 2 diabetes mellitus; complications of diabetes, including cataracts, micro- and macrovascular diseases, retinopathy, neuropathy, nephropathy and delayed wound healing, tissue ischaemia, diabetic foot, arteriosclerosis, myocardial infarction, acute coronary syndrome, unstable angina pectoris, stable
25 angina pectoris, stroke, peripheral arterial occlusive disease, cardiomyopathy, heart failure, heart rhythm disorders and vascular restenosis; diabetes-related diseases such as insulin resistance (impaired glucose homeostasis), hyperglycemia, hyperinsulinemia, elevated blood levels of fatty acids or glycerol, obesity, dyslipidemia, hyperlipidemia including hypertriglyceridemia, metabolic
30 syndrome (syndrome X), atherosclerosis and hypertension; and for increasing high density lipoprotein levels.

Gastrointestinal diseases and disorders include constipation (including chronic constipation, functional constipation, chronic idiopathic constipation (CIC), intermittent/sporadic constipation,
35 constipation secondary to diabetes mellitus, constipation secondary to stroke, constipation

secondary to chronic kidney disease, constipation secondary to multiple sclerosis, constipation secondary to Parkinson's disease, constipation secondary to systemic sclerosis, drug induced constipation, irritable bowel syndrome with constipation (IBS-C), irritable bowel syndrome mixed (IBS-M), pediatric functional constipation and opioid induced constipation); Crohn's disease; primary
 5 bile acid malabsorption; irritable bowel syndrome (IBS); inflammatory bowel disease (IBD); ileal inflammation; and reflux disease and complications thereof, such as Barrett's esophagus, bile reflux esophagitis and bile reflux gastritis.

A liver disease as defined herein is any disease in the liver and in organs connected therewith, such
 10 as the pancreas, portal vein, the liver parenchyma, the intrahepatic biliary tree, the extrahepatic biliary tree, and the gall bladder. In some cases, a liver disease a bile acid-dependent liver disease. Liver diseases and disorders include, but are not limited to, an inherited metabolic disorder of the liver; inborn errors of bile acid synthesis; congenital bile duct anomalies; biliary atresia; post-Kasai biliary atresia; post-liver transplantation biliary atresia; neonatal hepatitis; neonatal cholestasis;
 15 hereditary forms of cholestasis; cerebrotendinous xanthomatosis; a secondary defect of BA synthesis; Zellweger's syndrome; cystic fibrosis-associated liver disease; alpha1-antitrypsin deficiency; Alagilles syndrome (ALGS); Byler syndrome; a primary defect of bile acid (BA) synthesis; progressive familial intrahepatic cholestasis (PFIC) including PFIC-1, PFIC-2, PFIC-3 and non-specified PFIC, post-biliary diversion PFIC and post-liver transplant PFIC; benign recurrent intrahepatic
 20 cholestasis (BRIC) including BRIC1, BRIC2 and non-specified BRIC, post-biliary diversion BRIC and post-liver transplant BRIC; autoimmune hepatitis; primary biliary cirrhosis (PBC); liver fibrosis; non-alcoholic fatty liver disease (NAFLD); non-alcoholic steatohepatitis (NASH); portal hypertension; cholestasis; Down syndrome cholestasis; drug-induced cholestasis; intrahepatic cholestasis of pregnancy (jaundice during pregnancy); intrahepatic cholestasis; extrahepatic cholestasis; parenteral
 25 nutrition associated cholestasis (PNAC); low phospholipid-associated cholestasis; lymphedema cholestasis syndrome 1 (LCS1); primary sclerosing cholangitis (PSC); immunoglobulin G4 associated cholangitis; primary biliary cholangitis; cholelithiasis (gallstones); biliary lithiasis; choledocholithiasis; gallstone pancreatitis; Caroli disease; malignancy of bile ducts; malignancy causing obstruction of the biliary tree; biliary strictures; AIDS cholangiopathy; ischemic cholangiopathy; pruritus due to
 30 cholestasis or jaundice; pancreatitis; chronic autoimmune liver disease leading to progressive cholestasis; hepatic steatosis; alcoholic hepatitis; acute fatty liver; fatty liver of pregnancy; drug-induced hepatitis; iron overload disorders; congenital bile acid synthesis defect type 1 (BAS defect type 1); drug-induced liver injury (DILI); hepatic fibrosis; congenital hepatic fibrosis; hepatic cirrhosis; Langerhans cell histiocytosis (LCH); neonatal ichthyosis sclerosing cholangitis (NISCH); erythropoietic
 35 protoporphyria (EPP); idiopathic adulthood ductopenia (IAD); idiopathic neonatal hepatitis (INH); non

syndromic paucity of interlobular bile ducts (NS PILBD); North American Indian childhood cirrhosis (NAIC); hepatic sarcoidosis; amyloidosis; necrotizing enterocolitis; serum bile acid-caused toxicities, including cardiac rhythm disturbances (e.g., atrial fibrillation) in setting of abnormal serum bile acid profile, cardiomyopathy associated with liver cirrhosis (“cholecardia”), and skeletal muscle wasting associated with cholestatic liver disease; polycystic liver disease; viral hepatitis (including hepatitis A, hepatitis B, hepatitis C, hepatitis D and hepatitis E); hepatocellular carcinoma (hepatoma); cholangiocarcinoma; bile acid-related gastrointestinal cancers; and cholestasis caused by tumours and neoplasms of the liver, of the biliary tract and of the pancreas. Compounds of formula (I), or pharmaceutically acceptable salts thereof, are also useful in the enhancement of corticosteroid therapy in liver disease.

Other diseases that may be treated or prevented by the compounds of formula (I), or pharmaceutically acceptable salts thereof, include hyperabsorption syndromes (including abetalipoproteinemia, familial hypobetalipoproteinemia (FHBL), chylomicron retention disease (CRD) and sitosterolemia); hypervitaminosis and osteopetrosis; hypertension; glomerular hyperfiltration; polycystic kidney disease (PKD), including autosomal dominant polycystic kidney disease (ADPKD) and autosomal recessive polycystic kidney disease (ARPKD); and pruritus of renal failure. The compounds are also useful in the protection against liver- or metabolic disease-associated kidney injury.

The transport of bile acids in the human body is controlled by the action of the members of the SLC10 family of solute carrier proteins, in particular by the Na⁺-taurocholate cotransporting polypeptide (NTCP, also called liver bile acid transporter (LBAT); gene symbol *SLC10A1*), which is expressed in the sinusoidal membrane of hepatocytes, and by the apical sodium dependent bile acid transporter (ASBT, also called ileal bile acid transporter (IBAT), ISBT, ABAT or NTCP2; gene symbol *SLC10A2*), which is expressed in the apical membrane of ileal enterocytes, proximal renal tubule cells, biliary epithelium, large cholangiocytes and gallbladder epithelial cells. In the liver, bile acids are efficiently extracted from portal blood by the liver bile acid transporter (LBAT) and re-secreted across the canalicular membrane by the bile salt export pump (BSEP; gene symbol *ABCB11*).

The reabsorption of bile acids in the ileum is handled by the apical sodium-dependent bile acid transporter (ASBT), where it is commonly referred to as ileal bile acid transporter (IBAT). Both LBAT and ASBT function as electrogenic sodium-solute cotransporters that move two or more Na⁺ ions per molecule of solute.

Xenobiotics and endobiotics, including bile acids, are taken up by the liver from portal blood and secreted into bile by distinct transport proteins with individualized substrate specificities. Glycine- and taurine-conjugated bile acids exist in anionic form and are unable to cross membranes by diffusion, and thus, are completely dependent on membrane transport proteins to enter or exit the hepatocyte (Kosters and Karpen, *Xenobiotica* 2008, vol. 38, p. 1043-1071). ASBT and LBAT prefer glycine- and taurine-conjugated bile salts over their unconjugated counterparts and demonstrate a higher affinity for dihydroxy bile salts than for trihydroxy bile salts. No non-bile acid substrates have been identified for ASBT yet, however, LBAT was also found to transport a variety of steroid sulfates, hormones and xenobiotics.

LBAT is not as thoroughly characterized as ASBT in terms of drug inhibition requirements. Dong et al. have identified FDA approved drugs that inhibit human LBAT and compared LBAT and ASBT inhibition requirements. A series of LBAT inhibition studies were performed using FDA approved drugs, in concert with iterative computational model development. Screening studies identified 27 drugs as novel LBAT inhibitors, including irbesartan ($K_i = 11.9 \mu\text{M}$) and ezetimibe ($K_i = 25.0 \mu\text{M}$). The common feature pharmacophore indicated that two hydrophobes and one hydrogen bond acceptor were important for inhibition of LBAT. From 72 drugs screened in vitro, a total of 31 drugs inhibited LBAT, while 51 drugs (i.e. more than half) inhibited ASBT. Hence, while there was inhibitor overlap, ASBT unexpectedly was more permissive to drug inhibition than was LBAT, and this may be related to LBAT's possessing fewer pharmacophore features (Dong et al., *Mol. Pharm.* 2013, vol. 10, p. 1008–1019).

Vaz et al. describe the identification of LBAT deficiency as a new inborn error of metabolism with a relatively mild clinical phenotype. The identification of LBAT deficiency confirms that this transporter is the main import system for conjugated bile salts into the liver, but also indicates that auxiliary transporters are able to sustain the enterohepatic cycle in its absence (Vaz et al., *Hepatology* 2015, vol. 61, p. 260-267). These findings support the hypothesis that LBAT inhibition is a safe mechanism of action, as the hepatocytes still have the possibility to take up the necessary amount of bile acids.

Liu et al. describe the identification of a new type of hypercholanemia that is associated with homozygosity for the p.Ser267Phe mutation in *SLC10A1* (LBAT). The allele frequency of this mutation in gene *SLC10A1* varies in different populations, with the highest incidence occurring in Southern China (8% and 12% in Chinese Han and Dai respectively) and in Vietnam (11%). This "hidden" hypercholanemia was believed to affect 0.64% of the Southern Han, 1.44% of the Dai Chinese population, and 1.21% of the Vietnamese population. An increase in conjugated and unconjugated

serum BA levels in the homozygous individuals was also observed. Liu et al. suggest that this finding is most likely due to reduced BA transport from the portal circulation into hepatocytes. This supports the hypothesis that the physiological function of the enterohepatic circulation is not only to recycle bile acids but also to clear bile acids from the circulation to achieve homeostasis (Karpen and Dawson, *Hepatology* 2015, vol. 61, p. 24-27). Alternatively, the liver may synthesize increased levels of bile acids to compensate for the reduced enterohepatic recirculation in the homozygous carriers. As LBAT also transports unconjugated bile acids, the increase of the unconjugated bile acids in this study was not surprising (Liu et al., *Scientific Reports* 2017, 7: 9214, p. 1-7).

LBAT has been found to be downregulated in several forms of cholestatic liver injury and cholestasis, whereas ASBT has been found to be downregulated in a variety of gastrointestinal disorders such as Crohn's disease, primary bile acid malabsorption, inflammatory bowel disease, and ileal inflammation but upregulated in cholestasis. LBAT also functions as a cellular receptor for viral entry of the hepatitis B virus (HBV) and hepatitis D virus (HDV), which in turn is the major cause of liver disease and hepatocellular carcinoma.

ASBT inhibition has been investigated for decreasing plasma cholesterol levels and improving insulin resistance, as well as to relieving the hepatic bile acid burden in cholestatic liver disease. In addition, ASBT inhibition has been found to restore insulin levels and normoglycemia, thus establishing ASBT inhibition as a promising treatment for type 2 diabetes mellitus. ASBT inhibitors are also used for treatment of functional constipation.

As ASBT is predominantly expressed in the ileum (where it is often referred to as IBAT), ASBT inhibitors need not be systemically available. On the other hand, ASBT is also expressed in the proximal tubule cells of the kidneys. ASBT inhibitors that are systemically available may therefore also inhibit the reuptake of bile acids in the kidneys. It is believed that this would lead to increased levels of bile acids in urine, and to an increased removal of bile acids from the body via the urine. Systemically available ASBT inhibitors that exert their effect not only in the ileum but also in the kidneys are therefore expected to lead to a greater reduction of bile acid levels than non-systemically available ASBT inhibitors that only exert their effect in the ileum.

Compounds having a high ASBT inhibiting potency are particularly suitable for the treatment of liver diseases that cause cholestasis, such as progressive familial intrahepatic cholestasis (PFIC), Alagilles syndrome, biliary atresia and non-alcoholic steatohepatitis (NASH).

Biliary atresia is a rare pediatric liver disease that involves a partial or total blockage (or even absence) of large bile ducts. This blockage or absence causes cholestasis that leads to the accumulation of bile acids that damages the liver. In some embodiments, the accumulation of bile acids occurs in the extrahepatic biliary tree. In some embodiments, the accumulation of bile acids occurs in the intrahepatic biliary tree. The current standard of care is the Kasai procedure, which is a surgery that removes the blocked bile ducts and directly connects a portion of the small intestine to the liver. There are currently no approved drug therapies for this disorder.

Provided herein are methods for treating biliary atresia in a subject in need thereof, the methods comprising administration of a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the subject has undergone the Kasai procedure prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the subject is administered a compound of formula (I), or a pharmaceutically acceptable salt thereof, prior to undergoing the Kasai procedure. In some embodiments, the treatment of biliary atresia decreases the level of serum bile acids in the subject. In some embodiments, the level of serum bile acids is determined by, for example, an ELISA enzymatic assay or the assays for the measurement of total bile acids as described in Danese et al., PLoS One. 2017, vol. 12(6): e0179200, which is incorporated by reference herein in its entirety. In some embodiments, the level of serum bile acids can decrease by, for example, 10% to 40%, 20% to 50%, 30% to 60%, 40% to 70%, 50% to 80%, or by more than 90% of the level of serum bile acids prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the treatment of biliary atresia includes treatment of pruritus.

PFIC is a rare genetic disorder that is estimated to affect between one in every 50,000 to 100,000 children born worldwide and causes progressive, life-threatening liver disease.

One manifestation of PFIC is pruritus, which often results in a severely diminished quality of life. In some cases, PFIC leads to cirrhosis and liver failure. Current therapies include Partial External Biliary Diversion (PEBD) and liver transplantation, however, these options can carry substantial risk of post-surgical complications, as well as psychological and social issues.

Three alternative gene defects have been identified that correlate to three separate PFIC subtypes known as types 1, 2 and 3:

- 5 • PFIC, type 1, which is sometimes referred to as “Byler disease,” is caused by impaired bile secretion due to mutations in the ATP8B1 gene, which codes for a protein that helps to maintain an appropriate balance of fats known as phospholipids in cell membranes in the bile ducts. An imbalance in these phospholipids is associated with cholestasis and elevated bile acids in the liver. Subjects affected by PFIC, type 1 usually develop cholestasis in the first months of life and, in the absence of surgical treatment, progress to cirrhosis and end-stage liver disease before the end of the first decade of life.

- 10 • PFIC, type 2, which is sometimes referred to as “Byler syndrome,” is caused by impaired bile salt secretion due to mutations in the ABCB11 gene, which codes for a protein, known as the bile salt export pump, that moves bile acids out of the liver. Subjects with PFIC, type 2 often develop liver failure within the first few years of life and are at increased risk of developing a type of liver cancer known as hepatocellular carcinoma.

- 15 • PFIC, type 3, which typically presents in the first years of childhood with progressive cholestasis, is caused by mutations in the ABCB4 gene, which codes for a transporter that moves phospholipids across cell membranes.

20 In addition, TJP2 gene, NR1H4 gene or Myo5b gene mutations have been proposed to be causes of PFIC. In addition, some subjects with PFIC do not have a mutation in any of the ATP8B1, ABCB11, ABCB4, TJP2, NR1H4 or Myo5b genes. In these cases, the cause of the condition is unknown.

25 Exemplary mutations of the ATP8B1 gene or the resulting protein are listed in Tables 2 and 3, with numbering based on the human wild type ATP8B1 protein (e.g., SEQ ID NO: 1) or gene (e.g., SEQ ID NO: 2). Exemplary mutations of the ABCB11 gene or the resulting protein are listed in Tables 4 and 5, with numbering based on the human wild type ABCB11 protein (e.g., SEQ ID NO: 3) or gene (e.g., SEQ ID NO: 4).

30 As can be appreciated by those skilled in the art, an amino acid position in a reference protein sequence that corresponds to a specific amino acid position in SEQ ID NO: 1 or 3 can be determined by aligning the reference protein sequence with SEQ ID NO: 1 or 3 (e.g., using a software program, such as ClustalW2). Changes to these residues (referred to herein as “mutations”) may include single or multiple amino acid substitutions, insertions within or flanking the sequences, and deletions within or flanking the sequences. As can be appreciated by those skilled in the art, a nucleotide

position in a reference gene sequence that corresponds to a specific nucleotide position in SEQ ID NO: 2 or 4 can be determined by aligning the reference gene sequence with SEQ ID NO: 2 or 4 (e.g., using a software program, such as ClustalW2). Changes to these residues (referred to herein as “mutations”) may include single or multiple nucleotide substitutions, insertions within or flanking the sequences, and deletions within or flanking the sequences. See also Kooistra, et al., “KLIFS: A structural kinase-ligand interaction database,” *Nucleic Acids Res.* 2016, vol. 44, no. D1, pp. D365-D371, which is incorporated by reference in its entirety herein.

Canonical protein sequence of ATP8B1 (SEQ ID NO: 1) – Uniprot ID O43520

```

10  MSTERDSETT  FEEDSQPNDE  VVPYSDDETE  DELDDQGSVAV  EPEQNRVNRE  AEENREPFRK
    ECTWQVKAND  RKYHEQPHFM  NTKFLCIKES  KYANNAIKTY  KYNAFTFIPM  NLFEQFKRAA
    NLYFLALLIL  QAVPQISTLA  WYTTLVPLLV  VLGVTAIKDL  VDDVARHKMD  KEINNRTCEV
    IKDGRFKVAK  WKEIQVGDVI  RLKKNDFVPA  DILLLSSEP   NSLCYVETAE  LDGETNLKFK
    MSLEITDQYL  QREDTLATFD  GFIECEEPNN  RLDKFTGTLF  WRNTSFPLDA  DKILLRGCVI
15  RNTDFCHGLV  IFAGADTKIM  KNSGKTRFKR  TKIDYLMNYM  VYTI FVVLIL  LSAGLAIGHA
    YWEAQVGNSS  WYLYDGEDDT  PSYRGFLIFW  GYIIVLNTMV  PISLYVSVEV  IRLGQSHFIN
    WDLQMYAEEK  DTPAKARTTT  LNEQLGQIHY  IFSDKTGTLT  QNIMTFKKCC  INGQIYGDHR
    DASQHNHNKI  EQVDFSWNTY  ADGKLAFYDH  YLIEQIQSGK  EPEVRQFFFL  LAVCHTVMVD
    RTDGQLNYQA  ASPDEGALVN  AARNFGFAFL  ARTQNTITIS  ELGTERTYNV  LAILD FNSDR
20  KRMSIIVRTP  EGNIKLYCKG  ADTVIYERLH  RMNPTKQETQ  DALDI FANET  LRTLCLCYKE
    IEEKEFTewn  KKFMAASVAS  TNRDEALDKV  YEEIEKDLIL  LGATAIEDKL  QDGV PETISK
    LAKADIKIwV  LTGDKKETAE  NIGFACELLT  EDTTICYGED  INSLHARME  NQRNRGGVYA
    KFAPPVQESF  FPPGGNRALI  ITGSWLNEIL  LEKKTKRNKI  LKLKFP RTEE  ERRMRTQSKR
    RLEAKKEQRQ  KNFVDLACEC  SAVICCRVTP  KQKAMVVDLV  KRYKKAITLA  IGDGANDVNM
25  IKTAHIGVGI  SGQEGMQAVM  SSDYSFAQFR  YLQRLLLVHG  RWSYIRMCKF  LRYFFYKNFA
    FTLVHFWYSF  FNGYSAQTAY  EDWFITLYNV  LYTSLPVLLM  GLLDQDVSDK  LSLRFPGLYI
    VGQRDLLFNY  KRFFVSL LHG  VLTSMILFFI  PLGAYLQTVG  QDGEAPSDYQ  SFAVTIASAL
    VITVNFQIGL  DTSYWTFVNA  FSIFGSIALY  FGIMFDFHSA  GIHVLFPSAF  QFTGTASNAL
    RQPYIWL TII  LAVAVCLLPV  VAIRFLSMTI  WPSESDKIQK  HRKRLKAE EQ  WQRRQQVFRR
30  GVSTRRSAYA  FSHQRGYADL  ISSGRSIRKK  RSPLDAIVAD  GTAEYRRTGD  S
    
```

Canonical DNA Sequence for ATP8B1 (SEQ ID NO: 2)

```

    ATG AGT ACA GAA AGA GAC TCA GAA ACG ACA TTT GAC GAG GAT TCT CAG CCT
    AAT GAC GAA GTG GTT CCC TAC AGT GAT GAT GAA ACA GAA GAT GAA CTT GAT
35  GAC CAG GGG TCT GCT GTT GAA CCA GAA CAA AAC CGA GTC AAC AGG GAA GCA
    GAG GAG AAC CGG GAG CCA TTC AGA AAA GAA TGT ACA TGG CAA GTC AAA GCA
    AAC GAT CGC AAG TAC CAC GAA CAA CCT CAC TTT ATG AAC ACA AAA TTC TTG
    'G'I 'A'I' AAG GAG AG'I' AAA 'T'A'I' GCG AAT' AA'I' GCA A'I'I' AAA ACA 'T'AC AAG 'T'AC
    AAC GCA TTT ACC TTT ATA CCA ATG AAT CTG TTT GAG CAG TTT AAG AGA GCA
40  GCC AAT TTA TAT TTC CTG GCT CTT CTT ATC TTA CAG GCA GTT CCT CAA ATC
    TCT ACC CTG GCT TGG TAC ACC ACA CTA GTG CCC CTG CTT GTG GTG CTG GGC
    GTC ACT GCA ATC AAA GAC CTG GTG GAC GAT GTG GCT CGC CAT AAA ATG GAT
    AAG GAA ATC AAC AAT AGG ACG TGT GAA GTC ATT AAG GAT GGC AGG TTC AAA
    GTT GCT AAG TGG AAA GAA ATT CAA GTT GGA GAC GTC ATT CGT CTG AAA AAA
45  AAT GAT TTT GTT CCA GCT GAC ATT CTC CTG CTG TCT AGC TCT GAG CCT AAC
    AGC CTC TGC TAT GTG GAA ACA GCA GAA CTG GAT GGA GAA ACC AAT TTA AAA
    
```

TTT AAG ATG TCA CTT GAA ATC ACA GAC CAG TAC CTC CAA AGA GAA GAT ACA
 TTG GCT ACA TTT GAT GGT TTT ATT GAA TGT GAA GAA CCC AAT AAC AGA CTA
 GAT AAG TTT ACA GGA ACA CTA TTT TGG AGA AAC ACA AGT TTT CCT TTG GAT
 GCT GAT AAA ATT TTG TTA CGT GGC TGT GTA ATT AGG AAC ACC GAT TTC TGC
 5 CAC GGC TTA GTC ATT TTT GCA GGT GCT GAC ACT AAA ATA ATG AAG AAT AGT
 GGG AAA ACC AGA TTT AAA AGA ACT AAA ATT GAT TAC TTG ATG AAC TAC ATG
 GTT TAC ACG ATC TTT GTT GTT CTT ATT CTG CTT TCT GCT GGT CTT GCC ATC
 GGC CAT GCT TAT TGG GAA GCA CAG GTG GGC AAT TCC TCT TGG TAC CTC TAT
 10 GAT GGA GAA GAC GAT ACA CCC TCC TAC CGT GGA TTC CTC ATT TTC TGG GGC
 TAT ATC ATT GTT CTC AAC ACC ATG GTA CCC ATC TCT CTC TAT GTC AGC GTG
 GAA GTG ATT CGT CTT GGA CAG AGT CAC TTC ATC AAC TGG GAC CTG CAA ATG
 TAC TAT GCT GAG AAG GAC ACA CCC GCA AAA GCT AGA ACC ACC ACA CTC AAT
 GAA CAG CTC GGG CAG ATC CAT TAT ATC TTC TCT GAT AAG ACG GGG ACA CTC
 15 ACA CAA AAT ATC ATG ACC TTT AAA AAG TGC TGT ATC AAC GGG CAG ATA TAT
 GGG GAC CAT CGG GAT GCC TCT CAA CAC AAC CAC AAC AAA ATA GAG CAA GTT
 GAT TTT AGC TGG AAT ACA TAT GCT GAT GGG AAG CTT GCA TTT TAT GAC CAC
 TAT CTT ATT GAG CAA ATC CAG TCA GGG AAA GAG CCA GAA GTA CGA CAG TTC
 TTC TTC TTG CTC GCA GTT TGC CAC ACA GTC ATG GTG GAT AGG ACT GAT GGT
 20 CAG CTC AAC TAC CAG GCA GCC TCT CCC GAT GAA GGT GCC CTG GTA AAC GCT
 GCC AGG AAC TTT GGC TTT GCC TTC CTC GCC AGG ACC CAG AAC ACC ATC ACC
 ATC AGT GAA CTG GGC ACT GAA AGG ACT TAC AAT GTT CTT GCC ATT TTG GAC
 TTC AAC AGT GAC CGG AAG CGA ATG TCT ATC ATT GTA AGA ACC CCA GAA GGC
 AAT ATC AAG CTT TAC TGT AAA GGT GCT GAC ACT GTT ATT TAT GAA CGG TTA
 25 CAT CGA ATG AAT CCT ACT AAG CAA GAA ACA CAG GAT GCC CTG GAT ATC TTT
 GCA AAT GAA ACT CTT AGA ACC CTA TGC CTT TGC TAC AAG GAA ATT GAA GAA
 AAA GAA TTT ACA GAA TGG AAT AAA AAG TTT ATG GCT GCC AGT GTG GCC TCC
 ACC AAC CGG GAC GAA GCT CTG GAT AAA GTA TAT GAG GAG ATT GAA AAA GAC
 TTA ATT CTC CTG GGA GCT ACA GCT ATT GAA GAC AAG CTA CAG GAT GGA GTT
 30 CCA GAA ACC ATT TCA AAA CTT GCA AAA GCT GAC ATT AAG ATC TGG GTG CTT
 ACT GGA GAC AAA AAG GAA ACT GCT GAA AAT ATA GGA TTT GCT TGT GAA CTT
 CTG ACT GAA GAC ACC ACC ATC TGC TAT GGG GAG GAT ATT AAT TCT CTT CTT
 CAT GCA AGG ATG GAA AAC CAG AGG AAT AGA GGT GGC GTC TAC GCA AAG TTT
 GCA CCT CCT GTG CAG GAA TCT TTT TTT CCA CCC GGT GGA AAC CGT GCC TTA
 35 ATC ATC ACT GGT TCT TGG TTG AAT GAA ATT CTT CTC GAG AAA AAG ACC AAG
 AGA AAT AAG ATT CTG AAG CTG AAG TTC CCA AGA ACA GAA GAA GAA AGA CGG
 ATG CGG ACC CAA AGT AAA AGG AGG CTA GAA GCT AAG AAA GAG CAG CGG CAG
 AAA AAC TTT GTG GAC CTG GCC TGC GAG TGC AGC GCA GTC ATC TGC TGC CGC
 GTC ACC CCC AAG CAG AAG GCC ATG GTG GTG GAC CTG GTG AAG AGG TAC AAG
 40 AAA GCC ATC ACG CTG GCC ATC GGA GAT GGG GCC AAT GAC GTG AAC ATG ATC
 AAA ACT GCC CAC ATT GGC GTT GGA ATA AGT GGA CAA GAA GGA ATG CAA GCT
 GTC ATG TCG AGT GAC TAT TCC TTT GCT CAG TTC CGA TAT CTG CAG AGG CTA
 CTG CTG GTG CAT GGC CGA TGG TCT TAC ATA AGG ATG TGC AAG TTC CTA CGA
 TAC TTC TTT TAC AAA AAC TTT GCC TTT ACT TTG GTT CAT TTC TGG TAC TCC
 TTC TTC AAT GGC TAC TCT GCG CAG ACT GCA TAC GAG GAT TGG TTC ATC ACC
 45 CTC TAC AAC GTG CTG TAC ACC AGC CTG CCC GTG CTC CTC ATG GGG CTG CTC
 GAC CAG GAT GTG AGT GAC AAA CTG AGC CTC CGA TTC CCT GGG TTA TAC ATA
 GTG GGA CAA AGA GAC TTA CTA TTC AAC TAT AAG AGA TTC TTT GTA AGC TTG
 TTG CAT GGG GTC CTA ACA TCG ATG ATC CTC TTC TTC ATA CCT CTT GGA GCT
 TAT CTG CAA ACC GTA GGG CAG GAT GGA GAG GCA CCT TCC GAC TAC CAG TCT
 50 TTT GCC GTC ACC ATT GCC TCT GCT CTT GTA ATA ACA GTC AAT TTC CAG ATT
 GGC TTG GAT ACT TCT TAT TGG ACT TTT GTG AAT GCT TTT TCA ATT TTT GGA
 AGC ATT GCA CTT TAT TTT GGC ATC ATG TTT GAC TTT CAT AGT GCT GGA ATA

5 CAT GTT CTC TTT CCA TCT GCA TTT CAA TTT ACA GGC ACA GCT TCA AAC GCT
 CTG AGA CAG CCA TAC ATT TGG TTA ACT ATC ATC CTG GCT GTT GCT GTG TGC
 TTA CTA CCC GTC GTT GCC ATT CGA TTC CTG TCA ATG ACC ATC TGG CCA TCA
 GAA AGT GAT AAG ATC CAG AAG CAT CGC AAG CGG TTG AAG GCG GAG GAG CAG
 TGG CAG CGA CGG CAG CAG GTG TTC CGC CGG GGC GTG TCA ACG CGG CGC TCG
 GCC TAC GCC TTC TCG CAC CAG CGG GGC TAC GCG GAC CTC ATC TCC TCC GGG
 CGC AGC ATC CGC AAG AAG CGC TCG CCG CTT GAT GCC ATC GTG GCG GAT GGC
 ACC GCG GAG TAC AGG CGC ACC GGG GAC AGC TGA

10 **Table 2. Exemplary ATP8B1 Mutations**

Amino acid position 3 (e.g., T3K) ²⁷
Amino acid position 23 (e.g., P23L) ⁵
Amino acid position 45 (e.g., N45T) ^{5,8,9}
Amino acid position 46 (e.g., R46X) ^{A,25}
Amino acid position 62 (e.g., C62R) ²⁸
Amino acid position 63 (e.g., T63T) ⁴¹
Amino acid position 70 (e.g., D70N) ^{1,6}
Amino acid position 71 (e.g., R71H) ⁴³
Amino acid position 78 (e.g., H78Q) ¹⁹
Amino acid position 82 (e.g., T82T) ⁴¹
Amino acid position 92 (e.g., Y92Y) ⁴¹
Amino acid position 93 (e.g., A93A) ⁶
Amino acid position 96 (e.g., A96G) ²⁷
Amino acid position 114 (e.g., E114Q) ⁸
Amino acid position 127 (e.g., L127P ⁶ , L127V ³⁶)
Amino acid position 177 (e.g., T177T) ⁶
Amino acid position 179 (e.g., E179X) ²⁹
Δ Amino acid positions 185-282 ⁴⁴
Amino acid position 197 (e.g., G197Lfs*10) ²²
Amino acid position 201 (e.g., R201S ²⁷ , R201H ³⁵)
Amino acid position 203 (e.g., K203E ^{5,8} , K203R ⁹ , K203fs ²⁵)
Amino acid position 205 (e.g., N205fs ⁶ , N205Kfs*2 ³⁵)
Amino acid position 209 (e.g., P209T) ⁴
Amino acid position 217 (e.g., S217N) ⁴³
Amino acid position 232 (e.g., D232D) ³⁰
Amino acid position 233 (e.g., G233R) ³⁸

Amino acid position 243 (e.g., L243fs*28) ³³
Amino acid position 265 (e.g., C265R) ²⁵
Amino acid position 271 (e.g., R271X ¹³ , R271R ³⁰)
Amino acid position 288 (e.g., L288S) ⁶
Amino acid position 294 (e.g., L294S) ⁴³
Amino acid position 296 (e.g., R296C) ¹¹
Amino acid position 305 (e.g., F305I) ²⁸
Amino acid position 306 (e.g., C306R) ²³
Amino acid position 307 (e.g., H307L) ³⁵
Amino acid position 308 (e.g., G308V ¹ , G308D ⁶ , G308S ³⁵)
Amino acid position 314 (e.g., G314S) ¹³
Amino acid position 320 (e.g., M320Vfs*13) ¹¹
Amino acid position 337 (e.g., M337R) ¹⁸
Amino acid position 338 (e.g., N338K) ¹⁸
Amino acid position 340 (e.g., M340V) ¹⁸
Amino acid position 344 (e.g., I344F) ^{6,20}
Amino acid position 349 (e.g., I349T) ⁴¹
Amino acid position 358 (e.g., G358R) ²⁸
Amino acid position 367 (e.g., G367G) ⁴¹
Amino acid position 368 (e.g., N368D) ⁴¹
Amino acid position 393 (e.g., I393V) ²⁷
Amino acid position 403 (e.g., S403Y) ⁶
Amino acid position 407 (e.g., S407N) ⁴⁰
Amino acid position 412 (e.g., R412P) ⁶
Amino acid position 415 (e.g., Q415R) ²⁷
Amino acid position 422 (e.g., D422H) ³⁵
Amino acid position 429 (e.g., E429A) ⁶
Amino acid position 446 (e.g., G446R) ^{4,11}
Amino acid position 453 (e.g., S453Y) ⁶
Amino acid position 454 (e.g., D454G) ⁶
Amino acid position 455 (e.g., K455N) ⁴³
Amino acid position 456 (e.g., T456M ^{3,6} , T456K ³⁵)
Amino acid position 457 (e.g., G457G ⁶ , G457fs*6 ³³)

Amino acid position 469 (e.g., C469G) ⁴¹
Amino acid position 478 (e.g., H478H) ⁴¹
Amino acid position 500 (e.g., Y500H) ⁶
Amino acid position 525 (e.g., R525X) ⁴
Δ Amino acid position 529 ⁶
Amino acid position 535 (e.g., H535L ⁶ , H535N ⁴¹)
Amino acid position 553 (e.g., P553P) ⁴³
Amino acid position 554 (e.g., D554N ^{1,6} , D554A ³⁵)
Δ Amino acid positions 556-628 ⁴⁴
Δ Amino acid positions 559-563 ³⁵
Amino acid position 570 (e.g., L570L) ⁴¹
Amino acid position 577 (e.g., I577V) ¹⁹
Amino acid position 581 (e.g., E581K) ³⁵
Amino acid positions 554 and 581 (e.g., D554A+E581K) ³⁵
Amino acid position 585 (e.g., E585X) ²¹
Amino acid position 600 (e.g., R600W ^{2,4} , R600Q ⁶)
Amino acid position 602 (e.g., R602X) ^{3,6}
Amino acid position 628 (e.g., R628W) ⁵
Amino acid position 631 (e.g., R631Q) ²⁸
Δ Amino acid positions 645-699 ⁴
Amino acid position 661 (e.g., I661T) ^{1,4,6}
Amino acid position 665 (e.g., E665X) ^{4,6}
Amino acid position 672 (e.g., K672fs ⁶ , K672Vfs*1 ³⁵)
Amino acid position 674 (e.g., M674T) ¹⁹
Amino acid positions 78 and 674 (e.g., H78Q/M674T) ¹⁹
Amino acid position 684 (e.g., D684D) ⁴¹
Amino acid position 688 (e.g., D688G) ⁶
Amino acid position 694 (e.g., I694T ⁶ , I694N ¹⁷)
Amino acid position 695 (e.g., E695K) ²⁷
Amino acid position 709 (e.g., K709fs ⁶ , K709Qfs*41 ¹³)
Amino acid position 717 (e.g., T717N) ⁴
Amino acid position 733 (e.g., G733R) ⁶
Amino acid position 757 (e.g., Y757X) ⁴

Amino acid position 749 (e.g., L749P) ²¹
Amino acid position 792 (e.g., P792fs) ⁶
Δ Amino acid position 795-797 ⁶
Amino acid position 809 (e.g., I809L) ²⁷
Amino acid position 814 (e.g., K814N) ²⁸
Amino acid position 833 (e.g., R833Q ²⁷ , R833W ⁴¹)
Amino acid position 835 (e.g., K835Rfs*36) ³⁵
Amino acid position 845 (e.g., K845fs) ²⁵
Amino acid position 849 (e.g., R849Q) ²⁴
Amino acid position 853 (e.g., F853S, F853fs) ⁶
Amino acid position 867 (e.g., R867C ¹ , R867fs ⁶ , R867H ²³)
Amino acid position 885 (e.g., K885T) ⁴¹
Amino acid position 888 (e.g., T888T) ⁴¹
Amino acid position 892 (e.g., G892R) ⁶
Amino acid position 912 (e.g., G912R) ³⁵
Amino acid position 921 (e.g., S921S) ⁴¹
Amino acid position 924 (e.g., Y924C) ²⁸
Amino acid position 930 (e.g., R930X ⁶ , R930Q ²⁸)
Amino acid position 941 (e.g., R941X) ³⁵
Amino acid position 946 (e.g., R946T) ⁴¹
Amino acid position 952 (e.g., R952Q ^{5,9,15} , R952X ⁶)
Amino acid position 958 (e.g., N958fs) ⁶
Amino acid position 960 (e.g., A960A) ⁴¹
Δ Amino acid position 971 ⁴³
Amino acid position 976 (e.g., A976E ⁴¹ , A976A ⁴³)
Amino acid position 981 (e.g., E981K) ²⁰
Amino acid position 994 (e.g., S994R) ⁴
Amino acid position 1011 (e.g., L1011fs*18) ³³
Amino acid position 1012 (e.g., S1012I) ¹⁰
Amino acid position 1014 (e.g., R1014X) ^{6,11}
Amino acid position 1015 (e.g., F1015L) ²⁷
Amino acid position 1023 (e.g., Q1023fs) ⁶
Amino acid position 1040 (e.g., G1040R) ^{1,6}

Amino acid position 1044 (e.g., S0144L) ³⁴
Amino acid position 1047 (e.g., L1047fs) ⁶
Amino acid position 1050 (e.g., I1050K) ³¹
Amino acid position 1052 (e.g., L1052R) ²⁸
Amino acid position 1095 (e.g., W1095X) ¹¹
Amino acid position 1098 (e.g., V1098X) ³⁵
Amino acid position 1131 (e.g., Q1131X) ⁴⁴
Amino acid position 1142 (e.g., A1142Tfs*35) ⁴³
Amino acid position 1144 (e.g., Y1144Y) ⁴³
Amino acid position 1150 (e.g., I1150T) ⁴¹
Amino acid position 1152 (e.g., A1152T) ³⁰
Amino acid position 1159 (e.g., P1159P) ^{25,43}
Amino acid position 1164 (e.g., R1164X) ⁶
Amino acid position 1193 (e.g., R1193fs*39) ³³
Amino acid position 1197 (e.g., V1197L) ⁴¹
Amino acid position 1208 (e.g., A1208fs) ⁶
Amino acid position 1209 (e.g., Y1209Lfs*28) ⁴
Amino acid position 1211 (e.g., F1211L) ²⁷
Amino acid position 1219 (e.g., D1219H ⁵ , D1219G ²⁷)
Amino acid position 1223 (e.g., S1223S) ⁴¹
Amino acid position 1233 (e.g., P1233P) ⁴¹
Amino acid position 1241 (e.g., G1241fs) ⁶
Amino acid position 1248 (e.g., T1248T) ⁴³
Splice site mutation IVS3+1_+3delGTG ⁶
Splice site mutation IVS3-2A>G ⁶
IVS6+5T>G ^{17,25}
Splice site mutation IVS8+1G>T ⁶
IVS9-G>A ²⁶
IVS12+1G>A ²⁵
Splice site mutation IVS17-1G>A ⁶
Splice site mutation IVS18+2T>C ⁶
Splice site mutation IVS20-4CT>AA
Splice site mutation IVS21+5G>A ⁶

Splice site mutation IVS23-3C>A ⁶
Splice site mutation IVS26+2T>A ⁶
g.24774-42062del ⁴
c.-4C>G ⁴¹
c.145C>T ¹²
c.181-72G>A ⁹
c.182-5T>A ⁴¹
c.182-72G>A ⁴¹
c.246A>G ⁹
c.239G>A ³⁹
c.279+1_279+3delGTG ⁴⁶
c.280-2A>G ⁴⁶
c.625_62715delinsACAGTAAT ⁴⁶
c.554+122C>T ⁹
c.555-3T>C ²⁷
c.625+5 G>T ⁴
Amino acid position 209 (e.g., P209T) and c.625+5 G>T ⁴
c.628-30G>A ⁴¹
c.628-31C>T ⁴¹
c.698+1G>T ⁴⁶
c.698+20C>T ⁴¹
c.782-1G>A ⁴⁶
c.782-34G>A ⁴¹
Δ795-797 ¹⁴
c.782 -1G>A ⁴
c.852A>C ²⁷
c.941-1G>A ⁴⁶
c.1014C>T ⁹
c.1029+35G>A ⁹
c.1221-8C.G ⁴¹
1226delA ¹⁶
c.1429+1G>A ⁴⁶
c.1429+2T>G ¹³

c.1429+49G>A ⁴¹
c.1430-42A>G ⁴¹
c.1493T>C ¹²
c.1587_1589delCTT ⁴⁶
c.1630+2T>G ²⁷
c.1631-10T>A ⁴¹
c.1637-37T>C ⁴¹
1660 G>A ¹⁴
1798 C>T ¹⁴
1799 G>A ¹⁴
c.1819-39_41delAA ⁹
c.1819+1G>A ³¹
c.1820-27G>A ⁴¹
c.1918+8C>T ²⁷
c.1933-1G>AK46
c.2097+2T>C ³²
c.2097+60T>G ⁴¹
c.2097+89T>C ⁴¹
c.2097+97T>G ⁴¹
c.2210-114T>C ⁹
2210delA ¹⁶
c.2210-45_50dupATAAAA ⁹
c.2285+29C.T ⁴¹
c.2285+32A>G ⁴¹
c.2286-4_2286-3delinsAA ⁴⁶
c.2418+5G>A ⁴⁶
c.2707+3G>C ²⁷
c.2707+9T>G ⁴¹
c.2707+43A>G ⁴¹
c.2709-59T>C ⁴¹
c.2931+9A>G ⁴¹
c.2931+59T>A ⁴¹
c.2932-3C>A ⁴⁶

c.2932+59T>A ⁹
c.2937A>C ²⁷
c.3016-9C>A ³¹
c.3033-3034del ¹⁹
3122delTCCTA/ insACATCGATGTTGATGTTAGG ⁴⁵
3318 G>A ¹⁴
c.3400+2T>A ¹⁶
c.3401-175C>T ⁹
c.3401-167C>T ⁹
c.3401-108C>T ⁹
c.3531+8G>T ^{9,15}
c.3532-15C>T ⁹
Δ Phe ex 15 ⁴
Ex1_Ex13del ⁶
Ex2_Ex6del ³³
Ex12_Ex14del ²⁷
Skipped Exon 24 ⁴⁵
del5'UTR-ex18 ¹¹
c.*11C>T ⁴¹
c.*1101 + 366G > A ⁷
g.92918del565 ³¹
GC preceding exon 16 (e.g., resulting in a 4 bp deletion) ⁴²
Frameshift from the 5' end of exon 16 ⁴²
5' 1.4 kb deletion ⁴⁶

Table 3. Selected ATP8B1 Mutations Associated with PFIC-1

Amino acid position 23 (e.g., P23L) ⁵
Amino acid position 78 (e.g., H78Q) ¹⁹
Amino acid position 93 (e.g., A93A) ⁶
Amino acid position 96 (e.g., A96G) ²⁷
Amino acid position 127 (e.g., L127P) ⁶
Amino acid position 197 (e.g., G197Lfs*10) ²²

Amino acid position 205 (e.g., N205fs) ⁶
Amino acid position 209 (e.g., P209T) ⁴
Amino acid position 233 (e.g., G233R) ³⁸
Amino acid position 243 (e.g., L243fs*28) ³³
Amino acid position 288 (e.g., L288S) ⁶
Amino acid position 296 (e.g., R296C) ¹¹
Amino acid position 308 (e.g., G308V ^{1,6})
Amino acid position 320 (e.g., M320Vfs*13) ¹¹
Amino acid position 403 (e.g., S403Y) ⁶
Amino acid position 407 (e.g., S407N) ⁴⁰
Amino acid position 412 (e.g., R412P) ⁶
Amino acid position 415 (e.g., Q415R) ²⁷
Amino acid position 429 (e.g., E429A) ⁶
Amino acid position 446 (e.g., G446R) ⁴
Amino acid position 456 (e.g., T456M) ^{3,6}
Amino acid position 457 (e.g., G457G ⁶ , G457fs*6 ³³)
Amino acid position 500 (e.g., Y500H) ⁶
Amino acid position 525 (e.g., R525X) ⁴
Δ Amino acid position 529 ⁶
Amino acid position 535 (e.g., H535L) ⁶
Amino acid position 554 (e.g., D554N) ^{1,6}
Amino acid position 577 (e.g., I577V) ¹⁹
Amino acid position 585 (e.g., E585X) ²¹
Amino acid position 600 (e.g., R600W) ⁴
Amino acid position 602 (e.g., R602X) ^{3,6}
Amino acid position 661 (e.g., I661T) ^{4,6}
Amino acid position 665 (e.g., E665X) ^{4,6}
Δ Amino acid positions 645-699 ⁴
Amino acid position 672 (e.g., K672fs) ⁶
Amino acid position 674 (e.g., M674T) ¹⁹
Amino acid positions 78 and 674 (e.g., H78Q/M674T) ¹⁹
Amino acid position 688 (e.g., D688G) ⁶
Amino acid position 694 (e.g., I694N) ¹⁷

Amino acid position 695 (e.g., E695K) ²⁷
Amino acid position 709 (e.g., K709fs) ⁶
Amino acid position 717 (e.g., T717N) ⁴
Amino acid position 733 (e.g., G733R) ⁶
Amino acid position 749 (e.g., L749P) ²¹
Amino acid position 757 (e.g., Y757X) ⁴
Amino acid position 792 (e.g., P792fs) ⁶
Amino acid position 809 (e.g., I809L) ²⁷
Amino acid position 853 (e.g., F853S, F853fs) ⁶
Amino acid position 867 (e.g., R867fs) ⁶
Amino acid position 892 (e.g., G892R) ⁶
Amino acid position 930 (e.g., R930X ⁶ , R952Q ¹⁵)
Amino acid position 952 (e.g., R952X) ⁶
Amino acid position 958 (e.g., N958fs) ⁶
Amino acid position 981 (e.g., E981K) ²⁰
Amino acid position 994 (e.g., S994R) ⁴
Amino acid position 1014 (e.g., R1014X) ^{6,11}
Amino acid position 1015 (e.g., F1015L) ²⁷
Amino acid position 1023 (e.g., Q1023fs) ⁶
Amino acid position 1040 (e.g., G1040R) ^{1,6}
Amino acid position 1047 (e.g., L1047fs) ⁶
Amino acid position 1095 (e.g., W1095X) ¹¹
Amino acid position 1208 (e.g., A1208fs) ⁶
Amino acid position 1209 (e.g., Y1209Lfs*28) ⁴
Amino acid position 1211 (e.g., F1211L) ²⁷
Amino acid position 1219 (e.g., D1219H ⁵ , D1219G ²⁷)
Splice site mutation IVS3+1_+3delGTG ⁶
Splice site mutation IVS3-2A>G ⁶
IVS6+5T>G ¹⁷
Splice site mutation IVS8+1G>T ⁶
IVS9-G>A ²⁶
Splice site mutation IVS17-1G>A ⁶
Splice site mutation IVS18+2T>C ⁶

Splice site mutation IVS21+5G>A ⁶
g.24774-42062del ⁴
c.145C>T ¹²
c.239G>A ³⁹
c.625+5 G>T ⁴
Amino acid position 209 (e.g., P209T) and c.625+5 G>T ⁴
c.782 -1G>A ⁴
c.1493T>C ¹²
c.1630+2T>G ²⁷
1660 G>A ¹⁴
c.2707+3G>C ²⁷
c.2097+2T>C ³²
c.3033-3034del ¹⁹
3318 G>A ¹⁴
c.3158+8G>T ¹⁵
Δ Phe ex 15 ⁴
Ex1_Ex13del ⁶
Ex2_Ex6del ³³
Ex12_Ex14del ²⁷
del5'UTR-ex18 ¹¹
c.*1101 + 366G > A ⁷
GC preceding exon 16 (e.g., resulting in a 4 bp deletion) ⁴²
Frameshift from the 5' end of exon 16 ⁴²

^A A mutation to 'X' denotes an early stop codon

References for Tables 2 and 3

- ¹ Folmer et al., *Hepatology*. 2009, vol. 50(5), p. 1597-1605.
- 5 ² Hsu et al., *Hepatology Res.* 2009, vol. 39(6), p. 625-631.
- ³ Alvarez et al., *Hum Mol Genet.* 2004, vol. 13(20), p. 2451-2460.
- ⁴ Davit-Spraul et al., *Hepatology* 2010, vol. 51(5), p. 1645-1655.
- ⁵ Vitale et al., *J Gastroenterol.* 2018, vol. 53(8), p. 945-958.
- ⁶ Klomp et al., *Hepatology* 2004, vol. 40(1), p. 27-38.
- 10 ⁷ Zarenezhad et al., *Hepatitis Monthly*: 2017, vol. 17(2); e43500.
- ⁸ Dixon et al., *Scientific Reports* 2017, vol. 7, 11823.

- ⁹ Painter et al., *Eur J Hum Genet.* 2005, vol. 13(4), p. 435-439.
- ¹⁰ Deng et al., *World J Gastroenterol.* 2012, vol. 18(44), p. 6504-6509.
- ¹¹ Giovannoni et al., *PLoS One.* 2015, vol. 10(12): e0145021.
- ¹² Li et al., *Hepatology International* 2017, vol. 11, No. 1, Supp. Supplement 1, pp. S180. Abstract
5 Number: OP284.
- ¹³ Togawa et al., *Journal of Pediatric Gastroenterology and Nutrition* 2018, vol. 67, Supp. Supplement
1, pp. S363. Abstract Number: 615.
- ¹⁴ Miloh et al., *Gastroenterology* 2006, vol. 130, No. 4, Suppl. 2, pp. A759-A760. Meeting Info.:
Digestive Disease Week Meeting/107th Annual Meeting of the American-Gastroenterological-
10 Association. Los Angeles, CA, USA. May 19.
- ¹⁵ Dröge et al., *Zeitschrift fur Gastroenterologie* 2015, vol. 53, No. 12. Abstract Number: A3-27.
Meeting Info: 32. Jahrestagung der Deutschen Arbeitsgemeinschaft zum Studium der Leber.
Dusseldorf, Germany. 22 Jan 2016-23 Jan 2016
- ¹⁶ Mizuochi et al., *Clin Chim Acta.* 2012, vol. 413(15-16), p. 1301-1304.
- ¹⁷ Liu et al., *Hepatology International* 2009, vol. 3, No. 1, p. 184-185. Abstract Number: PE405.
15 Meeting Info: 19th Conference of the Asian Pacific Association for the Study of the Liver. Hong
Kong, China. 13 Feb 2009-16 Feb 2009
- ¹⁸ McKay et al., *Version 2. F1000Res.* 2013; 2: 32. DOI: 10.12688/f1000research.2-32.v2
- ¹⁹ Hasegawa et al., *Orphanet J Rare Dis.* 2014, vol. 9:89.
- ²⁰ ²⁰ Stone et al., *J Biol Chem.* 2012, vol. 287(49), p. 41139-51.
- ²¹ Kang et al., *J Pathol Transl Med.* 2019 May 16. doi: 10.4132/jptm.2019.05.03. [Epub ahead of print]
- ²² Sharma et al., *BMC Gastroenterol.* 2018, vol. 18(1), p. 107.
- ²³ Uegaki et al., *Intern Med.* 2008, vol. 47(7), p. 599-602.
- ²⁴ Goldschmidt et al., *Hepatol Res.* 2016, vol. 46(4), p. 306-311.
- ²⁵ ²⁵ Liu et al., *J Pediatr Gastroenterol Nutr.* 2010, vol. 50(2), p. 179-183.
- ²⁶ Jung et al., *J Pediatr Gastroenterol Nutr.* 2007, vol. 44(4), p. 453-458.
- ²⁷ Bounford. University of Birmingham. *Dissertation Abstracts International*, (2016) Vol. 75, No. 1C.
Order No.: AAI10588329. ProQuest Dissertations & Theses.
- ²⁸ Stolz et al., *Aliment Pharmacol Ther.* 2019, vol. 49(9), p. 1195-1204.
- ³⁰ ²⁹ Ivashkin et al., *Hepatology International* 2016, vol. 10, No. 1, Supp. SUPPL. 1, pp. S461. Abstract
Number: LBO-38. Meeting Info: 25th Annual Conference of the Asian Pacific Association for the
Study of the Liver, APASL 2016. Tokyo, Japan. 20 Feb 2016-24 Feb 2016
- ³⁰ Blackmore et al., *J Clin Exp Hepatol.* 2013, vol. 3(2), p. 159-161.
- ³¹ Matte et al., *J Pediatr Gastroenterol Nutr.* 2010, vol. 51(4), p. 488-493.
- ³⁵ ³² Squires et al., *J Pediatr Gastroenterol Nutr.* 2017, vol. 64(3), p. 425-430.

³³ Hayshi et al., EBioMedicine. 2018, vol. 27, p. 187-199.

³⁴ Nagasaka et al., J Pediatr Gastroenterol Nutr. 2007, vol. 45(1), p. 96-105.

³⁵ Wang et al., PLoS One. 2016; vol. 11(4): e0153114.

³⁶ Narchi et al., Saudi J Gastroenterol. 2017, vol. 23(5), p. 303–305.

5 ³⁷ Alashkar et al., Blood 2015, vol. 126, No. 23. Meeting Info.: 57th Annual Meeting of the American-Society-of-Hematology. Orlando, FL, USA. December 05 -08, 2015. Amer Soc Hematol.

³⁸ Ferreira et al., Pediatric Transplantation 2013, vol. 17, Suppl. SUPPL. 1, pp. 99. Abstract Number: 239. Meeting Info: IPTA 7th Congress on Pediatric Transplantation. Warsaw, Poland. 13 Jul 2013-16 Jul 2013.

10 ³⁹ Pauli-Magnus et al., J Hepatol. 2005, vol. 43(2), p. 342-357.

⁴⁰ Jericho et al., Journal of Pediatric Gastroenterology and Nutrition 2015, vol. 60(3), p. 368-374.

⁴¹ van der Woerd et al., PLoS One. 2013, vol. 8(11): e80553.

⁴² Copeland et al., J Gastroenterol Hepatol. 2013, vol. 28(3), p. 560-564.

⁴³ Dröge et al., J Hepatol. 2017, vol. 67(6), p. 1253-1264.

15 ⁴⁴ Chen et al., Journal of Pediatrics 2002, vol. 140(1), p. 119-124.

⁴⁵ Jirsa et al., Hepatol Res. 2004, vol. 30(1), p. 1-3.

⁴⁶ van der Woerd et al., Hepatology 2015, vol. 61(4), p. 1382-1391.

In some embodiments, the mutation in ATP8B1 is selected from L127P, G308V, T456M, D554N,
 20 F529del, I661T, E665X, R930X, R952X, R1014X, and G1040R.

Canonical Protein Sequence of ABCB11 (SEQ ID NO: 3) – Uniprot ID O95342

```

MSDSVILRSI KKFGEENDGF ESDKSYNNDK KSRLQDEKKG DGVRVGGFFQL FRFSSSTDIW
LMFVGSGLCAF LHGIAQPGVL LIFGTMDFV IDYDVELQEL QIPGKACVNN TIVWTNSSLN
25 QNMTNGTRCG LLNIESEMIK FASYAGIAV AVLITGYIQI CFWVIAAARQ IQKMRKFYFR
RIMRMEIGWF DCNSVGELNT RFSDDINKIN DAIADQMALF IQRMTSTICG FLLGGFFRGWK
LTLVIISVSP LIGIGAATIG LSVSKFTDYE LKAYAKAGVV ADEVISSMRT VAAFGEKRE
VERYEKNLVE AQRWGIRKGI VMGFFTGFVW CLIFLCYALA FWYGSTLVLD EGEYTPGTLV
QIFLSVIVGA LNLGNASPCL EAFATGRAAA TSIFETIDRK PIIDCMSEDG YKLDRIKGEI
30 EFHNVTFHYP SRPEVKILND LNMVIKPGEM TALVGPSGAG KSTALQLIQR FYDPCCEGMVT
VDGHDIRSLN IQWLRDQIGI VEQEPVLFST TIAENIRYGR EDATMEDIVQ AAKEANAYNF
IMDLPQQFDT LVGEGGGQMS GGQKQRVAIA RALIRNPKIL LLDMATSALD NESEAMVQEV
LSKIQHGHGTI ISVAHRLSTV RAADTIIGFE HGTAVERGTH EELLERKGVY FTLVTLQSQG
NQALNEEDIK DATEDDMLAR TFSRGSYQDS LRASIRQRSK SQLSYLVHEP PLAVVDHKST
35 YEEDRKDKDI PVQEEVEPAP VRRILKFSAP EWPYMLVGSV GAAVNGTVTP LYAFLFSQIL
GTFSSIPDKEE QRSQINGVCL LFMVAMGCVSL FTQFLQGYAF AKSGELLTKR LRKFGFRAML
GQDIAWFDDL RNSPGALTTR LATDASQVQG AAGSQIGMIV NSFTNVTVM IIAFSFSWKL
SLVILCFFPF LALSGATQTR MLTGFASRDK QALEMVGQIT NEALSNIRTV AGIGKERRFI
EALETELEKEP FKTAIQKANI YGFCFAFAQC IMFIANASAY RYGGYLISNE GLHFSYVFRV
40 ISAVVLSATA LGRAFSYTPS YAKAKISAAR FFQLLDRQPP ISVYNTAGEK WDNFQGGKIDF
VDCKFTYPSR PDSQVLNGLS VSISPGQTLA FVGSSGCGKS TSIQLLERFY DPDQGVKVID
    
```

GHDSKKNVNVQ FLRSNIGIVS QEPVLFACSI MDNIKYGDNT KEIPMERVIA AAKQAQLHDF
 VMSLPEKYET NVGSQGSQLS RGEKQRIAIA RAIVRDPKIL LLDEATSALD TESEKTVQVA
 LDKAREGRTC IVIAHRLSTI QNADIIVMA QGVVIEKGT H EELMAQKGAY YKLVTTGSP I S

5 Canonical DNA Sequence of ABCB11 (SEQ ID NO: 4)

ATG TCT GAC TCA GTA ATT CTT CGA AGT ATA AAG AAA TTT GGA GAG GAG AAT
 GAT GGT TTT GAG TCA GAT AAA TCA TAT AAT AAT GAT AAG AAA TCA AGG TTA
 CAA GAT GAG AAG AAA GGT GAT GGC GTT AGA GTT GGC TTC TTT CAA TTG TTT
 CGG TTT TCT TCA TCA ACT GAC ATT TGG CTG ATG TTT GTG GGA AGT TTG TGT
 10 GCA TTT CTC CAT GGA ATA GCC CAG CCA GGC GTG CTA CTC ATT TTT GGC ACA
 ATG ACA GAT GTT TTT ATT GAC TAC GAC GTT GAG TTA CAA GAA CTC CAG ATT
 CCA GGA AAA GCA TGT GTG AAT AAC ACC ATT GTA TGG ACT AAC AGT TCC CTC
 AAC CAG AAC ATG ACA AAT GGA ACA CGT TGT GGG TTG CTG AAC ATC GAG AGC
 GAA ATG ATC AAA TTT GCC AGT TAC TAT GCT GGA ATT GCT GTC GCA GTA CTT
 15 ATC ACA GGA TAT ATT CAA ATA TGC TTT TGG GTC ATT GCC GCA GCT CGT CAG
 ATA CAG AAA ATG AGA AAA TTT TAC TTT AGG AGA ATA ATG AGA ATG GAA ATA
 GGG TGG TTT GAC TGC AAT TCA GTG GGG GAG CTG AAT ACA AGA TTC TCT GAT
 GAT ATT AAT AAA ATC AAT GAT GCC ATA GCT GAC CAA ATG GCC CTT TTC ATT
 CAG CGC ATG ACC TCG ACC ATC TGT GGT TTC CTG TTG GGA TTT TTC AGG GGT
 20 TGG AAA CTG ACC TTG GTT ATT ATT TCT GTC AGC CCT CTC ATT GGG ATT GGA
 GCA GCC ACC ATT GGT CTG AGT GTG TCC AAG TTT ACG GAC TAT GAG CTG AAG
 GCC TAT GCC AAA GCA GGG GTG GTG GCT GAT GAA GTC ATT TCA TCA ATG AGA
 ACA GTG GCT GCT TTT GGT GGT GAG AAA AGA GAG GTT GAA AGG TAT GAG AAA
 AAT CTT GTG TTC GCC CAG CGT TGG GGA ATT AGA AAA GGA ATA GTG ATG GGA
 25 TTC TTT ACT GGA TTC GTG TGG TGT CTC ATC TTT TTG TGT TAT GCA CTG GCC
 TTC TGG TAC GGC TCC ACA CTT GTC CTG GAT GAA GGA GAA TAT ACA CCA GGA
 ACC CTT GTC CAG ATT TTC CTC AGT GTC ATA GTA GGA GCT TTA AAT CTT GGC
 AAT GCC TCT CCT TGT TTG GAA GCC TTT GCA ACT GGA CGT GCA GCA GCC ACC
 AGC ATT TTT GAG ACA ATA GAC AGG AAA CCC ATC ATT GAC TGC ATG TCA GAA
 30 GAT GGT TAC AAG TTG GAT CGA ATC AAG GGT GAA ATT GAA TTC CAT AAT GTG
 ACC TTC CAT TAT CCT TCC AGA CCA GAG GTG AAG ATT CTA AAT GAC CTC AAC
 ATG GTC ATT AAA CCA GGG GAA ATG ACA GCT CTG GTA GGA CCC AGT GGA GCT
 GGA AAA AGT ACA GCA CTG CAA CTC ATT CAG CGA TTC TAT GAC CCC TGT GAA
 GGA ATG GTG ACC GTG GAT GGC CAT GAC ATT CGC TCT CTT AAC ATT CAG TGG
 35 CTT AGA GAT CAG ATT GGG ATA GTG GAG CAA GAG CCA GTT CTG TTC TCT ACC
 ACC ATT GCA GAA AAT ATT CGC TAT GGC AGA GAA GAT GCA ACA ATG GAA GAC
 ATA GTC CAA GCT GCC AAG GAG GCC AAT GCC TAC AAC TTC ATC ATG GAC CTG
 CCA CAG CAA TTT GAC ACC CTT GTT GGA GAA GGA GGA GGC CAG ATG AGT GGT
 CGC CAC AAA CAA ACC GTA CCT ATC CCC AGA CCC CTC ATC CGA AAT CCC AAC
 40 ATT CTG CTT TTG GAC ATG GCC ACC TCA GCT CTG GAC AAT GAG AGT GAA GCC
 ATG GTC CAA GAA GTG CTG AGT AAG ATT CAG CAT GGG CAC ACA ATC ATT TCA
 GTT GCT CAT CGC TTG TCT ACG GTC AGA GCT GCA GAT ACC ATC ATT GGT TTT
 GAA CAT GGC ACT GCA GTG GAA AGA GGG ACC CAT GAA GAA TTA CTG GAA AGG
 AAA GGT GTT TAC TTC ACT CTA GTG ACT TTG CAA AGC CAG GGA AAT CAA GCT
 45 CTT AAT GAA GAG GAC ATA AAG GAT GCA ACT GAA GAT GAC ATG CTT GCG AGG
 ACC TTT AGC AGA GGG AGC TAC CAG GAT AGT TTA AGG GCT TCC ATC CGG CAA
 CGC TCC AAG TCT CAG CTT TCT TAC CTG GTG CAC GAA CCT CCA TTA GCT GTT
 GTA GAT CAT AAG TCT ACC TAT GAA GAA GAT AGA AAG GAC AAG GAC ATT CCT
 GTG CAG GAA GAA GTT GAA CCT GCC CCA GTT AGG AGG ATT CTG AAA TTC AGT
 50 GCT CCA GAA TGG CCC TAC ATG CTG GTA GGG TCT GTG GGT GCA GCT GTG AAC

5 GGG ACA GTC ACA CCC TTG TAT GCC TTT TTA TTC AGC CAG ATT CTT GGG ACT
 TTT TCA ATT CCT GAT AAA GAG GAA CAA AGG TCA CAG ATC AAT GGT GTG TGC
 CTA CTT TTT GTA GCA ATG GGC TGT GTA TCT CTT TTC ACC CAA TTT CTA CAG
 GGA TAT GCC TTT GCT AAA TCT GGG GAG CTC CTA ACA AAA AGG CTA CGT AAA
 10 TTT GGT TTC AGG GCA ATG CTG GGG CAA GAT ATT GCC TGG TTT GAT GAC CTC
 AGA AAT AGC CCT GGA GCA TTG ACA ACA AGA CTT GCT ACA GAT GCT TCC CAA
 GTT CAA GGG GCT GCC GGC TCT CAG ATC GGG ATG ATA GTC AAT TCC TTC ACT
 AAC GTC ACT GTG GCC ATG ATC ATT GCC TTC TCC TTT AGC TGG AAG CTG AGC
 CTG GTC ATC TTG TGC TTC TTC CCC TTC TTG GCT TTA TCA GGA GCC ACA CAG
 15 ACC AGG ATG TTG ACA GGA TTT GCC TCT CGA GAT AAG CAG GCC CTG GAG ATG
 GTG GGA CAG ATT ACA AAT GAA GCC CTC AGT AAC ATC CGC ACT GTT GCT GGA
 ATT GGA AAG GAG AGG CGG TTC ATT GAA GCA CTT GAG ACT GAG CTG GAG AAG
 CCC TTC AAG ACA GCC ATT CAG AAA GCC AAT ATT TAC GGA TTC TGC TTT GCC
 TTT GCC CAG TGC ATC ATG TTT ATT GCG AAT TCT GCT TCC TAC AGA TAT GGA
 20 GGT TAC TTA ATC TCC AAT GAG GGG CTC CAT TTC AGC TAT GTG TTC AGG GTG
 ATC TCT GCA GTT GTA CTG AGT GCA ACA GCT CTT GGA AGA GCC TTC TCT TAC
 ACC CCA AGT TAT GCA AAA GCT AAA ATA TCA GCT GCA CGC TTT TTT CAA CTG
 CTG GAC CGA CAA CCC CCA ATC AGT GTA TAC AAT ACT GCA GGT GAA AAA TGG
 GAC AAC TTC CAG GGG AAG ATT GAT TTT GTT GAT TGT AAA TTT ACA TAT CCT
 25 TCT CGA CCT GAC TCG CAA GTT CTG AAT GGT CTC TCA GTG TCG ATT AGT CCA
 GGG CAG ACA CTG GCG TTT GTT GGG AGC AGT GGA TGT GGC AAA AGC ACT AGC
 ATT CAG CTG TTG GAA CGT TTC TAT GAT CCT GAT CAA GGG AAG GTG ATG ATA
 GAT GGT CAT GAC AGC AAA AAA GTA AAT GTC CAG TTC CTC CGC TCA AAC ATT
 GGA ATT GTT TCC CAG GAA CCA GTG TTG TTT GCC TGT AGC ATA ATG GAC AAT
 30 ATC AAG TAT GGA GAC AAC ACC AAA GAA ATT CCC ATG GAA AGA GTC ATA GCA
 GCT GCA AAA CAG GCT CAG CTG CAT GAT TTT GTC ATG TCA CTC CCA GAG AAA
 TAT GAA ACT AAC GTT GGG TCC CAG GGG TCT CAA CTC TCT AGA GGG GAG AAA
 CAA CGC ATT GCT ATT GCT CGG GCC ATT GTA CGA GAT CCT AAA ATC TTG CTA
 CTA GAT GAA GCC ACT TCT GCC TTA GAC ACA GAA AGT GAA AAG ACG GTG CAG
 GTT GCT CTA GAC AAA GCC AGA GAG GGT CGG ACC TGC ATT GTC ATT GCC CAT
 CGC TTG TCC ACC ATC CAG AAC GCG GAT ATC ATT GCT GTC ATG GCA CAG GGG
 GTG GTG ATT GAA AAG GGG ACC CAT GAA GAA CTG ATG GCC CAA AAA GGA GCC
 TAC TAC AAA CTA GTC ACC ACT GGA TCC CCC ATC AGT TGA

35 **Table 4. Exemplary ABCB11 Mutations**

Amino acid position 1 (e.g., M1V) ⁹
Amino acid position 4 (e.g., S4X) ^{A,64}
Amino acid position 8 (e.g., R8X) ⁸⁸
Amino acid position 19 (e.g., G19R) ⁵⁶
Amino acid position 24 (e.g., K24X) ³⁵
Amino acid position 25 (e.g., S25X) ^{5,14}
Amino acid position 26 (e.g., Y26Ifs*7) ³⁸
Amino acid position 36 (e.g., D36D) ²⁷
Amino acid position 38 (e.g., K38Rfs*24) ⁷³
Amino acid position 43 (e.g., V43I) ⁵⁷

Amino acid position 49 (e.g., Q49X) ⁷³
Amino acid position 50 (e.g., L50S, L50W) ⁵⁷
Amino acid position 52 (e.g., R52W ²⁶ , R52R ²⁸)
Amino acid position 56 (e.g., S56L) ⁵⁸
Amino acid position 58 (e.g., D58N) ⁶²
Amino acid position 62 (e.g., M62K) ⁹
Amino acid position 66 (e.g., S66N) ¹⁷
Amino acid position 68 (e.g., C68Y) ⁴¹
Amino acid position 50 (e.g., L50S) ^{5,7}
Amino acid position 71 (e.g., L71H) ⁷³
Amino acid position 74 (e.g., I74R) ⁷¹
Amino acid position 77 (e.g., P77A) ⁷³
Amino acid position 87 (e.g., T87R) ⁶⁷
Amino acid position 90 (e.g., F90F) ^{7,27}
Amino acid position 93 (e.g., Y93S ¹³ , Y93X ⁸⁸)
Amino acid position 96 (e.g., E96X) ⁸⁸
Amino acid position 97 (e.g., L97X) ³⁹
Amino acid position 101 (e.g., Q101Dfs*8) ⁹
Amino acid position 107 (e.g., C107R) ³⁶
Amino acid position 112 (e.g., I112T) ⁹
Amino acid position 114 (e.g., W114R) ^{2,9}
Amino acid position 123 (e.g., M123T) ⁶⁷
Amino acid position 127 (e.g., T127Hfs*6) ⁵
Amino acid position 129 (e.g., C129Y) ²⁵
Amino acid position 130 (e.g., G130G) ⁷⁷
Amino acid position 134 (e.g., I134I) ²⁸
Amino acid position 135 (e.g., E135K ^{7,13} , E135L ¹⁷)
Amino acid position 137 (e.g., E137K) ⁷
Amino acid position 157 (e.g., Y157C) ⁵
Amino acid position 161 (e.g., C161X) ³⁹
Amino acid position 164 (e.g., V164Gfs*7 ³⁰ , V164I ⁸⁵)
Amino acid position 167 (e.g., A167S ⁴ , A167V ⁷ , A167T ^{9,17})
Amino acid position 181 (e.g., R181I) ³⁵

Amino acid position 182 (e.g., I182K) ⁹
Amino acid position 183 (e.g., M183V ⁸ , M183T ⁹)
Amino acid position 185 (e.g., M185I) ⁷³
Amino acid position 186 (e.g., E186G) ^{2,7,22}
Amino acid position 188 (e.g., G188W) ⁷³
Amino acid position 194 (e.g., S194P) ⁷
Amino acid position 198 (e.g., L198P) ⁷
Amino acid position 199 (e.g., N199Ifs*15X) ⁸⁸
Amino acid position 206 (e.g., I206V) ²⁸
Amino acid position 212 (e.g., A212T) ⁷³
Amino acid position 217 (e.g., M217R) ⁸⁸
Amino acid position 225 (e.g., T225P) ⁵⁷
Amino acid position 226 (e.g., S226L) ⁹
Amino acid position 232 (e.g., L232Cfs*9) ⁹
Amino acid position 233 (e.g., L233S) ⁸⁶
Amino acid position 238 (e.g., G238V) ^{2,7}
Amino acid position 242 (e.g., T242I) ^{5,7}
Amino acid position 245 (e.g., I245Tfs*26) ⁵⁷
Amino acid position 256 (e.g., A256G) ⁹
Amino acid position 260 (e.g., G260D) ⁷
Amino acid position 269 (e.g., Y269Y) ²⁷
Amino acid position 277 (e.g., A277E) ⁷⁷
Amino acid position 283 (e.g., E283D) ⁷³
Amino acid positions 212 and 283 (e.g., A212T+E283D) ⁷³
Amino acid position 284 (e.g., V284L ^{7,39} , V284A ⁷ , V284D ²³)
Amino acid position 297 (e.g., E297G ^{1,2,5,7} , E297K ⁷)
Amino acid position 299 (e.g., R299K) ²⁸
Amino acid position 303 (e.g., R303K ⁸ , R303M ⁶³ R303fsX321 ⁸³)
Amino acid position 304 (e.g., Y304X) ²⁶
Amino acid position 312 (e.g., Q312H) ⁷
Amino acid position 313 (e.g., R313S) ^{5,7}
Amino acid position 314 (e.g., W314X) ⁵⁷
Amino acid position 318 (e.g., K318Rfs*26) ²⁹

Amino acid position 319 (e.g., G319G) ⁷
Amino acid position 327 (e.g., G327E) ^{5,7}
Amino acid position 330 (e.g., W330X) ²⁴
Amino acid position 336 (e.g., C336S) ^{2,7}
Amino acid position 337 (e.g., Y337H) ^{21,27}
Amino acid position 342 (e.g., W342G) ⁵⁰
Amino acid position 354 (e.g., R354X) ⁹
Amino acid position 361 (e.g., Q361X ⁵⁷ , Q361R ⁷⁴)
Amino acid position 366 (e.g., V366V ²⁸ , V366D ⁵⁷)
Amino acid position 368 (e.g., V368Rfs*27) ⁵
Amino acid position 374 (e.g., G374S) ³
Amino acid position 380 (e.g., L380Wfs*18) ⁵
Amino acid position 382 (e.g., A382G) ⁸⁸
Δ Amino acid positions 382-388 ⁵
Δ Amino acid positions 383-389 ⁵⁷
Amino acid position 387 (e.g., R387H) ⁹
Amino acid position 390 (e.g., A390P) ^{5,7}
Amino acid position 395 (e.g., E395E) ²⁸
Amino acid position 404 (e.g., D404G) ⁹
Amino acid position 410 (e.g., G410D) ^{5,7}
Amino acid position 413 (e.g., L413W) ^{5,7}
Amino acid position 415 (e.g., R415X) ⁴²
Amino acid position 416 (e.g., I416I) ²⁷
Amino acid position 420 (e.g., I420T) ⁹
Amino acid position 423 (e.g., H423R) ¹³
Amino acid position 432 (e.g., R432T) ^{1,2,7}
Amino acid position 436 (e.g., K436N) ⁴⁰
Amino acid position 440 (e.g., D440E) ⁸⁸
Amino acid position 444 (e.g., V444A) ²
Amino acid position 454 (e.g., V454X) ⁴⁹
Amino acid position 455 (e.g., G455E) ⁹
Amino acid position 457 (e.g., S457Vfs*23) ⁸⁸
Amino acid position 461 (e.g., K461E) ^{2,7}

Amino acid position 462 (e.g., S462R) ⁸⁸
Amino acid position 463 (e.g., T463I) ^{5,7}
Amino acid position 466 (e.g., Q466K) ^{5,7}
Amino acid position 470 (e.g., R470Q ^{5,7} , R470X ⁹)
Amino acid position 471 (e.g., Y472X) ⁵
Amino acid position 472 (e.g., Y472C ^{5,27} , Y472X ¹⁴)
Amino acid position 473 (e.g., D473Q ³⁵ , D473V ⁸⁸)
Amino acid position 475 (e.g., C475X) ²⁹
Amino acid position 481 (e.g., V481E) ^{5,7}
Amino acid position 482 (e.g., D482G) ^{2,5,7}
Amino acid position 484 (e.g., H484Rfs*5) ⁹
Amino acid position 487 (e.g., R487H ² , R487P ⁵)
Amino acid position 490 (e.g., N490D) ^{5,7}
Amino acid position 493 (e.g., W493X) ⁸
Amino acid position 496 (e.g., D496V) ⁸⁸
Amino acid position 498 (e.g., I498T) ^{2,7}
Amino acid position 499 (e.g., G499E) ⁷³
Amino acid position 501 (e.g., V501G) ⁶⁸
Amino acid position 504 (e.g., E504K) ⁷⁹
Amino acid position 510 (e.g., T510T) ⁷
Amino acid position 512 (e.g., I512T) ^{5,7}
Amino acid position 515 (e.g., N515T ^{5,7} , N515D ⁶⁴)
Amino acid position 516 (e.g., I516M) ¹⁷
Amino acid position 517 (e.g., R517H) ^{5,7}
Amino acid position 520 (e.g., R520X) ⁵
Amino acid position 523 (e.g., A523G) ¹³
Amino acid position 528 (e.g., I528Sfs*21 ⁵ , I528X ⁹ , I528T ⁷³)
Amino acid position 535 (e.g., A535A ⁷ , A535X ⁸⁹)
Amino acid position 540 (e.g., F540L) ⁴⁶
Amino acid position 541 (e.g., I541L ^{5,7} , I541T ^{5,17})
Amino acid position 546 (e.g., Q546K ³⁹ , Q546H ⁷³)
Amino acid position 548 (e.g., F548Y) ^{5,7}
Amino acid position 549 (e.g., D549V) ⁹

Amino acid position 554 (e.g., E554K) ²¹
Amino acid position 556 (e.g., G556R) ⁶⁷
Amino acid position 558 (e.g., Q558H) ²³
Amino acid position 559 (e.g., M559T) ⁵⁷
Amino acid position 562 (e.g., G562D ^{5,7} , G562S ⁷³)
Amino acid position 570 (e.g., A570T ^{2,5,7} , A570V ²⁶)
Amino acid position 575 (e.g., R575X ^{2,5} , R575Q ²¹)
Amino acid position 580 (e.g., L580P) ⁵⁷
Amino acid position 586 (e.g., T586I) ⁷
Amino acid position 587 (e.g., S587X) ⁷³
Amino acid position 588 (e.g., A588V ^{5,7} , A588P ⁷³)
Amino acid position 591 (e.g., N591S) ^{2,7}
Amino acid position 593 (e.g., S593R) ^{2,7}
Amino acid position 597 (e.g., V597V ⁹ , V597L ¹³)
Amino acid position 603 (e.g., K603K) ⁵⁵
Amino acid position 609 (e.g., H609Hfs*46) ²⁶
Amino acid position 610 (e.g., I610Gfs*45 ⁹ , I610T ⁵⁷) ⁹
Amino acid position 615 (e.g., H615R) ²⁶
Amino acid position 616 (e.g., R616G ²⁸ , R616H ⁷³)
Amino acid position 619 (e.g., T619A) ²⁸
Amino acid position 623 (e.g., A623A) ²⁸
Amino acid position 625 (e.g., T625Nfs*5) ²⁶
Amino acid position 627 (e.g., I627T) ⁷
Amino acid position 628 (e.g., G628Wfs*3) ⁷⁰
Amino acid position 636 (e.g., E636G) ²
Amino acid position 648 (e.g., G648Vfs*6 ⁵ , G648V ⁵⁰)
Amino acid position 655 (e.g., T655I) ⁷
Amino acid position 669 (e.g., I669V) ²⁶
Amino acid position 676 (e.g., D676Y) ¹¹
Amino acid position 677 (e.g., M677V) ^{7,13}
Amino acid position 679 (e.g., A679V) ⁵⁸
Amino acid position 685 (e.g., G685W) ⁶⁰
Amino acid position 696 (e.g., R696W ²⁷ , R696Q ⁵⁸)

Amino acid position 698 (e.g., R698H ^{7,9} , R698K ⁶¹ , R698C ⁸⁸)
Amino acid position 699 (e.g., S699P) ⁹
Amino acid position 701 (e.g., S701P) ⁵⁸
Amino acid position 702 (e.g., Q702X) ⁸⁹
Amino acid position 709 (e.g., E709K) ⁷
Amino acid position 710 (e.g., P710P) ⁷
Amino acid position 712 (e.g., L712L) ²⁸
Amino acid position 721 (e.g., Y721C) ⁸⁸
Amino acid position 729 (e.g., D724N) ³⁹
Amino acid position 731 (e.g., P731S) ²³
Amino acid position 740 (e.g., P740Qfs*6) ⁷³
Amino acid position 758 (e.g., G758R) ⁵
Amino acid position 766 (e.g., G766R) ^{5,24}
Amino acid position 772 (e.g., Y772X) ⁵
Amino acid position 804 (e.g., A804A) ⁷
Amino acid position 806 (e.g., G806D ⁴⁴ , G806G ⁵⁵)
Amino acid position 809 (e.g., S809F) ⁸¹
Amino acid position 817 (e.g., G817G) ⁸⁸
Amino acid position 818 (e.g., Y818F) ⁷
Amino acid position 824 (e.g., G824E) ⁴²
Amino acid position 825 (e.g., G825G) ⁷³
Amino acid position 830 (e.g., R830Gfs*28) ⁷³
Amino acid position 832 (e.g., R832C ^{7,26} , R832H ⁴¹)
Amino acid position 842 (e.g., D842G) ²
Amino acid position 848 (e.g., D848N) ⁷³
Amino acid position 855 (e.g., G855R) ¹¹
Amino acid position 859 (e.g., T859R) ^{5,7}
Amino acid position 865 (e.g., A865V) ²⁷
Amino acid position 866 (e.g., S866A) ⁵⁷
Amino acid position 868 (e.g., V868D) ⁷³
Amino acid position 869 (e.g., Q869P) ⁷³
Amino acid position 875 (e.g., Q875X) ⁷³
Amino acid position 877 (e.g., G877R) ⁵⁶

Amino acid position 879 (e.g., I879R) ⁸⁸
Amino acid position 893 (e.g., A893V) ⁵⁷
Amino acid position 901 (e.g., S901R ¹⁷ , S901I ⁷³)
Amino acid position 903 (e.g., V903G) ⁵⁷
Δ Amino acid position 919 ¹²
Amino acid position 923 (e.g., T923P) ^{2,7}
Amino acid position 926 (e.g., A926P) ^{2,7}
Amino acid position 928 (e.g., R928X ¹⁵ , R928Q ⁴⁰)
Amino acid position 930 (e.g., K930X ⁵ , K930Efs*79 ^{5,10} , K930Efs*49 ²⁶)
Amino acid position 931 (e.g., Q931P) ²⁷
Amino acid position 945 (e.g., S945N) ⁵⁷
Amino acid position 948 (e.g., R948C) ^{5,7,26}
Amino acid position 958 (e.g., R958Q) ²⁸
Amino acid position 969 (e.g., K969K) ⁸⁸
Δ Amino acid positions 969-972 ⁵
Amino acid position 973 (e.g., T973I) ⁵⁷
Amino acid position 976 (e.g., Q976R ⁵⁸ , Q976X ⁸⁸)
Amino acid position 979 (e.g., N979D) ^{5,7}
Amino acid position 981 (e.g., Y981Y) ²⁸
Amino acid position 982 (e.g., G982R) ^{2,5,7}
Amino acid positions 444 and 982 (e.g., V444A+G982R) ³⁸
Amino acid position 995 (e.g., A995A) ²⁸
Amino acid position 1001 (e.g., R1001R) ⁹
Amino acid position 1003 (e.g., G1003R) ²⁴
Amino acid position 1004 (e.g., G1004D) ^{2,7}
Amino acid position 1027 (e.g., S1027R) ²⁶
Amino acid position 1028 (e.g., A1028A ^{7,10,88} , A1028E ⁸⁸)
Amino acid position 1029 (e.g., T1029K) ⁵
Amino acid position 1032 (e.g., G1032R) ¹²
Amino acid position 1041 (e.g., Y1041X) ⁹
Amino acid position 1044 (e.g., A1044P) ⁸⁸
Amino acid position 1050 (e.g., R1050C) ^{2,7,57}
Amino acid position 1053 (e.g., Q1053X) ⁵⁷

Amino acid position 1055 (e.g., L1055P) ³⁶
Amino acid position 1057 (e.g., R1057X ² , R1057Q ⁵⁸)
Amino acid position 1058 (e.g., Q1058Hfs*38 ⁹ , Q1058fs*38 ¹⁷ , Q1058X ⁷³)
Amino acid position 1061 (e.g., I1061Vfs*34) ⁹
Amino acid position 1083 (e.g., C1083Y) ⁴⁷
Amino acid position 1086 (e.g., T1086T) ²⁸
Amino acid position 1090 (e.g., R1090X) ^{2,5}
Amino acid position 1099 (e.g., L1099Lfs*38) ²⁶
Amino acid position 1100 (e.g., S1100Qfs*38) ¹³
Amino acid position 1110 (e.g., A1110E) ^{5,7}
Amino acid position 1112 (e.g., V1112F) ⁷⁰
Amino acid position 1116 (e.g., G1116R ⁷ , G1116F ^{9,17} , G1116E ³⁶)
Amino acid position 1120 (e.g., S1120N) ⁸⁸
Amino acid position 1128 (e.g., R1128H ^{2,7} , R1128C ^{5,7,13})
Amino acid position 1131 (e.g., D1131V) ²⁷
Amino acid position 1144 (e.g., S1144R) ⁷
Amino acid position 1147 (e.g., V1147X) ⁵
Amino acid position 1153 (e.g., R1153C ^{2,5,7} , R1153H ⁵)
Amino acid position 1154 (e.g., S1154P) ^{5,7}
Amino acid position 1162 (e.g., E1162X) ³⁹
Δ Amino acid position 1165 ⁸⁸
Amino acid position 1164 (e.g., V1164Gfs*7)
Amino acid position 1173 (e.g., N1173D) ⁵⁷
Amino acid position 1175 (e.g., K1175T) ⁵⁸
Amino acid position 1186 (e.g., E1186K) ⁷
Amino acid position 1192 (e.g., A1192Efs*50) ⁹
Amino acid position 1196 (e.g., Q1196X) ⁸⁸
Amino acid position 1197 (e.g., L1197G) ⁷
Amino acid position 1198 (e.g., H1198R) ²⁷
Amino acid position 1204 (e.g., L1204P) ⁸⁸
Amino acid position 1208 (e.g., Y1208C) ⁷³
Amino acid position 1210 (e.g., T1210P ^{5,7} , T1210F ⁵⁷)
Amino acid position 1211 (e.g., N1211D) ⁷

Amino acid position 1212 (e.g., V1212F) ³⁶
Amino acid position 1215 (e.g., Q1215X) ⁵
Amino acid position 1221 (e.g., R1221K) ⁵³
Amino acid position 1223 (e.g., E1223D) ⁷
Amino acid position 1226 (e.g., R1226P) ⁷³
Amino acid position 1228 (e.g., A1228V) ⁷
Amino acid position 1231 (e.g., R1231W ^{5,7} , R1231Q ^{5,7})
Amino acid position 1232 (e.g., A1232D) ¹⁷
Amino acid position 1235 (e.g., R1235X) ^{5,12}
Amino acid position 1242 (e.g., L1242I) ^{5,7}
Amino acid position 1243 (e.g., D1243G) ⁶⁷
Amino acid position 1249 (e.g., L1249X) ⁷³
Amino acid position 1256 (e.g., T1256fs*1296) ⁸³
Amino acid position 1268 (e.g., R1268Q) ^{2,7}
Amino acid position 1276 (e.g., R1276H) ³⁰
Amino acid position 1283 (e.g., A1283A ²⁸ , A1283V ⁸⁸)
Amino acid position 1292 (e.g., G1292V) ⁷³
Amino acid position 1298 (e.g., G1298R) ⁵
Amino acid position 1302 (e.g., E1302X) ⁵
Amino acid position 1311 (e.g., Y1311X) ⁵⁷
Amino acid position 1316 (e.g., T1316Lfs*64) ¹⁵
Amino acid position 1321 (e.g., S1321N) ⁵⁷
Intron 4 ((+3)A>C) ¹
IVS4-74A>T ⁸⁹
Splice site mutation 3' Intron 5 c.3901G>A ⁵
Splice site mutation 5; Intron 7 c.6111G>A ⁵
Splice site mutation IVS7+1G>A ¹⁴
IVS7+5G>A ⁴⁰
IVS8+1G>C ⁷⁶
Splice site mutation 5' Intron 9 c.9081delG ⁵
Splice site mutation 5' Intron 9 c.9081G>T ⁵
Splice site mutation 5' Intron 9 c.9081G>A ⁵
Splice site mutation IVS9+1G>T ¹⁴

Splice site mutation 3' Intron 13 c.143513_1435-8del ⁵
Splice site mutation IVS13del-13 ⁸
Splice site mutation 3' Intron 16 c.20128T>G ⁵
Splice site mutation IVS16-8T>G ¹⁴
Splice site mutation 5' Intron 18 c.21781G>T ⁵
Splice site mutation 5' Intron 18 c.21781G>A ⁵
Splice site mutation 5' Intron 18 c.21781G>C ⁵
Splice site mutation 3' Intron 18 c.21792A>G ⁵
Splice site mutation IVS18+1G>A ¹⁴
Splice site mutation 5' Intron 19 c.2343+1G>T ⁵
Splice site mutation 5' Intron 19 c.2343+2T>C ⁵
Splice site mutation IVS19+2T>C ¹⁴
Splice site mutation IVS19+1G>A ²²
Splice site mutation 3' Intron 21 c.26112A>T ⁵
IVS22+3A>G ⁸⁹
IVS 23-8 G-A ³⁶
IVS24+5G>A ⁵¹
Splice site mutation 5' Intron 24 c.32131delG ⁵
IVS35-6C>G ⁸⁹
Putative splice mutation 1198-1G>C ¹⁷
Putative splice mutation 1810-3C>G ¹⁷
Putative splice mutation 2178+1G>A ¹⁷
Putative splice mutation 2344-1G>T ¹⁷
Putative splice mutation c.2611-2A>T ³⁹
Putative splice mutation 3213+1_3213+2delinsA ¹⁷
c.-24C>A ^{44,78}
c.76 13 G>T ⁹
c.77-19T>A ⁵²
c.90_93delGAAA ¹⁸
c.124G>A ⁶⁹
c.150 +3 A>C ¹⁰
174C>T ⁵⁴
c.245T>C ⁸⁷

c.249_250insT ¹⁸
270T>C ⁵⁴
402C>T ⁵⁴
585G>C ⁵⁴
c.611+1G>A ⁷⁰
c.611+4A>G ³⁶
c.612-15_-6del10bp ⁵⁵
c.625A>C ³¹
c.627+5G>T ³¹
c.625A>C/ c.627+5G>T ³¹
696G>T ⁵⁴
c. 784+1G>C ⁴⁹
807T>C ⁵⁴
c.886C>T ³¹
c.890A>G ⁵⁹
c.908+1G>A ⁵⁷
c.908+5G>A ⁵⁵
c.908delG ⁵⁹
c.909-15A>G ⁶⁶
957A>G ⁵⁴
c.1084-2A>G ⁵⁷
1145 1bp deletion ⁹⁰
1281C>T ^{54,57}
c.1309-165C > T ¹⁹
c.1434 + 174G > A ¹⁹
c.1434 + 70C > T ¹⁹
c.1530C>A ⁵⁷
c.1587-1589delCTT ³¹
c.1621A>C ^{33,59}
c.1638+32T>C ⁶⁶
c.1638+80C>T ⁶⁶
1671C>T ⁵⁴
1791G>T ⁵⁴

1939delA ¹⁴
c.2075+3A>G ⁵³
c.2081T>A ³¹
c.2093G>A ⁶⁵
2098delA ¹⁶
c.2138-8T>G ⁶⁷
2142A>G ⁵⁴
c.2178+1G>T ^{36,39}
c.2179-17C>A ⁶⁶
c.2344-157T>G ⁶⁶
c.2344-17T>C ⁶⁶
c.2417G>A ⁷⁸
c.2541delG ⁸⁷
c.2620C>T ^{32,33}
c.2815-8A>G ⁵⁵
c.3003A>G ³⁷
c.3084A>G ^{48,54}
c.3213 +4 A>G ^{9,37}
c.3213 +5 G>A ⁹
c.3268C>T ⁷⁵
3285A>G ⁵⁴
c.3382C>T ⁷⁵
3435A>G ⁵⁴
c.3491delT ⁷²
c.3589C>T ⁵⁷
c.3765(+1 +5)del5 ⁴²
c.3766-34A>G ⁶⁶
c.3767-3768insC ⁶
c.3770delA ⁶⁷
c.3826C>T ⁷²
c.3846C>T ⁵⁷
c.3929delG ⁶⁷
c.*236A>G ⁶⁶

1145delC ⁸
Ex13_Ex17del ⁸²

Table 5. Selected ABCB11 Mutations Associated with PFIC-2

Amino acid position 1 (e.g., M1V) ⁹
Amino acid position 4 (e.g., S4X) ⁶⁴
Amino acid position 19 (e.g., G19R) ⁵⁶
Amino acid position 25 (e.g., S25X) ¹⁴
Amino acid position 26 (e.g., Y26Ifs*7) ³⁸
Amino acid position 50 (e.g., L50S) ^{7,57}
Amino acid position 52 (e.g., R52W) ²⁶
Amino acid position 58 (e.g., D58N) ⁶²
Amino acid position 62 (e.g., M62K) ⁹
Amino acid position 66 (e.g., S66N) ¹⁷
Amino acid position 68 (e.g., C68Y) ⁴¹
Amino acid position 93 (e.g., Y93S) ¹³
Amino acid position 101 (e.g., Q101Dfs*8) ⁹
Amino acid position 107 (e.g., C107R) ³⁶
Amino acid position 112 (e.g., I112T) ⁹
Amino acid position 114 (e.g., W114R) ^{2,9}
Amino acid position 129 (e.g., C129Y) ²⁵
Amino acid position 135 (e.g., E135K ¹³ , E135L ¹⁷)
Amino acid position 167 (e.g., A167V ⁷ , A167T ^{9,17})
Amino acid position 182 (e.g., I182K) ⁹
Amino acid position 183 (e.g., M183V ⁸ , M183T ⁹)
Amino acid position 225 (e.g., T225P) ⁵⁷
Amino acid position 226 (e.g., S226L) ⁹
Amino acid position 232 (e.g., L232Cfs*9) ⁹
Amino acid position 233 (e.g., L233S) ⁸⁶
Amino acid position 238 (e.g., G238V) ^{2,7}
Amino acid position 242 (e.g., T242I) ⁷
Amino acid position 245 (e.g., I245Tfs*26) ⁵⁷
Amino acid position 256 (e.g., A256G) ⁹

Amino acid position 260 (e.g., G260D) ⁵⁷
Amino acid position 284 (e.g., V284L) ⁷
Amino acid position 297 (e.g., E297G) ^{2,7}
Amino acid position 303 (e.g., R303K ⁸ , R303M ⁶³ , R303fsX321 ⁸³)
Amino acid position 304 (e.g., Y304X) ²⁶
Amino acid position 312 (e.g., Q312H) ⁷
Amino acid position 313 (e.g., R313S) ⁷
Amino acid position 314 (e.g., W314X) ⁵⁷
Amino acid position 318 (e.g., K318Rfs*26) ²⁹
Amino acid position 327 (e.g., G327E) ⁷
Amino acid position 330 (e.g., V330X) ²⁴
Amino acid position 336 (e.g., C336S) ^{2,7}
Amino acid position 337 (e.g., Y337H) ²¹
Amino acid position 342 (e.g., W342G) ⁵⁰
Amino acid position 354 (e.g., R354X) ⁹
Amino acid position 361 (e.g., Q361X) ⁵⁷
Amino acid position 366 (e.g., V366D) ⁵⁷
Amino acid position 386 (e.g., G386X) ³⁴
Δ Amino acid positions 383-389 ⁵⁷
Amino acid position 387 (e.g., R387H) ⁹
Amino acid position 390 (e.g., A390P) ⁷
Amino acid position 410 (e.g., G410D) ⁷
Amino acid position 413 (e.g., L413W) ⁷
Amino acid position 415 (e.g., R415X) ⁴²
Amino acid position 420 (e.g., I420T) ⁹
Amino acid position 454 (e.g., V454X) ⁴⁹
Amino acid position 455 (e.g., G455E) ⁹
Amino acid position 461 (e.g., K461E) ^{2,7}
Amino acid position 463 (e.g., T463I) ⁷
Amino acid position 466 (e.g., Q466K) ⁷
Amino acid position 470 (e.g., R470Q ⁷ , R470X ⁹)
Amino acid position 472 (e.g., Y472X ¹⁴ , Y472C ²⁷)
Amino acid position 475 (e.g., C475X) ²⁹

Amino acid position 481 (e.g., V481E) ⁷
Amino acid position 482 (e.g., D482G) ^{2,7}
Amino acid position 484 (e.g., H484Rfs*5) ⁹
Amino acid position 487 (e.g., R487H ² , R487P ⁸⁴)
Amino acid position 490 (e.g., N490D) ⁷
Amino acid position 493 (e.g., W493X) ⁸
Amino acid position 498 (e.g., I498T) ⁷
Amino acid position 501 (e.g., V501G) ⁶⁸
Amino acid position 512 (e.g., I512T) ⁷
Amino acid position 515 (e.g., N515T ⁷ , N515D ⁶⁴)
Amino acid position 516 (e.g., I516M) ¹⁷
Amino acid position 517 (e.g., R517H) ⁷
Amino acid position 520 (e.g., R520X) ⁵⁷
Amino acid position 523 (e.g., A523G) ¹³
Amino acid position 528 (e.g., I528X) ⁹
Amino acid position 540 (e.g., F540L) ⁴⁶
Amino acid position 541 (e.g., I541L ⁷ , I541T ¹⁷)
Amino acid position 548 (e.g., F548Y) ⁷
Amino acid position 549 (e.g., D549V) ⁹
Amino acid position 554 (e.g., E554K) ²¹
Amino acid position 559 (e.g., M559T) ⁵⁷
Amino acid position 562 (e.g., G562D) ⁷
Amino acid position 570 (e.g., A570T ⁷ , A570V ²⁶)
Amino acid position 575 (e.g., R575X ² , R575Q ²¹)
Amino acid position 588 (e.g., A588V) ⁷
Amino acid position 591 (e.g., N591S) ^{9,17}
Amino acid position 593 (e.g., S593R) ^{2,7}
Amino acid position 597 (e.g., V597V ⁹ , V597L ¹³)
Amino acid positions 591 and 597 (e.g., N591S+V597V) ⁹
Amino acid position 603 (e.g., K603K) ⁵⁵
Amino acid position 609 (e.g., H609Hfs*46) ²⁶
Amino acid position 610 (e.g., I610Gfs*45) ⁹
Amino acid position 615 (e.g., H615R) ²⁶

Amino acid position 625 (e.g., T625Nfs*5) ²⁶
Amino acid position 627 (e.g., I627T) ⁷
Amino acid position 636 (e.g., E636G) ²
Amino acid position 669 (e.g., I669V) ²⁶
Amino acid position 698 (e.g., R609H) ⁹
Amino acid positions 112 and 698 (e.g., I112T+R698H) ⁹
Amino acid position 699 (e.g., S699P) ⁹
Amino acid position 766 (e.g., G766R) ²⁴
Amino acid position 806 (e.g., G806G) ⁵⁵
Amino acid position 824 (e.g., G824E) ⁴²
Amino acid position 832 (e.g., R832C ^{7,26} , R832H ⁴¹)
Amino acid position 842 (e.g., D842G) ²
Amino acid position 859 (e.g., T859R) ⁷
Amino acid position 865 (e.g., A865V) ⁴⁵
Amino acid position 877 (e.g., G877R) ⁵⁶
Amino acid position 893 (e.g., A893V) ⁵⁷
Amino acid position 901 (e.g., S901R) ¹⁷
Amino acid position 903 (e.g., V903G) ⁵⁷
Δ Amino acid position 919 ¹²
Amino acid position 928 (e.g., R928X) ^{15,21}
Amino acid position 930 (e.g., K930Efs*79 ¹⁰ , K930Efs*49 ²⁶)
Amino acid position 948 (e.g., R948C) ^{7,26}
Amino acid position 979 (e.g., N979D) ⁷
Amino acid position 982 (e.g., G982R) ^{2,7}
Amino acid positions 444 and 982 (e.g., V444A+G982R) ³⁸
Amino acid position 1001 (e.g., R1001R) ⁹
Amino acid position 1003 (e.g., G1003R) ²⁴
Amino acid position 1004 (e.g., G1004D) ^{2,7}
Amino acid position 1027 (e.g., S1027R) ²⁶
Amino acid position 1028 (e.g., A1028A) ¹⁰
Amino acid position 1032 (e.g., G1032R) ¹²
Amino acid position 1041 (e.g., Y1041X) ⁹
Amino acid position 1050 (e.g., R1050C) ⁵⁷

Amino acid position 1053 (e.g., Q1053X) ⁵⁷
Amino acid position 1055 (e.g., L1055P) ³⁶
Amino acid position 1057 (e.g., R1057X) ²
Amino acid position 1058 (e.g., Q1058Hfs*38 ⁹ , Q1058fs*38 ¹⁷)
Amino acid position 1061 (e.g., I1061Vfs*34) ⁹
Amino acid position 1083 (e.g., C1083Y) ⁴⁷
Amino acid position 1090 (e.g., R1090X) ²
Amino acid position 1099 (e.g., L1099Lfs*38) ²⁶
Amino acid position 1100 (e.g., S1100Qfs*38) ¹³
Amino acid position 1110 (e.g., A1110E) ⁷
Amino acid position 1116 (e.g., G1116R ⁷ , G1116F ^{9,17} , G1116E ³⁶)
Amino acid position 1128 (e.g., R1128C) ^{7,13}
Amino acid position 1131 (e.g., D1131V) ²⁷
Amino acid position 1144 (e.g., S1144R) ⁷
Amino acid position 1153 (e.g., R1153C ^{2,7} , R1153H ^{7,26})
Amino acid position 1154 (e.g., S1154P) ⁷
Amino acid position 1173 (e.g., N1173D) ⁵⁷
Amino acid position 1192 (e.g., A1192Efs*50) ⁹
Amino acid position 1198 (e.g., H1198R) ²⁷
Amino acid position 1210 (e.g., T1210P ⁷ , T1210F ⁵⁷)
Amino acid position 1211 (e.g., N1211D) ⁷
Amino acid position 1212 (e.g., V1212F) ³⁶
Amino acid position 1231 (e.g., R1231W ⁷ , R1223Q ⁷)
Amino acid position 1232 (e.g., A1232D) ¹⁷
Amino acid position 1235 (e.g., R1235X) ¹²
Amino acid position 1242 (e.g., L1242I) ⁷
Amino acid position 1256 (e.g., T1256fs*1296) ⁸³
Amino acid position 1268 (e.g., R1268Q) ^{2,7}
Amino acid position 1302 (e.g., E1302X) ⁵⁷
Amino acid position 1311 (e.g., Y1311X) ⁵⁷
Amino acid position 1316 (e.g., T1316Lfs*64) ¹⁵
Intron 4 ((+3)A>C) ¹
Splice site mutation IVS7+1G>A ¹⁴

IVS8+1G>C ⁷⁶
Splice site mutation IVS9+1G>T ¹⁴
Splice site mutation IVS13del-13 ⁸ -8 ¹⁴
Splice site mutation IVS16-8T>G ¹⁴
Splice site mutation IVS18+1G>A ¹⁴
Splice site mutation IVS19+2T>C ¹⁴
IVS 23-8 G-A ³⁶
IVS24+5G>A ⁵¹
Putative splice mutation 1198-1G>C ¹⁷
Putative splice mutation 1810-3C>G ¹⁷
Putative splice mutation 2178+1G>A ¹⁷
Putative splice mutation 2344-1G>T ¹⁷
Putative splice mutation 3213+1_3213+2delinsA ¹⁷
c.-24C>A ⁷⁸
c.76 13 G>T ⁹
c.77-19T>A ⁵²
c.90_93delGAAA ¹⁸
c.124G>A ⁶⁹
c.150 +3 A>C ¹⁰
c.249_250insT ¹⁸
c.611+1G>A ⁸⁴
c.611+4A>G ³⁶
c.612-15_-6del10bp ⁵⁵
c.625A>C ³¹
c.627+5G>T ³¹
c.625A>C/ c.627+5G>T ³¹
c.886C>T ³¹
c.890A>G ⁵⁹
c.908+1G>A ⁵⁷
c.908+5G>A ⁵⁵
c.908delG ⁵⁹
1273 1bp deletion ⁹¹
c.1084-2A>G ⁵⁷

c.1445A>G ⁵⁹
c.1587-1589delCTT ³¹
c.1621A>C ⁵⁹
1939delA ¹⁴
c.2081T>A ³¹
2098delA ¹⁶
c.2343+1 G>T ⁸⁰
c.2178+1G>T ³⁶
c.2417G>A ⁷⁸
c.2620C>T ³²
c.2815-8A>G ⁵⁵
c.3003A>G ³⁷
c.3213 +4 A>G ^{9,37}
c.3213 +5 G>A ⁹
c.3268C>T ⁷⁵
c.3382C>T ⁷⁵
c.3765(+1 +5)del5 ⁴²
c.3767-3768insC ⁶
1145delC ⁸
Ex13_Ex17del ⁸²

^A A mutation to 'X' denotes an early stop codon

References for Tables 4 and 5

- ¹ Noe et al., J Hepatol. 2005, vol. 43(3), p. 536-543.
- 5 ² Lam et al., Am J Physiol Cell Physiol. 2007, vol. 293(5), p. C1709-16.
- ³ Stindt et al., Liver Int. 2013, vol. 33(10), p. 1527-1735.
- ⁴ Gao et al., Shandong Yiyao 2012, vol. 52(10), p. 14-16.
- ⁵ Strautnieks et al., Gastroenterology. 2008, vol. 134(4), p. 1203-1214.
- ⁶ Kagawa et al., Am J Physiol Gastrointest Liver Physiol. 2008, vol. 294(1), p. G58-67.
- 10 ⁷ Byrne et al., Hepatology. 2009, vol. 49(2), p. 553-567.
- ⁸ Chen et al., J Pediatr. 2008, vol. 153(6), p. 825-832.
- ⁹ Davit-Spraul et al., Hepatology 2010, vol. 51(5), p. 1645-1655.
- ¹⁰ Dröge et al., Sci Rep. 2016, vol. 6: 24827.
- ¹¹ Lang et al., Pharmacogenet Genomics. 2007, vol. 17(1), p. 47-60.

- ¹² Ellinger et al., *World J Gastroenterol.* 2017, vol. 23(29), p. :5295-5303.
- ¹³ Vitale et al., *J Gastroenterol.* 2018, vol. 53(8), p. 945-958.
- ¹⁴ Knisely et al., *Hepatology.* 2006, vol. 44(2), p. 478-86.
- ¹⁵ Ellis et al., *Hepatology.* 2018, vol. 67(4), p. 1531-1545.
- 5 ¹⁶ Lam et al., *J Hepatol.* 2006, vol. 44(1), p. 240-242.
- ¹⁷ Varma et al., *Hepatology* 2015, vol. 62(1), p. 198-206.
- ¹⁸ Treepongkaruna et al., *World J Gastroenterol.* 2009, vol. 15(34), p. 4339-4342.
- ¹⁹ Zarenezhad et al., *Hepatitis Monthly: 2017*, vol. 17(2); e43500.
- ²⁰ Hayashi et al., *Hepatol Res.* 2016, vol. 46(2), p. 192-200.
- 10 ²¹ Guorui et al., *Linchuang Erke Zazhi* 2013, vol. 31(10), 905-909.
- ²² van Mil et al., *Gastroenterology.* 2004, vol. 127(2), p. 379-384.
- ²³ Anzivino et al., *Dig Liver Dis.* 2013, vol. 45(3), p. 226-232.
- ²⁴ Park et al., *World J Gastroenterol.* 2016, vol. 22(20), p. 4901-4907.
- ²⁵ Imagawa et al., *J Hum Genet.* 2018, vol. 63(5), p. 569-577.
- 15 ²⁶ Giovannoni et al., *PLoS One.* 2015, vol. 10(12): e0145021.
- ²⁷ Hu et al., *Mol Med Rep.* 2014, vol. 10(3), p. 1264–1274.
- ²⁸ Lang et al., *Drug Metab Dispos.* 2006, vol. 34(9), p. 1582-1599.
- ²⁹ Masahata et al., *Transplant Proc.* 2016, vol. 48(9), p. 3156-3162.
- ³⁰ Holz et al., *Hepatol Commun.* 2018, vol. 2(2), p. 152-154.
- 20 ³¹ Li et al., *Hepatology International* 2017, vol. 11, No. 1, Supp. Supplement 1, pp. S180. Abstract Number: OP284.
- ³² Francalanci et al., *Laboratory Investigation* 2011, vol. 91, Supp. SUPPL. 1, pp. 360A. Abstract Number: 1526.
- ³³ Francalanci et al., *Digestive and Liver Disease* 2010, vol. 42, Supp. SUPPL. 1, pp. S16. Abstract
- 25 Number: T.N.5.
- ³⁴ Shah et al., *J Pediatr Genet.* 2017, vol. 6(2), p. 126-127.
- ³⁵ Gao et al., *Hepatitis Monthly* 2017, vol. 17(10), e55087/1-e55087/6.
- ³⁶ Evason et al., *Am J Surg Pathol.* 2011, vol. 35(5), p. 687-696.
- ³⁷ Davit-Spraul et al., *Mol Genet Metab.* 2014, vol. 113(3), p. 225-229.
- 30 ³⁸ Maggiore et al., *J Hepatol.* 2010, vol. 53(5), p. 981-6.
- ³⁹ McKay et al., *Version 2. F1000Res.* 2013; 2: 32. DOI: 10.12688/f1000research.2-32.v2
- ⁴⁰ Liu et al., *Pediatr Int.* 2013, vol. 55(2), p. 138-144.
- ⁴¹ Waisbourd-Zinman et al., *Ann Hepatol.* 2017, vol. 16(3), p. 465-468.

- ⁴² Griffin, et al., Canadian Journal of Gastroenterology and Hepatology 2016, vol. 2016. Abstract Number: A200. Meeting Info: 2016 Canadian Digestive Diseases Week, CDDW 2016. Montreal, QC, United States. 26 Feb 2016-29 Feb 2016
- ⁴³ Qiu et al., Hepatology 2017, vol. 65(5), p. 1655-1669.
- 5 ⁴⁴ Imagawa et al., Sci Rep. 2017, 7:41806.
- ⁴⁵ Kang et al., J Pathol Transl Med. 2019 May 16. doi: 10.4132/jptm.2019.05.03. [Epub ahead of print]
- ⁴⁶ Takahashi et al., Eur J Gastroenterol Hepatol. 2007, vol. 19(11), p. 942-6.
- ⁴⁷ Shimizu et al., Am J Transplant. 2011, vol. 11(2), p. 394-398.
- ⁴⁸ Krawczyk et al., Ann Hepatol. 2012, vol. 11(5), p. 710-744.
- 10 ⁴⁹ Sharma et al., BMC Gastroenterol. 2018, vol. 18(1), p. 107.
- ⁵⁰ Sattler et al., Journal of Hepatology 2017, vol. 66, No. 1, Suppl. S, pp. S177. Meeting Info.: International Liver Congress / 52nd Annual Meeting of the European-Association-for-the-Study-of-the-Liver. Amsterdam, NETHERLANDS. April 19 -23, 2017. European Assoc Study Liver.
- ⁵¹ Jung et al., J Pediatr Gastroenterol Nutr. 2007, vol. 44(4), p. 453-458.
- 15 ⁵² Sciveres. Digestive and Liver Disease 2010, vol. 42, Suppl. SUPPL. 5, pp. S329. Abstract Number: CO18. Meeting Info: 17th National Congress SIGENP. Pescara, Italy. 07 Oct 2010-09 Oct 2010
- ⁵³ Sohn et al., Pediatr Gastroenterol Hepatol Nutr. 2019, vol. 22(2), p. 201-206.
- ⁵⁴ Ho et al., Pharmacogenet Genomics. 2010, vol. 20(1), p. 45-57.
- ⁵⁵ Wang et al., Hepatol Res. 2018, vol. 48(7), p. 574-584.
- 20 ⁵⁶ Shaprio et al., J Hum Genet. 2010, vol. 55(5), p. 308-313.
- ⁵⁷ Bounford. University of Birmingham. Dissertation Abstracts International, (2016) Vol. 75, No. 1C. Order No.: AAI10588329. ProQuest Dissertations & Theses.
- ⁵⁸ Stolz et al., Aliment Pharmacol Ther. 2019, vol. 49(9), p. 1195-1204.
- ⁵⁹ Jankowska et al., J Pediatr Gastroenterol Nutr. 2014, vol. 58(1), p. 92-95.
- 25 ⁶⁰ Kim. Journal of Pediatric Gastroenterology and Nutrition 2016, vol. 62, Suppl. SUPPL. 1, pp. 620. Abstract Number: H-P-045. Meeting Info: 49th Annual Meeting of the European Society for Paediatric Gastroenterology, Hepatology and Nutrition, ESPGHAN 2016. Athens, Greece. 25 May 2016-28 May 2016.
- ⁶¹ Pauli-Magnus et al., Hepatology 2003, vol. 38, No. 4 Suppl. 1, pp. 518A. print. Meeting Info.: 54th
30 Annual Meeting of the American Association for the Study of Liver Diseases. Boston, MA, USA. October 24-28, 2003. American Association for the Study of Liver Diseases.
- ⁶² Li et al., Hepatology International 2017, vol. 11, No. 1, Suppl. Supplement 1, pp. S362. Abstract Number: PP0347. Meeting Info: 26th Annual Conference of the Asian Pacific Association for the Study of the Liver, APASL 2017. Shanghai, China. 15 Feb 2017-19 Feb 2017.

- ⁶³ Rumbo et al., *Transplantation* 2018, vol. 102, No. 7, Supp. Supplement 1, pp. S848. Abstract Number: P.752. Meeting Info: 27th International Congress of The Transplantation Society, TTS 2018. Madrid, Spain. 30 Jun 2018-05 Jul 2018.
- ⁶⁴ Lee et al., *Pediatr Gastroenterol Hepatol Nutr.* 2017, vol. 20(2), p. 114-123.
- 5 ⁶⁵ Sherrif et al., *Liver international: official journal of the International Association for the Study of the Liver* 2013, vol. 33, No. 8, pp. 1266-1270.
- ⁶⁶ Blackmore et al., *J Clin Exp Hepatol.* 2013, vol. 3(2), p. 159-161.
- ⁶⁷ Matte et al., *J Pediatr Gastroenterol Nutr.* 2010, vol. 51(4), p. 488-493.
- ⁶⁸ Lin et al., *Zhongguo Dang Dai Er Ke Za Zhi.* 2018, vol. 20(9), p. 758-764.
- 10 ⁶⁹ Harmanci et al., *Experimental and Clinical Transplantation* 2015, vol. 13, Supp. SUPPL. 2, pp. 76. Abstract Number: P62. Meeting Info: 1st Congress of the Turkic World Transplantation Society. Astana, Kazakhstan. 20 May 2015-22 May 2015.
- ⁷⁰ Herbst et al., *Mol Cell Probes.* 2015, vol. 29(5), p. 291-298.
- ⁷¹ Moghadamrad et al., *Hepatology.* 2013, vol. 57(6), p. 2539-2541.
- 15 ⁷² Holz et al., *Zeitschrift fur Gastroenterologie* 2016, vol. 54, No. 8. Abstract Number: KV275. Meeting Info: Viszeralmedizin 2016, 71. Jahrestagung der Deutschen Gesellschaft fur Gastroenterologie, Verdauungs- und Stoffwechselkrankheiten mit Sektion Endoskopie - 10. Herbsttagung der Deutschen Gesellschaft fur Allgemein- und Viszeralchirurgie. Hamburg, Germany. 21 Sep 2016-24 Sep 2016.
- 20 ⁷³ Wang et al., *PLoS One.* 2016; vol. 11(4): e0153114.
- ⁷⁴ Hao et al., *International Journal of Clinical and Experimental Pathology* 2017, vol. 10(3), p. 3480-3487.
- ⁷⁵ Arnell et al., *J Pediatr Gastroenterol Nutr.* 2010, vol. 51(4), p. 494-499.
- ⁷⁶ Sharma et al., *Indian Journal of Gastroenterology* 2017, vol. 36, No. 1, Supp. Supplement 1, pp. 25 A99. Abstract Number: M-20. Meeting Info: 58th Annual Conference of the Indian Society of Gastroenterology, ISGCON 2017. Bhubaneswar, India. 14 Dec 2017-17 Dec 2017.
- ⁷⁷ Beauséjour et al., *Can J Gastroenterol.* 2011, vol. 25(6), p. 311-314.
- ⁷⁸ Imagawa et al., *Journal of Pediatric Gastroenterology and Nutrition* 2016, vol. 63, Supp. Supplement 2, pp. S51. Abstract Number: 166. Meeting Info: World Congress of Pediatric Gastroenterology, Hepatology and Nutrition 2016. Montreal, QC, Canada. 05 Oct 2016-08 Oct 2016.
- 30 ⁷⁹ Peng et al., *Zhonghua er ke za zhi (Chinese journal of pediatrics)* 2018, vol. 56, No. 6, pp. 440-444.
- ⁸⁰ Tibesar et al., *Case Rep Pediatr.* 2014, vol. 2014: 185923.
- ⁸¹ Ng et al., *Journal of Pediatric Gastroenterology and Nutrition* 2018, vol. 66, Supp. Supplement 2, 35 pp. 860. Abstract Number: H-P-127. Meeting Info: 51st Annual Meeting European Society for

Paediatric Gastroenterology, Hepatology and Nutrition, ESPGHAN 2018. Geneva, Switzerland. 09
May 2018-12 May 2018.

⁸² Wong et al., Clin Chem. 2008, vol. 54(7), p. 1141-1148.

⁸³ Pauli-Magnus et al., J Hepatol. 2005, vol. 43(2), p. 342-357.

5 ⁸⁴ Jericho et al., Journal of Pediatric Gastroenterology and Nutrition. 60, vol. 3, p. 368-374.

⁸⁵ Scheimann et al., Gastroenterology 2007, vol. 132, No. 4, Suppl. 2, pp. A452. Meeting Info.:
Digestive Disease Week Meeting/108th Annual Meeting of the American-Gastroenterological-
Association. Washington, DC, USA. May 19 -24, 2007. Amer Gastroenterol Assoc; Amer Assoc
Study Liver Dis; Amer Soc Gastrointestinal Endoscopy; Soc Surg Alimentary Tract.

10 ⁸⁶ Jaquotot-Haerranz et al., Rev Esp Enferm Dig. 2013, vol. 105(1), p. 52-54.

⁸⁷ Khosla et al., American Journal of Gastroenterology 2015, vol. 110, No. Suppl. 1, pp. S397. Meeting
Info.: 80th Annual Scientific Meeting of the American-College-of-Gastroenterology. Honolulu, HI,
USA. October 16 -21, 2015.

⁸⁸ Dröge et al., J Hepatol. 2017, vol. 67(6), p. 1253-1264.

15 ⁸⁹ Liu et al., Liver International 2010, vol. 30(6), p. 809-815.

⁹⁰ Chen et al., Journal of Pediatrics 2002, vol. 140(1), p. 119-124.

⁹¹ U.S. Patent No. 9,295,677

In some embodiments, the mutation in ABCB11 is selected from A167T, G238V, V284L, E297G,
20 R470Q, R470X, D482G, R487H, A570T, N591S, A865V, G982R, R1153C, and R1268Q.

Provided are methods of treating PFIC (e.g., PFIC-1 and PFIC-2) in a subject that includes performing
an assay on a sample obtained from the subject to determine whether the subject has a mutation
associated with PFIC (e.g., a ATP8B1, ABCB11, ABCB4, TJP2, NR1H4 or Myo5b mutation), and
25 administering (e.g., specifically or selectively administering) a therapeutically effective amount of a
compound of formula (I), or a pharmaceutically acceptable salt thereof, to the subject determined to
have a mutation associated with PFIC. In some embodiments, the mutation is a ATP8B1 or ABCB11
mutation. For example, a mutation as provided in any one of Tables 1-4. In some embodiments, the
mutation in ATP8B1 is selected from L127P, G308V, T456M, D554N, F529del, I661T, E665X, R930X,
30 R952X, R1014X, and G1040R. In some embodiments, the mutation in ABCB11 is selected from A167T,
G238V, V284L, E297G, R470Q, R470X, D482G, R487H, A570T, N591S, A865V, G982R, R1153C, and
R1268Q.

Also provided are methods for treating PFIC (e.g., PFIC-1 and PFIC-2) in a subject in need thereof, the
35 method comprising: (a) detecting a mutation associated with PFIC (e.g., a ATP8B1, ABCB11, ABCB4,

TJP2, NR1H4 or Myo5b mutation) in the subject; and (b) administering to the subject a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, methods for treating PFIC can include administering a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof, to a
5 subject having a mutation associated with PFIC (e.g., a ATP8B1, ABCB11, ABCB4, TJP2, NR1H4 or Myo5b mutation). In some embodiments, the mutation is a ATP8B1 or ABCB11 mutation. For example, a mutation as provided in any one of Tables 1-4. In some embodiments, the mutation in ATP8B1 is selected from L127P, G308V, T456M, D554N, F529del, I661T, E665X, R930X, R952X, R1014X, and G1040R. In some embodiments, the mutation in ABCB11 is selected from A167T,
10 G238V, V284L, E297G, R470Q, R470X, D482G, R487H, A570T, N591S, A865V, G982R, R1153C, and R1268Q.

In some embodiments, the subject is determined to have a mutation associated with PFIC in a subject or a biopsy sample from the subject through the use of any art recognized tests, including
15 next generation sequencing (NGS). In some embodiments, the subject is determined to have a mutation associated with PFIC using a regulatory agency-approved, e.g., FDA-approved test or assay for identifying a mutation associated with PFIC in a subject or a biopsy sample from the subject or by performing any of the non-limiting examples of assays described herein. Additional methods of diagnosing PFIC are described in Gunaydin, M. et al., *Hepat Med.* 2018, vol. 10, p. 95-104,
20 incorporated by reference in its entirety herein.

In some embodiments, the treatment of PFIC (e.g., PFIC-1 or PFIC-2) decreases the level of serum bile acids in the subject. In some embodiments, the level of serum bile acids is determined by, for example, an ELISA enzymatic assay or the assays for the measurement of total bile acids as described
25 in Danese et al., *PLoS One.* 2017, vol. 12(6): e0179200, which is incorporated by reference herein in its entirety. In some embodiments, the level of serum bile acids can decrease by, for example, 10% to 40%, 20% to 50%, 30% to 60%, 40% to 70%, 50% to 80%, or by more than 90% of the level of serum bile acids prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the treatment of PFIC includes treatment of pruritus.

30

Since LBAT is expressed on hepatocytes, LBAT and dual ASBT/LBAT inhibitor substances need to have at least some bioavailability and free fraction in blood. Because LBAT inhibitor compounds only need to survive from the intestine to the liver, it is expected that a relatively low systemic exposure of such compounds will be sufficient, thereby minimizing the potential risk for any side effects in the rest of
35 the body. It is expected that inhibition of LBAT and ASBT will have at least additive effects in

decreasing the intrahepatic bile acid concentration. It is also expected that a dual ASBT/LBAT inhibitor may be able to reduce bile acid levels without inducing diarrhoea, as is sometimes observed with ASBT inhibitors.

- 5 Compounds having a high LBAT inhibiting potency and sufficient bioavailability are expected to be particularly suitable for the treatment of hepatitis. Compounds having a dual ASBT/LBAT inhibiting potency and sufficient bioavailability are expected to be particularly suitable for the treatment of non-alcoholic steatohepatitis (NASH).
- 10 NASH is a common and serious chronic liver disease that resembles alcoholic liver disease, but that occurs in people who drink little or no alcohol. In NASH patients, fat accumulation in the liver, known as nonalcoholic fatty liver disease (NAFLD) or steatosis, and other factors such as high LDL cholesterol and insulin resistance induce chronic inflammation in the liver and may lead to progressive scarring of tissue, known as fibrosis, and cirrhosis, followed eventually by liver failure and death. Patients
- 15 with NASH have been found to have significantly higher total serum bile acid concentrations than healthy subjects under fasting conditions (2.2- to 2.4-fold increase in NASH) and at all post-prandial time points (1.7- to 2.2-fold increase in NASH). These are driven by increased taurine- and glycine-conjugated primary and secondary bile acids. Patients with NASH exhibited greater variability in their fasting and post-prandial bile acid profile. These results indicate that patients with NASH have higher
- 20 fasting and post-prandial exposure to bile acids, including the more hydrophobic and cytotoxic secondary species. Increased bile acid exposure may be involved in liver injury and the pathogenesis of NAFLD and NASH (Ferslew et al., *Dig Dis Sci.* 2015, vol. 60, p. 3318–3328). It is therefore likely that ASBT and/or LBAT inhibition will be beneficial for the treatment of NASH.
- 25 NAFLD is characterized by hepatic steatosis with no secondary causes of hepatic steatosis including excessive alcohol consumption, other known liver diseases, or long-term use of a steatogenic medication (Chalasani et al., *Hepatology* 2018, vol. 67(1), p. 328-357). NAFLD can be categorized into non-alcoholic fatty liver (NAFL) and non-alcoholic steatohepatitis (NASH). According to Chalasani et al., NAFL is defined as the presence of $\geq 5\%$ hepatic steatosis without evidence of hepatocellular
- 30 injury in the form of hepatocyte ballooning. NASH is defined as the presence of $\geq 5\%$ hepatic steatosis and inflammation with hepatocyte injury (e.g., ballooning), with or without any liver fibrosis. NASH is also commonly associated with hepatic inflammation and liver fibrosis, which can progress to cirrhosis, end-stage liver disease, and hepatocellular carcinoma. While liver fibrosis is not always present in NASH, the severity of the fibrosis, when present, can be linked to long-term outcomes.

35

There are many approaches used to assess and evaluate whether a subject has NAFLD and if so, the severity of the disease, including differentiating whether the NAFLD is NAFL or NASH. In some embodiments, the severity of NAFLD can be assessed using the NAS. In some embodiments, treatment of NAFLD can be assessed using the NAS. In some embodiments, the NAS can be determined as described in Kleiner et al., *Hepatology*. 2005, 41(6):1313-1321, which is hereby incorporated by reference in its entirety. See, for example, Table 6 for a simplified NAS scheme adapted from Kleiner.

Table 6. Example of the NAFLD Activity Score (NAS) with Fibrosis Stage

Feature	Degree	Score
Steatosis	<5%	0
	5-33%	1
	>33-66%	2
	>66%	3
Lobular Inflammation	No foci	0
	<2 foci/200x	1
	2-4 foci/200x	2
	>4 foci/200x	3
Ballooning degeneration	None	0
	Few	1
	Many cells/Prominent ballooning	2
Fibrosis	None	0
	Perisinusoidal or periportal	1
	Perisinusoidal & portal/periportal	2
	Bridging fibrosis	3
	Cirrhosis	4

10

In some embodiments, the NAS is determined non-invasively, for example, as described in U.S. Application Publication No. 2018/0140219, which is incorporated by reference herein in its entirety. In some embodiments, the NAS is determined for a sample from the subject prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments,

the NAS is determined during the period of time or after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, a lower NAS score during the period of time or after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof indicates treatment of NAFLD (e.g., NASH). For example, a decrease in the NAS by 1, by 2, by 3, by 4, by 5, by 6, or by 7 indicates treatment of NAFLD (e.g., NASH). In some embodiments, the NAS following administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is 7 or less. In some embodiments, the NAS during the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is 5 or less, 4 or less, 3 or less, or 2 or less. In some embodiments, the NAS during the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is 7 or less. In some embodiments, the NAS during the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is 5 or less, 4 or less, 3 or less, or 2 or less. In some embodiments, the NAS after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is 7 or less. In some embodiments, the NAS after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is 5 or less, 4 or less, 3 or less, or 2 or less.

Additional approaches of assessing and evaluating NASH in a subject include determining one or more of hepatic steatosis (e.g., accumulation of fat in the liver); hepatic inflammation; biomarkers indicative of one or more of liver damage, hepatic inflammation, liver fibrosis, and/or liver cirrhosis (e.g., serum markers and panels). Further examples of physiological indicators of NASH can include liver morphology, liver stiffness, and the size or weight of the subject's liver.

In some embodiments, NASH in the subject is evidenced by an accumulation of hepatic fat and detection of a biomarker indicative of liver damage. For example, elevated serum ferritin and low titers of serum autoantibodies can be common features of NASH.

In some embodiments, methods to assess NASH include magnetic resonance imaging, either by spectroscopy or by proton density fat fraction (MRI-PDFF) to quantify steatosis, transient elastography (FIBROSCAN®), hepatic venous pressure gradient (HPVG), hepatic stiffness measurement with MRE for diagnosing significant liver fibrosis and/or cirrhosis, and assessing histological features of liver biopsy. In some embodiments, magnetic resonance imaging is used to detect one or more of steatohepatitis (NASH-MRI), liver fibrosis (Fibro-MRI), and steatosis. See, for

example, U.S. Application Publication Nos. 2016/146715 and 2005/0215882, each of which are incorporated herein by reference in their entireties.

5 In some embodiments, treatment of NASH can include a decrease of one or more symptoms associated with NASH; reduction in the amount of hepatic steatosis; a decrease in the NAS; a decrease in hepatic inflammation; a decrease in the level of biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis; and a reduction in fibrosis and/or cirrhosis, a lack of further progression of fibrosis and/or cirrhosis, or a slowing of the progression of fibrosis and/or cirrhosis in the subject following administration of one or more doses of a compound
10 of formula (I), or a pharmaceutically acceptable salt thereof.

In some embodiments, treatment of NASH comprises a decrease of one or more symptoms associated with NASH in the subject. Exemplary symptoms can include one or more of an enlarged liver, fatigue, pain in the upper right abdomen, abdominal swelling, enlarged blood vessels just
15 beneath the skin's surface, enlarged breasts in men, enlarged spleen, red palms, jaundice, and pruritus. In some embodiments, the subject is asymptomatic. In some embodiments, the total body weight of the subject does not increase. In some embodiments, the total body weight of the subject decreases. In some embodiments, the body mass index (BMI) of the subject does not increase. In some embodiments, the body mass index (BMI) of the subject decreases. In some embodiments, the
20 waist and hip (WTH) ratio of the subject does not increase. In some embodiments, the waist and hip (WTH) ratio of the subject decreases.

In some embodiments, treatment of NASH can be assessed by measuring hepatic steatosis. In some embodiments, treatment of NASH comprises a reduction in hepatic steatosis following administration
25 of a compound of formula (I), or a pharmaceutically acceptable salt thereof, as described herein. In some embodiments, hepatic steatosis is determined by one or more methods selected from the group consisting of ultrasonography, computed tomography (CT), magnetic resonance imaging, magnetic resonance spectroscopy (MRS), magnetic resonance elastography (MRE), transient elastography (TE) (e.g., FIBROSCAN®), measurement of liver size or weight, or by liver biopsy (see,
30 e.g., Di Lascio et al., *Ultrasound Med Biol.* 2018, vol. 44(8), p. 1585-1596; Lv et al., *J Clin Transl Hepatol.* 2018, vol. 6(2), p. 217-221; Reeder et al., *J Magn Reson Imaging.* 2011, vol. 34(4), spcone; and de Lédinghen V, et al., *J Gastroenterol Hepatol.* 2016, vol. 31(4), p. 848-855, each of which are incorporated herein by reference in their entireties). A subject diagnosed with NASH can have greater than about 5% hepatic steatosis, for example, greater than about 5% to about 25%, about 25% to
35 about 45%, about 45% to about 65%, or greater than about 65% hepatic steatosis. In some

embodiments, a subject with greater than about 5% to about 33% hepatic steatosis has stage 1 hepatic steatosis, a subject with about 33% to about 66% hepatic steatosis has stage 2 hepatic steatosis, and a subject with greater than about 66% hepatic steatosis has stage 3 hepatic steatosis.

5 In some embodiments, the amount of hepatic steatosis is determined prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the amount of hepatic steatosis is determined during the period of time or after the period of time of administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof. In some
10 embodiments, a reduction in the amount of hepatic steatosis during the period of time or after the period of time of administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof, indicates treatment of NASH. For example, a reduction in the amount of hepatic steatosis by about 1% to about 50%, about 25% to about 75%, or about 50% to about 100% indicates treatment of NASH. In some embodiments, a reduction in the amount of
15 hepatic steatosis by about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95% indicates treatment of NASH.

In some embodiments, the presence of hepatic inflammation is determined by one or more methods
20 selected from the group consisting of biomarkers indicative of hepatic inflammation and a liver biopsy sample(s) from the subject. In some embodiments, the severity of hepatic inflammation is determined from a liver biopsy sample(s) from the subject. For example, hepatic inflammation in a liver biopsy sample can be assessed as described in Kleiner et al., *Hepatology* 2005, vol. 41(6), p. 1313-1321 and Brunt et al., *Am J Gastroenterol* 1999, vol. 94, p. 2467-2474, each of which are hereby
25 incorporated by reference in their entireties. In some embodiments, the severity of hepatic inflammation is determined prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the severity of hepatic inflammation is determined during the period of time or after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, a decrease in the
30 severity of hepatic inflammation during the period of time or after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof, indicates treatment of NASH. For example, a decrease in the severity of hepatic inflammation by about 1% to about 50%, about 25% to about 75%, or about 50% to about 100% indicates
35 treatment of NASH. In some embodiments, a decrease in the severity of hepatic inflammation by

about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, or about 95% indicates treatment of NASH.

- 5 In some embodiments, treatment of NASH comprises treatment of fibrosis and/or cirrhosis, e.g., a decrease in the severity of fibrosis, a lack of further progression of fibrosis and/or cirrhosis, or a slowing of the progression of fibrosis and/or cirrhosis. In some embodiments, the presence of fibrosis and/or cirrhosis is determined by one or more methods selected from the group consisting of
- 10 transient elastography (e.g., FIBROSCAN®), non-invasive markers of hepatic fibrosis, and histological features of a liver biopsy. In some embodiments, the severity (e.g., stage) of fibrosis is determined by one or more methods selected from the group consisting of transient elastography (e.g., FIBROSCAN®), a fibrosis-scoring system, biomarkers of hepatic fibrosis (e.g., non-invasive
- 15 biomarkers), and hepatic venous pressure gradient (HVPG). Non-limiting examples of fibrosis scoring systems include the NAFLD fibrosis scoring system (see, e.g., Angulo et al., *Hepatology* 2007, vol. 45(4), p. 846-54), the fibrosis scoring system in Brunt et al., *Am. J. Gastroenterol.* 1999, vol. 94, p. 2467-2474, the fibrosis scoring system in Kleiner et al., *Hepatology* 2005, vol. 41(6), p. 1313-1321, and the ISHAK fibrosis scoring system (see Ishak et al., *J. Hepatol.* 1995, vol. 22, p. 696-699), the contents of each of which are incorporated by reference herein in their entireties.
- 20 In some embodiments, the severity of fibrosis is determined prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the severity of fibrosis is determined during the period of time or after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, a decrease in the severity of fibrosis during the period of time or after the period of time of
- 25 administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof, indicates treatment of NASH. In some embodiments, a decrease in the severity of fibrosis, a lack of further progression of fibrosis and/or cirrhosis, or a slowing of the progression of fibrosis and/or cirrhosis indicates treatment of NASH. In some embodiments, the severity of fibrosis is
- 30 determined using a scoring system such as any of the fibrosis scoring systems described herein, for example, the score can indicate the stage of fibrosis, e.g., stage 0 (no fibrosis), stage 1, stage 2, stage 3, and stage 4 (cirrhosis) (see, e.g., Kleiner et al). In some embodiments, a decrease in the stage of the fibrosis is a decrease in the severity of the fibrosis. For example, a decrease by 1, 2, 3, or 4 stages is a decrease in the severity of the fibrosis. In some embodiments, a decrease in the stage, e.g., from
- 35 stage 4 to stage 3, from stage 4 to stage 2, from stage 4 to stage 1, from stage 4 to stage 0, from stage

3 to stage 2, from stage 3 to stage 1, from stage 3 to stage 0, from stage 2 to stage 1, from stage 2 to stage 0, or from stage 1 to stage 0 indicates treatment of NASH. In some embodiments, the stage of fibrosis decreases from stage 4 to stage 3, from stage 4 to stage 2, from stage 4 to stage 1, from stage 4 to stage 0, from stage 3 to stage 2, from stage 3 to stage 1, from stage 3 to stage 0, from stage 2 to stage 1, from stage 2 to stage 0, or from stage 1 to stage 0 following administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the stage of fibrosis decreases from stage 4 to stage 3, from stage 4 to stage 2, from stage 4 to stage 1, from stage 4 to stage 0, from stage 3 to stage 2, from stage 3 to stage 1, from stage 3 to stage 0, from stage 2 to stage 1, from stage 2 to stage 0, or from stage 1 to stage 0 during the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the stage of fibrosis decreases from stage 4 to stage 3, from stage 4 to stage 2, from stage 4 to stage 1, from stage 4 to stage 0, from stage 3 to stage 2, from stage 3 to stage 1, from stage 3 to stage 0, from stage 2 to stage 1, from stage 2 to stage 0, or from stage 1 to stage 0 after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof.

In some embodiments, the presence of NASH is determined by one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis or scoring systems thereof. In some embodiments, the severity of NASH is determined by one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis or scoring systems thereof. The level of the biomarker can be determined by, for example, measuring, quantifying, and monitoring the expression level of the gene or mRNA encoding the biomarker and/or the peptide or protein of the biomarker. Non-limiting examples of biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis and/or scoring systems thereof include the aspartate aminotransferase (AST) to platelet ratio index (APRI); the aspartate aminotransferase (AST) and alanine aminotransferase (ALT) ratio (AAR); the FIB-4 score, which is based on the APRI, alanine aminotransferase (ALT) levels, and age of the subject (see, e.g., McPherson et al., Gut 2010, vol. 59(9), p. 1265-9, which is incorporated by reference herein in its entirety); hyaluronic acid; pro-inflammatory cytokines; a panel of biomarkers consisting of α 2-macroglobulin, haptoglobin, apolipoprotein A1, bilirubin, gamma glutamyl transpeptidase (GGT) combined with a subject's age and gender to generate a measure of fibrosis and necroinflammatory activity in the liver (e.g., FIBROTEST[®], FIBROSURE[®]), a panel of biomarkers consisting of bilirubin,

gamma-glutamyltransferase, hyaluronic acid, α 2-macroglobulin combined with the subject's age and sex (e.g., HEPASCOPE®; see, e.g., Adams et al., Clin. Chem. 2005, vol. 51(10), p. 1867-1873), and a panel of biomarkers consisting of tissue inhibitor of metalloproteinase-1, hyaluronic acid, and α 2-macroglobulin (e.g., FIBROSPECT®); a panel of biomarkers consisting of tissue inhibitor of metalloproteinases 1 (TIMP-1), amino-terminal propeptide of type III procollagen (PIIINP) and hyaluronic acid (HA) (e.g., the Enhanced Liver Fibrosis (ELF) score, see, e.g., Lichtinghagen R, et al., J Hepatol. 2013 Aug;59(2):236-42, which is incorporated by reference herein in its entirety). In some embodiments, the presence of fibrosis is determined by one or more of the FIB-4 score, a panel of biomarkers consisting of α 2-macroglobulin, haptoglobin, apolipoprotein A1, bilirubin, gamma glutamyl transpeptidase (GGT) combined with a subject's age and gender to generate a measure of fibrosis and necroinflammatory activity in the liver (e.g., FIBROTEST®, FIBROSURE®), a panel of biomarkers consisting of bilirubin, gamma-glutamyltransferase, hyaluronic acid, α 2-macroglobulin combined with the subject's age and sex (e.g., HEPASCOPE®; see, e.g., Adams et al., Clin. Chem. 2005, vol. 51(10), p. 1867-1873), and a panel of biomarkers consisting of tissue inhibitor of metalloproteinase-1, hyaluronic acid, and α 2-macroglobulin (e.g., FIBROSPECT®); and a panel of biomarkers consisting of tissue inhibitor of metalloproteinases 1 (TIMP-1), amino-terminal propeptide of type III procollagen (PIIINP) and hyaluronic acid (HA) (e.g., the Enhanced Liver Fibrosis (ELF) score). In some embodiments, the level of aspartate aminotransferase (AST) does not increase. In some embodiments, the level of aspartate aminotransferase (AST) decreases. In some embodiments, the level of alanine aminotransferase (ALT) does not increase. In some embodiments, the level of alanine aminotransferase (ALT) decreases. In some embodiments, the "level" of an enzyme refers to the concentration of the enzyme, e.g., within blood. For example, the level of AST or ALT can be expressed as Units/L.

In some embodiments, the severity of fibrosis is determined by one or more of the FIB-4 score, a panel of biomarkers consisting of α 2-macroglobulin, haptoglobin, apolipoprotein A1, bilirubin, gamma glutamyl transpeptidase (GGT) combined with a subject's age and gender to generate a measure of fibrosis and necroinflammatory activity in the liver (e.g., FIBROTEST®, FIBROSURE®), a panel of biomarkers consisting of bilirubin, gamma-glutamyltransferase, hyaluronic acid, α 2-macroglobulin combined with the subject's age and sex (e.g., HEPASCOPE®; see, e.g., Adams et al., Clin. Chem. 2005, vol. 51(10), p. 1867-1873, which is incorporated by reference herein in its entirety), and a panel of biomarkers consisting of tissue inhibitor of metalloproteinase-1, hyaluronic acid, and α 2-macroglobulin (e.g., FIBROSPECT®); and a panel of biomarkers consisting of tissue inhibitor of metalloproteinases 1 (TIMP-1), amino-terminal propeptide of type III procollagen (PIIINP) and hyaluronic acid (HA) (e.g., the Enhanced Liver Fibrosis (ELF) score).

In some embodiments, hepatic inflammation is determined by the level of liver inflammation biomarkers, e.g., pro-inflammatory cytokines. Non-limiting examples of biomarkers indicative of liver inflammation include interleukin-(IL) 6, interleukin-(IL) 1 β , tumor necrosis factor (TNF)- α ,
5 transforming growth factor (TGF)- β , monocyte chemotactic protein (MCP)-1, C-reactive protein (CRP), PAI-1, and collagen isoforms such as Col1a1, Col1a2, and Col4a1 (see, e.g., Neuman, et al., Can. J. Gastroenterol. Hepatol. 2014, vol. 28(11), p. 607-618 and U.S. Patent No. 9,872,844, each of which are incorporated by reference herein in their entireties). Liver inflammation can also be assessed by change of macrophage infiltration, e.g., measuring a change of CD68 expression level. In some
10 embodiments, liver inflammation can be determined by measuring or monitoring serum levels or circulating levels of one or more of interleukin-(IL) 6, interleukin-(IL) 1 β , tumor necrosis factor (TNF)- α , transforming growth factor (TGF)- β , monocyte chemotactic protein (MCP)-1, and C-reactive protein (CRP).

15 In some embodiments, the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis is determined for a sample from the subject prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis is determined during the period of time or
20 after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, a decrease in the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis during the period of time or after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, compared to prior to administration of the compound of
25 formula (I), or a pharmaceutically acceptable salt thereof, indicates treatment of NASH. For example, a decrease in the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least
30 about 90%, at least about 95%, or at least about 99% indicates treatment of NASH. In some embodiments, the decrease in the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis following administration of the compound of formula (I), or a pharmaceutically acceptable salt thereof, is by at least about 5%, at least about
35 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about

35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99%. In some embodiments, the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis during the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99%. In some embodiments, the level of one or more biomarkers indicative of one or more of liver damage, inflammation, liver fibrosis, and/or liver cirrhosis after the period of time of administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof, is by at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 60%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99%.

In some embodiments, the treatment of NASH decreases the level of serum bile acids in the subject. In some embodiments, the level of serum bile acids is determined by, for example, an ELISA enzymatic assay or the assays for the measurement of total bile acids as described in Danese et al., PLoS One. 2017, vol. 12(6): e0179200, which is incorporated by reference herein in its entirety. In some embodiments, the level of serum bile acids can decrease by, for example, 10% to 40%, 20% to 50%, 30% to 60%, 40% to 70%, 50% to 80%, or by more than 90% of the level of serum bile acids prior to administration of a compound of formula (I), or a pharmaceutically acceptable salt thereof. In some embodiments, the NASH is NASH with attendant cholestasis. In cholestasis, the release of bile, including bile acids, from the liver is blocked. Bile acids can cause hepatocyte damage (see, e.g., Perez MJ, Briz O. World J. Gastroenterol. 2009, vol. 15(14), p. 1677-1689) likely leading to or increasing the progression of fibrosis (e.g., cirrhosis) and increasing the risk of hepatocellular carcinoma (see, e.g., Sorrentino P et al., *Dig. Dis. Sci.* 2005, vol. 50(6), p. 1130-1135 and Satapathy SK and Sanyal AJ. *Semin. Liver Dis.* 2015, vol. 35(3), p. 221-235, each of which are incorporated by reference herein in their entireties). In some embodiments, the treatment of NASH includes treatment of pruritus. In some embodiments, the treatment of NASH with attendant cholestasis includes treatment of pruritus. In some embodiments, a subject with NASH with attendant cholestasis has pruritus.

35

Exemplary biomarkers for NASH are provided in Table 7.

Table 7. Exemplary NASH biomarkers

Liver Fibrosis Biomarkers

Aspartate aminotransferase (AST) to platelet ratio index (APRI)

Aspartate aminotransferase (AST) and alanine aminotransferase (ALT) ratio (AAR)

FIB-4 score¹

Hyaluronic acid

Pro-inflammatory cytokines

A panel including α 2-macroglobulin, haptoglobin, apolipoprotein A1, bilirubin, gamma glutamyl transpeptidase (GGT) combined with a subject's age and gender to generate a measure of fibrosis and necroinflammatory activity in the liver (e.g., FIBROTEST[®], FIBROSURE[®])

A panel including bilirubin, gamma-glutamyltransferase, hyaluronic acid, α 2-macroglobulin combined with the subject's age and sex (e.g., HEPAScore^{®2})

A panel including tissue inhibitor of metalloproteinase-1, hyaluronic acid, and α 2-macroglobulin (e.g., FIBROSPECT[®])

A panel including tissue inhibitor of metalloproteinases 1 (TIMP-1), amino-terminal propeptide of type III procollagen (PIIINP) and hyaluronic acid (HA) (e.g., the Enhanced Liver Fibrosis (ELF) score³)

Liver inflammation biomarkers^{4,5}

Interleukin-(IL) 6

Interleukin-(IL) 1 β

Tumor necrosis factor (TNF)- α

Transforming growth factor (TGF)- β

Monocyte chemotactic protein (MCP)-1

C-reactive protein (CRP)

PAI-1

Collagen isoforms (e.g., Col1a1, Col1a2, and Col4a1)

Change of macrophage infiltration (e.g., a change of CD68 expression level)

References for Table 7

¹ McPherson et al., Gut. 2010, vol. 59(9), p. 1265-1269.

² Adams, et al. Clin Chem. 2005, vol. 51(10), p. 1867-1873.

5 ³ Lichtinghagen, et al. J Hepatol. 2013, vol. 59(2), p. 236-242.

⁴ Neuman, et al. Can J Gastroenterol Hepatol. 2014, vol. 28(11), p. 607–618.

⁵ U.S. Patent No. 9,872,844

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may show a higher
10 free fraction in plasma. In some embodiments, the free fraction is greater than about 0.2%, such as
greater than about 0.4%, such as greater than about 0.6%, such as greater than about 0.8%, such as
greater than about 1.0%, such as greater than about 1.25%, such as greater than about 1.5%, such as
greater than about 1.75%, such as greater than about 2.0%, such as greater than about 2.5%, such as
15 greater than about 3%, such as greater than about 4%, such as greater than about 5%, such as
greater than about 7.5%, such as greater than about 10%, or such as greater than about 20%.

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may be excreted in
urine. In some embodiments, the fraction of the compound that is excreted in urine is greater than
about 0.2%, such as greater than about 0.4%, such as greater than about 0.6%, such as greater than
20 about 0.8%, such as greater than about 1.0%, such as greater than about 2%, such as greater than
about 3%, such as greater than about 5%, such as greater than about 7.5%, such as greater than
about 10%, such as greater than about 15%, such as greater than about 20%, such as greater than
about 30%, or such as greater than about 50%.

25 Following absorption from the intestine, some compounds of formula (I), or pharmaceutically
acceptable salts thereof, may be circulated via the enterohepatic circulation. In some embodiments,
the fraction of the compound that is circulated via the enterohepatic circulation is greater than about
0.1%, such as greater than about 0.2%, such as greater than about 0.3%, such as greater than about
0.5%, such as greater than about 1.0%, such as greater than about 1.5%, such as greater than about
30 2%, such as greater than about 3%, such as greater than about 5%, such as greater than about 7%,
such as greater than about 10%, such as greater than about 15%, such as greater than about 20%,
such as greater than about 30% or such as greater than about 50%.

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may cause renal excretion of bile salts. In some embodiments, the fraction of circulating bile acids that is excreted by the renal route is greater than about 1 %, such as greater than about 2%, such as greater than about 5%, such as greater than about 7%, such as greater than about 10%, such as greater than about 15%, such as greater than about 20%, or such as greater than about 25%.

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may show improved or optimal permeability. The permeability may be measured in Caco2 cells, and values are given as Papp (apparent permeability) values in cm/s. In some embodiments, the permeability is greater than at least about 0.1×10^{-6} cm/s, such as greater than about 0.2×10^{-6} cm/s, such as greater than about 0.4×10^{-6} cm/s, such as greater than about 0.7×10^{-6} cm/s, such as greater than about 1.0×10^{-6} cm/s, such as greater than about 2×10^{-6} cm/s, such as greater than about 3×10^{-6} cm/s, such as greater than about 5×10^{-6} cm/s, such as greater than about 7×10^{-6} cm/s, such as greater than about 10×10^{-6} cm/s, such as greater than about 15×10^{-6} cm/s.

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may show an improved or optimal bioavailability. In some embodiments, the oral bioavailability is greater than about 5%, such as greater than about 7%, such as greater than about 10%, such as greater than about 15%, such as greater than about 20%, such as greater than about 30%, such as greater than about 40%, such as greater than about 50 %, such as greater than about 60 %, such as greater than about 70% or such as greater than about 80%. In other embodiments, the oral bioavailability is between about 10 and about 90%, such as between about 20 and about 80%, such as between about 30 and about 70% or such as between about 40 and about 60%.

25

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may be a substrate to relevant transporters in the kidney.

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may give rise to concentrations of bile acids in the intestine, the liver and in serum that do not cause adverse gastrointestinal effects.

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may decrease the concentration of bile acids in the liver without causing gastrointestinal disorders such as diarrhoea.

35

As used herein, the terms "treatment", "treat" and "treating" refer to reversing, alleviating, delaying the onset of, or inhibiting the progress of a disease or disorder, or one or more symptoms thereof, as described herein. In some embodiments, treatment may be administered after one or more symptoms have developed. In other embodiments, treatment may be administered in the absence of symptoms. For example, treatment may be administered to a susceptible individual prior to the onset of symptoms (e.g., in light of a history of symptoms and/or in light of genetic or other susceptibility factors). Treatment may also be continued after symptoms have resolved, for example to prevent or delay their recurrence.

10 A suitable pharmaceutically acceptable salt of a compound of the invention is, for example, a base-addition salt of a compound of the invention which is sufficiently acidic, such as an alkali metal salt (e.g., a sodium or potassium salt), an alkaline earth metal salt (e.g., a calcium or magnesium salt), an ammonium salt, or a salt with an organic base which affords a physiologically acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-
15 hydroxyethyl)amine.

Some compounds of formula (I), or pharmaceutically acceptable salts thereof, may have chiral centres and/or geometric isomeric centres (E- and Z-isomers). It is to be understood that the invention encompasses all such optical isomers, diastereoisomers and geometric isomers that possess ASBT and/or LBAT inhibitory activity. The invention also encompasses any and all tautomeric forms of compounds of formula (I), or pharmaceutically acceptable salts thereof, that possess ASBT and/or LBAT inhibitory activity. Certain compounds of formula (I), or pharmaceutically acceptable salts thereof, may exist in unsolvated as well as solvated forms, such as, for example, hydrated forms. It is to be understood that the invention encompasses all such solvated forms that possess ASBT and/or LBAT inhibitory activity.

In another aspect, the invention relates to a pharmaceutical composition comprising a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable excipients. The excipients may e.g. include fillers, binders, disintegrants, glidants and lubricants. In general, pharmaceutical compositions may be prepared in a conventional manner using conventional excipients.

Examples of suitable fillers include, but are not limited to, dicalcium phosphate dihydrate, calcium sulfate, lactose (such as lactose monohydrate), sucrose, mannitol, sorbitol, cellulose, microcrystalline cellulose, dry starch, hydrolyzed starches and pregelatinized starch.

Examples of suitable binders include, but are not limited to, starch, pregelatinized starch, gelatin, sugars (such as sucrose, glucose, dextrose, lactose and sorbitol), polyethylene glycol, waxes, natural and synthetic gums (such as acacia gum and tragacanth gum), sodium alginate, cellulose derivatives
5 (such as hydroxypropylmethylcellulose (or hypromellose), hydroxypropylcellulose and ethylcellulose) and synthetic polymers (such as acrylic acid and methacrylic acid copolymers, methacrylic acid copolymers, methyl methacrylate copolymers, aminoalkyl methacrylate copolymers, polyacrylic acid/polymethacrylic acid copolymers and polyvinylpyrrolidone (povidone)).

10 Examples of suitable disintegrants include, but are not limited to, dry starch, modified starch (such as (partially) pregelatinized starch, sodium starch glycolate and sodium carboxymethyl starch), alginic acid, cellulose derivatives (such as sodium carboxymethylcellulose, hydroxypropyl cellulose, and low substituted hydroxypropyl cellulose (L-HPC)) and cross-linked polymers (such as carmellose, croscarmellose sodium, carmellose calcium and cross-linked PVP (crospovidone)).

15 Examples of suitable glidants and lubricants include, but are not limited to, talc, magnesium stearate, calcium stearate, stearic acid, glyceryl behenate, colloidal silica, aqueous silicon dioxide, synthetic magnesium silicate, fine granulated silicon oxide, starch, sodium lauryl sulfate, boric acid, magnesium oxide, waxes (such as carnauba wax), hydrogenated oil, polyethylene glycol, sodium benzoate,
20 polyethylene glycol, and mineral oil.

The pharmaceutical composition may be conventionally coated with one or more coating layers. Enteric coating layers or coating layers for delayed or targeted release of the compound of formula (I), or pharmaceutically acceptable salts thereof, are also contemplated. The coating layers may
25 comprise one or more coating agents, and may optionally comprise plasticizers and/or pigments (or colorants).

Example of suitable coating agents include, but are not limited to, cellulose-based polymers (such as ethylcellulose, hydroxypropylmethylcellulose (or hypromellose), hydroxypropylcellulose, cellulose
30 acetate phthalate, cellulose acetate succinate, hydroxypropyl methylcellulose acetate succinate and hydroxypropyl methylcellulose phthalate), vinyl-based polymers (such as polyvinyl alcohol) and polymers based on acrylic acid and derivatives thereof (such as acrylic acid and methacrylic acid copolymers, methacrylic acid copolymers, methyl methacrylate copolymers, aminoalkyl methacrylate copolymers, polyacrylic acid/polymethacrylic acid copolymers).

35

Examples of suitable plasticizers include, but are not limited to, triethyl citrate, glyceryl triacetate, tributyl citrate, diethyl phthalate, acetyl tributyl citrate, dibutyl phthalate, dibutyl sebacate and polyethylene glycol.

- 5 Examples of suitable pigments include, but are not limited to, titanium dioxide, iron oxides (such as yellow, brown, red or black iron oxides) and barium sulfate.

The pharmaceutical composition may be in a form that is suitable for oral administration, for parenteral injection (including intravenous, subcutaneous, intramuscular and intravascular injection),
10 for topical administration or for rectal administration. In a preferred embodiment, the pharmaceutical composition is in a form that is suitable for oral administration, such as a tablet or a capsule.

The dosage required for the therapeutic or prophylactic treatment will depend on the route of
15 administration, the severity of the disease, the age and weight of the patient and other factors normally considered by the attending physician, when determining the appropriate regimen and dosage level for a particular patient.

The amount of the compound to be administered will vary for the patient being treated, and may
20 vary from about 1 µg/kg of body weight to about 50 mg/kg of body weight per day. A unit dose form, such as a tablet or capsule, will usually contain about 1 to about 250 mg of active ingredient, such as about 1 to about 100 mg, or such as about 1 to about 50 mg, or such as about 1 to about 20 mg, e.g. about 2.5 mg, or about 5 mg, or about 10 mg, or about 15 mg. The daily dose can be administered as a single dose or divided into one, two, three or more unit doses. An orally administered daily dose of
25 a bile acid modulator is preferably within about 0.1 to about 250 mg, more preferably within about 1 to about 100 mg, such as within about 1 to about 5 mg, such as within about 1 to about 10 mg, such as within about 1 to about 15 mg, or such as within about 1 to about 20 mg.

In another aspect, the invention relates to a compound of formula (I), or a pharmaceutically
30 acceptable salt thereof, for use as a medicament. The invention also relates to the use of a compound of formula (I), or a pharmaceutically acceptable salt thereof, as a medicament.

In another aspect, the invention relates to a compound of formula (I), or a pharmaceutically acceptable salt thereof, for use in the treatment or prevention of any of the diseases recited herein.
35 The invention also relates to the use of a compound of formula (I), or a pharmaceutically acceptable

salt thereof, in the manufacture of a medicament for the treatment or prevention of any of the diseases recited herein. The invention also relates to a method of treating or preventing any of the diseases recited herein in a subject, such as man, comprising administering to the subject in need of such treatment or prevention a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof.

Combination therapy

In one aspect of the invention, the compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with at least one other therapeutically active agent, such as with one, two, three or more other therapeutically active agents. The compound of formula (I), or a pharmaceutically acceptable salt thereof, and the at least one other therapeutically active agent may be administered simultaneously, sequentially or separately. Therapeutically active agents that are suitable for combination with the compounds of formula (I) include, but are not limited to, known active agents that are useful in the treatment of any of the aforementioned conditions, disorders and diseases.

In one embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with another ASBT inhibitor. Suitable ASBT inhibitors are disclosed in WO 93/16055, WO 94/18183, WO 94/18184, WO 96/05188, WO 96/08484, WO 96/16051, WO 97/33882, WO 98/03818, WO 98/07449, WO 98/40375, WO 99/35135, WO 99/64409, WO 99/64410, WO 00/47568, WO 00/61568, WO 00/38725, WO 00/38726, WO 00/38727, WO 00/38728, WO 00/38729, WO 01/66533, WO 01/68096, WO 02/32428, WO 02/50051, WO 03/020710, WO 03/022286, WO 03/022825, WO 03/022830, WO 03/061663, WO 03/091232, WO 03/106482, WO 2004/006899, WO 2004/076430, WO 2007/009655, WO 2007/009656, WO 2011/137135, WO 2019/234077, WO 2020/161216, WO 2020/161217, WO 2021/110883, WO 2021/110884, WO 2021/110885, WO 2021/110886, WO 2021/110887, WO 2022/029101, DE 19825804, EP 864582, EP 489423, EP 549967, EP 573848, EP 624593, EP 624594, EP 624595, EP 624596, EP 0864582, EP 1173205, EP 1535913 and EP 3210977, all of which are incorporated herein by reference in their entireties. Particular examples of suitable ASBT inhibitors include 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)-1'-phenyl-1'-[*N*'-(carboxymethyl)carbamoyl]-methyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine (elobixibat) and 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N*'-(*S*)-1-carboxypropyl] carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine (odevixibat).

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a bile acid binder (also referred to as a bile acid sequestrant, or a resin), such as colestevlam, cholestyramine or cholestipol. In a preferred embodiment of such a combination, the bile acid binder is formulated for colon release. Examples of such formulations are disclosed in e.g. WO 2017/138877, WO 2017/138878, WO 2019/032026 and WO 2019/032027, all of which are incorporated herein by reference in their entireties.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a DPP-IV inhibitor, including gliptins such as sitagliptin, vildagliptin, saxagliptin, linagliptin, gemigliptin, anagliptin, teneligliptin, alogliptin, trelagliptin, omarigliptin, evogliptin, gosogliptin and dutogliptin, or a pharmaceutically acceptable salt thereof.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an HMG CoA reductase inhibitor, such as fluvastatin, lovastatin, pravastatin, simvastatin, atorvastatin, pitavastatin cerivastatin, mevastatin, rosuvastatin, bervastatin or dalvastatin, or a pharmaceutically acceptable salt thereof.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a cholesterol absorption inhibitor such as ezetimibe, or a pharmaceutically acceptable salt thereof.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a PPAR alpha agonist, including fibrates such as clofibrate, bezafibrate, ciprofibrate, clinofibrate, clofibrade, fenofibrate, gemfibrozil, ronifibrate and simfibrate, or a pharmaceutically acceptable salt thereof.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a PPAR gamma agonist, including thiazolidinediones such as pioglitazone, rosiglitazone and lobeglitazone, or a pharmaceutically acceptable salt thereof.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a dual PPAR alpha/gamma agonist, including glitazars such as saroglitazar, aleglitazar, muraglitazar or tesaglitazar, or a pharmaceutically acceptable salt thereof.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a dual PPAR alpha/delta agonist, such as elafibranor.

5 In yet another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a pan PPAR agonist (i.e. a PPAR agonist that has activity across all subtypes: α , γ and δ), such as IVA337.

10 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a farnesoid X receptor (FXR) modulators, including FXR agonists such as cafestol, chenodeoxycholic acid, 6 α -ethyl-chenodeoxycholic acid (obeticholic acid; INT-747), fexaramine, tropifexor, cilofexor and MET409.

15 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a TGR5 receptor modulator, including TGR5 agonists such as 6 α -ethyl-23(S)-methylcholic acid (INT-777).

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a dual FXR/TGR5 agonist such as INT-767.

20 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with ursodeoxycholic acid (UDCA). In yet another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with nor-ursodeoxycholic acid (nor-UDCA).

25 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an FGF19 modulator, such as NGM282.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an FGF21 agonist, such as BMS-986036.

30

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an integrin inhibitor, such as PLN-74809 and PLN-1474.

35 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a CCR2/CCR5 inhibitor, such as cenicriviroc.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a caspase protease inhibitor, such as emricasan.

- 5 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a galectin-3 inhibitor, such as GR-MD-02.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a stearyl-CoA desaturase (SCD) Inhibitor, such as aramchol
10 (arachidyl amido cholanoic acid).

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an apoptosis signal-regulating kinase 1 (ASK1) inhibitor, such as selonsertib.

15

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an LOXL2 inhibitor, such as simtuzumab.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are
20 administered in combination with an ACC inhibitor, such as GS-0976.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a thyroid hormone receptor- β agonist, such as MGL3196.

25 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a GLP-1 agonist such as liraglutide.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a dual glucagon-like peptide and glucagon receptor agonists, such
30 as SAR425899.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a mitochondrial pyruvate carrier inhibitor, such as MSDC-0602K.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an anti-oxidant agent, such as vitamin E.

5 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an SGLT1 inhibitor, an SGLT2 inhibitor or a dual SGLT1 and SGLT2 inhibitor. Examples of such compounds are dapagliflozin, sotagliflozin, canagliflozin, empagliflozin, LIK066 and SGL5213.

10 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a diacylglycerol O-Acyltransferase 2 (DGAT2) inhibitor, such as DGAT2RX and PF-06865571.

15 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a fatty acid synthase (FASN) Inhibitor, such as TVB-2640.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an AMP-activated protein kinase (AMPK) activator, such as PXL-770.

20 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a glucocorticoid receptor antagonist (GR), a mineralocorticoid receptor antagonist (MR), or a dual GR/MR antagonist. Examples of such compounds are MT-3995 and CORT-118335.

25 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a cannabinoid receptor 1 (CB1) antagonist, such as IM102.

30 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a Klotho β (KLB) and fibroblast growth factor receptor (FGFR) activator, such as MK-3655 (previously known as NGM-313).

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a chemokine (c-c motif) ligand 24 (CCL24) inhibitor, such as CM101.

35

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an A3 antagonist, such as PBF-1650.

5 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a P2x7 receptor antagonist, such as SGM 1019.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with P2Y13 receptor agonists, such as CER-209.

10 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a sulfated oxysterol, such as Dur-928.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a leukotriene D4 (LTD4) receptor antagonist, such as MN-001.

15

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a type 1 natural killer T cell (NKT1) inhibitor, such as GRI-0621.

20 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an anti-lipopolysaccharide (LPS) compound, such as IMM-124E.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a VAP1 inhibitor, such as BI1467335.

25 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an A3 adenosine receptor agonist, such as CF-102.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a SIRT-1 activator, such as NS-20.

30

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a nicotinic acid receptor 1 agonist, such as ARI-3037MO.

35 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a TLR4 antagonist, such as JKB-121.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a ketohexokinase inhibitor, such as PF-06835919.

- 5 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an adiponectin receptor agonist, such as ADP-335.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with an autotaxin inhibitor, such as PAT-505 and PF8380.

10

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a chemokine (c-c motif) receptor 3 (CCR3) antagonist, such as bertilimumab.

- 15 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a chloride channel stimulator, such as cobiprostone and lubiprostone.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a heat shock protein 47 (HSP47) inhibitor, such as ND-L02-s0201.

20

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a sterol regulatory element-binding protein (SREBP) transcription factor inhibitor, such as CAT-2003 and MDV-4463.

25

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a biguanidine, such as metformin.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with insulin.

30

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a glycogen phosphorylase inhibitor and/or a glucose-6-phosphatase inhibitor.

35

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a sulfonylurea, such as glipizid, glibenklamid and glimepirid.

5 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a meglitinide, such as repaglinide, nateglinide and ormglinide.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a glucosidase inhibitor, such as acarbose or miglitol.

10 In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a squalene synthase inhibitor, such as TAK-475.

In another embodiment, compounds of formula (I), or pharmaceutically acceptable salts thereof, are administered in combination with a PTPB1 inhibitor, such as trodusquemine, ertiprotafib, JTT-551
15 and claramine.

Preparation of compounds

The compounds of the invention can be prepared as a free acid or a pharmaceutically acceptable salt
20 thereof by the processes described below. Throughout the following description of such processes it is understood that, where appropriate, suitable protecting groups will be added to, and subsequently removed from the various reactants and intermediates in a manner that will be readily understood by one skilled in the art of organic synthesis. Conventional procedures for using such protecting groups as well as examples of suitable protecting groups are for example described in *Greene's*
25 *Protective Groups in Organic Synthesis* by P.G.M Wutz and T.W. Greene, 4th Edition, John Wiley & Sons, Hoboken, 2006.

General methods

30 All solvents used were of analytical grade. Commercially available anhydrous solvents were routinely used for reactions. Starting materials were available from commercial sources or prepared according to literature procedures. 3-Butyl-8-hydroxy-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepine 1,1-dioxide and 3-ethyl-8-hydroxy-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepine 1,1-dioxide may be prepared as described in WO 2021/110883 (Intermediates 7 and

18, respectively). Room temperature refers to 20 - 25 °C. Solvent mixture compositions are given as volume percentages or volume ratios.

LCMS:

- 5 **Instrument name:** Agilent 1290 infinity II.
- Method A:** Mobile phase: A: 0.1% HCOOH in water: ACN (95:5), B: ACN; flow rate: 1.5 mL/min; column: ZORBAX XDB C-18 (50 x 4.6 mm) 3.5 µM.
- Method B:** Mobile phase: A: 10 mM NH₄HCO₃ in water, B: ACN; flow rate: 1.2 mL/min; column: XBridge C8 (50 x 4.6 mm), 3.5 µM.
- 10 **Method C:** Mobile phase: A: 0.1% HCOOH in water: ACN (95:5), B: ACN; flow rate: 1.5 mL/min; column: ATLANTIS dC18 (50 x 4.6 mm), 5 µM.
- Method D:** Mobile phase: A: 10 mM NH₄OAc in water, B: ACN; flow rate: 1.2 mL/min; column: Zorbax Extend C18 (50 x 4.6mm) 5 µM.
- Method E:** Mobile Phase: A: 0.1% TFA in water: ACN (95:5), B: 0.1% TFA in ACN; flow rate: 1.5 mL/min; Column: XBridge C8 (50 x 4.6 mm), 3.5 µM.
- 15 **Method F:** Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; flow Rate: 0.8 mL/min; column: ZORBAX ECLIPSE PLUS C18 (50 x 2.1 mm), 1.8 µm.
- Method G:** Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; flow rate: 0.8 mL/min; column: Acquity UPLC BEH C18 (2.1 x 50 mm), 1.7 µm.
- 20 **Method H:** Mobile phase: A: 10 mM NH₄OAc, B: 100% ACN; flow rate: 0.8 mL/min; Column: Acquity UPLC BEH C18 (2.1 x 50) mm; 1.7 µm.
- Method I:** Mobile phase: A: 0.1% HCOOH in water: ACN (95:5), B: ACN; flow rate: 0.8 mL/min; Column: ZORBAX ECLIPSE PLUS C18 (2.1 x 50) mm, 1.8 µm.
- Method J:** Mobile phase: A: 0.1% TFA in water, B: ACN; Flow rate: 1.0 mL/min; Column: Zorbax Extend C18 (50 x 4.6 mm), 5 µM.
- 25 **Method K:** Mobile Phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; Flow Rate: 1.5 mL/min; Column: XBridge C8 (50 x 4.6 mm), 3.5 µM.

UPLC:

- 30 **Instrument name:** waters Acquity I Class
- Method A:** Mobile Phase: A: 0.1% HCOOH in water, B: 0.1% HCOOH in ACN; Flow Rate: 0.8 mL/min; Column: Acquity UPLC HSS T3 (2.1 x 50) mm; 1.8 µm.

Instrument Name : Shimadzu Nexera X2 LC / 2020 MSD

Method B: Mobile Phase: A: 0.1% HCOOH in water, B: ACN; Flow Rate: 0.8 mL/min; Column: Acquity UPLC BEH C18 (2.1 x 50) mm; 1.7 µm.

5 **HPLC:**

Instrument name: Agilent 1260 Infinity II series instruments as followed using % with UV detection (maxplot).

Method A: Mobile phase: A: 10 mM NH₄HCO₃ in water, B: ACN; flow rate: 1.0 mL/min; column: XBridge C8 (50 x 4.6 mm, 3.5 µm).

10 **Method B:** Mobile phase: A: 0.1% TFA in water, B: 0.1% TFA in ACN; flow rate: 2.0 mL/min; column: XBridge C8 (50 x 4.6 mm, 3.5 µm).

Method C: Mobile phase: A: 10 mM NH₄OAc in milli-q water, B: ACN; flow rate: 1.0 ml/min; column: Phenomenex Gemini C18 (150 x 4.6 mm, 3.0 µm).

Method D: Mobile phase: A: 0.1% TFA in water, B: ACN; flow rate: 1.0 mL/min; column: ATLANTIS dC18 (250 x 4.6 mm, 5.0 µm).

15 **Method E:** Mobile phase: A: 0.1% TFA in water, B: ACN, flow rate: 2.0 mL/ min; column: X-Bridge C8 (50 X 4.6 mm, 3.5 µm).

Chiral SFC:

20 **Instrument name:** PIC SFC 10 (analytical)

Ratio between CO₂ and co-solvent is ranging between 60:40 and 80:20

Method A: Mobile phase: 0.5% isopropylamine in IPA; flow rate: 3 mL/min; column: YMC Amylose-SA (250 x 4.6 mm, 5 µm).

Method B: Mobile Phase: 0.5% isopropylamine in IPA; flow rate: 3 mL/min; column: Chiralpak AD-H (250 x 4.6 mm, 5 µm).

Method C: Mobile Phase: 20 mM ammonia in methanol; flow rate: 3 mL/min; column: YMC Cellulose-SC (250 x 4.6 mm, 5 µm).

Method D: Mobile Phase: methanol; flow rate: 3 mL/min; column: Lux A1 (250 x 4.6 mm, 5 µm).

Method E: Mobile Phase: 0.5% isopropylamine in methanol; flow rate: 5 mL/min; column: Lux C4.

30 **Method F:** Mobile Phase: 0.5% isopropylamine in methanol; flow rate: 3 mL/min; column: YMC Cellulose-SC.

Method G: Mobile Phase: 0.5% isopropylamine in methanol; flow rate: 3 mL/min; column: Lux A1.

Method H: Mobile Phase: 0.5% isopropylamine in IPA; flow rate: 3 mL/min; column: Lux A1 (250 x 4.6 mm, 5 µm).

Method I: Mobile phase: 0.5% isopropylamine in methanol; flow rate: 3 mL/min; column: Chiral CCS (250 x 4.6 mm, 5 µm).

Method J: Mobile Phase: 0.5% isopropylamine in IPA; flow rate: 5 mL/min; column: YMC Cellulose-SC AD-H (250 x 4.6 mm, 5 µm).

5 **Method K:** Mobile phase: 0.5% Isopropylamine in methanol; flow rate: 4 mL/min; column: (R,R)-Whelk-01 (250 x 4.6 mm, 5 µm).

Method L: Mobile phase: 0.5% Isopropylamine in IPA; flow rate: 3 mL/min; column: Chiralcel OX-H (250 x 4.6 mm, 5 µm).

10 **Method M:** Mobile phase: 0.5% Isopropylamine in IPA; flow rate: 5 mL/min; column: YMC Cellulose-SC (250 x 4.6 mm, 5 µm).

Method N: Mobile phase: methanol, flow rate: 5 mL/min; column: Chiralcel OX-H (250 x 4.6 mm, 5 µm).

Method O: Mobile phase: 0.1% Isopropylamine in IPA:methanol (1:1), flow rate: 3 mL/min; column: Chiralpak AS-H (250 x 4.6 mm, 5 µm).

15 **Method P:** Mobile phase: 0.5% Isopropylamine in methanol, flow rate: 3 mL/min; column: Chiralpak AS-H (250 x 4.6 mm, 5 µm).

Method Q: Mobile phase: IPA, flow rate: 3 mL/min; column: Lux A1 (250 x 4.6 mm, 5 µm).

Method R: Mobile phase: 0.1% Isopropylamine in IPA:methanol (1:1), flow rate: 3 mL/min; column: Lux A1 (250 x 4.6 mm, 5 µm).

20 **Method S:** Mobile phase: 0.5% Isopropylamine in methanol, flow rate: 3 mL/min; column: Chiralpak OX-H (250 x 4.6 mm, 5 µm).

Method T: Mobile phase: 0.5% Isopropylamine in IPA, flow rate: 4 mL/min; column: YMC Cellulose-SB (250 x 4.6 mm, 5 µm).

25 **Method U:** Mobile phase: 0.5% Isopropylamine in IPA, flow rate: 3 mL/min; column: Chiralpak AS-H (250 x 4.6 mm, 5 µm).

Prep-HPLC:

Instrument name: Agilent 1290 Infinity II

30 **Method A:** Mobile phase: A: 0.1% TFA in water; Mobile phase; B: 0.1% TFA in ACN; flow rate: 2.0 mL/min; Column: X-Bridge C8 (50 X 4.6 mm, 3.5 µM).

Method B: Mobile phase: A: 10 mM NH₄OAc in water; B: ACN; flow rate: 35 mL/min; column: X select C18 (30 x 150 mm, 5 µm).

Method C: Mobile phase: A: 10 mM NH₄HCO₃ in water; B: ACN; flow rate: 1.0 mL/min; column: XBridge C8 (50 x 4.6 mm, 3.5 µm).

Method D: Mobile phase: A: 0.1% HCOOH in water; B: ACN; flow rate: 1.0 mL/min; column: X-select C18 (30 x 150 mm, 5 µm).

Chiral Preparative SFC:

- 5 **Instrument name:** PIC SFC 100 and PSC SFC 400
Ratio between CO₂ and co-solvent is ranging between 60:40 and 80:20
- Method A:** Mobile phase: 0.5% isopropylamine in IPA; flow rate: 3 mL/min; column: YMC Amylose-SA (250 x 30 mm, 5 µm).
- Method B:** Mobile Phase: 0.5% isopropylamine in IPA; flow rate: 3 mL/min; column: Chiralpak AD-H
10 (250 x 30 mm, 5 µm).
- Method C:** Mobile phase: 20 mM ammonia in methanol; flow rate: 3 mL/min; column: YMC Cellulose-SC (250 x 30 mm, 5 µm).
- Method D:** Mobile phase: methanol; flow rate: 3 mL/min; column: Chiral CCS (250 x 30 mm, 5 µm).
- Method E:** Mobile phase: methanol; flow rate: 3 mL/min; column: Lux A1 (250 x 30 mm, 5 µm).
- 15 **Method F:** Mobile Phase: 0.5% isopropylamine in IPA; flow rate: 3 mL/min; column: Lux A1 (250 x 30 mm, 5 µm).
- Method G:** Mobile phase: 0.5% isopropylamine in methanol; flow rate: 3 mL/min; column: Chiral CCS (250 x 30 mm, 5 µm).
- Method H:** Mobile phase: 0.5% Isopropylamine in methanol; flow rate: 4 mL/min; column: (R,R)-
20 Whelk-01 (250 x 30 mm, 5 µm).
- Method I:** Mobile phase: 0.5% Isopropylamine in IPA; flow rate: 5 mL/min; column: YMC Cellulose-SC (250 x 30 mm, 5 µm).
- Method J:** Mobile phase: 0.5% Isopropylamine in IPA; flow rate: 3 mL/min; column: Chiralcel OX-H (250 x 30 mm, 5 µm).
- 25 **Method K:** Mobile phase: 0.5% Isopropylamine in methanol; flow rate: 5 mL/min; column: YMC Cellulose-SC (250 x 30 mm, 5 µm).
- Method L:** Mobile phase: Methanol; flow rate: 5 mL/min; column: Chiralcel OX-H (250 x 30 mm, 5 µm).

30 **Abbreviations**


ACN	acetonitrile
BOC	<i>tert</i> -butoxycarbonyl
DBAD	di- <i>tert</i> -butyl azodicarboxylate
35 DCM	dichloromethane

	DMF	dimethylformamide
	HPLC	high-performance liquid chromatography
	IPA	isopropyl alcohol
	LCMS	liquid chromatography - mass spectrometry
5	NMP	<i>N</i> -methyl-2-pyrrolidone
	PE	petroleum ether
	SFC	supercritical fluid chromatography
	TFA	trifluoroacetic acid
	THF	tetrahydrofuran
10	TLC	thin layer chromatography
	UPLC	ultra performance liquid chromatography

The invention will now be described by the following examples which do not limit the invention in any respect. All cited documents and references are incorporated by reference.

15

EXAMPLES

Intermediate 1**2-((*tert*-Butoxycarbonyl)amino)hexanoic acid**


20

To a solution of 2-aminohexanoic acid (25 g, 0.19 mol) in water (250 mL) and THF (250 mL), NaHCO₃ (48 g, 0.57 mol) and Boc-anhydride (52.2 mL, 0.23 mol) were added and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was cooled, quenched and acidified using 1.5N HCl. The reaction mixture was

25 extracted with EtOAc (2 x 150 mL). The combined organic layer was washed with ice-cold water (150 mL) and brine (150 mL), dried over anhydrous Na₂SO₄ and concentrated to afford the title compound. **Yield:** 45.5 g (crude, white solid).


¹H NMR (400 MHz, DMSO-*d*₆): δ 5.10-4.98 (m, 1H), 4.33-4.30 (m, 1H), 1.95-1.81 (m, 1H), 1.75-1.55 (m, 1H), 1.46 (s, 10H), 1.45-1.31 (m, 4H), 0.93 (t, *J* = 7.2 Hz, 3H). **LCMS:** (Method E) 132.2 (M⁺-Boc+H),

30 Rt. 2.36 min, 99.98% (max).

Intermediate 2***tert*-Butyl (1-oxo-1-(phenylamino)hexan-2-yl)carbamate**

To a stirred solution of 2-((*tert*-butoxycarbonyl)amino)hexanoic acid (Intermediate 1; 45.5 g, 0.196 mol) in DMF (150 mL), triethylamine (54.83 mL, 0.39 mol) was added and the reaction mixture was cooled to 0 °C. 1-Propanephosphonic anhydride solution (50% in EtOAc; 75.1 g, 0.23 mol) and aniline (18 g, 0.196 mol) were added and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with ice-cold water (50 mL) and diluted with EtOAc (200 mL). The aqueous layer was washed with ice-cold water (200 mL) and brine (200 mL), and then dried over anhydrous Na₂SO₄ and filtered off. The organic part was concentrated under vacuum to afford the title compound. **Yield:** 82% (49.7 g, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 9.93 (s, 1H), 7.60 (d, *J* = 7.6 Hz, 2H), 7.30 (t, *J* = 8.0 Hz, 2H), 7.06-6.99 (m, 2H), 4.07-4.02 (m, 1H), 1.63-1.56 (m, 2H), 1.46-1.28 (m, 13H), 0.86 (t, *J* = 6.8 Hz, 3H). **LCMS:** (Method E) 207.0 (M⁺-Boc+H), Rt. 2.69 min, 91.20% (max).


Intermediate 3**2-Amino-N-phenylhexanamide**

To a solution of *tert*-butyl (1-oxo-1-(phenylamino)hexan-2-yl)carbamate (Intermediate 2; 49.7 g, 0.162 mol) in 1,4-dioxane (500 mL) at 0 °C, a solution of HCl in 1,4-dioxane (4 M, 199 mL) was added and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was concentrated under vacuum and the obtained residue was quenched with saturated NaHCO₃ solution. The aqueous layer was extracted with EtOAc (2 x 200 mL). The combined organic layer was washed brine (200 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum to afford the title compound. **Yield:** 95% (32 g, colourless gum).

¹H NMR (400 MHz, DMSO-*d*₆): δ 10.20-9.66 (m, 1H), 7.64 (dd, *J* = 4.8, 2.4 Hz, 2H), 7.32-7.27 (m, 2H), 7.06-7.02 (m, 1H), 3.29-3.26 (m, 1H), 1.67-1.61 (m, 1H), 1.46-1.27 (m, 5H), 0.89-0.85 (m, 3H). LCMS: (Method B) 207.2 (M⁺+H), Rt. 2.03 min, 84.36% (max).

5 Intermediate 4


N1-Phenylhexane-1,2-diamine



To a solution of 2-amino-N-phenylhexanamide (Intermediate 3; 32 g, 0.15 mol) in THF (320 mL) at 0 °C was added borane dimethylsulfide (2M solution in THF, 117 mL, 0.23 mol) and the reaction mixture was heated for 16 hours at 75 °C. After completion of the reaction (monitored by TLC), the reaction mixture was cooled to 0 °C, quenched with methanol (150 mL) and then heated for 2 hours at 60 °C. The reaction mixture was cooled to room temperature and concentrated under vacuum. The obtained residue was partitioned between water (50 mL) and EtOAc (50 mL). The aqueous layer was extracted with EtOAc (2 x 250 mL). The combined organic layer was washed with water (250 mL) and brine (250 mL). The organic part was dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 10% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 93% (28 g, yellow oil). **LCMS:** (Method B) 193.3 (M⁺+H), Rt. 1.98 min, 76.9% (max).

20 Intermediate 5

2,4-Dibromo-5-methoxy-N-(1-(phenylamino)hexan-2-yl)benzenesulfonamide




To a solution of 2,4-dibromo-5-methoxybenzenesulfonyl chloride (3.2 g, 8.79 mmol) in THF (50 mL) at 0 °C were added N1-phenylhexane-1,2-diamine (Intermediate 4; 1.3 g, 6.76 mmol) and triethylamine (2.8 mL, 20.3 mmol) and the reaction mixture was stirred for 4 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was concentrated under vacuum and the obtained residue was dissolved in EtOAc (60 mL). The organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The

resulting crude material was purified by Isolera column chromatography (eluent: 18% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 97% (3.5 g, brown gum). **LCMS:** (Method E) 521.0 (M^+H), Rt. 3.14 min, 93.57% (max).

5 Intermediate 6


7-Bromo-3-butyl-8-methoxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide



To a solution of 2,4-dibromo-5-methoxy-N-(1-(phenylamino)hexan-2-yl)benzenesulfonamide (Intermediate 5; 3.5 g, 6.72 mmol) in DMF (30 mL), K_2CO_3 (1.73 g, 12.5 mmol) and copper powder (0.42 g, 6.72 mmol) were added. The reaction mixture was degassed for 5 minutes under N_2 atmosphere and the reaction mixture was then heated for 16 hours at 115 °C. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with ice-cold water (25 mL) and the aqueous layer was extracted with a 1:1 mixture of EtOAc and PE (2 x 50 mL). The combined organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 25% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 51% (1.5 g, brown gum). **LCMS:** (Method A) 439.0 (M^+H), Rt. 2.83 min, 82.06% (max).

Intermediate 7

20 **7-Bromo-3-butyl-8-methoxy-2-methyl-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide**




To a stirred solution of 7-bromo-3-butyl-8-methoxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 6; 4.5 g, 10.24 mmol) in N-methyl-2-pyrrolidone (20 mL) were added Cs_2CO_3 (6.7 g, 20.5 mmol) and then iodomethane (3.2 mL, 51.2 mmol) and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with water (25 mL) and the aqueous layer was extracted

with a mixture of EtOAc and PE (30%, 2 x 75 mL). The combined organic layer was washed with brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude material was forwarded as such to the next step without any further purification. **Yield:** 4.5 g (crude, pale brown solid).

5 **LCMS:** (Method E) 452.8 (M⁺+H), Rt. 3.19 min, 95.26% (max).

Intermediate 8

3-Butyl-8-hydroxy-2-methyl-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide



10


To a solution of 7-bromo-3-butyl-8-methoxy-2-methyl-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 7; 4.5 g, 9.92 mmol) in DMF (50 mL), sodium thiomethoxide (3.5 g, 49.6 mmol) was added and the reaction mixture was heated for 16 hours at 80 °C. After completion of the reaction (monitored by UPLC), the reaction mixture was quenched with ice-cold water (25 mL) and the aqueous layer was extracted with EtOAc (4 x 50 mL). The combined organic layer was washed with water (2 x 100 mL) and brine (100 mL) and dried over anhydrous Na₂SO₄. The organic part was concentrated under vacuum and the resulting crude material was purified by Isolera column chromatography (eluent: 26% EtOAc PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 77% (3.1 g, off-white solid).

15

20 **¹H NMR** (400 MHz, DMSO-*d*₆): δ 10.73 (s, 1H), 7.29 (s, 1H), 7.12-7.16 (m, 2H), 6.99 (s, 1H), 6.67 (t, *J* = 7.2 Hz, 1H), 6.54 (d, *J* = 8.0 Hz, 2H), 4.06-3.89 (m, 2H), 3.21-3.09 (m, 1H), 2.42 (s, 3H), 2.34 (s, 3H), 1.68-1.55 (m, 1H), 1.54-1.42 (m, 1H), 1.41-1.25 (m, 4H), 0.93-0.90 (m, 3H). **LCMS:** (Method A) 406.9 (M⁺+H), Rt. 2.65 min, 92.05% (max).

25 Intermediate 9

(S)-3-Butyl-8-hydroxy-2-methyl-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide and (R)-3-butyl-8-hydroxy-2-methyl-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide




The two enantiomers of racemic 3-butyl-8-hydroxy-2-methyl-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 8; 10.0 g, 24.59 mmol) were separated by SFC Instrument (method E). The material was concentrated under vacuum at 40 °C. The first eluting fraction corresponded to enantiomer 1 and the second eluting fraction corresponded to enantiomer 2. The absolute configuration of the two enantiomers is not known.

Enantiomer 1: **Yield:** 42% (4.3 g, off-white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 10.69 (s, 1H), 7.28 (s, 1H), 7.13 (t, *J* = 10.8 Hz, 2H), 6.99 (s, 1H), 6.66 (t, *J* = 9.6 Hz, 1H), 6.53 (d, *J* = 10.8 Hz, 2H), 4.00-3.89 (m, 2H), 3.19-3.14 (m, 1H), 2.42 (s, 3H), 2.34 (s, 3H), 1.68-1.55 (m, 1H), 1.55-1.45 (m, 1H), 1.40-1.25 (m, 4H), 0.95-0.85 (m, 3H). **LCMS:** (Method A) 407.1 (M⁺+H), Rt. 2.56 min, 98.41% (Max). **HPLC:** (Method E) Rt. 5.43 min, 98.58% (Max). **Chiral SFC:** (method D) Rt. 1.83 min, 100% (Max).

Enantiomer 2: **Yield:** 34% (4.2 g, off-white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 10.58 (s, 1H), 7.27 (s, 1H), 7.13 (t, *J* = 10.0 Hz, 2H), 6.98 (s, 1H), 6.66 (t, *J* = 10.0 Hz, 1H), 6.53 (d, *J* = 10.8 Hz, 2H), 4.10-3.80 (m, 2H), 3.25-3.05 (m, 1H), 2.42 (s, 3H), 2.34 (s, 3H), 1.68-1.45 (m, 2H), 1.45-1.25 (m, 4H), 0.98-0.85 (m, 3H). **LCMS:** (Method A) 407.2 (M⁺+H), Rt. 2.56 min, 98.25% (Max). **HPLC:** (Method E) Rt. 5.43 min, 97.49% (Max). **Chiral SFC:** (method D) Rt. 3.06 min, 99.76% (Max).

Intermediate 10

Methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoate



To a stirred solution of 3-butyl-8-hydroxy-2-methyl-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 8; 0.2 g, 0.49 mmol) in THF (5 mL) at 0 °C, methyl 3-hydroxy-2,2-dimethylpropanoate (0.07 g, 0.49 mmol) and triphenylphosphine (0.19 g, 0.73 mmol) were added and the reaction mixture was stirred for 10 minutes. DBAD (0.22 g, 0.98 mmol) was then added at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After


completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 5 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 20-22% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.

Yield: 60% (0.15 g, white solid).

LCMS: (Method K) 521.1 (M⁺+H), Rt. 3.23 min, 75.58% (Max).

Intermediate 11

- 10 **Methyl (S)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoate and methyl (R)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoate**



- 15 To a stirred solution of enantiomer 1 of 3-butyl-8-hydroxy-2-methyl-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 9; 0.1 g, 0.24 mmol) in THF (5 mL) at 0 °C, methyl 3-hydroxy-2,2-dimethylpropanoate (0.04 g, 0.24 mmol) and triphenylphosphine (0.09 g, 0.36 mmol) were added and the reaction mixture was stirred for 10 minutes. DBAD (0.11 g, 0.48 mmol) was then added and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 20% EtOAc/PE; silica gel: 230-400 mesh) to afford enantiomer 1 of title compound.


Enantiomer 2 of the title compound was obtained following the same procedure, starting from 0.3 g of enantiomer 2 of Intermediate 9. After completion of the reaction, the reaction mixture was concentrated under vacuum and the resulting crude was purified by Isolera column chromatography (eluent: 8% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. The absolute configuration of the two enantiomers is not known.

- 30 **Enantiomer 1: Yield:** 80% (80 mg, white solid). **LCMS:** (Method K) 521.0 (M⁺+H), Rt. 3.24 min, 91.11% (Max).

Enantiomer 2: **Yield:** 99% (0.4 g, yellow solid). **LCMS:** (Method K) 521.0 (M^+H), Rt. 3.26 min, 95.22% (Max).

Intermediate 12


- 5 **Ethyl 1-(((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate**



To a stirred solution of 3-butyl-8-hydroxy-2-methyl-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 8; 0.1 g, 0.25 mmol) in THF (5 mL) at 0 °C were added ethyl 1-(hydroxymethyl)cyclopropane-1-carboxylate (0.04 g, 0.25 mmol) and triphenylphosphine (0.09 g, 0.37 mmol) and the reaction mixture was stirred for 10 minutes. DBAD (0.09 g, 0.37 mmol) was then added at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (5 mL) and the aqueous layer extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL), dried over anhydrous Na_2SO_4 and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 30% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 46% (0.12 g, white solid). **LCMS:** (Method K) 533.3 (M^+H), Rt. 3.10 min, 39.60%.

20 Intermediate 13

Ethyl (S)-1-(((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate and ethyl (R)-1-(((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate



25 Enantiomer 1 of the title compound was obtained following the procedure as described for intermediate 12 above, starting from 0.1 g of enantiomer 1 of Intermediate 9. Enantiomer 2 of the


title compound was obtained following the same procedure, but starting from 0.1 g of enantiomer 2 of Intermediate 9. The absolute configuration of the two enantiomers is not known.

Enantiomer 1: **Yield:** 72% (0.12 g, white solid). **LCMS:** (Method K) 533.0 (M^+H), Rt. 3.29 min, 78.91% (Max).

Enantiomer 2: **Yield:** 66% (0.13 g, white solid). **LCMS:** (Method K) 533.0 (M^+H), Rt. 3.29 min, 66.72% (Max).

Intermediate 14

3-Butyl-7-(ethylthio)-8-hydroxy-2-methyl-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide




To a solution of 7-bromo-3-butyl-8-methoxy-2-methyl-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 7; 0.3 g, 0.66 mmol) in N-methyl-2-pyrrolidone (5 mL), sodium ethanethiolate (0.27 g, 3.31 mmol) was added and the reaction mixture was stirred for 12 hours at 100 °C. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with ice-cold water (25 mL) and the aqueous layer was extracted with EtOAc (2 x 50 mL). The combined organic layer was washed with ice-cold water (50 mL) and brine (50 mL) and dried over anhydrous Na_2SO_4 . The organic part was concentrated under vacuum and the resulting crude material was purified by Isolera column chromatography (eluent: 30% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 69% (0.2 g, off-white solid).

1H NMR (400 MHz, $DMSO-d_6$): δ 7.63 (s, 1H), 7.42 (s, 1H), 7.23-7.21 (m, 2H), 6.83 (t, $J = 7.6$ Hz, 1H), 6.68-6.64 (m, 3H), 4.05-4.01 (m, 2H), 3.23-3.21 (m, 1H), 2.80 (q, $J = 7.2$ Hz, 2H), 2.60 (s, 3H), 1.57-1.31 (m, 6H), 1.27 (t, $J = 7.20$ Hz, 3H), 0.96 (t, $J = 7.20$ Hz, 3H). **LCMS:** (Method E) 421.0 (M^+H), Rt. 3.04 min, 95.23% (max).

Intermediate 15


Methyl 3-((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoate



To a stirred solution of 3-butyl-7-(ethylthio)-8-hydroxy-2-methyl-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 14; 0.15 g, 0.36 mmol) in THF (5 mL) at 0 °C, methyl 3-hydroxy-2,2-dimethylpropanoate (0.05 g, 0.36 mmol) and triphenylphosphine (0.14 g, 0.54 mmol) were added and the reaction mixture was stirred for 10 minutes. DBAD (0.12 g, 0.54 mmol) was then added at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 5 mL). The combined organic layer was washed with brine (5 mL), dried over anhydrous Na₂SO₄, and then filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 22% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 32% (0.16 g, white solid). **LCMS:** (Method A) 535.2 (M⁺H), Rt. 3.08 min, 38.01% (Max).

Intermediate 16

Ethyl 1-(((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate




To a stirred solution of 3-butyl-7-(ethylthio)-8-hydroxy-2-methyl-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 14; 0.15 g, 0.36 mmol) in THF (10 mL) at 0 °C were added ethyl 1-(hydroxymethyl)cyclopropane-1-carboxylate (0.05 g, 0.36 mmol) and triphenylphosphine (0.14 g, 0.54 mmol) and the reaction mixture was stirred for 10 minutes. DBAD (0.12 g, 0.54 mmol) was then added at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 5 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified

by Isolera column chromatography (eluent: 22% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 81% (0.16 g, white solid).

LCMS: (Method A) 547.2 (M⁺+H), Rt. 3.06 min, 98.17% (Max).

5 Intermediate 17

3-Butyl-7-chloro-5-(4-fluorophenyl)-8-methoxy-2,3-dihydro-1,5-benzothiazepin-4(5H)-one




To a stirred solution of 3-butyl-7-chloro-8-methoxy-2,3-dihydro-1,5-benzothiazepin-4(5H)-one (6 g, 20.01 mmol) in 1-bromo-4-fluorobenzene (88 g, 500 mmol), copper (I) iodide (0.38 g, 2.00 mmol) and K₂CO₃ (5.53 g, 40.0 mmol) were added and the solution was purged with nitrogen for 20 minutes for degasification. Tris[2-(2-methoxyethoxy)ethyl]amine (1.30 g, 4.00 mmol) was then added under nitrogen atmosphere and the resulting reaction mixture was heated for 40 h at 135 °C. After completion of the reaction (monitored by UPLC), the reaction mixture was filtered through celite and the celite pad was washed with EtOAc (100 mL). The filtrate was concentrated under vacuum and the resulting crude material was purified by Isolera column chromatography (eluent: 15-20% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 64% (5.5 g, off-white solid).

LCMS: (Method E) 394.0 (M⁺+H), Rt. 3.19 min, 91.57%.

Intermediate 18

20 3-Butyl-7-chloro-5-(4-fluorophenyl)-8-methoxy-2,3,4,5-tetrahydro-1,5-benzothiazepine




To a stirred solution of 3-butyl-7-chloro-5-(4-fluorophenyl)-8-methoxy-2,3-dihydro-1,5-benzothiazepin-4(5H)-one (Intermediate 17; 5.5 g, 13.96 mmol) in THF (60 mL) at 0 °C, borane dimethylsulfide (140 mL, 140 mmol) was added dropwise and the reaction mixture was refluxed for 40 hours at 75 °C. After completion of the reaction (monitored by UPLC), the reaction mixture was cooled to 0 °C and quenched with methanol (100 mL). The resulting solution was heated for 2 hours at 65 °C, then cooled to room temperature and concentrated under vacuum. The resulting crude

material was purified by Isolera column chromatography (eluent: 15-20% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 77% (4.5 g, colourless liquid).

LCMS: (Method A) 380.0 (M⁺) Rt. 3.61 min, 90.76% (Max).

5 Intermediate 19


3-Butyl-7-chloro-5-(4-fluorophenyl)-8-methoxy-2,3,4,5-tetrahydro-1,5-benzothiazepine 1,1-dioxide



To a stirred solution of 3-butyl-7-chloro-5-(4-fluorophenyl)-8-methoxy-2,3,4,5-tetrahydro-1,5-benzothiazepine (Intermediate 18; 4.5 g, 11.84 mmol) in THF (45 mL) and water (10 mL), oxone (36.4 g, 59.2 mmol) was added and the reaction mixture was stirred for 24 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was filtered through a Büchner funnel and the filtrate was extracted with EtOAc (2 x 200 mL). The combined organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The crude material was purified by Isolera column chromatography (eluent: 10-13% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 84% (4.1 g, colourless solid). **LCMS:** (Method A) 412.1 (M⁺+H), Rt. 2.83 min, 96.78% (Max).

Intermediate 20

3-Butyl-5-(4-fluorophenyl)-8-hydroxy-7-(methylthio)-2,3,4,5-tetrahydro-1,5-benzothiazepine 1,1-dioxide



To a stirred solution of 3-butyl-7-chloro-5-(4-fluorophenyl)-8-methoxy-2,3,4,5-tetrahydro-1,5-benzothiazepine 1,1-dioxide (Intermediate 19; 1 g, 2.42 mmol) in DMF (12 mL), sodium thiomethoxide (0.85 g, 12.14 mol) was added at room temperature and the resulting mixture was

stirred overnight at 100 °C. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with water (20 mL) and the aqueous layer was extracted with EtOAc (2 x 20 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 30-35% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.


Yield: 30% (0.300 g, off-white solid).

¹H NMR (400 MHz, CDCl₃): δ 7.66 (s, 1H), 7.27 (s, 1H), 6.94-6.94 (m, 2H), 6.62-6.64 (m, 2H), 4.14-4.16 (m, 1H), 3.41-3.42 (m, 1H), 2.96-2.96 (m, 1H), 2.50 (s, 1H), 2.40 (s, 3H), 1.37-1.37 (m, 4H), 1.29-1.30 (m, 4H), 0.95 (t, *J* = 7.20 Hz, 3H). **LCMS:** (Method A) 408.2 (M⁺-H), Rt. 2.54 min, 98.39% (Max).

10

Intermediate 21

Methyl 3-((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)-2,2-dimethylpropanoate



To a stirred solution of 3-butyl-5-(4-fluorophenyl)-8-hydroxy-7-(methylthio)-2,3,4,5-tetrahydro-1,5-benzothiazepine 1,1-dioxide (Intermediate 20; 0.1 g, 0.25 mmol) in THF (5 mL) at 0° C, methyl 3-hydroxy-2,2-dimethylpropanoate (0.03 g, 0.25 mmol) and triphenylphosphine (0.09 g, 0.37 mmol) were added and the reaction mixture was stirred for 10 minutes. DBAD (0.08 g, 0.37 mmol) was then added at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 5 mL). The combined organic layer was washed with brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by Isolera column chromatography (eluent: 22% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.


Yield: 73% (0.12 g, white solid).

LCMS: (Method A) 524.1 (M⁺+H), Rt. 3.33 min, 77.91% (Max).

Intermediate 22

Ethyl 1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate

30




To a stirred solution of 3-butyl-5-(4-fluorophenyl)-8-hydroxy-7-(methylthio)-2,3,4,5-tetrahydro-1,5-benzothiazepine 1,1-dioxide (Intermediate 20; 0.1 g, 0.25 mmol) in THF (5 mL) at 0 °C, ethyl 1-(hydroxymethyl)cyclopropane-1-carboxylate (0.04 g, 0.25 mmol) and triphenylphosphine (0.09 g, 0.37 mmol) were added and the reaction mixture was stirred for 10 minutes. DBAD (0.08 g, 0.37 mmol) was then added at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 5 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 20-22% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 82% (0.11 g, white solid).

LCMS: (Method A) 536.2 (M⁺+H), Rt. 3.35 min, 97.37% (Max).

15 Intermediate 23

tert-Butyl (1-((4-fluorophenyl)amino)-1-oxohexan-2-yl)carbamate




To a stirred solution of 2-((tert-butoxycarbonyl)amino)hexanoic acid (93.0 g, 402 mmol) in DMF (930 mL) at 0 °C, triethylamine (112 mL, 804 mmol) was added dropwise. 2,4,6-Tripropyl-1,3,5,2,4,6-trioxatriphosphinane 2,4,6-trioxide (307 g, 483 mmol) was then added and the reaction mixture was stirred for 10 minutes at 0 °C. 4-Fluoroaniline (44.7 g, 402 mmol) was then added dropwise and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with ice-cold water (500 mL) and a solid product was precipitated. The obtained solid was filtered, washed with water (2 x 100 mL) and dried under vacuum to afford the title compound. The resulting crude was forwarded as such to the next step without any further purification. **Yield:** 69% (90 g, light yellow solid).

LCMS: (Method K) 225.2 (M⁺+H-Boc), Rt. 2.66 min, 99.41% (max).

Intermediate 24

2-Amino-N-(4-fluorophenyl)hexanamide



5

To a stirred solution of *tert*-butyl (1-((4-fluorophenyl)amino)-1-oxohexan-2-yl)carbamate (Intermediate 23; 90 g, 277 mmol) in 1,4-dioxane (900 mL), HCl in dioxane (4M in 1,4-dioxane, 360 mL, 1280 mmol) was added at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was concentrated under vacuum and the obtained residue was basified with 10% NaHCO₃ solution (400 mL). The aqueous layer was extracted with EtOAc (2 x 500 mL) and the combined organic layer was washed with water (2 x 100 mL) and dried over anhydrous Na₂SO₄. The organic part was concentrated under vacuum and the resulting crude was forwarded as such to the next step without any further purification. **Yield:** 91% (60 g, light-brown liquid).


10

15

LCMS: (Method K) 225.1 (M⁺+H), Rt. 1.97 min, 94.23% (max).

Intermediate 25

N1-(4-fluorophenyl)hexane-1,2-diamine




20

To a stirred solution of 2-amino-N-(4-fluorophenyl)hexanamide (Intermediate 24; 60 g, 268 mmol) in THF (300 mL), borane dimethylsulfide (1M in THF, 401 mL, 401 mmol) was added at 0 °C and the reaction mixture was heated for 24 hours at 75 °C. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with methanol (200 mL) at 0 °C and the reaction mixture was heated for 2 hours at 75 °C. The reaction mixture was concentrated under vacuum and the obtained residue was dissolved in DCM (500 mL). The organic layer was washed with water (2 x 200 mL), dried over anhydrous Na₂SO₄ and then concentrated under vacuum. The resulting crude was purified by Isolera column chromatography (eluent: 5-20% MeOH in DCM; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 87% (53 g, brown liquid).

25


LCMS: (Method K) 211.2 (M⁺+H), Rt. 1.97 min, 92.44% (max).

30

Intermediate 26**2,4-Dibromo-N-(1-((4-fluorophenyl)amino)hexan-2-yl)-5-methoxybenzenesulfonamide**

To a stirred solution of *N*1-(4-fluorophenyl)hexane-1,2-diamine (Intermediate 25; 20 g, 95 mmol) in THF (200 mL) were added 2,4-dibromo-5-methoxybenzenesulfonyl chloride (41.6 g, 114 mmol) and triethylamine (39.8 mL, 285 mmol) at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with EtOAc (400 mL). The organic layer was washed with water (2 x 100 mL) and dried over anhydrous Na₂SO₄. The organic part was concentrated under vacuum and the resulting crude was purified by Isolera column chromatography (eluent: 15% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 86% (45.5 g, brown solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 8.02-7.90 (m, 2H), 7.55 (s, 1H), 6.82 (m, 2H), 6.31-6.26 (m, 2H), 5.45-5.30 (m, 1H), 3.90 (s, 3H), 3.30-3.15 (m, 1H), 3.05-2.82 (m, 2H), 1.60-1.40 (m, 1H), 1.40-1.30 (m, 1H), 1.30-1.20 (m, 1H), 1.15-0.98 (m, 3H), 0.73 (t, *J* = 8.80 Hz, 3H). **LCMS:** (Method A) 538.9 (M⁺+H), Rt. 3.04 min, 85.57% (max).

Intermediate 27**7-Bromo-3-butyl-5-(4-fluorophenyl)-8-methoxy-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide**

20


To a stirred solution of 2,4-dibromo-*N*-(1-((4-fluorophenyl)amino)hexan-2-yl)-5-methoxybenzenesulfonamide (Intermediate 26; 45 g, 84 mmol) in DMF (450 mL), potassium carbonate (23.11 g, 167 mmol) and copper iodide (1.59 g, 8.36 mmol) were added at room temperature. The reaction mixture was degassed for 15 minutes with nitrogen. Tris[2-(2-methoxyethoxy)ethyl]amine (5.41 g,

16.72 mmol) was then added at room temperature and the reaction mixture was heated for 16 hours at 130 °C. After completion of the reaction (monitored by TLC), the reaction mixture was filtered through a celite pad and the celite pad was washed with EtOAc (50 mL). The filtrate part was concentrated under vacuum. The obtained residue was diluted with EtOAc (400 mL) and the organic layer was washed with water (2 x 100 mL). The organic part was dried over anhydrous Na₂SO₄ and the resulting crude was purified by Isolera column chromatography (eluent: 12% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 55% (22.0 g, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 7.57 (s, 1H), 7.50-7.42 (d, *J* = 6.6 Hz, 2H), 7.02 (t, *J* = 9.0 Hz, 2H), 6.70-6.50 (m, 2H), 4.38-4.22 (m, 1H), 3.95 (s, 3H), 3.33 (m, 1H), 2.95-2.80 (m, 1H), 1.65-1.15 (m, 6H), 0.89 (t, *J* = 9.2 Hz, 3H). **LCMS:** (Method A) 455.0 (M⁺-H), Rt. 2.71 min, 95.55% (max).


Intermediate 28

7-Bromo-3-butyl-5-(4-fluorophenyl)-8-methoxy-2-methyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide



To a stirred solution of 7-bromo-3-butyl-5-(4-fluorophenyl)-8-methoxy-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 27; 5.8 g, 12.68 mmol) in N-methyl-2-pyrrolidone (24 mL), Cs₂CO₃ (8.26 g, 25.4 mmol) was added at 0 °C and reaction mixture was stirred for 15 minutes. Methyl iodide (1.58 mL, 25.4 mmol) was then added dropwise at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with EtOAc (100 mL) and the organic layer was washed with water (50 mL). The organic part was dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude was purified by Isolera column chromatography (eluent: 15% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 75% (5.1 g, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 7.47 (s, 1H), 7.41 (s, 1H), 6.98-6.94 (m, 2H), 6.72-6.69 (m, 2H), 3.99-3.92 (m, 5H), 3.40-3.22 (m, 1H), 2.63 (s, 3H), 1.68-1.60 (m, 1H), 1.50-1.35 (m, 5H), 0.95 (t, *J* = 7.2 Hz, 3H). **LCMS:** (Method A) 471.0 (M⁺+H), Rt. 2.89 min, 88.26% (max).


Intermediate 29**3-Butyl-5-(4-fluorophenyl)-8-hydroxy-2-methyl-7-(methylthio)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide**

5 To a stirred solution of 7-bromo-3-butyl-5-(4-fluorophenyl)-8-methoxy-2-methyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 28; 2.0 g, 4.24 mmol) in DMF (20 mL), sodium thiomethoxide (1.48 g, 21.21 mmol) was added at room temperature and the reaction mixture was stirred for 16 hours at 100 °C. After completion of the reaction (monitored by TLC), the reaction mixture was cooled to room temperature and quenched with water (20 mL). The aqueous layer was
 10 extracted with EtOAc (2 x 30 mL). The combined organic layer was washed with brine (10 mL) and dried over anhydrous Na₂SO₄. The organic part was concentrated under vacuum and the resulting crude material was purified by Isolera column chromatography (eluent: 15-20% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 81% (1.6 g, light-yellow solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 10.72 (s, 1H), 7.27 (s, 1H), 6.97 (t, *J* = 11.2 Hz, 3H), 6.55-6.50 (m, 2H),
 15 3.98-3.86 (m, 2H), 3.30-3.05 (m, 1H), 2.42 (s, 3H), 2.34 (s, 3H), 1.70-1.45 (m, 2H), 1.40-1.25 (m, 4H), 1.00-0.80 (m, 3H). **LCMS:** (Method A) 425.2 (M⁺+H), Rt. 2.56 min, 90.64% (max).

Intermediate 30

Ethyl 1-(((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate
 20



To a stirred solution of 3-butyl-5-(4-fluorophenyl)-8-hydroxy-2-methyl-7-(methylthio)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 29; 0.06 g, 0.14 mmol) in THF (3 mL) at 0 °C were added ethyl 1-(hydroxymethyl)cyclopropane-1-carboxylate (0.02 g, 0.14 mmol) and


triphenylphosphine (0.056 g, 0.212 mmol) and the reaction mixture was stirred for 10 minutes. DBAD (0.05 g, 0.21 mmol) was then added at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 5 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 22-25% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 69% (70 mg, off-white solid).

LCMS: (Method A) 551.2 (M⁺+H), Rt. 2.96 min, 77.38% (Max).

10

Intermediate 31

Methyl 3-((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoate




15 To a stirred solution of 3-butyl-5-(4-fluorophenyl)-8-hydroxy-2-methyl-7-(methylthio)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 29; 0.05 g, 0.12 mmol) in THF (3 mL) at 0 °C, methyl 3-hydroxy-2,2-dimethylpropanoate (0.02 g, 0.12 mmol) and triphenylphosphine (0.05 g, 0.18 mmol) were added and the reaction mixture was stirred for 30 minutes. DBAD (0.04 g, 0.18 mmol) was then added at 0 °C and the reaction mixture was stirred for 12 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 5 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 30% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 57% (60 mg, brown solid).

25

LCMS: (Method A) 538.8 (M⁺), Rt. 3.03 min, 60.90% (Max).

Intermediate 32

2-((tert-Butoxycarbonyl)amino)hexanoic acid




To a solution of 2-aminohexanoic acid (25 g, 0.19 mol) in water (250 mL) and THF (250 mL), NaHCO₃ (48 g, 0.57 mol) and Boc-anhydride (52.2 mL, 0.23 mol) were added and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was cooled, quenched and acidified using 1.5N HCl. The reaction mixture was extracted with EtOAc (2 x 150 mL). The combined organic layer was washed with ice-cold water (150 mL) and brine (150 mL), dried over anhydrous Na₂SO₄ and concentrated to afford the title compound. **Yield:** 45.5 g (crude, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 5.10-4.98 (m, 1H), 4.33-4.30 (m, 1H), 1.95-1.81 (m, 1H), 1.75-1.55 (m, 1H), 1.46 (s, 10H), 1.45-1.31 (m, 4H), 0.93 (t, *J* = 7.2 Hz, 3H). **LCMS:** (Method E) 132.2 (M⁺-Boc+H), Rt. 2.36 min, 99.98% (max).


Intermediate 33

tert-Butyl (1-oxo-1-(phenylamino)hexan-2-yl)carbamate




To a stirred solution of 2-((*tert*-butoxycarbonyl)amino)hexanoic acid (Intermediate 32; 45.5 g, 0.196 mol) in DMF (150 mL), triethylamine (54.83 mL, 0.39 mol) was added and the reaction mixture was cooled to 0 °C. 1-Propanephosphonic anhydride solution (50% in EtOAc; 75.1 g, 0.23 mol) and aniline (18 g, 0.196 mol) were added and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with ice-cold water (50 mL) and diluted with EtOAc (200 mL). The aqueous layer was washed with ice-cold water (200 mL) and brine (200 mL), and then dried over anhydrous Na₂SO₄ and filtered off. The organic part was concentrated under vacuum to afford the title compound. **Yield:** 82% (49.7 g, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 9.93 (s, 1H), 7.60 (d, *J* = 7.6 Hz, 2H), 7.30 (t, *J* = 8.0 Hz, 2H), 7.06-6.99 (m, 2H), 4.07-4.02 (m, 1H), 1.63-1.56 (m, 2H), 1.46-1.28 (m, 13H), 0.86 (t, *J* = 6.8 Hz, 3H). **LCMS:** (Method E) 207.0 (M⁺-Boc+H), Rt. 2.69 min, 91.20% (max).


Intermediate 34**2-Amino-N-phenylhexanamide**

5 To a solution of *tert*-butyl (1-oxo-1-(phenylamino)hexan-2-yl)carbamate (Intermediate 33; 49.7 g, 0.162 mol) in 1,4-dioxane (500 mL) at 0 °C, a solution of HCl in 1,4-dioxane (4 M, 199 mL) was added and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was concentrated under vacuum and the obtained residue was quenched with saturated NaHCO₃ solution. The aqueous layer was extracted with EtOAc
10 (2 x 200 mL). The combined organic layer was washed brine (200 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum to afford the title compound. **Yield:** 95% (32 g, colourless gum).

¹H NMR (400 MHz, DMSO-*d*₆): δ 10.20-9.66 (m, 1H), 7.64 (dd, *J* = 4.8, 2.4 Hz, 2H), 7.32-7.27 (m, 2H), 7.06-7.02 (m, 1H), 3.29-3.26 (m, 1H), 1.67-1.61 (m, 1H), 1.46-1.27 (m, 5H), 0.89-0.85 (m, 3H). **LCMS:**
15 (Method B) 207.2 (M⁺+H), Rt. 2.03 min, 84.36% (max).

Intermediate 35**N1-Phenylhexane-1,2-diamine**


20 To a solution of 2-amino-N-phenylhexanamide (Intermediate 34; 32 g, 0.15 mol) in THF (320 mL) at 0 °C was added borane dimethylsulfide (2M solution in THF, 117 mL, 0.23 mol) and the reaction mixture was heated for 16 hours at 75 °C. After completion of the reaction (monitored by TLC), the reaction mixture was cooled to 0 °C, quenched with methanol (150 mL) and then heated for 2 hours at 60 °C. The reaction mixture was cooled to room temperature and concentrated under vacuum.
25 The obtained residue was partitioned between water (50 mL) and EtOAc (50 mL). The aqueous layer was extracted with EtOAc (2 x 250 mL). The combined organic layer was washed with water (250 mL) and brine (250 mL). The organic part was dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 10% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 93% (28 g, yellow oil).
30 **LCMS:** (Method B) 193.3 (M⁺+H), Rt. 1.98 min, 76.9% (max).

Intermediate 36**2,4-Dibromo-5-methoxy-N-(1-(phenylamino)hexan-2-yl)benzenesulfonamide**


To a solution of 2,4-dibromo-5-methoxybenzenesulfonyl chloride (3.2 g, 8.79 mmol) in THF (50 mL) at 5 0 °C were added N1-phenylhexane-1,2-diamine (Intermediate 35; 1.3 g, 6.76 mmol) and triethylamine (2.8 mL, 20.3 mmol) and the reaction mixture was stirred for 4 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was concentrated under vacuum and the obtained residue was dissolved in EtOAc (60 mL). The organic layer was washed with water (50 mL) and brine (50 mL), dried over anhydrous Na₂SO₄ and 10 concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 18% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 97% (3.5 g, brown gum). **LCMS:** (Method E) 521.0 (M⁺+H), Rt. 3.14 min, 93.57% (max).

Intermediate 37

15 **7-Bromo-3-butyl-8-methoxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide**




To a solution of 2,4-dibromo-5-methoxy-N-(1-(phenylamino)hexan-2-yl)benzenesulfonamide (Intermediate 36; 3.5 g, 6.72 mmol) in DMF (30 mL), K₂CO₃ (1.73 g, 12.5 mmol) and copper powder (0.42 g, 6.72 mmol) were added. The reaction mixture was degassed for 5 minutes under N₂ 20 atmosphere and the reaction mixture was then heated for 16 hours at 115 °C. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with ice-cold water (25 mL) and the aqueous layer was extracted with a 1:1 mixture of EtOAc and PE (2 x 50 mL). The combined organic layer was washed with brine (30 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 25% 25 EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 51% (1.5 g, brown gum). **LCMS:** (Method A) 439.0 (M⁺+H), Rt. 2.83 min, 82.06% (max).

Intermediate 38**7-bromo-3-butyl-8-methoxy-2-(4-methoxybenzyl)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide**

5

To a stirred solution of 7-bromo-3-butyl-8-methoxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 37; 2 g, 4.55 mmol) in NMP (15 mL), Cs₂CO₃ (1.78 g, 5.46 mmol) was added at 0 °C and the reaction mixture was stirred for 15 minutes. Then 1-(bromomethyl)-4-methoxybenzene (1.37 g, 6.83 mmol) was added dropwise at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with EtOAc (50 mL) and the organic layer was washed with water (2 x 10 mL). The organic part was dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude was purified by Isolera column chromatography (eluent: 20-22% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 46% (2.1 g, white solid).

15 **LCMS:** (Method K) 559.1 (M⁺), Rt. 2.99 min, 56.12% (Max).

Intermediate 39**3-butyl-7-(ethylthio)-8-hydroxy-2-(4-methoxybenzyl)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide**

20


To a stirred solution of 7-bromo-3-butyl-8-methoxy-2-(4-methoxybenzyl)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 38; 1.0 g, 1.79 mmol) in NMP (10 mL), sodium ethanethiolate (0.75 g, 8.94 mmol) was added at room temperature and the reaction mixture was heated for 12 hours at 100 °C. After completion of the reaction (monitored by TLC), the reaction mixture was cooled to room temperature and quenched with water (20 mL). The aqueous layer was extracted with EtOAc (2 x 30 mL) and the combined organic layer was washed with brine (10 mL) and then dried over anhydrous Na₂SO₄. The organic part was concentrated under vacuum and the resulting crude material was purified by Isolera column chromatography (eluent: 20-22% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 29% (0.3 g, white solid).

5

10 **LCMS:** (Method A) 527.1 (M⁺+H), Rt. 2.89 min, 91.85% (Max).

Intermediate 40

3-Butyl-7-(ethylthio)-8-hydroxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide



To a stirred solution of 3-butyl-7-(ethylthio)-8-hydroxy-2-(4-methoxybenzyl)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 39; 0.4 g, 0.76 mmol) in toluene (5 mL) at 0°C were added triphenylamine (0.09 g, 0.38 mmol) and 2,2,2-trifluoroacetic acid (1.73 g, 15.19 mmol). The reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with water (10 mL) and the aqueous layer was extracted with EtOAc (2 x 20 mL). The combined organic layer was washed with brine (10 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 25% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 30% (0.2 g, white solid).


15

20 **LCMS:** (Method B) 407.1 (M⁺+H), Rt. 2.58 min, 47.18% (Max).

25

Intermediate 41

Ethyl 1-(((3-butyl-7-(ethylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate




To a stirred solution of 3-butyl-7-(ethylthio)-8-hydroxy-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 40; 0.05 g, 0.12 mmol) in THF (5 mL) at 0 °C, ethyl 1-(hydroxymethyl)cyclopropane-1-carboxylate (0.03 g, 0.18 mmol) and triphenylphosphine (0.05 g, 0.18 mmol) were added and the reaction mixture was stirred for 10 minutes. DBAD (0.04 g, 0.19 mmol) was then added at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 5 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 20-22% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 76% (50 mg, white solid).

LCMS: (Method A) 532.7 (M⁺), Rt. 3.00 min, 98.06% (Max).

15 Intermediate 42

7-Bromo-3-butyl-5-(4-fluorophenyl)-8-methoxy-2-(4-methoxybenzyl)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide



To a stirred solution of 7-bromo-3-butyl-5-(4-fluorophenyl)-8-methoxy-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 27; 5 g, 10.93 mmol) in NMP (50 mL), Cs₂CO₃ (7.12 g, 21.86 mmol) was added and the reaction mixture was stirred for 30 minutes at 0 °C. Then 1-(bromomethyl)-4-methoxybenzene (3.30 g, 16.40 mmol) was added and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the


reaction mixture was filtered through a celite bed and the celite bed was washed with EtOAc (2 x 20 mL). The filtrate was concentrated under vacuum and the crude was purified by Isolera column chromatography (eluent: 15-20% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.

Yield: 73% (6 g, off-white solid).

5 **LCMS:** (Method A) 578.1 (M⁺+H), Rt. 3.49 min, 75.44%.

Intermediate 43

3-Butyl-5-(4-fluorophenyl)-8-hydroxy-2-(4-methoxybenzyl)-7-(methylthio)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide



10

To a stirred solution of 7-bromo-3-butyl-5-(4-fluorophenyl)-8-methoxy-2-(4-methoxybenzyl)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 42; 1 g, 1.23 mmol) in DMF (20 mL), sodium thiomethoxide (0.43 g, 6.14 mmol) was added at room temperature and the resulting mixture was stirred for 16 hours at 100 °C. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with water (50 mL) and the aqueous layer was extracted with EtOAc (2 x 50 mL). The combined organic layer was washed with brine (50 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 10-15% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.


15

Yield: 66% (500 g, off-white solid).

20 **LCMS:** (Method B) 529.1 (M⁺-H), Rt. 3.20 min, 85.70% (Max).

Intermediate 44


3-Butyl-5-(4-fluorophenyl)-8-hydroxy-7-(methylthio)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide



To a stirred solution of 3-butyl-5-(4-fluorophenyl)-8-hydroxy-2-(4-methoxybenzyl)-7-(methylthio)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 43; 0.2 g, 0.37 mmol) in toluene (10 ml), triphenylamine (0.09 g, 0.37 mmol) and 2,2,2-trifluoroacetic acid (0.04 g, 0.37 mmol) was added at 0 °C and the reaction mixture was stirred for 4 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with water (10 mL) and the aqueous layer was extracted with EtOAc (2 x 20 mL). The combined organic layer was washed with brine (10 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 20-25% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 90% (0.15 g, white solid). **LCMS:** (Method A) 411.0 (M⁺+H), Rt. 2.26 min, 93.21% (Max).

Intermediate 45

Ethyl 1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate



To a stirred solution of 3-butyl-5-(4-fluorophenyl)-8-hydroxy-7-(methylthio)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 44; 0.05 g, 0.12 mmol) in THF (5 mL) at 0 °C, ethyl 1-(hydroxymethyl)cyclopropane-1-carboxylate (0.02 g, 0.12 mmol) and triphenylphosphine (0.05 g, 0.18 mmol) were added and the reaction mixture was stirred for 10 minutes. DBAD (0.04 g, 0.18 mmol) was then added at 0 °C and the reaction mixture was stirred for 1 hour at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 5 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part


was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 20-22% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 51% (0.1 g, white solid).

LCMS: (Method B) 536.2 (M⁺), Rt. 2.79 min, 96.56% (Max).

5

Intermediate 46

3-Butyl-7-(ethylthio)-5-(4-fluorophenyl)-8-hydroxy-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide



10 To a stirred solution of 7-bromo-3-butyl-5-(4-fluorophenyl)-8-methoxy-2-(4-methoxybenzyl)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 42; 0.3 g, 0.52 mmol) in DMF (5 mL), sodium ethanethiolate (0.22 g, 2.60 mmol) was added at room temperature and the reaction mixture was stirred for 12 hours at 100 °C. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with water (20 mL) and the aqueous layer was extracted with EtOAc (2 x 30

15 mL). The combined organic layer was washed with brine (10 mL), dried over anhydrous Na₂SO₄ and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 20% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.


Yield: 97% (0.23 g, white solid).

LCMS: (Method A) 425.1 (M⁺+H), Rt. 2.97 min, 92.88% (Max).

20

Intermediate 47

Ethyl 1-(((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate




To a stirred solution of 3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-8-hydroxy-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 46; 0.1 g, 0.24 mmol) in THF (5 mL) at 0 °C, ethyl 1-(hydroxymethyl)cyclopropane-1-carboxylate (0.102 g, 0.71 mmol) and triphenylphosphine (0.19 g, 0.71 mmol) were added and the reaction mixture was stirred for 10 minutes. DBAD (0.16 g, 0.71 mmol) was then added at 0 °C and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 5 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 22-25% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 49% (70 mg, brown solid).

LCMS: (Method A) 551.1 (M⁺+H), Rt. 2.93 min, 90.19% (Max).

Intermediate 48


Ethyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-difluoropropanoate



To a stirred solution of 3-butyl-8-hydroxy-2-methyl-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 8; 0.1 g, 0.25 mmol) in DMF (3 mL) at 0 °C, Cs₂CO₃ (0.12 g, 0.37 mmol) was added and the reaction mixture was stirred for 30 minutes at room temperature. Then ethyl 3-bromo-2,2-difluoropropanoate (0.07 g, 0.30 mmol) was added and the reaction mixture was stirred for 16 hours at room temperature. After TLC revealed that the starting material had not been fully consumed, the reaction mixture was heated for 2 hours at 60 °C. The reaction mixture was then diluted with water (5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 8-10% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 33% (0.1 g, brown solid).

UPLC: (Method B) 543.2 (M⁺+H), Rt. 2.55 min, 43.90% (Max).


30

Intermediate 49**Methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxy-2-methylpropanoate**

5 To a stirred solution of 3-butyl-8-hydroxy-2-methyl-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 8; 0.5 g, 1.22 mmol) in DMF (5 mL), Cs₂CO₃ (0.78 g, 2.4 mmol) and methyl 2-methyloxirane-2-carboxylate (0.42 g, 3.6 mmol) were added at 0 °C. The reaction mixture was stirred for 72 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with dilute HCl (1.5 N, 5 mL). The aqueous layer was extracted with ethyl acetate (2 x 20 mL) and the combined organic layer was washed with water (10 mL) and brine (10 mL). The organic part was dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 20% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.

Yield: 27% (0.17 g, brown solid).

15 ¹H NMR (400 MHz, CDCl₃): δ 7.36 (s, 1H), 7.29-7.25 (m, 2H), 6.97 (s, 1H), 6.83-6.80 (m, 1H), 6.68-6.65 (m, 2H), 4.37-4.36 (m, 1H), 4.15 (m, 1H), 4.04 (s, 1H), 4.00 (s, 1H), 3.88 (s, 3H), 3.65 (bs, 1H), 3.15 (bs, 1H), 2.59 (s, 3H), 2.33 (s, 3H), 1.66 (m, 2H), 1.6 (m, 6H), 1.3 (m, 1H), 0.98-0.96 (m, 3H). **LCMS:** (Method K) 523.1 (M⁺+H), Rt. 2.97 min, 94.53% (Max).

20 Intermediate 50**Methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxy-2-methylpropanoate**


The title compound was obtained following the same procedure as described for Intermediate 49 above, starting from 0.15 g of enantiomer 2 of 3-butyl-8-hydroxy-2-methyl-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 9). The crude material was purified by Isolera column chromatography (eluent: 10-15% EtOAc/PE; silica gel: 230-400 mesh) to

afford the title compound. The absolute configuration of the compound is not known. **Yield:** 47% (90 mg, brown solid).

¹H NMR (400 MHz, CDCl₃): δ 7.37 (s, 1H), 7.29 (m, 2H), 6.98 (s, 1H), 6.84 (t, *J* = 4.4 Hz, 1H), 6.68-6.66 (m, 2H), 4.37-4.36 (m, 1H), 4.13 (m, 1H), 4.04 (m, 2H), 3.88 (s, 3H), 3.30 (bs, 1H), 2.98 (s, 2H), 2.59 (s, 3H), 2.33 (s, 3H), 1.60-1.57 (m, 3H), 1.56-1.49 (m, 4H), 0.98-0.96 (m, 3H). **LCMS:** (Method K) 522.9 (M⁺+H), Rt. 2.78 min, 92.89% (Max).

Intermediate 51

Methyl (S)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxy-2-methylpropanoate and methyl (R)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxy-2-methylpropanoate (individual diastereomers)



The two diastereoisomers of Intermediate 50 (0.3 g, 0.58 mmol) were separated by chiral SFC Instrument (method N). The material was concentrated under vacuum at 40 °C. The first eluting fraction corresponded to diastereomer 1 and the second eluting fraction corresponded to diastereomer 2. Each of the two fractions was then individually treated for further purification. The obtained residue was diluted with ethyl acetate (30 ml) and washed with dilute HCl (1.5 N, 10 mL) and water (10 mL). The organic part was then dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum to afford a purified diastereomer of the title compound. The absolute configuration of the two diastereomers is not known.


Diastereoisomer 1: Yield: 43% (0.13 g, off-white gum). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 7.30 (s, 1H), 7.16 (t, *J* = 8.4 Hz, 2H), 7.05 (s, 1H), 6.70 (t, *J* = 7.2 Hz, 1H), 6.57 (d, *J* = 8.0 Hz, 2H), 5.74 (s, 1H), 4.21 (m, 2H), 3.88 (bs, 1H), 3.69 (s, 3H), 3.20 (bs, 1H), 2.46 (s, 3H), 2.34 (s, 3H), 1.64 (m, 1H), 1.50 (m, 1H), 1.49 (s, 3H), 1.36 (m, 4H), 0.93-0.92 (m, 3H). **LCMS:** (Method H) 523.0 (M⁺+H), Rt. 2.50 min, 96.90% (Max). **HPLC:** (Method E) Rt. 5.62 min, 98.27% (Max). **Chiral SFC:** (method L) Rt. 3.73 min, 98.36% (Max).

Diastereoisomer 2: Yield: 41% (0.13 g, off-white gum). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 7.30 (s, 1H), 7.16 (t, *J* = 8.4 Hz, 2H), 7.04 (s, 1H), 6.69 (t, *J* = 7.2 Hz, 1H), 6.57 (d, *J* = 8.0 Hz, 2H), 5.73 (s, 1H), 4.20

(m, 2H), 4.04 (bs, 2H), 3.90 (bs, 1H), 3.69 (s, 4H), 1.99 (m, 1H), 1.44 (m, 4H), 1.35-1.30 (m, 7H), 0.91 (m, 4H). **LCMS:** (Method H) 523.0 (M⁺+H), Rt. 2.50 min, 93.08% (Max). **HPLC:** (Method E) Rt. 5.63 min, 94.12% (Max). **Chiral SFC:** (method L) Rt. 4.68 min, 98.75% (Max).

5 Intermediate 52

Methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoate




To a stirred solution of methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxy-2-methylpropanoate (Intermediate 49; 0.17 g, 0.32 mmol) in DMF (5 mL) at 0 °C, sodium hydride (6.5 mg, 0.16 mmol) was added and the reaction mixture was stirred for 10 minutes. Then methyl iodide (0.23 g, 1.62 mmol) was added dropwise and the reaction mixture was stirred for 3 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with dilute HCl (1.5 N, 5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 20 mL). The combined organic layer was washed with water (10 mL) and brine (10 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 10-12% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.

Yield: 75% (0.13 g, brown solid).

LCMS: (Method K) 537.2 (M⁺+H), Rt. 2.98 min, 96.48% (Max).

Intermediate 53

Methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoate




25

The title compound was obtained following the same procedure as described for Intermediate 52, starting from 0.08 g of Intermediate 50. After work-up of the reaction mixture, the crude material was purified by Isolera column chromatography (eluent: 10-15% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. The absolute configuration of the compound is not known. **Yield:** 57% (50 mg, off-white solid).

LCMS: (Method K) 537.0 (M⁺+H), Rt. 3.12 min, 93.30% (Max).

Intermediate 54

Methyl (S)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoate and methyl (R)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoate (individual diastereomers)




Diastereoisomers 1 and 2 of the title compound were prepared from diastereoisomer 1 (0.13 g) and diastereoisomer 2 (0.13 g) of Intermediate 51, respectively, following the same procedure as described for Intermediate 52. After work-up of the reaction mixtures, the crude materials were purified by Isolera column chromatography (eluent: 22% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compounds. The absolute configuration of the two diastereomers is not known.

Diastereoisomer 1: Yield: 69% (95 mg, pale yellow solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 7.32 (s, 1H), 7.16 (t, *J* = 8.0 Hz, 2H), 7.05 (s, 1H), 6.71 (t, *J* = 7.2 Hz, 1H), 6.58 (d, *J* = 8.0 Hz, 2H), 4.36 (m, 2H), 4.04-4.03 (m, 1H), 3.90 (bs, 1H), 3.73 (s, 3H), 3.30 (s, 3H), 2.46 (s, 3H), 2.35 (s, 4H), 1.61 (m, 1H), 1.49 (s, 4H), 1.36 (m, 4H), 0.94-0.92 (m, 3H). **LCMS:** (Method A) 537.1 (M⁺+H), Rt. 2.89 min, 96.63% (Max).

Diastereoisomer 2: Yield: 64% (90 mg, pale yellow solid). **LCMS:** (Method A) 537.1 (M⁺+H), Rt. 2.89 min, 95.04% (Max).

Intermediate 55

Methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate



To a solution of 3-butyl-8-hydroxy-2-methyl-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 8; 0.5 g, 1.22 mmol) in DMF (10 mL), Cs₂CO₃ (0.79 g, 2.45 mmol) was added and the reaction mixture was stirred for 15 minutes at room temperature.

5 Methyl oxirane-2-carboxylate (0.37 g, 3.68 mmol) was then added and the reaction mixture was stirred for 72 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with dilute HCl (1.5 N, 15 mL) and diluted with water (10 mL). The aqueous layer was extracted with EtOAc (2 X 15 mL) and the combined organic layer was washed with water (15 mL) and brine (15 mL). The organic part was dried over anhydrous Na₂SO₄, filtered and concentrated under vacuum. The resulting crude material was purified by Isolera column chromatography (eluent: 10% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.


10 **Yield:** 29% (0.18 g, colourless gum).

¹H NMR (400 MHz, DMSO-*d*₆): δ 7.33 (s, 1H), 7.17-7.05 (m, 3H), 6.72-6.56 (m, 2H), 5.91-5.89 (m, 1H), 4.51-4.49 (m, 1H), 4.32-4.27 (m, 1H), 4.04-3.88 (m, 3H), 3.70 (s, 2H), 3.32 (m, 3H) 2.57-2.19 (m,

15 6H), 1.42-1.39 (m, 6H), 0.7-0.81 (m, 3H). **LCMS:** (Method B) 509.0 (M⁺+H), Rt. 2.57 min, 99.63% (Max).

Intermediate 56

Methyl 3-(((S)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate and methyl 3-(((R)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate



Stereoisomers 1 and 2 of the title compound were prepared from enantiomer 1 (0.3 g) and enantiomer 2 (0.3 g) of Intermediate 9, respectively, following the same procedure as described for Intermediate 55. The absolute configuration of the two stereoisomers is not known.

25


Stereoisomer 1: **Yield:** 47.1% (0.18 g, brown solid). **¹H-NMR** (400 MHz, CDCl₃): δ 7.39 (s, 1H), 7.29-7.20 (m, 2H), 6.98 (s, 1H), 6.83-6.78 (m, 1H), 6.67-6.65 (m, 2H), 4.60-4.59 (m, 1H), 4.44-4.42 (m, 2H), 4.04-4.00 (m, 2H), 3.90-3.85 (m, 3H), 3.26-3.24 (m, 1H), 2.99-2.90 (m, 1H), 2.58 (s, 3H), 2.33 (s, 3H), 1.49-1.24 (m, 6H), 0.98-0.94 (m, 3H). **LCMS:** (Method A) 509.1 (M⁺+H) Rt. 2.92 min, 98.11% (Max).

5 Stereoisomer 2: **Yield:** 44.8% (0.17 g, off white solid). **¹H-NMR** (400 MHz, DMSO-*d*₆): δ 7.39 (s, 1H), 7.29-7.21 (m, 2H), 6.98 (s, 1H), 6.85-6.83 (m, 1H), 6.68-6.67 (m, 2H), 4.60-4.58 (m, 1H), 4.46-4.42 (m, 2H), 4.04-4.00 (m, 2H), 3.90-3.85 (m, 3H), 3.26-3.24 (m, 1H), 2.98-2.91 (m, 1H), 2.58 (s, 3H), 2.33 (s, 3H), 1.49-1.24 (m, 6H), 0.98-0.94 (m, 3H). **LCMS:** (Method E) 508.9 (M⁺+H) Rt. 2.72 min, 98.85% (Max).

10

Intermediate 57

Methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoate



15 To a stirred solution of methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate (Intermediate 55; 180 mg, 0.35 mmol) in DMF (2 mL) at 0 °C, a suspension of sodium hydride (7.06 mg, 0.17 mmol) in DMF (2 mL) at 0 °C was added dropwise and the reaction mixture was stirred for 10 minutes at room temperature. Methyl iodide (109 mL, 1.76 mmol) in DMF (1 mL) was then added dropwise at 0 °C and the reaction

20 mixtures was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with dilute HCl (1.5 N, 1 mL), diluted with water (5 mL) and the aqueous layer was extracted with EtOAc (2 x 10 mL). The combined organic layer was washed with water (10 mL), brine (10 mL), and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by Isolera column


25 chromatography (eluent: 20-30% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 31.4% (55 mg, brown solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 7.34 (s, 1H), 7.15 (t, *J* = 7.6 Hz, 2H), 7.13 (s, 1H), 6.69 (t, *J* = 7.2 Hz, 1H), 6.58 (d, *J* = 8.4 Hz, 2H), 4.44-4.42 (m, 1H), 4.37-4.35 (m, 2H), 4.00-3.95 (m, 1H), 3.73-3.73 (m, 1H), 3.43 (s, 3H), 3.41 (s, 3H), 3.20-3.19 (m, 1H), 2.37 (s, 3H), 2.28 (s, 3H), 1.36-1.34 (m, 2H), 1.27-1.24

30 (m, 4H), 0.9 (t, *J* = 6.8 Hz, 3H). **LCMS:** (Method E) 522.8 (M⁺+H), Rt. 2.87 min, 95.53% (Max).

Intermediate 58

Methyl 3-(((S)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoate and methyl 3-(((R)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoate




Stereoisomers 1 and 2 of the title compound were prepared from stereoisomer 1 (0.25 g) and stereoisomer 2 (0.24 g) of Intermediate 56, respectively, following the same procedure as described for Intermediate 57. The absolute configuration of the two stereoisomers is not known.

Stereoisomer 1: **Yield:** 54.9% (0.15 g, white solid). **¹H-NMR** (400 MHz, CDCl₃): δ 7.39 (s, 1H), 7.22-7.20 (m, 2H), 6.98 (s, 1H), 6.83-6.78 (m, 1H), 6.67-6.65 (m, 2H), 4.44-4.38 (m, 2H), 4.27-4.25 (m, 1H), 4.09-4.04 (m, 2H), 3.87-3.80 (m, 5H), 3.61 (s, 3H), 3.23 (s, 1H), 2.57 (s, 3H), 2.33 (s, 3H), 1.75-1.70 (m, 1H), 1.58 (s, 3H), 1.50-1.28 (m, 3H). **LCMS:** (Method B) 523.1(M⁺H) Rt. 2.71 min, 94.97% (Max).

Stereoisomer 2: **Yield:** 38.6% (0.1 g, white solid). **LCMS:** (Method B) 523.0(M⁺H) Rt. 2.71 min, 95.14% (Max).

Intermediate 59

Methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-ethoxypropanoate



To a stirred solution of methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate (Intermediate 55; 140 mg, 0.28 mmol) in DMF (3 mL) at 0 °C, sodium hydride (5.5 mg, 0.14 mmol) was added and the reaction mixture was stirred for 5 minutes. Ethyl iodide (0.11 mL, 1.38 mmol) was then added dropwise and


the reaction mixture was stirred for 1 hour at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with dilute HCl (1.5 N, 1 mL) and the aqueous layer was extracted with EtOAc (2 x 10 mL). The combined organic layer was washed with water (5 mL), brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by Isolera column chromatography (eluent: 30-45% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.

Yield: 20% (0.03 g, off-white gum.)

LCMS: (Method E) 537.0 (M⁺+H) Rt. 3.12 min, 73.10% (Max).


10 Intermediate 60

Methyl 3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)-2-hydroxypropanoate



To a stirred solution of 3-butyl-8-hydroxy-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepine 1,1-dioxide (0.2 g, 0.51 mmol) in DMF (5 mL) at 0 °C were added methyl oxirane-2-carboxylate (0.16 g, 1.53 mmol) and Cs₂CO₃ (0.33 g, 1.02 mmol), and the reaction mixture was stirred for 16 hours at room temperature. As TLC showed incomplete conversion, more methyl oxirane-2-carboxylate (0.16 g, 1.53 mmol) was added and the reaction mixture was stirred for 16 hours at room temperature. Even though TLC showed that not all starting material had been consumed, the reaction mixture was quenched with dilute HCl (1.5 N, 5 mL) and the aqueous layer was extracted with EtOAc (2 x 15 mL). The combined organic layer was washed with water (5 mL), brine (10 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by Isolera column chromatography (eluent: 40% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 67% (0.17 g, brown gum).


¹H NMR (400 MHz, DMSO-*d*₆): δ 7.42 (s, 1H), 7.28 (t, *J* = 6.04 Hz, 2H), 7.17 (s, 1H), 6.68 (t, *J* = 7.32 Hz, 1H), 6.55 (d, *J* = 8.04 Hz, 2H), 5.91 (m, 1H), 4.52-4.49 (m, 1H), 4.37-4.31 (m, 2H), 4.29-4.26 (m, 1H), 3.72 (s, 3H), 3.69-3.68 (m, 1H), 3.34-3.32 (m, 1H), 3.18-3.01 (m, 1H), 2.34 (s, 3H), 1.39-1.18 (m, 7H), 0.92-0.89 (m, 3H). **LCMS:** (Method E) 494.1 (M⁺+H) Rt. 2.44 min, 95.67% (Max).

Intermediate 61**3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-8-hydroxy-2-methyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide**

5 To a stirred solution of 7-bromo-3-butyl-5-(4-fluorophenyl)-8-methoxy-2-methyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 28; 0.500 g, 1.06 mmol) in NMP (4 mL), sodium ethanethiolate (0.45 g, 5.30 mmol) was added and the reaction mixture was heated for 16 h at 100 °C. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (20 mL) and the aqueous layer was extracted with EtOAc (2 x 20 mL). The combined organic layer was
 10 washed with water (5 mL), brine (5 mL), and the organic part was dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude was purified by Isolera column chromatography (eluent: 30-50% EtOAc/PE; silica gel: 230-400 mesh) afford the title compound. **Yield:** 86% (0.4 g, yellowish solid).

LCMS: (Method A) 439.1(M⁺+H) Rt. 2.69 min, 54.66% (Max).

15

Intermediate 62**Methyl 3-((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-2-methyl-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate**

20 To a stirred solution of 3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-8-hydroxy-2-methyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine 1,1-dioxide (Intermediate 61; 100 mg, 0.23 mmol) in DMF (1 mL), Cs₂CO₃ (153 mg, 0.47 mmol) was added and the reaction mixture was stirred for 10 minutes at room temperature. Methyl oxirane-2-carboxylate (68.4 mg, 0.69 mmol) was then added dropwise and the reaction mixture was stirred for 24 hours at room temperature. After completion of the


reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 2 mL) and the aqueous layer was extracted with EtOAc (2 x 10 mL). The combined organic layer was washed with water (5 mL), brine (5 mL), and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by Isolera column chromatography (eluent: 0-20% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.

Yield: 79.6% (55 mg, white solid).

LCMS: (Method B) 541.2 (M⁺+H) Rt. 2.65 min, 79.63% (Max).

Example 1

10 3-((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid




To a stirred solution of methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoate (Intermediate 10; 0.15 g, 0.28 mmol) in a mixture of 1,4-dioxane and water (3:2; 5 mL) at 0 °C, lithium hydroxide (24 mg, 0.50 mmol) was added and the reaction mixture was stirred for 3 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude was purified by prep HPLC purification (Method A) to afford the title compound. **Yield:** 7% (10 mg, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 12.46 (s, 1H), 7.28 (s, 1H), 7.15 (t, *J* = 8.8 Hz, 2H), 7.05 (s, 1H), 6.69 (t, *J* = 7.2 Hz, 1H), 6.57 (d, *J* = 8.4 Hz, 2H), 4.12 (s, 2H), 4.08-3.92 (m, 1H), 3.92-3.82 (m, 1H), 3.25-3.12 (m, 1H), 2.46 (s, 3H), 2.34 (s, 3H), 1.70-1.60 (m, 1H), 1.58-1.42 (m, 1H), 1.42-1.32 (m, 4H), 1.28-1.25 (m, 6H), 0.9 (t, *J* = 7.2 Hz, 3H). **LCMS:** (Method K) 506.9 (M⁺+H), Rt. 2.85 min, 95.20% (Max). **HPLC:** (Method E) Rt. 5.85 min, 97.92% (Max).

Example 2

30 (S)-3-((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid and (R)-3-((3-butyl-2-methyl-7-

(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid



To a stirred solution of enantiomer 1 of methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoate (Intermediate 11; 80 mg, 0.06 mmol) in a mixture of 1,4-dioxane and water (3:2; 5 mL) at 0 °C, lithium hydroxide (15 mg, 0.11 mmol) was added and the reaction mixture was stirred for 3 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 20 mL). The combined organic layer was washed with water (10 mL) and brine (10 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude was purified by prep HPLC purification (Method A) to afford enantiomer 1 of the title compound.

Enantiomer 2 of the title compound was obtained following the same procedure, starting from 0.4 g of enantiomer 2 of Intermediate 11. The resulting crude was purified by prep-HPLC purification (Method B) to afford of the title compound. The absolute configuration of the two enantiomers is not known.


Enantiomer 1: **Yield:** 75% (15 mg, white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 12.45 (s, 1H), 7.28 (s, 1H), 7.16 (t, *J* = 7.6 Hz, 2H), 7.05 (s, 1H), 6.70 (t, *J* = 7.2 Hz, 1H), 6.58 (d, *J* = 8.4 Hz, 2H), 4.12 (s, 2H), 4.1-3.98 (m, 1H), 3.98-3.8 (m, 1H), 3.3-3.1 (m, 1H), 2.46 (s, 3H), 2.34 (s, 3H), 1.65-1.64 (m, 1H), 1.51-1.49 (m, 1H), 1.28 (m, 10H), 0.94 - 0.90 (m, 3H). **LCMS:** (Method K) 507.0 (M⁺+H), Rt. 3.04 min, 98.00% (Max). **HPLC:** (Method E) Rt. 6.02 min, 99.04% (Max). **Chiral Purity:** (Method D) Rt. 1.90 min, 100.00% (Max).

Enantiomer 2: **Yield:** 13% (50 mg, off-white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 12.46 (s, 1H), 7.28 (s, 1H), 7.16 (t, *J* = 7.6 Hz, 2H), 7.05 (s, 1H), 6.70 (t, *J* = 7.6 Hz, 1H), 6.58 (d, *J* = 8.0 Hz, 2H), 4.12 (s, 2H), 4.1-3.98 (m, 1H), 3.98-3.8 (m, 1H), 3.25-3.1 (bs, 1H), 2.46 (s, 3H), 2.34 (s, 3H), 1.7-1.45 (m, 2H), 1.4-1.2 (m, 10H), 0.94-0.92 (m, 3H). **LCMS:** (Method H) 507.0 (M⁺+H), Rt. 2.82 min, 96.15% (Max). **HPLC:** (Method D) Rt. 5.94 min, 95.57% (Max). **Chiral Purity:** (Method D) Rt. 2.61 min, 97.52% (Max).

30

Example 3


1-(((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid



5 To a stirred solution of ethyl 1-(((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate (Intermediate 12; 0.12 g, 0.23 mmol) in a mixture of 1,4-dioxane and water (3:2, 5 mL) at 0 °C, lithium hydroxide (10.79 mg, 0.45 mmol) was added and the reaction mixture was stirred for 2 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 3 mL) and the aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by prep-HPLC purification (Method A) to afford the title compound. **Yield:** 22% (25 mg, white solid). ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.40 (s, 1H), 7.28 (s, 1H), 7.16 (t, *J* = 8.4 Hz, 2H), 7.05 (s, 1H), 6.70 (t, *J* = 7.6 Hz, 1H), 6.58 (d, *J* = 8.0 Hz, 2H), 4.27 (dd, *J* = 3.6, 10.0 Hz, 2H), 4.04-3.99 (m, 1H), 3.90 (bs, 1H), 3.15 (bs, 1H), 2.40 (s, 3H), 2.29 (s, 3H), 1.65-1.62 (m, 1H), 1.51-1.50 (m, 1H), 1.47-1.45 (m, 4H), 1.24-1.23 (m, 2H), 1.15-1.12 (m, 2H), 0.94-0.92 (m, 3H). **LCMS:** (Method K) 505.3 (M⁺+H), Rt. 2.76 min, 99.10% (Max). **HPLC:** (Method E) Rt. 5.70 min, 98.59% (Max).

20 Example 4

(S)-1-(((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid and (R)-1-(((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid



25

To a stirred solution of enantiomer 1 of ethyl 1-(((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate

(Intermediate 13; 0.12 g, 0.23 mmol) in a mixture of 1,4-dioxane and water (3:2; 5 mL), lithium hydroxide (10.79 mg, 0.45 mmol) was added at 0 °C. The reaction mixture was stirred for 2 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 3 mL). The aqueous layer was extracted with ethyl acetate (2 x 10 mL) and the combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by prep HPLC purification (Method A) to afford the title compound.


Enantiomer 2 of the title compound was obtained following the same procedure, starting from 0.13 g of enantiomer 2 of Intermediate 13. The absolute configuration of the two enantiomers is not known.

Enantiomer 1: **Yield:** 26% (30 mg, white solid). **¹H NMR** (400 MHz, DMSO-*d*₆) δ 12.47 (s, 1H), 7.28 (s, 1H), 7.16 (t, *J* = 8.4 Hz, 2H), 7.05 (s, 1H), 6.70 (t, *J* = 7.2 Hz, 1H), 6.57 (d, *J* = 8.0 Hz, 2H), 4.28 - 4.26 (m, 2H), 4.02 (m, 1H), 3.89 (bs, 1H), 3.47 (bs, 1H), 2.46 (s, 3H), 2.34 (s, 3H), 1.63-1.62 (m, 1H), 1.51-1.50 (m, 1H), 1.36-1.35 (m, 4H), 1.23-1.22 (m, 2H), 1.10-1.09 (m, 2H), 0.94-0.92 (m, 3H). **LCMS:** (Method K) 505.0 (M⁺+H), Rt. 3.01 min, 96.72% (Max). **HPLC:** (Method E) Rt. 5.61 min, 97.75% (Max). **Chiral Purity:** (Method H) Rt. 3.19 min, 99.69% (Max).

Enantiomer 2: **Yield:** 20% (25 mg, white solid). **¹H NMR** (400 MHz, DMSO-*d*₆) δ 12.44 (s, 1H), 7.28 (s, 1H), 7.16 (t, *J* = 7.2 Hz, 2H), 7.05 (s, 1H), 6.71 (t, *J* = 7.2 Hz, 1H), 6.58 (d, *J* = 8.4 Hz, 2H), 4.27 (m, 2H), 4.02 (m, 1H), 3.90 (bs, 1H), 3.21 (bs, 1H), 2.46 (s, 3H), 2.34 (s, 3H), 1.65-1.62 (m, 1H), 1.50-1.49 (m, 1H), 1.37-1.35 (m, 4H), 1.24-1.23 (m, 2H), 1.11-1.10 (m, 2H), 0.94-0.92 (m, 3H). **LCMS:** (Method K) 505.0 (M⁺+H), Rt. 3.01 min, 97.62% (Max). **HPLC:** (Method E) Rt. 5.61 min, 98.84% (Max). **Chiral Purity:** (Method H) Rt. 4.18 min, 99.55% (Max).

Example 5

3-((3-Butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid




To a stirred solution of methyl 3-((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoate (Intermediate 15; 0.16 g, 0.30

mmol) in a mixture of 1,4-dioxane and water (3: 2, 5 mL) at 0 °C, lithium hydroxide (0.02 g, 0.60 mmol) was added and the reaction mixture was stirred for 2 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (10 mL) and brine (10 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by prep HPLC purification (Method A) to afford the title compound. **Yield:** 38% (60 mg, white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 12.45 (s, 1H), 7.29 (s, 1H), 7.16 (t, *J* = 7.6 Hz, 2H), 7.08 (s, 1H), 6.71 (t, *J* = 7.6 Hz, 1H), 6.59 (d, *J* = 7.6 Hz, 2H), 4.11 (s, 2H), 4.05-3.85 (m, 2H), 3.25-3.15 (m, 1H), 3.02-2.8 (m, 2H), 2.60 (m, 1H), 2.47 (s, 3H), 1.7-1.45 (m, 2H), 1.42-1.3 (m, 5H), 1.28 (m, 1H), 1.27 (s, 6H), 1.21 (t, *J* = 7.60 Hz, 3H), 0.92 (t, *J* = 6.8 Hz, 3H). **LCMS:** (Method B) 521.0 (M⁺+H), Rt. 2.05 min, 96.18% (Max). **HPLC:** (Method E) Rt. 5.94 min, 97.88% (Max).

Example 6

(S)-3-((3-Butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid and (R)-3-((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid



The two enantiomers of racemic 3-((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid (Example 5; 50 mg, 0.09 mmol) were separated by chiral SFC Instrument (Method F). The material was concentrated under vacuum at 40 °C. The first eluting fraction corresponded to enantiomer 1 and the second eluting fraction corresponded to enantiomer 2. The absolute configuration of the two enantiomers is not known.

25


Enantiomer 1: **Yield:** 37% (20 mg, off-white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 7.29 (s, 1H), 7.15 (t, *J* = 8.8 Hz, 2H), 7.08 (s, 1H), 6.70 (t, *J* = 7.2 Hz, 1H), 6.59 (d, *J* = 8.0 Hz, 2H), 4.11 (s, 2H), 4.05-3.83 (m, 2H), 3.28-3.12 (m, 1H), 2.95-2.82 (m, 2H), 2.47 (s, 3H), 1.68-1.55 (m, 1H), 1.55-1.45 (m, 1H), 1.42-1.32 (m, 4H), 1.32-1.22 (m, 6H), 1.20 (t, *J* = 7.20 Hz, 3H), 0.95-0.88 (m, 3H). **LCMS:** (Method K) 521.2 (M⁺+H), Rt. 3.13 min, 98.08% (Max). **HPLC:** (Method E) Rt. 5.91 min, 91.39% (Max). **Chiral SFC:** (Method H) Rt. 1.72 min, 95.81% (Max).

30

Enantiomer 2: **Yield:** 23% (15 mg, off-white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 12.45 (s, 1H), 7.29 (s, 1H), 7.15 (t, *J* = 8.8 Hz, 2H), 7.08 (s, 1H), 6.70 (t, *J* = 7.6 Hz, 1H), 6.59 (d, *J* = 8.0 Hz, 2H), 4.11 (s, 2H), 4.08-3.80 (m, 2H), 3.28-3.12 (m, 1H), 2.92-2.82 (m, 2H), 2.47 (s, 3H), 1.68-1.58 (m, 1H), 1.58-1.45 (m, 1H), 1.42-1.32 (m, 4H), 1.32-1.22 (m, 6H), 1.2 (t, *J* = 7.2 Hz, 3H), 0.95-0.88 (m, 3H). **LCMS:** (Method D) 521.0 (M⁺+H), Rt. 3.13 min, 94.38% (Max). **HPLC:** (Method E) Rt. 5.93 min, 94.93% (Max). **Chiral SFC:** (Method H) Rt. 2.30 min, 95.03% (Max).

Example 7


1-(((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid



To a stirred solution of ethyl 1-(((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate (Intermediate 16; 0.05 g, 0.091 mmol) in a mixture of 1,4-dioxane and water (3:2, 5 mL) at 0 °C, lithium hydroxide (4.38 mg, 0.18 mmol) was added and the reaction mixture was stirred for 2 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 3 mL) and the aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by prep HPLC purification (Method A) to afford the title compound. **Yield:** 30% (15 mg, white solid). **¹H NMR** (400 MHz, DMSO-*d*₆) δ 12.46 (s, 1H), 7.28 (s, 1H), 7.15 (t, *J* = 7.6 Hz, 2H), 7.08 (s, 1H), 6.70 (t, *J* = 7.2 Hz, 1H), 6.58 (d, *J* = 8.0 Hz, 2H), 4.29-4.20 (m, 2H), 4.04-4.00 (m, 1H), 3.88 (m, 1H), 3.22 (m, 1H), 2.88 (m, 2H), 2.46 (s, 3H), 1.65-1.60 (m, 1H), 1.54-1.49 (m, 1H), 1.35 (m, 4H), 1.24-1.18 (m, 5H), 1.09 (m, 2H), 0.93 (m, 3H). **LCMS:** (Method K) 519.2 (M⁺+H), Rt. 2.83 min, 94.83% (Max). **HPLC:** (Method E) Rt. 5.79 min, 94.74% (Max).

Example 8

(S)-1-(((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid and (R)-1-(((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid




The two enantiomers of racemic 1-(((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid (Example 7; 60 mg, 0.12 mmol) were separated by chiral SFC Instrument (Method H). The material was concentrated under vacuum at 40 °C. The first eluting fraction corresponded to enantiomer 1 and the second eluting fraction corresponded to enantiomer 2. The absolute configuration of the two enantiomers is not known.

Enantiomer 1: **Yield:** 24% (15 mg, off-white solid). ¹H NMR (400 MHz, DMSO-*d*₆): δ 12.43 (s, 1H), 7.28 (s, 1H), 7.15 (t, *J* = 8.8 Hz, 2H), 7.08 (s, 1H), 6.70 (t, *J* = 7.2 Hz, 1H), 6.58 (d, *J* = 8.4 Hz, 2H), 4.24 (m, 2H), 4.10-3.95 (m, 1H), 3.95-3.80 (m, 1H), 3.28-3.12 (m, 1H), 2.98-2.84 (m, 2H), 2.46 (s, 3H), 1.70-1.58 (m, 1H), 1.58-1.45 (m, 1H), 1.40-1.32 (m, 4H), 1.28-1.18 (m, 5H), 1.15-1.05 (m, 2H), 0.9 (t, *J* = 6.8 Hz, 3H). **LCMS:** (Method A) 519.2 (M⁺+H), Rt. 3.15 min, 97.05% (Max). **HPLC:** (Method E) Rt. 5.94 min, 97.46% (Max). **Chiral SFC:** (Method K) Rt. 2.87 min, 99.75% (Max).

Enantiomer 2: **Yield:** 23% (15 mg, off-white solid). ¹H NMR (400 MHz, DMSO-*d*₆): δ 12.43 (s, 1H), 7.28 (s, 1H), 7.15 (t, *J* = 7.6 Hz, 2H), 7.08 (s, 1H), 6.70 (t, *J* = 7.6 Hz, 1H), 6.58 (d, *J* = 8.0 Hz, 2H), 4.35-4.12 (m, 2H), 4.10-3.98 (m, 1H), 3.98-3.80 (m, 1H), 3.28-3.10 (m, 1H), 2.89 (m, 2H), 2.47 (s, 3H), 1.68-1.58 (m, 1H), 1.58-1.45 (m, 1H), 1.45-1.30 (m, 4H), 1.28-1.18 (m, 5H), 1.15-1.05 (m, 2H), 1.00-0.85 (m, 3H). **LCMS:** (Method A) 519.1 (M⁺+H), Rt. 2.70 min, 95.84% (Max). **HPLC:** (Method E) Rt. 5.79 min, 94.28% (Max). **Chiral SFC:** (Method K) Rt. 3.61 min, 99.46% (Max).

Example 9

3-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid




25

To a stirred solution of methyl 3-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)-2,2-dimethylpropanoate (Intermediate 21; 0.12 g, 0.23 mmol) in a mixture of 1,4-dioxane and water (3:2; 5 mL) at 0 °C, lithium hydroxide (10.97 mg, 0.46 mmol) was added and the reaction mixture was stirred for 2 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by prep HPLC purification (Method A) to afford the title compound. **Yield:** 17% (20 mg, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 12.43 (s, 1H), 7.35 (s, 1H), 7.03-6.97 (m, 3H), 6.58-6.54 (m, 2H), 4.25-4.08 (m, 3H), 3.50-3.35 (m, 1H), 3.25-3.02 (m, 2H), 2.34 (s, 3H), 2.32-2.20 (m, 1H), 1.38-1.35 (m, 4H), 1.32-1.27 (m, 8H), 0.91 (t, *J* = 7.20 Hz, 3H). **LCMS:** (Method A) 510.0 (M⁺+H), Rt. 3.03 min, 98.51% (Max). **HPLC:** (Method E) Rt. 5.67 min, 99.12% (Max).

15 Example 10

1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid




To a stirred solution of ethyl 1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate (Intermediate 22; 0.1 g, 0.19 mmol) in a mixture of 1,4-dioxane and water (5:1, 6 mL) at 0 °C, lithium hydroxide (8.94 mg, 0.373 mmol) was added and the reaction mixture was stirred for 2 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 3 mL) and the aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by prep HPLC purification (Method A) to afford the title compound. **Yield:** 15% (15 mg, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 12.43 (s, 1H), 7.35 (s, 1H), 7.03-6.97 (m, 3H), 6.57-6.54 (m, 2H), 4.32-4.12 (m, 3H), 3.50-3.38 (m, 1H), 3.25-3.00 (m, 2H), 2.34 (s, 3H), 2.32-2.20 (m, 1H), 1.45-1.20 (m, 8H),

1.12-1.08 (m, 2H), 0.90 (t, $J = 7.20$ Hz, 3H). **LCMS:** (Method A) 508.1 ($M^+ + H$), Rt. 2.96 min, 94.20% (Max). **HPLC:** (Method E) Rt. 5.53 min, 94.35% (Max).

Example 11

- 5 **1-(((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid**



- To a stirred solution of ethyl 1-(((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate
 10 (Intermediate 30; 80 mg, 0.15 mmol) in a mixture of 1,4-dioxane and water (3:2, 5 mL) at 0 °C, lithium hydroxide (6.96 mg, 0.29 mmol) was added and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 3 mL) and the aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over
 15 anhydrous Na_2SO_4 . The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by prep HPLC purification (Method A) to afford the title compound.


Yield: 29% (22 mg, white solid).

- $^1\text{H NMR}$** (400 MHz, $\text{DMSO}-d_6$): δ 12.45 (s, 1H), 7.27 (s, 1H), 7.10-6.82 (m, 3H), 6.72-6.48 (m, 2H), 4.25 (m, 2H), 4.05-3.92 (m, 1H), 3.92-3.80 (m, 1H), 3.28-3.10 (m, 1H), 2.45 (s, 3H), 2.34 (s, 3H), 1.68-1.55
 20 (m, 1H), 1.55-1.42 (m, 1H), 1.42-1.28 (m, 4H), 1.28-1.18 (m, 2H), 1.18-1.02 (m, 2H), 1.00-0.82 (m, 3H).

LCMS: (Method B) 523.0 ($M^+ + H$), Rt. 2.10 min, 98.06% (Max). **HPLC:** (Method E) Rt. 5.75 min, 99.00% (Max).

Example 12

- 25 **(S)-1-(((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid and (R)-1-(((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid**




The two enantiomers of racemic 1-(((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid (Example 11; 40 mg, 0.077 mmol) were separated by chiral SFC Instrument (Method A). The material was concentrated under vacuum at 40 °C. The first eluting fraction corresponded to enantiomer 1 and the second eluting fraction corresponded to enantiomer 2. The absolute configuration of the two enantiomers is not known.

Enantiomer 1: **Yield:** 25% (10 mg, white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 12.44 (s, 1H), 7.27 (s, 1H), 7.02-7.01 (m, 3H), 6.59 (m, 2H), 4.26 (m, 2H), 4.05-3.98 (m, 1H), 3.92-3.82 (m, 1H), 3.2-3.15 (m, 1H), 2.46 (s, 3H), 2.34 (s, 3H), 1.68-1.55 (m, 1H), 1.55-1.48 (m, 1H), 1.4-1.3 (m, 4H), 1.25-1.18 (m, 2H), 1.12-1.05 (m, 2H), 0.98-0.88 (m, 3H). **LCMS:** (Method A) 523.1 (M⁺+H), Rt. 3.05 min, 97.58% (Max). **HPLC:** (Method E) Rt. 5.66 min, 99.56% (Max). **Chiral SFC:** (Method U) Rt. 2.37 min, 99.39% (Max).

Enantiomer 2: **Yield:** 36% (15 mg, white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 12.45 (s, 1H), 7.27 (s, 1H), 7.10-6.90 (m, 3H), 6.65-6.48 (m, 2H), 4.32-4.18 (m, 2H), 4.05-3.95 (m, 1H), 3.90-3.82 (m, 1H), 3.25-3.15 (m, 1H), 2.47 (s, 3H), 2.34 (s, 3H), 1.65-1.55 (m, 1H), 1.55-1.45 (m, 1H), 1.42-1.28 (m, 4H), 1.28-1.20 (m, 2H), 1.15-1.05 (m, 2H), 0.9 (t, *J* = 6.8 Hz, 3H). **LCMS:** (Method A) 523.1 (M⁺+H), Rt. 3.05 min, 96.30% (Max). **HPLC:** (Method E) Rt. 5.62 min, 95.65% (Max). **Chiral SFC:** (Method U) Rt. 2.93 min, 99.67% (Max).

Example 13

3-((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid




To a stirred solution of methyl 3-((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoate (Intermediate 31; 0.04 g, 0.074 mmol) in a mixture of 1,4-dioxane and water (5:1, 6 mL) at 0 °C, lithium hydroxide (3.56 mg, 0.15 mmol) was added and the reaction mixture was stirred for 12 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude material was purified by prep HPLC purification (Method A) to afford the title compound. **Yield:** 25% (10 mg, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 12.46 (s, 1H), 7.27 (s, 1H), 7.13-6.82 (m, 3H), 6.70-6.40 (m, 2H), 4.11 (s, 2H), 4.05-3.95 (m, 1H), 3.95-3.75 (m, 1H), 3.30-3.15 (m, 1H), 2.46 (s, 3H), 2.34 (s, 3H), 1.72-1.55 (m, 1H), 1.55-1.42 (m, 1H), 1.40-1.30 (m, 4H), 1.30-1.25 (m, 6H), 0.95-0.88 (m, 3H). **LCMS:** (Method A) 524.8 (M⁺), Rt. 2.74 min, 95.76% (Max). **HPLC:** (Method E) Rt. 5.75 min, 96.44% (Max).

Example 14

(S)-3-((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid and (R)-3-((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid



The two enantiomers of racemic 3-((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid (Example 13; 0.025 g, 0.143 mmol) were separated by chiral SFC (Method G). The material was concentrated under vacuum at 40 °C. The first eluting fraction corresponded to enantiomer 1 and the second eluting fraction corresponded to enantiomer 2. The absolute configuration of the two enantiomers is not known.


Enantiomer 1: **Yield:** 33.2% (0.025 g, off-white solid). **¹H-NMR** (400 MHz, DMSO-*d*₆): δ 12.46 (s, 1H), 7.27 (s, 1H), 7.02-7.01 (m, 1H), 6.98-6.96 (m, 2H), 6.59-6.56 (m, 2H), 4.11 (s, 2H), 4.01-3.97 (m, 1H), 3.86 (s, 1H), 3.24-3.21 (m, 1H), 2.46 (s, 3H), 2.34 (s, 3H), 1.60-1.41 (m, 2H), 1.35-1.34 (m, 4H), 1.27 (s,

6H), 0.93-0.91 (m, 3H). **LCMS:** (Method A) 525.1 (M^+H), Rt.3.18 min, 99.48% (Max). **HPLC:** (Method B) Rt. 5.88 min, 99.62% (Max). **Chiral HPLC:** (Method G) Rt. 1.65 min, 100% (Max)

Enantiomer 2: **Yield:** 26.6% (0.020 g, off-white solid). **1H -NMR** (400 MHz, DMSO- d_6): δ 12.46 (s, 1H), 7.27 (s, 1H), 7.02-7.01 (m, 1H), 6.98-6.96 (m, 2H), 6.59-6.57 (m, 2H), 4.11 (s, 2H), 4.01-3.97 (m, 1H),
 5 3.87 (s, 1H), 3.24-3.23 (m, 1H), 2.46 (s, 3H), 2.34 (s, 3H), 1.60 (s, 2H), 1.35 (s, 4H), 1.27-1.24 (m, 6H), 0.93-0.91 (m, 3H). **LCMS:** (Method A) 525.2 (M^+H), Rt.3.17 min, 98.37% (Max). **HPLC:** (Method B) Rt. 5.88 min, 99.71% (Max). **Chiral HPLC** (Method G) Rt. 2.22 min, 99.76% (Max)

Example 15

10 **1-(((3-butyl-7-(ethylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid**




To a stirred solution of ethyl 1-(((3-butyl-7-(ethylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate (Intermediate 41; 0.05 g, 0.09 mmol)
 15 in a mixture of 1,4-dioxane and water (5:1, 6 mL) at 0 °C, lithium hydroxide (4.50 mg, 0.19 mmol) was added and the reaction mixture was stirred for 12 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 3 mL) and the aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was
 20 filtered and concentrated under vacuum. The resulting crude material was purified by prep HPLC purification (Method A) to afford the title compound. **Yield:** 20% (10 mg, white solid).

1H NMR (400 MHz, DMSO- d_6): δ 12.45 (s, 1H), 7.32 (s, 1H), 7.25 (d, J = 9.2 Hz, 1H), 7.14 (t, J = 8.4 Hz, 2H), 7.08 (s, 1H), 6.69 (t, J = 7.2 Hz, 1H), 6.57 (d, J = 8.4 Hz, 2H), 4.45-4.30 (m, 1H), 4.30-4.15 (m, 2H), 3.52-3.42 (m, 1H), 2.95-2.80 (m, 3H), 1.70-1.52 (m, 2H), 1.52-1.38 (m, 4H), 1.35-1.28 (m, 2H), 1.25-
 25 1.22 (m, 3H), 1.12-1.08 (m, 2H), 0.9 (t, J = 7.2 Hz, 3H). **LCMS:** (Method A) 505.1 (M^+H), Rt. 2.52 min, 92.00% (Max). **HPLC:** (Method E) Rt. 5.55 min, 92.07% (Max).

Example 16

30 **1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid**




To a stirred solution of ethyl 1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate (Intermediate 45; 0.03 g, 0.06 mmol) in a mixture of 1,4-dioxane and water (5:1, 6 mL) at 0 °C, lithium hydroxide (2.68 mg, 0.11 mmol) was added and the reaction mixture was stirred for 12 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 3 mL) and the aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude was purified by prep HPLC purification (Method A) to afford the title compound. **Yield:** 33% (10 mg, white solid). ¹H NMR (400 MHz, DMSO-*d*₆): δ 12.51 (s, 1H), 7.36 (s, 1H), 7.32 (d, *J* = 9.6 Hz, 1H), 7.12-6.98 (m, 3H), 6.72-6.55 (m, 2H), 4.42-4.22 (m, 3H), 3.52-3.48 (m, 1H), 2.92-2.88 (m, 1H), 2.39 (s, 3H), 1.70-1.25 (m, 10H), 1.00-0.92 (m, 3H). **LCMS:** (Method A) 509.1 (M⁺+H), Rt. 2.41 min, 91.15% (Max). **HPLC:** (Method E) Rt. 5.40 min, 93.38% (Max).

15

Example 17

(S)-1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid and (R)-1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid

20




The two enantiomers of racemic 1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid (Example 16; 0.1 g, 0.19 mmol) were separated by chiral SFC Instrument (Method M). The material was concentrated

under vacuum at 40 °C. The first eluting fraction corresponded to enantiomer 1 and the second eluting fraction corresponded to enantiomer 2. The absolute configuration of the two enantiomers is not known.

- 5 Enantiomer 1: **Yield:** 40% (40 mg, off-white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 12.45 (s, 1H), 7.30 (s, 1H), 7.26 (d, *J* = 9.2 Hz, 1H), 7.05-6.95 (m, 3H), 6.62-6.48 (m, 2H), 4.38-4.15 (m, 3H), 3.48-3.40 (m, 1H), 2.95-2.78 (m, 1H), 2.33 (s, 3H), 1.65-1.42 (m, 2H), 1.42-1.28 (m, 4H), 1.28-1.20 (m, 2H), 1.15-1.05 (m, 2H), 0.90 (t, *J* = 7.20 Hz, 3H). **LCMS:** (Method K) 509.2 (M⁺+H), Rt. 2.40 min, 98.25% (Max). **HPLC:** (Method E) Rt. 5.51 min, 99.22% (Max). **Chiral SFC:** (Method G) Rt. 1.41 min, 100% (Max).
- 10 Enantiomer 2: **Yield:** 40% (40 mg, off-white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 12.45 (s, 1H), 7.30 (s, 1H), 7.26 (d, *J* = 8.8 Hz, 1H), 7.10-7.92 (m, 3H), 6.65-6.48 (m, 2H), 4.38-4.18 (m, 3H), 3.52-3.42 (m, 1H), 2.95-2.78 (m, 1H), 2.33 (s, 3H), 1.65-1.45 (m, 2H), 1.45-1.30 (m, 4H), 1.28-1.20 (m, 2H), 1.15-1.05 (m, 2H), 0.90 (t, *J* = 6.80 Hz, 3H). **LCMS:** (Method B) 508.9 (M⁺+H), Rt. 1.88 min, 96.57% (Max). **HPLC:** (Method E) Rt. 5.51 min, 98.55% (Max). **Chiral SFC:** (Method G) Rt. 2.13 min, 99.05% (Max).

15


Example 18**1-(((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid**

- 20 To a stirred solution of ethyl 1-(((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylate (Intermediate 47; 0.07 g, 0.13 mmol) in a mixture of 1,4-dioxane and water (5:1, 6 mL) at 0 °C, lithium hydroxide (6.09 mg, 0.25 mmol) was added and the reaction mixture was stirred for 2 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute
- 25 HCl (1.5 N, 3 mL) and the aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude was purified by prep HPLC purification (Method A) to afford the title compound. **Yield:** 58% (40 mg, white solid).
- ¹H NMR** (400 MHz, DMSO-*d*₆): δ 12.45 (s, 1H), 7.31 (s, 1H), 7.26 (d, *J* = 9.2 Hz, 1H), 7.05 (s, 1H), 6.99 (t, *J* = 8.8 Hz, 2H), 6.68-6.45 (m, 2H), 4.42-4.22 (m, 2H), 4.22-4.10 (m, 1H), 3.50-3.40 (m, 1H), 2.87 (m,
- 30

3H), 1.65-1.52 (m, 1H), 1.48-1.22 (m, 7H), 1.22-1.18 (m, 3H), 1.15-1.05 (m, 2H), 0.90 (t, $J = 7.20$ Hz, 3H). **LCMS:** (Method A) 521.3 (M^+H), Rt. 3.02 min, 95.10% (Max). **HPLC:** (Method E) Rt. 5.61 min, 97.11% (Max).

5 Example 19

(S)-1-(((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid and (R)-1-(((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid



10

The two enantiomers of racemic 1-(((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid (Example 18; 30 mg, 0.06 mmol) were separated by chiral SFC Instrument (Method M). The material was concentrated under vacuum at 40 °C. The first eluting fraction corresponded to enantiomer 1 and the second eluting fraction corresponded to enantiomer 2. The absolute configuration of the two enantiomers is not known.

15


Enantiomer 1: **Yield:** 30% (9 mg, off-white solid). **1H NMR** (400 MHz, DMSO- d_6): δ 12.43 (s, 1H), 7.31 (s, 1H), 7.26 (d, $J = 9.6$ Hz, 1H), 7.05 (s, 1H), 6.99 (t, $J = 8.8$ Hz, 2H), 6.65-6.52 (m, 2H), 4.38-4.12 (m, 3H), 3.42-3.38 (m, 1H), 2.92-2.85 (m, 3H), 1.65-1.52 (m, 2H), 1.45-1.38 (m, 2H), 1.25-1.20 (m, 3H), 1.20-1.18 (m, 3H), 1.12-1.02 (m, 3H), 0.95-0.88 (m, 3H). **LCMS:** (Method A) 521.2 (M^+H), Rt. 3.01 min, 95.40% (Max). **HPLC:** (Method E) Rt. 5.63 min, 99.21% (Max). **Chiral SFC:** (Method G) Rt. 2.2 min, 99.74% (Max).

20

Enantiomer 2: **Yield:** 23% (7 mg, off-white solid). **1H NMR** (400 MHz, DMSO- d_6): δ 12.44 (s, 1H), 7.31 (s, 1H), 7.26 (d, $J = 9.6$ Hz, 1H), 7.05 (s, 1H), 6.99 (t, $J = 8.8$ Hz, 2H), 6.68-6.50 (m, 2H), 4.35-4.15 (m, 3H), 3.42-3.38 (m, 1H), 2.92-2.85 (m, 3H), 1.65-1.55 (m, 1H), 1.50-1.32 (m, 4H), 1.28-1.15 (m, 6H), 1.12-1.08 (m, 2H), 0.95-0.85 (m, 3H). **LCMS:** (Method A) 521.1 (M^+H), Rt. 3.00 min, 97.25% (Max). **HPLC:** (Method E) Rt. 5.70 min, 97.41% (Max). **Chiral SFC:** (Method G) Rt. 3.45 min, 99.49% (Max).


25

Example 20

3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-difluoropropanoic acid

5 To a stirred solution of ethyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-difluoropropanoate (Intermediate 48; 0.1 g, 0.18 mmol) in a mixture of 1,4-dioxane and water (3:1, 4 mL) at 0 °C, lithium hydroxide (8.83 mg, 0.37 mmol) was added and the reaction mixture was stirred for 16 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 10 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered and concentrated under vacuum. The resulting crude was purified by prep HPLC purification (Method A) to afford the title compound. **Yield:** 14% (14 mg, off-white solid).

15 ¹H NMR (400 MHz, DMSO-*d*₆): δ 7.33 (s, 1H), 7.15 (t, *J* = 8.4 Hz, 2H), 7.06 (s, 1H), 6.70 (t, *J* = 7.2 Hz, 1H), 6.58 (d, *J* = 8.4 Hz, 2H), 4.51 (t, *J* = 14.4 Hz, 2H), 4.08-3.98 (m, 1H), 3.95-3.85 (m, 1H), 3.25-3.15 (m, 1H), 2.46 (s, 3H), 2.36 (s, 3H), 1.68-1.58 (m, 1H), 1.55-1.45 (m, 1H), 1.40-1.28 (m, 4H), 0.95-0.88 (m, 3H). **LCMS:** (Method K) 515.2 (M⁺+H), Rt. 2.46 min, 95.16% (Max). **HPLC:** (Method E) Rt. 5.57 min, 96.09% (Max).

20 **Example 21****(S)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-difluoropropanoic acid and (R)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-difluoropropanoic acid**

25

The two enantiomers of racemic 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-difluoropropanoic acid (Example 20; 40 mg, 0.08

mmol) were separated by chiral SFC Instrument (Method F). The material was concentrated under vacuum at 40 °C. The first eluting fraction corresponded to enantiomer 1 and the second eluting fraction corresponded to enantiomer 2. The absolute configuration of the two enantiomers is not known.

5

Enantiomer 1: **Yield:** 12% (5 mg, off-white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 7.34 (s, 1H), 7.15 (t, *J* = 8.4 Hz, 2H), 7.06 (s, 1H), 6.70 (t, *J* = 7.2 Hz, 1H), 6.58 (d, *J* = 8.0 Hz, 2H), 4.52 (t, *J* = 14.0 Hz, 2H), 4.05-3.95 (m, 1H), 3.95-3.80 (m, 1H), 3.25-3.15 (m, 1H), 2.46 (s, 3H), 2.36 (s, 3H), 1.68-1.58 (m, 1H), 1.55-1.45 (m, 1H), 1.42-1.32 (m, 4H), 0.98-0.88 (m, 3H). **LCMS:** (Method B) 515.0 (M⁺+H), Rt. 2.09 min, 94.41% (Max). **HPLC:** (Method E) Rt. 5.52 min, 98.13% (Max). **Chiral SFC:** (Method H) Rt. 3.26 min, 100% (Max).

10


Enantiomer 2: **Yield:** 13% (5 mg, off-white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 7.34 (s, 1H), 7.15 (t, *J* = 8.8 Hz, 2H), 7.06 (s, 1H), 6.70 (t, *J* = 7.2 Hz, 1H), 6.58 (d, *J* = 8.4 Hz, 2H), 4.50 (t, *J* = 14.4 Hz, 2H), 4.05-3.98 (m, 1H), 3.95-3.82 (m, 1H), 3.25-3.12 (m, 1H), 2.46 (s, 3H), 2.36 (s, 3H), 1.68-1.58 (m, 1H), 1.50-1.45 (m, 1H), 1.42-1.30 (m, 4H), 0.92 (t, *J* = 6.80 Hz, 3H). **LCMS:** (Method B) 515.1 (M⁺+H), Rt. 2.09 min, 93.89% (Max). **HPLC:** (Method E) Rt. 5.53 min, 99.76% (Max). **Chiral SFC:** (Method H) Rt. 5.93 min, 100% (Max).

15

Example 22

20

3-((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid



To a stirred solution of methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoate (Intermediate 52; 0.13 g, 0.24 mmol) in a mixture of 1,4-dioxane and water (2:1, 3 mL) at 0 °C, lithium hydroxide (0.02 g, 0.48 mmol) was added and the reaction mixture was stirred for 3 hours at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, 5 mL) and the aqueous layer was extracted with ethyl acetate (2 x 20 mL). The combined organic layer was washed with water (10 mL) and brine (10 mL) and dried over anhydrous Na₂SO₄. The


25

organic part was filtered, concentrated under vacuum and the resulting crude was purified by prep-HPLC purification (Method A) to afford the title compound. **Yield:** 25% (30 mg, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 13.0 (s, 1H), 7.32 (s, 1H), 7.16 (t, *J* = 7.6 Hz, 2H), 7.05 (s, 1H), 6.71 (t, *J* = 7.2 Hz, 1H), 6.58 (d, *J* = 8.4 Hz, 2H), 4.36 (m, 1H), 4.27 (m, 1H), 4.02 (m, 1H), 3.95-3.85 (m, 1H), 3.31 (s, 3H), 3.25-3.15 (bs, 1H), 2.46 (s, 3H), 2.34 (s, 3H), 1.7-1.3 (m, 9H), 0.94-0.92 (m, 3H). **LCMS:** (Method K) 522.9 (M⁺+H), Rt. 2.94 min, 98.59% (Max). **HPLC:** (Method E) Rt. 5.54 min, 98.79% (Max).

Example 23

3-((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid




The title compound was obtained following the same procedure as described for Example 21, starting from 0.05 g of Intermediate 53. After work-up of the reaction mixture, the crude material was purified by Isolera column chromatography (eluent: 10-15% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound. The absolute configuration of the compound is not known. **Yield:** 50% (25 mg, off-white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 13.0 (s, 1H), 7.33 (s, 1H), 7.16 (t, *J* = 7.2 Hz, 2H), 7.05 (s, 1H), 6.70 (t, *J* = 7.6 Hz, 1H), 6.58 (d, *J* = 8.4 Hz, 2H), 4.36 (t, *J* = 9.6 Hz, 1H), 4.27 (t, *J* = 8.0 Hz, 1H), 4.02 (m, 1H), 3.89-3.88 (m, 1H), 3.30 (s, 3H), 3.25-3.15 (m, 1H), 2.46 (s, 3H), 2.35 (s, 3H), 1.7-1.6 (m, 1H), 1.6-1.42 (m, 4H), 1.42-1.28 (m, 4H), 1.0-0.85 (m, 3H). **LCMS:** (Method K) 522.9 (M⁺+H), Rt. 2.73 min, 99.48% (Max). **HPLC:** (Method E) Rt. 5.51 min, 99.35% (Max).

Example 24

(S)-3-((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid and (R)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid (individual diastereomers)




Diastereoisomers 1 and 2 of the title compound were prepared from diastereoisomer 1 (95 mg) and diastereoisomer 2 (90 mg) of Intermediate 54, respectively, following the same procedure as described for Example 21. After work-up of the reaction mixtures, the crude materials were purified by Isolera column chromatography (eluent: 4% MeOH/DCM; silica gel: 230-400 mesh) to afford the title compounds. The absolute configuration of the two diastereomers is not known.

Diastereoisomer 1: **Yield:** 57% (55 mg, white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 7.33 (s, 1H), 7.16 (t, *J* = 7.2 Hz, 2H), 7.05 (s, 1H), 6.71 (t, *J* = 7.6 Hz, 1H), 6.58 (d, *J* = 8.0 Hz, 2H), 4.42-4.32 (m, 1H), 4.3-4.2 (m, 1H), 4.1-3.8 (m, 2H), 3.31 (s, 3H), 3.25-3.15 (m, 1H), 2.46 (s, 3H), 2.35 (s, 3H), 1.7-1.58 (m, 1H), 1.55 - 1.28 (m, 8H), 0.95-0.88 (m, 3H). **LCMS:** (Method K) 523.3 (M⁺+H), Rt. 2.71 min, 98.96% (Max). **HPLC:** (Method E) Rt. 5.45 min, 97.51% (Max). **Chiral Purity:** (Method H) Rt. 9.02 min, 98.10% (Max).
 Diastereoisomer 2: **Yield:** 49% (45 mg, white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 13.08 (s, 1H), 7.33 (s, 1H), 7.16 (t, *J* = 8.0 Hz, 2H), 7.06 (s, 1H), 6.71 (t, *J* = 7.2 Hz, 1H), 6.58 (d, *J* = 8.4 Hz, 2H), 4.42-4.32 (m, 1H), 4.3-4.18 (m, 1H), 4.1-3.85 (m, 2H), 3.30 (s, 3H), 3.25-3.15 (m, 1H), 2.47 (s, 3H), 2.35 (s, 3H), 1.7-1.45 (m, 5H), 1.45-1.28 (m, 4H), 0.95-0.88 (m, 3H). **LCMS:** (Method K) 523.3 (M⁺+H), Rt. 2.71 min, 98.20% (Max). **HPLC:** (Method E) Rt. 5.45 min, 98.01% (Max). **Chiral Purity:** (Method H) Rt. 7.83 min, 98.92% (Max).

20 Example 25

3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)propanoic acid



To a stirred solution of 3-butyl-8-hydroxy-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepine 1,1-dioxide (0.2 g, 0.05 mmol) in THF (5 mL), potassium *tert*-butoxide (0.063 g, 0.06 mmol) and oxetan-2-one (0.04 g, 0.06 mmol) were added, and the reaction mixture was stirred for 3


hours at 50 °C. As TLC showed incomplete conversion, more oxetan-2-one (0.04 g, 0.06 mmol) was added and the reaction mixture was stirred for 16 hours at 50 °C. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (5 mL, 1.5 N HCl) and the aqueous layer was extracted with EtOAc (2 x 15 mL). The combined organic layer was washed with water (5 mL) and brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting material was purified by Isolera column chromatography (eluent: 20% EtOAc/PE; silica gel: 230-400 mesh) to afford the title compound.

Yield: 31% (75 mg, white solid).

¹H-NMR (400 MHz, DMSO-*d*₆): δ 7.51 (s, 1H), 7.18 (t, *J* = 8.00 Hz, 2H), 7.06 (s, 1H), 6.75 (t, *J* = 7.20 Hz, 1H), 6.67 (d, *J* = 8.00 Hz, 2H), 4.40 (s, 2H), 4.35-4.31 (m, 1H), 3.44-3.40 (m, 1H), 3.14-3.05 (m, 2H), 2.87-2.85 (m, 2H), 2.45 (s, 1H), 2.33 (s, 3H), 1.47-1.40 (m, 6H), 1.00-0.89 (m, 3H). **LCMS:** (Method E) 464.1 (M⁺+H) Rt. 4.81 min, 95.14% (Max). **HPLC:** (Method B) Rt. 5.36 min, 95.27% (Max).

Example 26

(S)-3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)propanoic acid and (R)-3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)propanoic acid




The two enantiomers of racemic 3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)propanoic acid (Example 25; 68 mg, 0.15 mmol) were separated by chiral SFC (Method D). The material was concentrated under vacuum at 40 °C. The first eluting fraction corresponded to enantiomer 1 and the second eluting fraction corresponded to enantiomer 2. The absolute configuration of the two enantiomers is not known.

Enantiomer 1: **Yield:** 35% (0.024 g, white solid). ¹H NMR (400 MHz, DMSO-*d*₆): δ 12.47 (s, 1H), 7.39 (s, 1H), 7.15 (t, *J* = 7.60 Hz, 2H), 7.06 (s, 1H), 6.68 (t, *J* = 7.20 Hz, 1H), 6.56 (d, *J* = 8.00 Hz, 2H), 4.34-4.21 (m, 3H), 3.34 (s, 1H), 3.14 (s, 1H), 3.10-3.04 (m, 1H), 2.72 (s, 2H), 2.34 (s, 4H), 1.40-1.30 (m, 6H), 0.93-0.89 (m, 3H). **LCMS:** (Method A) 464.1 (M⁺+H) Rt. 2.54 min, 94.45% (Max). **HPLC:** (Method B) Rt. 5.36 min, 97.37% (Max). **Chiral HPLC:** (Method D) Rt. 2.07min, 100% (Max).

Enantiomer 2: **Yield:** 12% (0.008 g, white solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 7.39 (s, 1H), 7.17-7.15 (m, 2H), 7.06 (s, 1H), 6.68 (t, *J* = 7.20 Hz, 1H), 6.55 (d, *J* = 8.00 Hz, 2H), 4.34-4.30 (m, 3H), 3.44-3.41 (m, 1H), 3.40-3.33 (m, 1H), 3.29-3.23 (m, 1H), 2.76-2.73 (m, 2H), 2.34 (s, 4H), 1.40-1.30 (m, 6H), 0.93-0.89 (m, 3H). **LCMS:** (Method A) 464.2 (M⁺+H) Rt. 2.55 min, 99.61% (Max). **HPLC:** (Method B) Rt. 5.35 min, 99.92% (Max). **Chiral HPLC:** (Method D) Rt. 3.28 min, 99.26% (Max).

Example 27

3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-ethoxypropanoic acid



10

To a stirred solution of methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-ethoxypropanoate (Intermediate 59; 30 mg, 0.06 mmol) in 1,4-dioxane (1 mL) at 0 °C, lithium hydroxide (4.69 mg, 0.11 mmol) was added and the reaction mixture was stirred for 30 minutes at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was quenched with dilute HCl (1.5 N, 1 mL) and the aqueous layer was extracted with EtOAc (2 x 5 mL). The combined organic layer was washed with water (5 mL), brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by Prep HPLC (Method A) to afford the title compound. **Yield:** 13% (4 mg, off-white solid).

15


20

¹H NMR (400 MHz, DMSO-*d*₆): δ 13.21 (s, 1H), 7.38 (s, 1H), 7.15 (t, *J* = 8.00 Hz, 2H), 7.05 (s, 1H), 6.70 (t, *J* = 4.00 Hz, 1H), 6.58 (d, *J* = 6.80 Hz, 2H), 4.43-4.41 (m, 1H), 4.33-4.25 (m, 2H), 4.03-3.87 (m, 2H), 3.74-3.70 (m, 1H), 3.58-3.54 (m, 1H), 3.23-3.11 (m, 1H), 2.51 (s, 3H), 2.34 (s, 3H), 1.61-1.48 (m, 2H), 1.36-1.30 (m, 4H), 1.17-1.13 (m, 3H), 0.94-0.90 (m, 3H). **LCMS:** (Method E) 523.0 (M⁺+H) Rt. 2.89 min, 95.56% (Max). **HPLC:** (Method B) Rt. 5.59 min, 93.21 % (Max).

25

Example 28


3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)-2-hydroxypropanoic acid



To a stirred solution of methyl 3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)-2-hydroxypropanoate (Intermediate 60; 0.17 g, 0.34 mmol) in 1,4-dioxane (2 mL) at 0 °C, dilute HCl (6 N, 2 mL) was added dropwise and the reaction mixture was heated for 3 hours at 80 °C. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with water (5 mL) and the aqueous layer was extracted with EtOAc (2 x 5 mL). The combined organic layer was washed with water (5 mL), brine (5 mL), and the organic part was dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by Isolera column chromatography (eluent: 3-10% MeOH in DCM; silica gel: 230-400 mesh) to afford the title compound. **Yield:** 30% (50 mg, light brown solid). **¹H NMR** (400 MHz, DMSO-*d*₆): δ 7.42 (s, 1H), 7.14 (t, *J* = 8.00 Hz, 2H), 7.04 (s, 1H), 6.67 (t, *J* = 7.60 Hz, 1H), 6.55 (d, *J* = 8.00 Hz, 2H), 4.37-4.32 (m, 1H), 4.27-4.19 (m, 3H), 3.43-3.39 (m, 1H), 3.33-3.20 (m, 1H), 3.08-3.02 (m, 1H), 2.34 (s, 3H), 2.33-2.27 (m, 1H), 1.38-1.35 (m, 4H), 1.33-1.25 (m, 3H), 0.91-0.88 (m, 3H). **LCMS:** (Method E) 480.1 (M⁺+H) Rt. 2.68 min, 98.95% (Max). **HPLC:** (Method B) Rt. 4.91 min, 97.58% (Max).

Example 29

3-((3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)propanoic acid




To a stirred solution of 3-ethyl-8-hydroxy-7-(methylthio)-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepine 1,1-dioxide (0.2 g, 0.55 mmol) in THF (5 mL) at 0 °C, KO^tBu (0.07 g, 0.60 mmol) and oxetan-2-one (0.04 g, 0.60 mmol) were added and the reaction mixture was stirred for 3 hours at 50 °C. As TLC showed incomplete conversion, more oxetan-2-one (0.04 g, 0.06 mmol) was added and the reaction mixture was stirred for 16 hours at 50 °C. After completion of the reaction, the reaction mixture was acidified with dilute HCl (1.5 N, 5 mL) and the aqueous layer was extracted with EtOAc (2

x 15 mL). The combined organic layer was washed with water (5 mL), brine (5 mL), and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by Isolera column chromatography (eluent: 45% EtOAc:PE; silica gel: 230-400 mesh). The obtained material was re-purified by prep-HPLC purification (Method A) to afford the title compound. **Yield:** 6.2% (15 mg, light brown solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 7.39 (s, 1H), 7.17-7.13 (m, 2H), 7.06 (s, 1H), 6.68 (t, *J* = 7.20 Hz, 1H), 6.57 (d, *J* = 8.00 Hz, 2H), 4.31-4.25 (m, 3H), 3.47-3.43 (m, 1H), 3.43-3.26 (m, 1H), 3.10-3.04 (m, 1H), 2.72-2.68 (m, 2H), 2.34 (s, 3H), 2.24 (s, 1H), 1.48-1.38 (m, 2H), 0.99 (t, *J* = 7.60 Hz, 3H). **LCMS:** (Method A) 436.1(M⁺+H) Rt. 2.10 min, 98.88% (Max). **HPLC:** (Method B) Rt. 4.33 min, 99.77% (Max).

Example 30

3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid




To a stirred solution of methyl 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoate (Intermediate 57; 55 mg, 0.10 mmol) in a mixture of 1,4-dioxane and water (2:1, 3 mL) at 0 °C, lithium hydroxide (10 mg, 0.21 mmol) was added and the reaction mixture was stirred for 1 hour at room temperature. After completion of the reaction (monitored by TLC), the reaction mixture was acidified with dilute HCl (1.5 N, pH~4) and diluted with ice-cold water (2 mL). The aqueous layer was extracted with EtOAc (2 x 5 mL). The combined organic layer was washed with water (5 mL), brine (5 mL) and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum to afford the title compound. **Yield:** 28% (15 mg, white solid).

¹H NMR (400 MHz, DMSO-*d*₆): δ 13.02 (bs, 1H), 7.34 (s, 1H), 7.15 (t, *J* = 8.0 Hz, 2H), 7.05 (s, 1H), 6.70 (t, *J* = 7.2 Hz, 1H), 6.58 (d, *J* = 8.0 Hz, 2H), 4.43 (t, *J* = 3.2 Hz, 1H), 4.37-4.31 (m, 1H), 4.19-4.15 (m, 1H), 4.04-4.01 (m, 1H), 4.00-3.90 (m, 1H), 3.43 (s, 3H), 3.20-3.19 (m, 1H), 2.45 (s, 3H), 2.35 (s, 3H), 1.62-1.60 (m, 1H), 1.52-1.50 (m, 1H), 1.36-1.31 (m, 4H), 0.93-0.92 (m, 3H). **LCMS:** (Method E) 508.9 (M⁺+H), Rt. 2.87 min, 96.57% (Max). **HPLC:** (Method B) Rt. 5.44 min, 98.31% (Max).

Example 31

3-(((S)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid and 3-(((R)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid




Stereoisomers 1 and 2 of the title compound were prepared from stereoisomer 1 (0.15 g) and stereoisomer 2 (0.1 g) of Intermediate 58, respectively, following the same procedure as described for Example 30. After work-up of the reaction mixtures, the crude materials were purified by Isolera column chromatography (eluent: 0 to 30% EtOAc/ PE; silica gel: 230-400 mesh) to afford the title compounds. The absolute configuration of the two stereoisomers is not known.

Stereoisomer 1: **Yield:** 71.3% (0.11 g, brown solid). **¹H-NMR** (400 MHz, DMSO-*d*₆): δ 12.64 (s, 1H), 7.35 (s, 1H), 7.15 (t, *J* = 8.00 Hz, 2H), 7.06 (s, 1H), 6.70 (t, *J* = 7.60 Hz, 1H), 6.58 (d, *J* = 8.40 Hz, 2H), 4.44-4.41 (m, 2H), 4.37-4.34 (m, 1H), 4.19-4.19 (m, 1H), 4.03-4.00 (m, 1H), 3.43 (s, 3H), 3.18-3.17 (m, 1H), 2.46 (s, 3H), 2.35-2.33 (m, 3H), 1.60-1.62 (m, 2H), 1.36-1.31 (m, 4H), 0.94-0.90 (m, 3H). **LCMS:** (Method E) 509.2 (M⁺+H) Rt. 2.64 min, 94.06% (Max). **HPLC:** (Method B) Rt. 5.33 min, 94.60% (Max).

Stereoisomer 2: **Yield:** 66.7% (0.07 g, white solid). **¹H-NMR** (400 MHz, DMSO-*d*₆): δ 13.03 (s, 1H), 7.35 (s, 1H), 7.15 (t, *J* = 8.00 Hz, 2H), 7.06 (s, 1H), 6.70 (t, *J* = 7.20 Hz, 1H), 6.58 (d, *J* = 8.40 Hz, 2H), 4.44-4.39 (m, 2H), 4.33-4.33 (m, 1H), 4.19-4.04 (m, 1H), 4.00-3.90 (m, 1H), 3.38 (s, 3H), 3.19-3.17 (m, 1H), 2.46 (s, 3H), 2.33 (s, 3H), 1.62-1.51 (m, 2H), 1.36-1.31 (m, 4H), 0.94-0.92 (m, 3H). **LCMS:** (Method A) 509.0 (M⁺+H) Rt. 2.90 min, 99.65% (Max). **HPLC:** (Method B) Rt. 5.29 min, 99.81% (Max).

Example 32

(S)-3-(((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid and (R)-3-(((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid (individual diastereomers)




The two diastereomers of stereoisomer 2 of Example 31 were separated by SFC (Method H). The material was concentrated under vacuum at 40 °C. The first eluting fraction corresponded to diastereomer 1 and the second eluting fraction corresponded to diastereomer 2. The absolute configuration of the two diastereomers is not known.

Diastereomer 1: **Yield:** 15.37% (0.01 g, white solid). **¹H-NMR** (400 MHz, DMSO-*d*₆): δ 7.34 (s, 1H), 7.15 (t, *J* = 7.20 Hz, 2H), 7.05 (s, 1H), 6.70 (t, *J* = 6.80 Hz, 1H), 6.58 (d, *J* = 8.00 Hz, 2H), 4.48-4.42 (m, 1H), 4.22 (m, 1H), 4.03-3.89 (m, 3H), 3.55-3.51 (m, 1H), 3.33 (s, 3H), 2.51 (s, 3H), 2.35 (s, 3H), 1.63-1.60 (m, 2H), 1.36-1.31 (m, 4H), 0.94-0.86 (m, 3H). **LCMS:** (Method E) 508.9 (M⁺+H) Rt. 2.91 min, 98.24% (Max). **HPLC:** (Method B) Rt. 5.30 min, 99.93% (Max). **Chiral HPLC** (Method H) Rt. 4.33 min, 100% (Max).

Diastereomer 2: **Yield:** 22.96% (0.02 g, white solid). **¹H-NMR** (400 MHz, DMSO-*d*₆): δ 7.33 (s, 1H), 7.15 (t, *J* = 7.60 Hz, 2H), 7.05 (s, 1H), 6.69 (t, *J* = 7.20 Hz, 1H), 6.58 (d, *J* = 8.40 Hz, 2H), 4.48-4.42 (m, 1H), 4.29-4.21 (m, 1H), 3.99-3.90 (m, 3H), 3.45-3.40 (m, 1H), 3.23 (s, 3H), 2.46 (s, 3H), 2.35 (s, 3H), 1.63-1.51 (m, 2H), 1.50-1.35 (m, 4H), 0.94-0.86 (m, 3H). **LCMS:** (Method A) 508.9 (M⁺+H) Rt. 2.91 min, 98.61% (Max). **HPLC:** (Method B) Rt. 5.30 min, 99.48% (Max). **Chiral HPLC** (Method H) Rt. 5.41 min, 99.23% (Max).

20 Example 33

3-((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-2-methyl-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid



To a stirred solution of methyl 3-((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-2-methyl-1,1-dioxido-

2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoate (Intermediate 63; 55 mg,

0.10 mmol) in 1,4-dioxane at 0 °C, dilute HCl (6 N, 3 mL) was added dropwise and the reaction mixture was heated for 16 hours at 80 °C. After completion of the reaction (monitored by TLC), the reaction mixture was diluted with cold water (5 mL) and the aqueous layer was extracted with EtOAc (2 x 5 mL). The combined organic layer was washed with water (5 mL), brine (5 mL), and dried over anhydrous Na₂SO₄. The organic part was filtered, concentrated under vacuum and the resulting crude material was purified by Prep HPLC (Method C) to afford the title compound. **Yield:** 18.11% (0.01 g, white solid).

¹H-NMR (400 MHz, DMSO-*d*₆): δ 12.84 (s, 1H), 7.34 (s, 1H), 7.06 (s, 1H), 6.99 (t, *J* = 8.80 Hz, 2H), 6.60-6.57 (m, 2H), 5.62-5.60 (m, 1H), 4.32-4.29 (m, 1H), 4.27-4.25 (m, 2H), 4.01-3.97 (m, 1H), 3.86-3.85 (m, 1H), 3.23-3.22 (m, 1H), 2.91-2.85 (m, 2H), 2.52 (s, 3H), 1.62-1.58 (m, 1H), 1.58-1.50 (m, 1H), 1.35-1.34 (m, 4H), 1.24-1.17 (m, 3H), 0.91 (t, *J* = 6.80 Hz, 3H). **LCMS:** (Method A) 527.1 (M⁺+H) Rt. 2.42 min, 93.98% (Max). **HPLC:** (Method B) Rt. 5.31 min, 95.21% (Max).

BIOLOGICAL ASSAYS

IBAT (h/m) assay protocol

10,000 cells (Human or Mouse IBAT-overexpressing cells) were seeded in 96-wells plate (Corning CLS3809) in 200 μL MEM-alpha medium (Gibco 12571-063) supplemented with 10% FBS (Gibco 10438026) containing Puromycin (Gibco A1113803) (10 μg/mL) and incubated at 37 °C in 5% CO₂ for 48 hours. After incubation, media was decanted from the wells and cells were washed two times with 300 μL of basal MEM-alpha medium (FBS-free). After decanting basal MEM-alpha medium each time, plates were tapped against paper towel to ensure maximum removal of residual media.

Test inhibitor dilutions (highest test concentration being 10 μM, 3-fold serial dilution, 10 points) prepared in DMSO (Sigma D2650) were added in incubation mix (maintaining 0.2% final DMSO concentration) containing 0.25 μM 3H-taurocholic acid (ARC ART-1368) and 5 μM of cold taurocholic acid (Sigma T4009). 50 μL of incubation mix containing test inhibitors was then added to the wells (in duplicate) and the plates were incubated for 20 minutes in a CO₂ incubator at 37 °C. After incubation, the reaction was stopped by keeping the plates on ice water mix for 2-3 minutes and then the incubation mix was aspirated completely from the wells. The wells were washed two times with 250 μL of chilled unlabelled 1 mM taurocholic acid dissolved in HEPES (Gibco 15630080)-buffered (10 mM) HBSS (Gibco 14175079) (pH 7.4). The plates were tapped against a paper towel after every wash to ensure maximum removal of blocking buffer.

100 μ L of MicroScint-20 (PerkinElmer 6013621) was added to the wells and kept overnight at room temperature before reading the plates in TopCount NXT™ Microplate Scintillation and Luminescence Counter from PerkinElmer under 3H Test protocol (set at 120 seconds reading time per well).

5 *LBAT (h/m) assay protocol*

20,000 cells (Human or Mouse LBAT-overexpressing cells) were seeded in 96-wells plate (Corning CLS3809) in 100 μ L MEM-alpha medium (Gibco 12571-063) supplemented with 10% FBS (Gibco 10438026) containing Geneticin (Gibco 10131-027) (1 mg/mL) and incubated at 37 °C in 5% CO₂ for
10 24 hours. After incubation, media was decanted from the wells and cells were washed two times with 300 μ L of basal MEM-alpha medium (FBS-free). After decanting basal MEM-alpha medium each time, plates were tapped against paper towel to ensure maximum removal of residual media. For human LBAT, incubation mix was prepared by adding test inhibitor dilutions (3-fold serial dilution in DMSO (Sigma D2650), 10 points) in MEM-alpha (without FBS) containing 0.3 μ M 3H-taurocholic
15 acid (ARC ART-1368) and 7.5 μ M cold taurocholic acid (Sigma T4009) (maintaining 0.2% final DMSO concentration). For mouse LBAT, incubation mix was prepared by adding test inhibitor dilutions (3-fold serial dilution in DMSO, 10 points) in MEM-alpha (without FBS) containing 0.3 μ M 3H-taurocholic acid and 25 μ M cold taurocholic acid maintaining 0.2% final DMSO concentration).

50 μ L of incubation mix containing test inhibitors was then added to the wells (in duplicate) and the
20 plates were incubated for 20 minutes in a CO₂ incubator at 37 °C. After incubation, the reaction was stopped by keeping the plates on ice water mix for 2-3 minutes and then the incubation mix was aspirated completely from the wells. The wells were washed two times with 250 μ L of chilled unlabelled 1 mM taurocholic acid dissolved in HEPES (Gibco 15630080)-buffered (10 mM) HBSS (Gibco 14175079) (pH 7.4). The plates were tapped against a paper towel after every wash to ensure
25 maximum removal of blocking buffer.

100 μ L of MicroScint-20 (PerkinElmer 6013621) was added to the wells and kept overnight at room temperature before reading the plates in TopCount NXT™ Microplate Scintillation and Luminescence Counter from PerkinElmer under 3H Test protocol (set at 120 seconds reading time per well, with normal plate orientation).

30

Bidirectional permeability assay (Caco-2 cells)

Caco-2 cells (Evotec) were seeded at a density of 70,000 cells/well in Millicell® 24-well cell culture insert plates and maintained in an incubator (37 °C, 5% CO₂, 95% RH) for 21 days with media change
35 on alternate days.

Stock solutions (10 mM) of the test compounds, atenolol (low permeability marker), propranolol (high permeability marker) and digoxin (substrate for P-gp transport pathway) were prepared in dimethylsulfoxide (DMSO). An intermediate stock solution (1 mM) was prepared by diluting 10 μ L of 10 mM master stock solution with 90 μ L of neat DMSO. A working stock solution (10 μ M) was prepared by diluting 50 μ L of 1 mM with 4950 μ L of FaSSIF buffer. Post addition of compounds to the FaSSIF, samples were subjected to sonication for 2 hours, and centrifuged at 4000 RPM for 30 minutes at 37 °C. The 4 mL of resultant supernatant was directly used in the assay. The final DMSO concentration in the transport experiments was 1%.

On the day of assay, Caco-2 monolayers were washed twice with transport buffer (HBSS, pH 7.4) and pre-incubated for 30 min (37 °C, 5% CO₂, 95% RH) in an incubator. The electrical resistance of the monolayers was measured with a Millicell® - ERS system. Monolayers with trans-epithelial electrical resistance (TEER) values greater than 350 ohm.cm² were selected for the assay.

The assay was conducted in absorptive direction (A2B) and secretory (B2A) directions. Transport experiments were initiated by addition of transport assay buffer (FaSSIF buffer prepared in HBSS) consisting of compounds to the donor compartment (apical chamber A-B; basolateral chamber B-A) in duplicate (n=2) wells. Drug free HBSS buffer (pH 7.4) containing 1% bovine serum albumin (BSA) was introduced to the receiver (A-B-basolateral; B-A- Apical) compartments. The volumes of apical and basolateral compartments were 0.4 and 0.8 mL, respectively. After adding dosing solution, plates were incubated in an incubator for 120 minutes at 37 °C. After 120 minutes, donor and receiver samples were collected and matrix matched (1:1, 30 μ L study sample + 30 μ L blank buffer) with the opposite buffer. Dosing samples matrix matched (1:1, 30 μ L study sample + 30 μ L blank buffer) with the opposite buffer. Samples were processed by adding acetonitrile containing internal standard (60 μ L study sample + 200 μ L acetonitrile containing internal standard -Tolbutamide, 500 ng/mL).

Samples were vortexed and centrifuged at 4000 rpm for 10 minutes. The obtained supernatant (100 μ L) was diluted with 100 μ L of water and transferred to fresh 96 well plates. The concentration of compounds in the samples was analyzed by liquid chromatography tandem mass spectrometry (LC-MS/MS) method using discovery grade bio-analytical method, as applicable.

The mean apparent permeability (P_{app} , $\times 10^{-6}$ cm/sec) of the test compounds, atenolol, propranolol and digoxin were calculated as follows:

$$P_{app} = \frac{dq}{dt} \times \frac{1}{C_0} \times \frac{1}{A}$$

where dq/dt = rate of transport (rate of transport of compound in the receiver compartment), C_0 = initial concentration in the donor compartment, A = surface area of the effective filter membrane.

HepaRG-based assay protocol

A cryopreserved vial of differentiated HepaRG cells (Biopredic International HPR116080) is thawed in HepaRG Thawing/Plating/General Purpose Medium (Biopredic International ADD670C) supplemented with 200 mM Glutamax (Gibco 35050061) following the protocol provided by Biopredic International. 70,000 cells per well are seeded in 96-wells plate (Corning CLS3809) in 100 μ L of HepaRG Thawing/Plating/General Purpose Medium supplemented with 200 mM Glutamax and incubated at 37 °C in 5% CO₂ for 24 hours. Post incubation, the seeding media is replaced by HepaRG Maintenance/Metabolism Medium (Biopredic International ADD620C) and incubated for 6 days, with fresh HepaRG Maintenance/Metabolism Medium replenishment every 48 hours. After 7 days incubation post seeding, incubation media is decanted from the wells and cells are washed two times with 250 μ L of William's E Basal Media (Gibco 12551032). After decanting William's E Basal Media each time, plates are tapped against paper towel to ensure maximum removal of residual media. Incubation mix is prepared by adding test inhibitor dilutions (3-fold serial dilution in DMSO (Sigma D2650)) in William's E media (basal) containing 0.3 μ M 3H-taurocholic acid (ARC ART-1368) and 7.5 μ M cold taurocholic acid (Sigma T4009) (maintaining 0.2% final DMSO concentration). 50 μ L of incubation mix containing test inhibitors is then added to the wells (in duplicate) and the plates are incubated for 30 minutes in 5% CO₂ incubator at 37 °C. After incubation, the reaction is stopped by keeping the plates on ice water mix for 2-3 minutes and the incubation mix is then aspirated completely from the wells. The wells are washed two times with 250 μ L of chilled unlabelled 1 mM taurocholic acid dissolved in HEPES (Gibco 15630080)-buffered (10mM) HBSS (Gibco 14175079) (pH 7.4). The plates are tapped against a paper towel after every wash to ensure maximum removal of blocking buffer. 100 μ L of MicroScint-20 (PerkinElmer 6013621) is added to the wells and kept overnight at room temperature before reading the plates in TopCount NXT™ Microplate Scintillation and Luminescence Counter from PerkinElmer under 3H Test protocol (set at 120 seconds reading time per well, with normal plate orientation).

Preparation of test compound dilutions

All test compounds were provided in powder form at room temperature. 10 mM DMSO stocks of the test compounds were prepared, aliquoted and stored at -20 °C. From the 10 mM DMSO stock of the compounds, a 3-fold serial dilution in DMSO was prepared to get a total of 10 dilutions of the test compounds. 0.5 μ L of this dilution in DMSO was added to 250 μ L of FBS-free basal media containing 3H-taurocholic acid and cold taurocholic acid to prepare the incubation mixture.

Bioavailability studies

Male mice (C57BL/6 or CD1) or Wistar rats of 8-9 weeks old were used. For each test compound, two groups of 3 animals each were used. One group was administered a single intravenous dose of 1 mg/kg (vehicle 100% DMSO) through the tail vein and the other group was administered a single oral dose of 10 mg/kg through gavage needle. The group that was administered an oral dose was fasted overnight. Blood samples were collected after 0.083, 0.25, 0.5, 1, 2, 4, 6, 8 and 24 hours following intravenous administration, and after 0.25, 0.5, 1, 2, 4, 6, 8 and 24 hours following oral administration. Blood samples were taken from saphenous vein. 0.2% EDTA was used as the anticoagulant. The samples were analyzed by a discovery grade bioanalytical method developed for the estimation of test compound in plasma, using an LC-MS/MS system.

Results

15

Biological data for the compounds of the examples is shown in Table 8 below.

Table 8

Example	hLBAT IC ₅₀ (nM)	hIBAT IC ₅₀ (nM)	Permeability (Caco-2)		Bioavailability (%)
			P _{app} A2B (x 10 ⁻⁶ cm/sec)	P _{app} B2A (x 10 ⁻⁶ cm/sec)	
1	9.8				
2, enantiomer 1	2278	10000			
2, enantiomer 2	2.3	682	13.2	15.2	100
3	13	274	20.8	12.9	
4, enantiomer 1	673	10000			
4, enantiomer 2	7.4	294	18.8	13.0	17
5	8.9	615	9.4	7.2	
6, enantiomer 1	536	10810			
6, enantiomer 2	6.2	290	8.6	6.5	
7	25	405	8.1	3.8	
8, enantiomer 1	5.0	166	15.8	5.0	40.0
8, enantiomer 2	591	4971			
9	5.3	2697			

Example	hLBAT IC ₅₀ (nM)	hIBAT IC ₅₀ (nM)	Permeability (Caco-2)		Bioavailability (%)
			P _{app} A2B (x 10 ⁻⁶ cm/sec)	P _{app} B2A (x 10 ⁻⁶ cm/sec)	
10	10	678			
11	4.5	167			
12, <i>enantiomer 1</i>	9.4	3007			
12, <i>enantiomer 2</i>	8.5	96	12.9	9.1	33
13	3.3	360	10.8	8.6	
14, <i>enantiomer 1</i>	918	1949			
14, <i>enantiomer 2</i>	3	193	12.2	7.3	100
15	4.2	>3333			
16	3.4	464			
17, <i>enantiomer 1</i>	2.8	234	7.7	10.6	11
17, <i>enantiomer 2</i>	131	1254			
18	3.3	460			
19, <i>enantiomer 1</i>	1.3	274	7.2	7.0	
19, <i>enantiomer 2</i>	106	>3333			
20	1.1	259			
21, <i>enantiomer 1</i>	>333	>10000			
21, <i>enantiomer 2</i>	0.6	203			
22	28	428	11.7	21.5	
23	14	257	10.6	30.2	
24, <i>diastereomer 1</i>	8.4	1030	8.6	15.9	
24, <i>diastereomer 2</i>	16	412	16.3	24.0	
25	93				
26, <i>enantiomer 1</i>	59				
28	187				
30	64				
31, <i>stereoisomer 1</i>	5682				
31, <i>stereoisomer 2</i>	22	2093			
32, <i>diastereomer 1</i>	22	2404	5.2	15.9	
32, <i>diastereomer 2</i>	32	4784			
33	6	117	2.2	4.3	

PD model: Evaluation of test compound on total bile acids levels in male C57BL6 mice.


C57BL/6N Tac mice of 8-9 weeks old are used to study the effect of bile acid modulators on bile acid levels. After completion of quarantine and acclimatization period, animals are randomized based on
5 bodyweight into x experimental groups: (i) vehicle control, and (ii) test compound y mg/kg po once daily. Animals are treated with test compound for 7 days. On day 5 of the study, animals are individually housed in fresh cages. On day 7, feces are collected from each cage, followed by blood withdrawal from each animal through retro-orbital route. Animals are euthanized to collect liver and terminal ileum from each animal for further analysis. Bodyweight and food consumption are
10 measured twice weekly. Serum lipid profiles are analyzed in serum samples of day 7. Total bile acids in serum is measured in the serum samples of day 7. Fecal bile excretion is measured in the fecal sample of day 7. Hepatic expression of CYP7A1 and SHP are quantified in the liver samples of day 7. Liver triglycerides and total cholesterol are analyzed in the liver samples of day 7.

15 *Urine bile acid model: Evaluation of test compounds on urine bile acid levels in male C57BL/6N mice.*

C57BL/6N Tac mice of 8-9 weeks old are used to study the effect of bile acid modulators on bile acid levels. After completion of quarantine and acclimatization period, animals are randomized based on
20 bodyweight into x experimental groups: (i) vehicle control, and (ii) test compound y mg/kg po once daily. Animals are treated with test compound for 7 days. On day 6 of the study, animals are transferred to a metabolic cage. On day 7, feces and urine are collected from each metabolic cage, followed by blood withdrawal from each animal through retro-orbital route. Animals are euthanized to collect kidney from each animal for further analysis. Bodyweight is measured twice weekly. Total
25 bile acids in serum is measured in serum samples of day 7. Fecal bile acid excretion is measured in the fecal sample of day 7. Urine excretion of bile acids is measured in the sample of day 7. Kidney expression of ASBT, OSTa, OSTAb and MRP2 is quantified in the samples of day 7.

CLAIMS

1. A compound of formula (I)



5

wherein

M is selected from $-CH_2-$ and $-NR^5-$;

10

R^1 is C_{1-4} alkyl;

R^2 is independently selected from the group consisting of hydrogen, halogen, hydroxy, C_{1-4} alkyl, C_{1-4} haloalkyl, C_{1-4} alkoxy, C_{1-4} haloalkoxy, cyano, nitro, amino, $N-(C_{1-4}$ alkyl)amino, N,N -di(C_{1-4} alkyl)amino, C_{1-6} alkylcarbonylamino, C_{3-6} cycloalkylcarbonylamino, $N-(C_{1-4}$ alkyl)-aminocarbonyl, N,N -di(C_{1-4} alkyl)aminocarbonyl, C_{1-4} alkyloxycarbonylamino, C_{3-6} cycloalkyl-oxycarbonylamino, C_{1-4} alkylsulfonamido and C_{3-6} cycloalkylsulfonamido;

15

n is an integer 1, 2 or 3;

R^3 is selected from the group consisting of hydrogen, halogen, cyano, C_{1-4} alkyl, C_{3-6} cycloalkyl, C_{1-4} alkoxy, C_{3-6} cycloalkyloxy, C_{1-4} alkylthio, C_{3-6} cycloalkylthio, amino, $N-(C_{1-4}$ alkyl)amino and N,N -di(C_{1-4} alkyl)amino;

20

R^{4A} and R^{4B} are each independently selected from the group consisting of hydrogen, halogen, hydroxy, C_{1-4} alkyl and C_{1-4} alkoxy; or R^{4A} and R^{4B} , together with the carbon atom to which they are attached, form a 3- to 5-membered saturated carbocyclic ring;

R^{4C} and R^{4D} are each independently selected from the group consisting of hydrogen and C_{1-4} alkyl; and

25

R^5 is selected from the group consisting of hydrogen and C_{1-4} alkyl; or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1, wherein R^1 is C_{2-4} alkyl.

3. A compound according to claim 1 or 2, wherein R¹ is *n*-propyl or *n*-butyl.
4. A compound according to any one of claims 1 to 3, wherein R² is independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, hydroxy, methoxy, amino,
5 methylamino and dimethylamino.
5. A compound according to any one of claims 1 to 4, wherein R³ is selected from the group consisting of fluoro, chloro, bromo, methyl, cyclopropyl, methoxy, ethoxy, methylthio, ethylthio, amino, methylamino and dimethylamino.
- 10 6. A compound according to any one of claims 1 to 5, wherein R^{4A} and R^{4B} are each independently selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄ alkyl and C₁₋₄ alkoxy, or R^{4A} and R^{4B}, together with the carbon atom to which they are attached, form a cyclopropyl ring.
- 15 7. A compound according to any one of claims 1 to 6, wherein R^{4C} and R^{4D} are each independently hydrogen or methyl.
8. A compound according to any one of claims 1 to 7, wherein R⁵ is hydrogen or methyl.
- 20 9. A compound according to claim 1, selected from the group consisting of:
3-((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;
(S)-3-((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;
25 (R)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;
1-(((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;
30 (S)-1-(((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;
(R)-1-(((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;
35 3-((3-Butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

(S)-3-((3-Butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

(R)-3-((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

5 1-(((3-Butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

(S)-1-(((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

10 (R)-1-(((3-butyl-7-(ethylthio)-2-methyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

3-((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

15 1-(((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

(S)-1-(((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

20 (R)-1-(((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

3-((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

(S)-3-((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

25 (R)-3-((3-butyl-5-(4-fluorophenyl)-2-methyl-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-dimethylpropanoic acid;

1-(((3-butyl-7-(ethylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

30 1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

(S)-1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

(R)-1-(((3-butyl-5-(4-fluorophenyl)-7-(methylthio)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

1-(((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

(S)-1-(((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

5 (R)-1-(((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)methyl)cyclopropane-1-carboxylic acid;

3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-difluoropropanoic acid;

10 (S)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-difluoropropanoic acid;

(R)-3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2,2-difluoropropanoic acid;

3-((3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid;

15 (S)-3-(((R)-3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid;

(S)-3-(((S)-3-Butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid;

20 (R)-3-(((R)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid;

(R)-3-(((S)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxy-2-methylpropanoic acid;

3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)propanoic acid;

25 (S)-3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)propanoic acid;

(R)-3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)propanoic acid;

30 3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-ethoxypropanoic acid;

3-((3-butyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)-2-hydroxypropanoic acid;

3-((3-ethyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,5-benzothiazepin-8-yl)oxy)propanoic acid;

3-((3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid;

3-(((S)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid;

5 3-(((R)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid;

(S)-3-(((R)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid;

10 (R)-3-(((R)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid;

(S)-3-(((S)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid;

(R)-3-(((S)-3-butyl-2-methyl-7-(methylthio)-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-methoxypropanoic acid; and

15 3-((3-butyl-7-(ethylthio)-5-(4-fluorophenyl)-2-methyl-1,1-dioxido-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepin-8-yl)oxy)-2-hydroxypropanoic acid;
or a pharmaceutically acceptable salt thereof.

10. A pharmaceutical composition comprising a therapeutically effective amount of a compound
20 according to any one of claims 1 to 9, and one or more pharmaceutically acceptable excipients.
11. The compound according to any one of claims 1 to 9, for use as a medicament.
12. The compound according to any one of claims 1 to 9, for use in the treatment or prevention of
25 a cardiovascular disease or a disorder of fatty acid metabolism or a glucose utilization disorder, such as hypercholesterolemia; disorders of fatty acid metabolism; type 1 and type 2 diabetes mellitus; complications of diabetes, including cataracts, micro- and macrovascular diseases, retinopathy, neuropathy, nephropathy and delayed wound healing, tissue ischaemia, diabetic foot, arteriosclerosis, myocardial infarction, acute coronary syndrome, unstable angina
30 pectoris, stable angina pectoris, stroke, peripheral arterial occlusive disease, cardiomyopathy, heart failure, heart rhythm disorders and vascular restenosis; diabetes-related diseases such as insulin resistance (impaired glucose homeostasis), hyperglycemia, hyperinsulinemia, elevated blood levels of fatty acids or glycerol, obesity, dyslipidemia, hyperlipidemia including hypertriglyceridemia, metabolic syndrome (syndrome X), atherosclerosis and hypertension;
35 and for increasing high density lipoprotein levels.

13. The compound according to any one of claims 1 to 9, for use in the treatment or prevention of a gastrointestinal disease or disorder, such as constipation (including chronic constipation, functional constipation, chronic idiopathic constipation (CIC), intermittent/sporadic constipation, constipation secondary to diabetes mellitus, constipation secondary to stroke, constipation secondary to chronic kidney disease, constipation secondary to multiple sclerosis, constipation secondary to Parkinson's disease, constipation secondary to systemic sclerosis, drug induced constipation, irritable bowel syndrome with constipation (IBS-C), irritable bowel syndrome mixed (IBS-M), pediatric functional constipation and opioid induced constipation); Crohn's disease; primary bile acid malabsorption; irritable bowel syndrome (IBS); inflammatory bowel disease (IBD); ileal inflammation; and reflux disease and complications thereof, such as Barrett's esophagus, bile reflux esophagitis and bile reflux gastritis.
14. The compound according to any one of claims 1 to 9, for use in the treatment or prevention of a liver disease or disorder, such as an inherited metabolic disorder of the liver; inborn errors of bile acid synthesis; congenital bile duct anomalies; biliary atresia; post-Kasai biliary atresia; post-liver transplantation biliary atresia; neonatal hepatitis; neonatal cholestasis; hereditary forms of cholestasis; cerebrotendinous xanthomatosis; a secondary defect of BA synthesis; Zellweger's syndrome; cystic fibrosis-associated liver disease; alpha1-antitrypsin deficiency; Alagilles syndrome (ALGS); Byler syndrome; a primary defect of bile acid (BA) synthesis; progressive familial intrahepatic cholestasis (PFIC) including PFIC-1, PFIC-2, PFIC-3 and non-specified PFIC, post-biliary diversion PFIC and post-liver transplant PFIC; benign recurrent intrahepatic cholestasis (BRIC) including BRIC1, BRIC2 and non-specified BRIC, post-biliary diversion BRIC and post-liver transplant BRIC; autoimmune hepatitis; primary biliary cirrhosis (PBC); liver fibrosis; non-alcoholic fatty liver disease (NAFLD); non-alcoholic steatohepatitis (NASH); portal hypertension; cholestasis; Down syndrome cholestasis; drug-induced cholestasis; intrahepatic cholestasis of pregnancy (jaundice during pregnancy); intrahepatic cholestasis; extrahepatic cholestasis; parenteral nutrition associated cholestasis (PNAC); low phospholipid-associated cholestasis; lymphedema cholestasis syndrome 1 (LCS1); primary sclerosing cholangitis (PSC); immunoglobulin G4 associated cholangitis; primary biliary cholangitis; cholelithiasis (gallstones); biliary lithiasis; choledocholithiasis; gallstone pancreatitis; Caroli disease; malignancy of bile ducts; malignancy causing obstruction of the biliary tree; biliary strictures; AIDS cholangiopathy; ischemic cholangiopathy; pruritus due to cholestasis or jaundice; pancreatitis; chronic autoimmune liver disease leading to progressive cholestasis; hepatic steatosis; alcoholic hepatitis; acute fatty liver; fatty liver of pregnancy;

drug-induced hepatitis; iron overload disorders; congenital bile acid synthesis defect type 1 (BAS defect type 1); drug-induced liver injury (DILI); hepatic fibrosis; congenital hepatic fibrosis; hepatic cirrhosis; Langerhans cell histiocytosis (LCH); neonatal ichthyosis sclerosing cholangitis (NISCH); erythropoietic protoporphyria (EPP); idiopathic adulthood ductopenia (IAD); idiopathic neonatal hepatitis (INH); non syndromic paucity of interlobular bile ducts (NSPILBD); North American Indian childhood cirrhosis (NAIC); hepatic sarcoidosis; amyloidosis; necrotizing enterocolitis; serum bile acid-caused toxicities, including cardiac rhythm disturbances (e.g., atrial fibrillation) in setting of abnormal serum bile acid profile, cardiomyopathy associated with liver cirrhosis ("cholecardia"), and skeletal muscle wasting associated with cholestatic liver disease; polycystic liver disease; viral hepatitis (including hepatitis A, hepatitis B, hepatitis C, hepatitis D and hepatitis E); hepatocellular carcinoma (hepatoma); cholangiocarcinoma; bile acid-related gastrointestinal cancers; and cholestasis caused by tumours and neoplasms of the liver, of the biliary tract and of the pancreas; or for use in the enhancement of corticosteroid therapy in liver disease.

- 15
15. The compound according to any one of claims 1 to 9, for use in the treatment or prevention of hyperabsorption syndromes (including abetalipoproteinemia, familial hypobetalipoproteinemia (FHBL), chylomicron retention disease (CRD) and sitosterolemia); hypervitaminosis and osteopetrosis; hypertension; glomerular hyperfiltration; polycystic kidney disease (PKD), including autosomal dominant polycystic kidney disease (ADPKD) and autosomal recessive polycystic kidney disease (ARPKD); and pruritus of renal failure; or for use in the protection against liver- or metabolic disease-associated kidney injury.
- 20