



HU000030888T2

(19) **HU****MAGYARORSZÁG****Szellemi Tulajdon Nemzeti Hivatala**(11) Lajstromszám: **E 030 888**(13) **T2**

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 13 166218**(22) A bejelentés napja: **2011. 03. 07.**(51) Int. Cl.: **A61K 35/74** (2006.01)**A61P 17/00** (2006.01)**A61P 37/00** (2006.01)

(96) Az európai bejelentés bejelentési száma:

EP 20110166218

(97) Az európai bejelentés közzétételi adatai:

EP 2626076 A1 **2013. 08. 14.**

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2626076 B1 **2016. 08. 24.**

(30) Elsőbbségi adatok:

MI20100375 **2010. 03. 08.** **IT**

(73) Jogosult(ak):

Probiotal S.p.a., 28100 Novara (NO) (IT)

(72) Feltaláló(k):

Mogna, Giovanni, 28100 Novara NO (IT)**Strozzi, Gian Paolo, 28100 Novara NO (IT)****Drago, Lorenzo, 28100 Novara NO (IT)****Mogna, Luca, 28100 Novara NO (IT)**

(74) Képviselő:

SBGK Szabadalmi Ügyvivői Iroda, Budapest

(54)

Lactobacillus salivarius (LS01) DSM 22775 terápiás alkalmazásai

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11)

EP 2 626 076 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
24.08.2016 Bulletin 2016/34

(51) Int Cl.:
A61K 35/74 ^(2015.01) **A61P 37/00** ^(2006.01)
A61P 17/00 ^(2006.01)

(21) Application number: **13166218.1**

(22) Date of filing: **07.03.2011**

(54) **Therapeutic uses of Lactobacillus salivarius LS01 DSM 22775**

Therapeutische Verwendungen von Lactobacillus salivarius LS01 DSM 22775

Utilisations thérapeutiques de Lactobacillus salivarius LS01 DSM 22775

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **08.03.2010 IT MI20100375**

(43) Date of publication of application:
14.08.2013 Bulletin 2013/33

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
11717732.9 / 2 544 698

(73) Proprietor: **Probiotal S.p.A.
28100 Novara (NO) (IT)**

(72) Inventors:
• **Mogna, Giovanni
28100 Novara NO (IT)**
• **Strozzi, Gian Paolo
28100 Novara NO (IT)**
• **Drago, Lorenzo
28100 Novara NO (IT)**
• **Mogna, Luca
28100 Novara NO (IT)**

(74) Representative: **Hoffmann Eitle
Patent- und Rechtsanwälte PartmbB
Arabellastraße 30
81925 München (DE)**

(56) References cited:
WO-A1-2008/153377 WO-A2-2005/039319

- **PROBIOTIC STUDY GROUP ET AL: "A differential effect of 2 probiotics in the prevention of eczema and atopy: A double-blind, randomized, placebo-controlled trial", JOURNAL OF ALLERGY AND CLINICAL IMMUNOLOGY, ELSEVIER, AMSTERDAM, NL, vol. 122, no. 4, 1 October 2008 (2008-10-01), pages 788-794, XP025493925, ISSN: 0091-6749, DOI: 10.1016/J.JACI.2008.07.011 [retrieved on 2008-08-31]**
- **KALLIOMAKIM ET AL: "Probiotics for prevention of atopic disease?", THE LANCET, LANCET LIMITED. LONDON, GB, vol. 362, no. 9382, 9 August 2003 (2003-08-09), page 496, XP004783945, ISSN: 0140-6736, DOI: 10.1016/S0140-6736(03)14087-1**
- **TAYLOR ET AL: "Probiotic supplementation for the first 6 months of life fails to reduce the risk of atopic dermatitis and increases the risk of allergen sensitization in high-risk children: A randomized controlled trial", JOURNAL OF ALLERGY AND CLINICAL IMMUNOLOGY, ELSEVIER, AMSTERDAM, NL, vol. 119, no. 1, 1 January 2007 (2007-01-01), pages 184-191, XP022128225, ISSN: 0091-6749, DOI: 10.1016/J.JACI.2006.08.036**
- **TANIUCHI S ET AL: "Administration of Bifidobacterium to infants with atopic dermatitis: Changes in fecal microflora and clinical symptoms", JOURNAL OF APPLIED RESEARCH 2005 US, vol. 5, no. 2, 2005, pages 387-396, XP002606412, ISSN: 1537-064X**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 626 076 B1

- SUNADAYOSUKE ET AL: "Effect of Lactobacillus acidophilus strain L-55 on the development of atopic dermatitis-like skin lesions in NC/Nga mice.", INTERNATIONAL IMMUNOPHARMACOLOGY 20 DEC 2008 LNKD-PUBMED:18790088, vol. 8, no. 13-14, 20 December 2008 (2008-12-20), pages 1761-1766, XP002606413, ISSN: 1567-5769
- BUYS LUCINDA M: "Treatment options for atopic dermatitis.", AMERICAN FAMILY PHYSICIAN 15 FEB 2007 LNKD- PUBMED:17323714, vol. 75, no. 4, 15 February 2007 (2007-02-15), pages 523-528, XP002606414, ISSN: 0002-838X
- L. NIERSE ET AL: "The effects of selected probiotic strains on the development of eczema (the Panda study)", ALLERGY, vol. 64, no. 9, 1 September 2009 (2009-09-01), pages 1349-1358, XP55069616, ISSN: 0105-4538, DOI: 10.1111/j.1398-9995.2009.02021.x
- CHRISTINA E. WEST ET AL: "Probiotics during weaning reduce the incidence of eczema", PEDIATRIC ALLERGY AND IMMUNOLOGY, vol. 20, no. 5, 1 August 2009 (2009-08-01) , pages 430-437, XP55069609, ISSN: 0905-6157, DOI: 10.1111/j.1399-3038.2009.00745.x
- JI YEUN KIM ET AL: "Effect of probiotic mix (Bifidobacterium bifidum, Bifidobacterium lactis, Lactobacillus acidophilus) in the primary prevention of eczema: a double-blind, randomized, placebo-controlled trial", PEDIATRIC ALLERGY AND IMMUNOLOGY, vol. 21, no. 2p2, 1 March 2010 (2010-03-01), pages e386-e393, XP55069608, ISSN: 0905-6157, DOI: 10.1111/j.1399-3038.2009.00958.x
- DATABASE BIOSIS [Online] BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US October 2011 DRAGO L ET AL: 'EFFECTS OF LACTOBACILLUS SALIVARIUS LS01 (DSM 22775) TREATMENT ON ADULT ATOPIC DERMATITIS: A RANDOMIZED PLACEBO-CONTROLLED STUDY' Database accession no. PREV201200185427
- DATABASE MEDLINE [Online] US NATIONAL LIBRARY OF MEDICINE (NLM), BETHESDA, MD, US October 2012 DRAGO LORENZO ET AL: 'Changing of fecal flora and clinical effect of L. salivarius LS01 in adults with atopic dermatitis.' Database accession no. NLM22955359
- DRAGO LORENZO ET AL: 'Strain-dependent release of cytokines modulated by Lactobacillus salivarius human isolates in an in vitro model' BMC RESEARCH NOTES, BIOMED CENTRAL LTD, GB vol. 3, no. 1, 44, 25 February 2010, pages 1 - 5, XP021070143 ISSN: 1756-0500

Description

[0001] The present invention is described in claims 1-12.

[0002] The present invention relates to a composition comprising probiotic bacteria for the treatment of pathologies associated with alterations of the immune system. In particular, the present invention relates to the use of selected probiotic bacteria for the preparation of a composition for the treatment of allergies, such as atopic dermatitis.

[0003] It is well known that Atopic Dermatitis (hereinafter AD for the sake of brevity) is a chronic relapsing skin disorder which begins in the neonatal period or childhood and may persist into adulthood. Therefore, the phases of AD are divided into neonatal, infantile and adult. AD is an inflammatory skin disease which, like asthma and allergic rhinitis, is associated with local infiltration of T lymphocytes that produce interleukin-4 (IL-4) and interleukin-5 (IL-5).

[0004] IL-4, as is well known, regulates the development of the T helper 2 (Th2) phenotype, with consequent hyperproduction of immunoglobulin (Ig) and eosinophilia. Elevated serum Ig E levels and skin tests positive for food and inhalant allergens can be demonstrated in 80-90% of subjects with atopic dermatitis.

[0005] AD is based both on genetic factors and on immunological factors, even though a large number of external factors can modify its expression. In 60% of the cases of AD a family history of atopy can be demonstrated; in other words, if one parent has an atopic diathesis there is a 60% likelihood that the children will be atopic.

[0006] Atopy is the genetic predisposition to developing localised anaphylactic reactions following contact, ingestion, inoculation or inhalation of allergens.

[0007] The immunological factors taken into consideration in the etiology of AD are: allergy to foods, allergy induced by contact allergens and irritants, allergy induced by aeroallergens and immunoregulatory abnormalities.

[0008] Among the multitude of treatments directed at curing and/or preventing allergies or skin inflammations, in particular, at treating atopic dermatitis, many drug-based treatments stand out. Drug-based treatments have several contraindications that limit their use. Moreover, these drugs do not seem well tolerated by some categories of patients.

[0009] However, at present there exists no treatment having absolute and definitive efficacy. It is possible only to adopt certain precautions against the onset of allergies or, as needed, remedies for mitigating the bothersome symptoms, such as itching and reddening of the skin.

[0010] Therefore, there remains a need to have a treatment which does not present the limitations of drug-based treatments and which can be administered to all categories of patients.

[0011] In particular, there remains a need to have a treatment which can serve to combat the onset of allergies and the development of skin inflammations and/or to reduce bothersome symptoms such as reddening of skin tissue, itching and eczema.

[0012] The Applicant has provided an answer to the above-mentioned needs since it has succeeded, after intense research activity, in selecting specific probiotic bacterial strains.

[0013] The subject matter of the present invention is a composition comprising a culture of probiotic bacteria, as claimed in the appended independent claim.

[0014] The subject matter of the present invention moreover relates to the use of one culture of probiotic bacteria, as claimed in the appended independent claim.

[0015] The Applicant set itself the objective of selecting specific bacterial strains for the purpose of evaluating their effect on the clinical course and quality of life of patients with moderate or severe AD and for the purpose of determining their influence on clinical and immunological parameters. Moreover, the Applicant set itself the objective of selecting bacterial strains with probiotic activity able to be tolerated by patients and capable of colonising the intestinal tract.

[0016] The diagnosis of moderate/severe AD was evaluated using a standardized severity index for Atopic Dermatitis - hereinafter indicated as AD - (SCORAD index = SCORing Atopic Dermatitis). Said index is known to those persons skilled in the art.

[0017] It was found that the bacterial strains of the present invention have a high immunomodulating effect and are thus useful for the treatment of pathologies associated with alterations of the immune system.

[0018] Following intense research activity, the Applicant selected, from among the many strains tested, the following probiotic bacterial strains:

1) *Lactobacillus salivarius* (LS01) DSM 22775, deposited on 23.07.2009, by the company Probiotical SpA of Novara - Italy;

2) *Bifidobacterium breve* (BR03) DSM 16604, deposited on 20.07.2004, by the company Probiotical SpA of Novara - Italy;

3) *Lactobacillus pentosus* (LPS01) DSM 21980, deposited on 14.11.2008, by the company Probiotical SpA of Novara - Italy;

4) *Streptococcus thermophilus* (FP4) DSM 18616, deposited on 13.09.2006, by the company Mofin Srl of Novara - Italy;

5) *Lactobacillus casei* ssp. *rhamnosus* (LR04) DSM 16605, deposited on 20.07.2004, by the company Probiotical

SpA of Novara - Italy; and

6) *Lactobacillus acidophilus* (LA02) DSM 21717, deposited on 06.08.2008, by the company Probiotal SpA of Novara - Italy.

[0019] The cultures listed above were all deposited with the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, (DSMZ) in Germany, in accordance with the Budapest Treaty.

[0020] The cultures of probiotic bacteria listed above can be in the form of live bacteria, dead bacteria or cellular components, cell extracts or lysates thereof.

[0021] In a preferred embodiment, the composition comprises the bacterial culture as claimed for use as an immunomodulating composition capable of modulating the immune system.

[0022] The expression "immunomodulating composition" means that the composition is capable of modulating the immune system in the sense that it is capable of stimulating/inducing several responses of the immune system, rendering it more reactive, for example by intervening through the production of specific cytokines. Advantageously, said immunomodulating composition induces the immune system toward the production of type 1 cytokines.

[0023] Advantageously, the bacterial strains and the corresponding bacterial cultures have demonstrated to be capable of improving clinical and immunological parameters. In particular, it was observed that they are capable of:

1) significantly improving the SCORAD index and DLQI;

2) reducing microbial translocation (plasma LPS levels) and activating CD8+ T lymphocytes;

3) increasing the percentage both of total regulatory T lymphocytes (Treg), and of the same expressing the markers TLR2- and TLR4;

4) improving the Th1/ Th2 and Th17/ Treg ratios.

[0024] Therefore, the composition of the present invention (immunomodulating composition) is capable of preventing, reducing and/or curing the pathologies associated with alterations of the immune system such as allergies, atopy, allergic rhinitis, food hypersensitivity, atopic dermatitis, eczema, asthma and immunodeficiencies.

[0025] In accordance with one embodiment, the present invention concerns a composition comprising or, alternatively, consisting of a culture of probiotic bacteria of *Lactobacillus salivarius* (LS01) DSM 22775.

[0026] In accordance with another preferred embodiment, the present invention concerns the use of a culture of probiotic bacteria of *Lactobacillus salivarius* (LS01) DSM 22775 for the preparation of a composition capable of preventing, reducing and/or curing the pathologies selected from the group consisting of allergies, atopy, allergic rhinitis, food hypersensitivity, atopic dermatitis, eczema, asthma and immunodeficiencies.

[0027] In accordance with another preferred embodiment, the present invention concerns the use of a composition comprising said culture of *Lactobacillus salivarius* (LS01) DSM 22775.

[0028] Preferably, the invention concerns the use of said culture of *Lactobacillus salivarius* (LS01) DSM 22775 for the preparation of a composition for the preventive and/or curative treatment of atopic dermatitis.

[0029] In accordance with another preferred embodiment, the present invention concerns the use of a composition comprising or, alternatively, consisting of said culture of *Lactobacillus salivarius* (LS01) DSM 22775.

[0030] The bacterial cultures of the present invention can be considered as "active ingredients" which can be blended, in the due proportions, with physiologically and/or pharmaceutically acceptable excipients.

[0031] In order to enhance their activity, the active ingredients can be administered systemically, advantageously orally, in the form of compositions in a solid state, prepared according to techniques known to a person skilled in the art. In a preferred embodiment, the compositions are in the form of lyophilizates, powders, granules, tablets, soft-gel capsules or suspensions.

[0032] The compositions of the present invention can be pharmaceutical, dietetic, nutritional or nutraceutical compositions.

[0033] The compositions of the present invention can moreover comprise dietary fibres with prebiotic activity, such as fructooligosaccharides (FOS), inulin and partially hydrolyzed guar gum (PHGG).

[0034] The pharmaceutical, dietetic, nutritional or nutraceutical compositions of the present invention comprise at least one culture of probiotic bacteria, as indicated above. The final concentration of the composition is comprised from 1×10^7 to 1×10^{11} CFU/g of composition, preferably from 1×10^8 to 1×10^{10} CFU/g of composition.

[0035] The compositions of the present invention, which comprise selected probiotic bacteria, are useful as immunomodulating compositions capable of modulating the immune system.

[0036] Said compositions are capable of curing and/or preventing the pathologies associated with alterations in the functions of the following cytokines: Th1 (IFN- γ and IL-12) and Th2 (IL-4, IL-5 and IL-10).

[0037] Furthermore, said compositions are capable of inducing the immune system, inducing the latter toward the production of type 1 cytokines.

[0038] Therefore, the compositions of the present invention have valid application for the preventive and/or curative

treatment of pathologies associated with alterations of the immune system, in particular for the preventive and/or curative treatment of allergies, atopy, allergic rhinitis, food hypersensitivity, atopic dermatitis, eczema, asthma or immunodeficiencies.

[0039] Advantageously, the compositions of the present invention have valid application in the preventive and/or curative treatment of atopic dermatitis.

[0040] Advantageously, the intake, by patients affected with atopic dermatitis AD, of a probiotic composition comprising the culture of *Lactobacillus salivarius* (LS01) DSM 22775 and the culture of *Bifidobacterium breve* (BR03) DSM 16604 at a concentration comprised from 1×10^7 to 1×10^{11} CFU, preferably 1×10^9 CFU/g, determines a decrease in the Scorad index (a lower Scorad value indicates a better quality of life).

[0041] Moreover, the administration of said probiotic composition for a period comprised from 12 to 24 weeks, preferably 16 weeks, is capable of reducing/blocking the secretion of the cytokine IL-4.

Experimental part (I)

[0042] The Applicant tested all six of the above-mentioned strains.

[0043] A first clinical study was conducted to test a composition comprising the culture of *Lactobacillus salivarius* (LS01) DSM 22775 and the culture of *Bifidobacterium breve* (BR03) DSM 16604.

[0044] A group of 40 patients (males and females, aged from 18 to 55 years) with moderate/severe AD were randomized (1:1) into two groups.

[0045] The diagnosis of moderate/severe AD was evaluated using a standardized AD severity index (SCORAD index = SCORing Atopic Dermatitis). Said index is known to those persons skilled in the art.

[0046] T0 represents time zero, as the starting point of the treatment, whereas T16 represents the 16-week period of treatment. At T16 an examination was performed, with Scorad and sampling.

[0047] Group A was treated with a composition comprising the culture of *Lactobacillus salivarius* (LS01) DSM 22775 and the culture of *Bifidobacterium breve* (BR03) DSM 16604, having a concentration of 1×10^9 CFU (corresponding to 0.01 g). The two cultures were blended with 1 g of com starch to give a final composition of 1.01 g in sachets. The 20 patients of group A received 2 sachets per day for 16 weeks.

[0048] Group B was treated with a placebo represented by 1 g of (only) com starch. The 20 patients of group B received 2 sachets of placebo per day for 16 weeks.

[0049] The following cytokines were tested for Th1 (IFN- γ and IL-12) and Th2 (IL-4, IL-5 and IL-10).

[0050] All data of the 40 patients were carefully collected and analysed.

[0051] Figure 1 shows a histogram relative to the Scorad index at T0 and T16.

[0052] The results obtained show that the intake, by the 20 patients of group A, of the probiotic composition comprising the culture of *Lactobacillus salivarius* (LS01) DSM 22775 and the culture of *Bifidobacterium breve* (BR03) DSM 16604, at a concentration of 1×10^9 CFU, determined a decrease in the Scorad in patients affected by AD (a lower Scorad value indicates a better quality of life). Moreover, the administration of said probiotic composition for 16 weeks was able to block secretion of the cytokine considered, IL-4, whose mean value was comparable to that obtained at the time of enrolment. Whereas in contrast the administration of the placebo was not able to block the release of the cytokine IL-4, which significantly increased after 16 weeks of treatment (evolution of the disease).

[0053] A series of pre-clinical in vitro studies were conducted on a group of bacteria comprising, among others, the culture of *Lactobacillus salivarius* (LS01) DSM 22775 and the culture of *Bifidobacterium breve* (BR03) DSM 16604.

[0054] Pre-clinical in vitro studies comparing the immunomodulating capacities of the tested bacterial strains demonstrated that the culture of *Lactobacillus salivarius* (LS01) DSM 22775 and the culture of *Bifidobacterium breve* (BR03) DSM 16604 show a superior capacity in inducing the production of type 1 cytokines.

[0055] The cultures of the bacteria *Lactobacillus pentosus* (LPS01) DSM 21980, *Streptococcus thermophilus* (FP4) DSM 18616, *Lactobacillus casei ssp. rhamnosus* (LR04) DSM 16605 and *Lactobacillus acidophilus* (LA02) DSM 21717 were also tested.

[0056] The culture of the strain *L. pentosus* (LPS01) DSM 22775 demonstrated in vitro to induce a very strong pro-Th1 cytokine profile.

[0057] Moreover, all the cultures of the strains *Lactobacillus pentosus* (LPS 01) DSM 21980, *Streptococcus thermophilus* (FP4) DSM 18616, *Lactobacillus casei ssp. rhamnosus* (LR04) DSM 16605 and *Lactobacillus acidophilus* (LA02) DSM 21717 were tested for their capacity to induce the synthesis of IL-12 by PBMCs in an in vivo study. IL-12 is a Th1-type cytokine and represents a key element in the treatment of allergies and other atopic pathologies.

[0058] By way of example, the data relative to *L. pentosus* (LPS01) DSM 21980, tested on PBMCs, are reported: Th-1 (pg/ml) = 1.754; Th-2 (pg/ml) = 326; Th1/Th2 = 5.38.

Experimental part (II)

[0059] The Applicant conducted a first clinical study using a composition comprising the cultures *Lactobacillus salivarius* (LS01) DSM 22775 and *Bifidobacterium breve* (BR03) and a second clinical study using a composition comprising the cultures *Lactobacillus salivarius* (LS01) DSM 22775, *Bifidobacterium breve* (BR03) and *Lactobacillus pentosus* (LPS01) DSM 21980. For the sake of simplicity, only the details of said first clinical study are reported, since said second clinical study was conducted with the same methods. Both clinical studies (first and second) gave comparable results.

[0060] The Applicant conducted said first clinical study (randomised, double-blind, on patients affected by severe or moderate atopic dermatitis), in order to evaluate the clinical efficacy of the administration of two probiotic bacterial strains, namely *Lactobacillus salivarius* (LS01), deposited with the DSMZ and having the deposit number DSM 22775, and *Bifidobacterium breve* (BR03), deposited with the DSMZ and having the deposit number DSM 16604.

Materials and methods

[0061] 48 patients affected by atopic dermatitis (hereinafter AD) were recruited into the study. Their characteristics are shown in Table 1.

Table 1

	Patients with probiotics (n=31)	Placebo patients (n=15)
Sex(M/F)	14/17	5/10
Age (years)	32.44 $\pm$ 1.47	30.91 $\pm$ 2.79
Respiratory allergy	23/31 (74.2%)	9/15 (60%)
Food allergy	2/31(6.4%)	2/15 (13.3%)
Other allergies	7/31 (22.6%)	2/15 (13.3%)
No allergies	5/31 (16%)	4/15 (26.6 %)
Years of AD	19.36 $\pm$ 2.38	15.08 $\pm$ 3.04

[0062] The present clinical study was conducted in the period of April-September 2010. 32 patients (treatment group) received a preparation containing the two probiotic bacterial strains, namely *L. salivarius* (LS01) DSM 22775 and *B. breve* (BR03) DSM 16604, at a dose of 1×10^9 CFU/g of each strain in maltodextrin.

[0063] Whereas 16 patients (placebo group) received a preparation of maltodextrin alone. Both groups were treated twice a day for 12 weeks. The compositions in lyophilised form were dissolved in water and administered orally to the patients.

[0064] None of the patients modified his or her diet and all were told not to consume fermented products containing live microorganisms during the clinical study. Patients who were affected by active allergic (respiratory or skin) disease, or infectious disease during pregnancy or breastfeeding, were excluded from the present study. Patients who had received a treatment with probiotics, antibiotics or immunomodulators during the six months prior to enrolment or who had received treatment with oral steroids a month prior to enrolment were excluded from the present study. All patients gave their informed consent at the time of enrolment.

[0065] The clinical evaluation was based on the SCORAD index (SCORing Atopic Dermatitis), on the DLQI (Dermatology Life Quality Index) and on microbial translocation and immunological parameters. For each individual patient, all evaluations were made at the time of enrolment (baseline), after 3 months of treatment (3ms) and two months after the end of the therapy (3+2ms).

Questionnaire on symptoms and quality of life

[0066] The diagnostic criteria for AD were set following the recently published clinical guide. Clinical severity was evaluated using the SCORAD index developed by the European Task Force for atopic dermatitis. In the course of the study, in order to measure variations in the symptoms tied to the pathology, all enrolled patients filled out the specific DLQI (Dermatology Life Quality Index) questionnaire. The SCORAD evaluation was performed at the start, end and two months after suspension of the treatment (probiotics or placebo) by an operator who did not know in which group the patient was enrolled. The DLQI was filled in by patients at the time of enrolment and at 12 and 20 weeks.

Separation of plasma and peripheral blood mononuclear cells (PBMCs)

[0067] For the separation of plasma and peripheral blood mononuclear cells (PBMCs), samples of peripheral venous blood were drawn from all study subjects at the time of enrolment, after 3 months of treatment and 2 months after suspension of the therapy. The plasma was stored until the time of analysis, whereas the PBMCs were separated by density gradient centrifugation. Cell viability was determined using an Adam-MC cell counter.

Stimulation of PBMCs

[0068] The PBMCs were incubated for 18 hours in the presence or absence of Lipopolysaccharide (LPS) bacterial stimulation. An analysis of the cytokines induced by the antigenic stimulus was performed by flow cytometry, adding Brefeldin A as an inhibitor of cell transport in the last hours of culture.

Immunophenotype analysis

[0069] The percentage of activated CD8⁺ T lymphocytes was calculated directly on samples of peripheral venous blood by cytofluorimetric analysis using monoclonal antibodies (mAb) conjugated with fluorochromes (CD8, CD38, CD45RO). After incubation with the mAb, the erythrocytes were lysed and the cells were fixed and analysed with an FC500 flow cytometer (Beckman Coulter).

Analysis of lymphocyte subpopulations

[0070] The identification of the different subpopulations of T lymphocytes was conducted on non-stimulated PBMC cultures and PMBC cultures stimulated for 18 hours with LPS, at all times (baseline, 3ms and 3+2ms). Cell marking was done by simultaneously using combinations of fluorescent antibodies capable of recognising molecules both on the cell surface and on intracellular components. Specifically, type 1 T helper lymphocytes (Th1) were identified as CD4⁺/IFN- γ ⁺/Tbet⁺; type 2 T helper lymphocytes (Th2) as CD4⁺/IL-4⁺/GATA3⁺; type 17 T helper lymphocytes (Th17) as CD4⁺/IL-17⁺/ROR γ T⁺ and, finally, regulatory T helper lymphocytes (Treg) as CD4⁺/CD25⁺/FoxP3⁺/IL-10⁺/TGF- β ⁺.

Evaluation of the expression of toll-like preceptors (TLR) on regulatory T lymphocytes

[0071] The analysis was performed on non-stimulated PBMCs, following the protocol provided by the manufacturer of the mAb. Specifically, the following mAb were necessary: CD4, CD25, FoxP3, TLR-4, TLR-2.

Plasma LPS assay

[0072] In order to determine the concentration of LPS present in the plasma samples, use was made of "LAL Chromogenic Endpoint Assay" kit sold by the company Hycult biotechnology, following the manufacturer's instructions. The test was conducted in a 96 well plate. After 45 minutes of incubation at room temperature, the absorbance at a wavelength of 405 nm was read with a spectrophotometer and the LPS concentration, expressed in pg/ml, was calculated by interpolation with a curve of known concentration.

Collection and storage of faecal samples

[0073] Faecal samples were collected at the time of enrolment, after 3 months of treatment and 2 months after the end of the study. The samples were immediately placed at 4°C, aliquoted and stored at - 80°C until the time of analysis.

Quantification of the bacteria present in the faecal samples

[0074] The quantification was conducted on all aerobic bacteria and on the following bacterial groups: Enterobacteria, Staphylococci, Lactobacilli (in particular *L.salivarius* LS01), and Bifidobacteria (in particular *B. breve* BR03). Faecal samples were diluted in physiological solution and the appropriate dilutions were incubated according to the following scheme:

- All AEROBIC bacteria: Tryptic soy agar with 5% sheep's blood (AS);
- Enterobacteria: MacConkey Agar (MC);
- Staphylococci: Mannitol salt agar (MSA);
- Lactobacilli: Man, Rogosa and Sharp agar (MRS); Bifidobacteria: Bifidobacterium selective medium (BSM).

[0075] The MC and BSM media were incubated at 37°C for 24 and 48 hours, respectively. The AS, MRS and BSM media were incubated at 37°C, in the presence of 10% carbon dioxide, for 24, 48 and 72 hours, respectively.

[0076] All colonies were identified through growth in selective media, Gram stain, cell morphology and tests for catalase and oxidase.

[0077] The composition of the intestinal microbiota was expressed as the count (mean $\pm$ standard deviation of the base 10 logarithm (log₁₀) per gram of faeces) for each bacterial group. The detection limits were 2 log₁₀ CFU/g for all microorganisms with the exception of *B. Breve*, whose limit was 5 log₁₀ CFU/g. The modifications of the counts of the different microbial groups were calculated as:

$$[(\text{Log}_{10} \text{ CFU/g at T12 or T16}) - (\text{Log}_{10} \text{ CFU/g at T0})]$$

Molecular identification of *L. salivarius* LS01 and of *B. breve* BR03

[0078] The two bacterial strains were initially identified based on their morphological peculiarities: *L. salivarius* LS01 cultivated on agarized BSM medium forms cream-white coloured colonies, roundish with an elongated shape and a diameter comprised from 2 to 4 mm; *B. breve* BR03 on agarized BSM medium forms fuchsia coloured colonies, roundish with an elongated shape and a diameter comprised from 1 to 2 mm.

[0079] In the cultures in which it was possible to assume the presence of the strains *L. salivarius* and *B. breve*, 10 random colonies were analysed by Polymerase Chain Reaction (PCR) in accordance with the guidelines published by the National Institute of Health. All of the colonies classified as *L. salivarius* or *B. breve* were further characterised by running pulsed field gel electrophoresis (PFGE) and comparing the profiles with suitable references.

Statistical analysis

[0080] The results were analysed using appropriate statistical tests. T-tests were used to compare patients during the treatments. Possible relationships were evaluated using the test of Pearson's correlation. The variations in the bacterial count were analysed using the Wilcoxon-Mann-Whitney test. The statistical analysis was performed using the SPSS statistical package.

Results

a) Clinical efficacy

[0081] 46 patients affected by AD completed the clinical study; 2 patients (1 patient per group) were excluded. The evaluation of the SCORAD (SCORAD index) and DLQI was made during the follow-up examinations. The patients treated with the probiotics showed a significant reduction in the SCORAD at the end of the treatment ($p=0.001$), which was also maintained after suspension of the treatment ($p=0.006$) (Table 2).

[0082] Analogously, the DLQI showed an improvement in the patients treated with the probiotic mixture (enrolment vs. 3m treatment: $p=0.024$; enrolment vs. 2ms after suspension: $p=0.001$). In the group of patients to whom the placebo was administered, no significant difference was observed either in terms of SCORAD or of DLQI (Table 2).

b) Plasma LPS concentration

[0083] The LPS concentration is an index of microbial translocation; an increase in the LPS concentration is associated with alterations in intestinal permeability. The treatment with the two probiotic strains induced a reduction in the plasma LPS concentration, which was maintained after interruption of the therapy (baseline vs. 3 months of treatment: $p=0.050$; baseline vs. 2 months after suspension of the treatment: $p<0.001$; 3 months of treatment vs. 2 months after suspension of the treatment: $p<0.001$), as shown below (Table 3).

Table 3

	Patients with probiotics	Placebo
Baseline	71.71	56.64
3ms of treatment	53.17	76.7
2ms after suspension	36	119.61

[0084] In patients treated with the placebo a significant increase was observed in the plasma LPS levels during the study period (baseline vs. 2 months after suspension of the treatment: $p=0.004$; 3 months of treatment vs. 2 months after suspension of the treatment: $p=0.016$) (Table 3). At the end of the study, the plasma LPS concentration was significantly higher in the control group (placebo) in comparison to the treatment group ($p<0.001$).

c) Activated T lymphocytes

[0085] The activated CD8+ T cells were analysed by evaluating the expression on the same of the CD38 and CD45 molecules. A progressive reduction was observed in said cells during the study period in the patients treated with the two probiotics (baseline vs. 2 months after suspension of the treatment $p<0.001$; 3 months of treatment vs. 2 months after suspension of the treatment: $p<0.001$) (Table 4). No significant difference was observed in the placebo group.

Table 4

	Patients with probiotics	Placebo
Baseline	2.51	1.8
3ms treatment	2.08	1.61
2ms after suspension	1.19	1.3

d) Regulatory T cells (Treg)

[0086] The regulatory T cells were evaluated both in the absence of stimulation and after stimulation with LPS. After treatment with the probiotic mixture, the percentage of non-stimulated Treg lymphocytes was significantly increased (baseline vs. 2 months after suspension of the treatment: $p=0.002$; 3 months of treatment vs. 2 months after suspension of the treatment: $p=0.034$) (Table 5A and Table 5B). Similar results were obtained from the analysis after stimulation with LPS.

Table 5A

	Patients with probiotics	Placebo
Baseline	18.57	33.22
3ms treatment	23.08	32.08
2ms after suspension	39.12	27.22

Table 5B

	Patients with probiotics	Placebo
Baseline	19.85	16.77
3ms treatment	24.99	28.31
2ms after suspension	31.8	22.38

e) Expression of Toll-like receptors on regulatory T lymphocytes

[0087] The expression of the Toll-like receptors 2 (TLR-2) and 4 (TLR-4) was evaluated directly in peripheral venous blood. As shown in Table 5C and Table 5D, the expression of both molecules on the Treg lymphocytes was significantly increased after 3 months of administration of probiotics; this result was also maintained 2 months after suspension of the treatment (TLR-2: baseline vs. 3 months of treatment: $p<0.001$; baseline vs. 2 months after suspension of the treatment: $p=0.010$; TLR-4: baseline vs. 3 months of treatment: $p<0.001$; baseline vs. 2 months after suspension of the treatment $p=0.011$). In the control group (placebo), a significant decrease was observed in the expression of the molecule TLR-2 by the Treg lymphocytes at the end of the study (3 months of treatment vs. 2 months after suspension of the treatment: $p=0.009$).

Table 5C

	Patients with probiotics	Placebo
Baseline	8.67	16.67
3ms treatment	30.89	23
2ms after suspension	23.78	12.51

Table 5D

	Patients with probiotics	Placebo
Baseline	9.27	17.02
3ms treatment	31.17	22.97
2ms after suspension	22.67	18.87

f) Lymphocyte subpopulations Th1, Th2 and Th17

[0088] The lymphocyte subpopulations Th1, Th2 and Th17 were evaluated in the absence of stimulation and after stimulation with LPS in both groups undergoing analysis: the percentage of the subpopulation Th1, identified on the basis of the expression of the CD4 and Tbet molecules and secreting the cytokine IFN- γ , was significantly increased in the subjects treated with the probiotic mixture, both in the absence of stimulation and after stimulation with LPS (non-stimulated: baseline vs. 3 months of treatment: $p=0.003$; LPS: baseline vs. 3 months of treatment: $p=0.025$; baseline vs. 2 months after suspension of the treatment: $p=0.019$) (Table 6A and Table 6B). In the patients treated with the probiotics, two months after the end of the treatment a further decrease in the Th1 cells was observed ($p<0.001$).

Table 6A

	Patients with probiotics	Placebo
Baseline	0.11	0.18
3ms treatment	0.23	0.2
2ms after suspension	0.16	0.14

Table 6B

	Patients with probiotics	Placebo
Baseline	0.09	0.13
3ms treatment	0.19	0.21
2ms after suspension	0.18	0.1

[0089] The percentage of the Th2 cell subpopulation, identified on the basis of the expression of the CD4 and GATA3 molecules and secreting the cytokine IL-4, was significantly decreased during the administration of probiotics but after suspension of the therapy it increased again (non-stimulated: baseline vs. 3 months of treatment: $p=0.016$; 3 months of treatment vs. 2 months after suspension of the treatment: $p=0.005$; LPS: baseline vs. 3 months of treatment: $p=ns$; 3 months of treatment vs. 2 months after suspension of the treatment: $p=0.0045$) (Table 6C and Table 6D).

Table 6C

	Patients With probiotics	Placebo
Baseline	0.85	0.45
T1	0.26	0.19
T 2	0.9	0.41

Table 6D

	Patients with probiotics	Placebo
Baseline	0.81	0.61
T 1	0.45	0.3
T 2	1.69	0.07

[0090] The percentage of the Th17 cell subpopulation, identified on the basis of the expression of the CD4 and ROR γ molecules and secreting the cytokine IL-17, was decreased in the patients treated with the probiotics after 3 months of treatment (non-stimulated: baseline vs. 3 months of treatment: $p=0.037$; LPS: baseline vs. 3 months of treatment: $p=0.046$) (Table 6E and Table 6F). Two months after suspension of the treatment significant differences were observed between the two groups undergoing analysis as regards the Th2 and Th17 lymphocyte subpopulations ($p=0.008$ and $p=0.031$, respectively). No significant difference was found in the Th1, Th2 and Th17 cell populations in the subjects of the control group (placebo).

Table 6E

	Patients with probiotics	Placebo
Baseline	0.18	0.14
3ms treatment	0.1	0.18
2ms after suspension	0.12	0.04

Table 6F

	Patients with probiotics	Placebo
Baseline	0.17	0.15
3ms treatment	0.1	0.12
2ms after suspension	0.11	0.1

g) Th1/Th2 and Th17/Treg ratios

[0091] After 3 months of administration of probiotics, a significant increase was observed in the Th1/Th2 ratio ($p=0.028$) (Table 7A). This effect was lost after interruption of the treatment ($p=0.002$). In contrast, in the group treated with the bacterial mixture, the Th17/Treg ratio progressively decreased and this result was also found 2 months after interruption of the therapy ($p=0.029$). During the study the subjects enrolled in the control group (placebo) did not show any significant variation in the Th1/Th2 and Th17/Treg ratios. The Th17/Treg ratio was significantly higher in the placebo group in comparison with the probiotic group after 3 months of treatment ($P=0.037$) (Table 7B).

Table 7A

	Patients with probiotics	Placebo
Baseline	0.48	0.74
3ms treatment	1.27	0.95
2ms after suspension	0.49	0.95

Table 7B

	Patients with probiotics	Placebo
Baseline	1.075	0.94

(continued)

	Patients with probiotics	Placebo
3ms treatment	0.57	1.19
2ms after suspension	0.62	0.333

h) Correlations

[0092] In the patients to whom the probiotics were administered, at the end of the treatment a negative correlation was observed between the SCORAD and the percentage of Treg lymphocytes after stimulation with LPS ($p=0.020$) and between the percentage of activated CD8+ T lymphocytes and that of the Th1 cells stimulated with LPS ($p=0.046$).

[0093] A significant positive correlation was observed between the plasma LPS concentration and the percentage of non-stimulated Th2 cells ($p=0.016$), between the plasma LPS concentration and the percentage of Th17 cells stimulated with LPS ($p=0.04$) and between the percentage of activated CD8+ T lymphocytes and that of the Th17 cells stimulated with LPS ($p=0.037$).

i) Modifications of the faecal microbiota and detection of the strains *L. salivarius* LS01 and *B. breve* BR03

[0094] The cell counts at the time of enrolment, after 3 months of treatment and 2 months after the end of administration are shown in Table 8.

Table 8

	baseline	3ms treatment	2ms after suspension
Patients with probiotics			
Total aerobes	7.7 ± 1.0	7.9 ± 0.9	8.2 ± 0.7
Enterobacteriaceae	6.8 ± 1.3	7.3 ± 1.2	6.7 ± 2.6
Staphylococci	4.0 ± 1.6	$2.8 \pm 1.5^{*\diamond}$	$2.3 \pm 1.6^{*\diamond}$
Lactobacilli	5.4 ± 2.0	5.1 ± 1.8	5.5 ± 0.4
Bifidobacteria	9.1 ± 0.6	8.8 ± 0.5	8.4 ± 0.6
Patients with placebo			
Total aerobes	7.1 ± 1.5	7.5 ± 1.2	7.7 ± 1.3
Enterobacteriaceae	5.8 ± 1.7	6.7 ± 1.6	6.2 ± 1.5
Staphylococci	4.1 ± 0.9	3.9 ± 1.1	4.0 ± 1.1
Lactobacilli	5.4 ± 1.9	5.2 ± 1.6	5.4 ± 1.8
Bifidobacteria	8.5 ± 1.1	8.6 ± 0.6	8.6 ± 0.9
*significance decreased ($P<0.05$) compared to baseline			
$\diamond$ Significant difference ($P<0.05$) compared to placebo at the same time.			

[0095] In the control group (placebo), no significant alterations were found in the faecal flora. In the group treated with probiotics, after 3 months a significant decrease in staphylococci was detected. This finding was significant also in comparison with the control group after 3 months of treatment. The decrease remained statistically significant even two months after suspension of the probiotics. No significant modifications were found in any other microbial group analysed.

[0096] The findings related to the probiotic strains in the faecal samples of the subjects enrolled in the study are shown in Table 9. All the colonies with a morphology corresponding to that of the strains considered, identified by PCR analysis and PFGE, were correspondent to those of the strains previously identified as *L. salivarius* LS01 or *B. breve* BR03.

[0097] The strain *L. salivarius* LS01 was found in all subjects after 3 months of treatment, in a quantity equal to 10^2 - 10^5 CFU/g. The strain *B. breve* BR03 was detected in 90% of the subjects, after three months of treatment, in a quantity equal to 10^6 - 10^8 CFU/g.

[0098] In some patients, both strains were also isolated 2 months after suspension of the treatment but in a smaller quantity compared to those obtained immediately after the end of administration.

[0099] The Applicant was able to verify that the compositions of the present invention, in particular those of said first clinical study (*Lactobacillus salivarius* (LS01) DSM 22775, and *Bifidobacterium breve* (BR03) DSM 16604) and said second clinical study (*Lactobacillus salivarius* (LS01) DSM 22775, *Bifidobacterium breve* (BR03) DSM 16604 and *Lactobacillus pentosus* (LPS01) DSM 21980), administered for 3 months to adult subjects affected by severe or moderate atopic dermatitis, demonstrated to be able to improve the clinical and immunological parameters thereof. In particular, at the end of the probiotic treatment, the following were observed:

- 1) a significant improvement in the SCORAD index and DLQI;
- 2) a reduction in microbial translocation (plasma LPS levels) and in the activation of CD8+ T lymphocytes;
- 3) an increase in the percentage both of the total regulatory T lymphocytes (Treg), and of the same expressing the markers TLR2- and TLR4;
- 4) an improvement in the Th1/ Th2 and Th17/ Treg ratios.

[0100] Advantageously, it was demonstrated that the probiotic strains used were capable of colonising the gastrointestinal tract of the patients and of reducing the staphylococcal load in faeces.

[0101] Advantageously, the clinical effect of the administration of probiotics, in terms of improving the symptoms of the pathology and quality of life, revealed itself after 3 months of treatment. Said effect showed to persist even 2 months after suspension of the treatment.

[0102] Advantageously, the persistence of the 3 probiotic bacterial strains (*Lactobacillus salivarius* (LS01) DSM 22775, *Bifidobacterium breve* (BR03) DSM 16604 and *Lactobacillus pentosus* (LPS01) DSM 21980) in the gastrointestinal tract was also observed after suspension of the treatment, suggesting the ability of the specific bacterial combinations to colonise the intestine.

[0103] Alterations in intestinal permeability play a key role in the pathogenesis of atopic dermatitis and in other diseases characterised by an inflammatory component. The results obtained from the clinical study demonstrate an important aspect, which is represented by the presence of a negative correlation between plasma LPS levels and atopy. Specifically, the presence of a significant improvement in microbial translocation both during and after the probiotic treatment demonstrated the immunomodulating effect possessed by the 3 bacterial strains, able to improve the intestinal barrier by acting both directly and indirectly.

[0104] Studies on HIV infections have shown an association between microbial translocation and immunological activation with an excessive T-cell turnover. The evaluation of the percentage of activated CD8+ T lymphocytes, considered an important prognostic factor in HIV infections, is one of the most widely accepted markers of LPS-mediated activation of the immune system. As expected, in the patients affected by atopic dermatitis enrolled in the present study, a reduction in the plasma LPS values was observed in association with a decrease in activated CD8+ T cells. These data demonstrate that the bacterial strains tested have a strong immunomodulating activity as a consequence of their capacity to increase the intestinal barrier. In the placebo group this correlation was not observed.

[0105] Four different subpopulations of CD4+ T-helper lymphocytes were identified on the basis of their capacity to secrete cytokines: Th1, main producers of IFN- γ and IL-2, are involved in cell-mediated immune responses; Th2, secreting IL-4, are responsible for triggering the humoral response; Th17, secreting IL-17 IL-21 and IL-22, are involved in immune responses directed against extracellular pathogens and in autoimmune diseases; regulatory T cells with an immunoregulatory function control the hyperactivation of immune responses.

[0106] A decrease in Treg cells has been reported in patients with atopic dermatitis and asthma, and their number is inversely correlatable to the secretion of IgE, eosinophilia, and IFN- γ levels.

[0107] The intake of the probiotic strains induced a significant increase in the percentage of total Treg lymphocytes and a significant increase in Treg lymphocytes expressing the receptors TLR2 and TLR4 after the treatment. These cell subpopulations perform an immunosuppressive activity as their main function. The colonisation of *S. aureus* is a crucial question for the progression of atopic dermatitis, as is the positive correlation between the severity of the disease and colonisation by *S. aureus*. The beneficial effects of the probiotics could be due to a Treg-cell mediated recovery of immunological tolerance toward *S. aureus* and Gram-negative bacteria. Moreover, a negative correlation was found between the SCORAD index and Treg cells, suggesting a positive role of the latter in the evolution of atopic dermatitis.

[0108] The Th2 subtype plays a critical role in the pathogenesis of allergic reactions and is responsible for the acute phase of atopic dermatitis, while the Th1 subpopulation has a protective function that is associated with the chronic phase of the disease. In patients affected by atopic dermatitis, Th17 cells are increased in acute skin lesions but not in chronic lesions and are positively correlated with the severity of the disease. The data obtained in this study have shown that this new probiotic combination increases the functionality of Th1 cells to the detriment of the Th2 and Th17 subpopulations, thus increasing the Th1/Th2 and Th17/Treg ratios.

[0109] The Th2 and Th17 subtypes are positively correlated with plasma LPS levels, suggesting a strong interaction

between microbial translocation and immune system modifications. This finding is further confirmed by the positive correlation between Th17 and activated CD8+. The role performed by microbial intestinal flora in the development of immunological tolerance and in the pathogenesis of allergic diseases has moreover been amply demonstrated. The alteration in intestinal microflora plays a role in the onset of atopic dermatitis and in the worsening of skin symptoms.

The same authors, when analysing the composition of the faecal microbiota of sensitized wheezy and non-wheezy children, observed that the lactobacteria and bifidobacteria were similar in both groups.

[0110] In the present study a significant difference was observed in the microbial composition of the two groups analysed, with respect to Staphylococci. Therefore, it is presumable that said genus could have a crucial role both at the level of the skin and in the intestines of patients affected by atopic dermatitis and that the administration of the present probiotic combination could rebalance it.

[0111] The present clinical study has shown that the bacterial strains of the present invention are well tolerated, and are capable of colonising the intestinal tract and of inducing beneficial clinical effects in subjects affected by atopic dermatitis through the modulation of some of the most important immunopathological alterations typical of this inflammatory skin disease.

[0112] The subject matter of the present invention is the use of the above tested bacterial strains in an additional therapy supporting the treatment of atopic dermatitis in adult subjects.

Herein described is

1. A composition comprising at least one culture of probiotic bacteria selected from the group comprising the following bacterial cultures deposited with the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, (DSMZ):

- *Lactobacillus salivarius* (LS01) DSM 22775, deposited on 23.07.2009;
- *Bifidobacterium breve* (BR03) DSM 16604, deposited on 20.07.2004;
- *Lactobacillus pentosus* (LPS01) DSM 21980, deposited on 14.11.2008;
- *Streptococcus thermophilus* (FP4) DSM 18616, deposited on 13.09.2006;
- *Lactobacillus casei ssp. rhamnosus* (LRO4) DSM 16605, deposited on 20.07.2004; and
- *Lactobacillus acidophilus* (LA02) DSM 21717, deposited on 06.08.2008,

for use in the treatment of pathologies associated with alterations of the immune system.

2. The composition for use in accordance with EM 1, in the preventive and/or curative treatment of pathologies associated with alterations of the immune system selected from the group comprising allergies, atopy, allergic rhinitis, food hypersensitivity, atopic dermatitis, eczema, asthma or immunodeficiencies.

3. The composition for use in accordance with EMs 1 or 2, in the preventive and/or curative treatment of atopic dermatitis.

4. The composition for use in accordance with one of the EMs 1-3, wherein said composition comprises at least one culture of prebiotic bacteria selected from the group comprising *Lactobacillus salivarius* (LS01) DSM 22775, *Bifidobacterium breve* (BR03) DSM 16604 and *Lactobacillus pentosus* (LPS 01) DSM 21980.

5. The composition for use in accordance with EM 4, wherein said composition consists of *Lactobacillus salivarius* (LS01) DSM 22775, *Bifidobacterium breve* (BR03) DSM 16604 and *L. pentosus* (LPS01) DSM 21980.

6. The composition for use in accordance with EM 4, wherein said composition consists of *Lactobacillus salivarius* (LS01) DSM 22775 and *Bifidobacterium breve* (BR03) DSM 16604.

7. The composition for use in accordance with any one of EMs 1-6, wherein said bacterial cultures are present in said compositions in the form of live bacteria, dead bacteria or cellular components, cell extracts or lysates thereof.

8. The composition for use in accordance with any one of EMs 1-7, wherein said composition is in solid form for oral administration.

9. The composition for use in accordance with any one of EMs 1-8, wherein said cultures of probiotic bacteria are present at a concentration comprised from 1×10^7 to 1×10^{11} CFU/g of composition, preferably from 1×10^8 to 1×10^{10} CFU/g of composition.

10. The composition for use in accordance with any one of EMs 1-9, wherein said composition further comprises dietary fibres with prebiotic activity selected from among fructooligosaccharides (FOS), inulin and partially hydrolyzed guar gum (PHGG).

11. The composition for use in accordance with any one of EMs 1-10, wherein said composition is a pharmaceutical, dietetic, nutritional or nutraceutical composition.

12. Use of at least one culture of probiotic bacteria selected from the group comprising the following cultures of bacteria deposited with the Deutsche Sammlung und Mikroorganismen und Zellkulturen GmbH, (DSMZ):

- *Lactobacillus salivarius* (LS01) DSM 22775, deposited on 23.07.2009;
- *Bifidobacterium breve* (BR03) DSM 16604, deposited on 20.07.2004;

- *Lactobacillus pentosus* (LPS01) DSM 21980, deposited on 14.11.2008;
- *Streptococcus thermophilus* (FP4) DSM 18616, deposited on 13.09.2006;
- *Lactobacillus casei ssp. rhamnosus* (LR04) DSM 16605, deposited on 20.07.2004; and
- *Lactobacillus acidophilus* (LA02) DSM 21717, deposited on 06.08.2008,

for use in the treatment of pathologies associated with alterations of the immune system.

13. The use according to EM 12, wherein said composition is for the preventive and/or curative treatment of pathologies associated with alterations of the immune system selected from the group comprising allergies, atopy, allergic rhinitis, food hypersensitivity, atopic dermatitis, eczema, asthma or immunodeficiency.

14. The use according to EM 13, wherein said composition is for the preventive and/or curative treatment of atopic dermatitis.

15. The use according to any one of EMs 12-14, wherein said composition is in accordance with any one of EMs 4-11.

Claims

1. A composition comprising a culture of probiotic bacteria *Lactobacillus salivarius* (LS01) DSM 22775, deposited on 23.07.2009 at Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH-DSMZ, for use in the preventive and/or curative treatment of pathologies selected from the group consisting of allergies, atopy, allergic rhinitis, food hypersensitivity, atopic dermatitis, eczema, asthma or immunodeficiency.
2. The composition for use in accordance with claim 1, wherein the composition comprises one culture of probiotic bacteria, wherein the culture of probiotic bacteria is *Lactobacillus salivarius* (LS01) DSM 22775, deposited on 23.07.2009 at Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH-DSMZ.
3. The composition for use in accordance with claim 1 or 2, in the preventive and/or curative treatment of atopic dermatitis and/or eczema.
4. The composition for use in accordance with claim 1 or 2, in the preventive and/or curative treatment of atopic dermatitis.
5. The composition for use in accordance with any one of claims 1-4, wherein said bacterial cultures are present in said compositions in the form of live bacteria, dead bacteria or cellular components, cell extracts or lysates thereof.
6. The composition for use in accordance with any one of claims 1-5, wherein said composition is in solid form for oral administration.
7. The composition for use in accordance with any one of claims 1-6, wherein said culture of probiotic bacteria is present at a concentration comprised from 1×10^7 to 1×10^{11} CFU/g of composition, preferably from 1×10^8 to 1×10^{10} CFU/g of composition.
8. The composition for use in accordance with any one of claims 1-7, wherein said composition further comprises dietary fibres with prebiotic activity selected from among fructooligosaccharides (FOS), inulin and partially hydrolyzed guar gum (PHGG).
9. The composition for use in accordance with any one of claims 1-8, wherein said composition is a pharmaceutical, dietetic, nutritional or nutraceutical composition.
10. A culture of probiotic bacteria of *Lactobacillus salivarius* (LS01) DSM 22775, deposited on 23.07.2009 at Deutsche Sammlung und Mikroorganismen und Zellkulturen GmbH - DSMZ, for use in the preventive and/or curative treatment of pathologies selected from the group consisting of allergies, atopy, allergic rhinitis, food hypersensitivity, atopic dermatitis, eczema, asthma or immunodeficiency.
11. The culture for use according to claim 10, wherein said culture is for the use in the preventive and/or curative treatment of atopic dermatitis and/or eczema.
12. The culture for use according to claim 10, wherein said culture is for the use in the preventive and/or curative treatment of atopic dermatitis.

Patentansprüche

1. Zusammensetzung, umfassend eine probiotische Bakterienkultur *Lactobacillus salivarius* (LS01) DSM 22775, hinterlegt am 23.07.2009 bei der Deutschen Sammlung von Mikroorganismen und Zellkulturen GmbH-DSMZ, zur Verwendung bei der präventiven und/oder kurativen Behandlung von Erkrankungen, ausgewählt aus der Gruppe, bestehend aus Allergien, Atopie, allergischer Rhinitis, Nahrungsmittelüberempfindlichkeit, atopischer Dermatitis, Ekzem, Asthma oder Immundefekt.
2. Zusammensetzung zur Verwendung nach Anspruch 1, wobei die Zusammensetzung eine einzige probiotische Bakterienkultur umfasst, wobei die probiotische Bakterienkultur *Lactobacillus salivarius* (LS01) DSM 22775 ist, hinterlegt am 23.07.2009 bei der Deutschen Sammlung von Mikroorganismen und Zellkulturen GmbH-DSMZ.
3. Zusammensetzung zur Anwendung nach Anspruch 1 oder 2 bei der präventiven und/oder kurativen Behandlung von atopischer Dermatitis und/oder Ekzem.
4. Zusammensetzung zur Anwendung nach Anspruch 1 oder 2, bei der präventiven und/oder kurativen Behandlung von atopischer Dermatitis.
5. Zusammensetzung zur Verwendung nach einem der Ansprüche 1-4, wobei diese Bakterienkulturen in diesen Zusammensetzungen in der Form von lebenden Bakterien, abgetöteten Bakterien oder Zellbestandteilen, Zellextrakten oder Lysaten davon vorliegen.
6. Zusammensetzung zur Verwendung nach einem der Ansprüche 1-5, wobei diese Zusammensetzung in fester Form zur oralen Verwendung vorliegt.
7. Zusammensetzung zur Verwendung nach einem der Ansprüche 1-6, wobei diese probiotischen Bakterienkulturen bei einer Konzentration vorliegen, die 1×10^7 bis 1×10^{11} KBE/g Zusammensetzung, bevorzugt 1×10^8 bis 1×10^{10} KBE/g Zusammensetzung, umfasst.
8. Zusammensetzung zur Verwendung nach einem der Ansprüche 1-7, wobei diese Zusammensetzung ferner diätetische Fasern mit präbiotischer Aktivität umfasst, ausgewählt aus Fructooligosacchariden (FOS), Inulin und teilweise hydrolysiertem Guargummi (PHGG).
9. Zusammensetzung zur Verwendung nach einem der Ansprüche 1-8, wobei diese Zusammensetzung eine pharmazeutische, diätetische, Ernährungs- oder nutraceuticalische Zusammensetzung ist.
10. Probiotische Bakterienkultur *Lactobacillus salivarius* (LS01) DSM 22775, hinterlegt am 23.07.2009 bei der Deutschen Sammlung von Mikroorganismen und Zellkulturen GmbH-DSMZ, zur Verwendung bei der präventiven und/oder kurativen Behandlung von Erkrankungen, ausgewählt aus der Gruppe, bestehend aus Allergien, Atopie, allergischer Rhinitis, Nahrungsmittelüberempfindlichkeit, atopischer Dermatitis, Ekzem, Asthma oder Immundefekt.
11. Kultur zur Verwendung nach Anspruch 10, wobei diese Kultur zur Anwendung bei der präventiven und/oder kurativen Behandlung von atopischer Dermatitis und/oder Ekzem bestimmt ist.
12. Kultur zur Verwendung nach Anspruch 10, wobei diese Kultur zur Anwendung bei der präventiven und/oder kurativen Behandlung von atopischer Dermatitis bestimmt ist.

Revendications

1. Composition comprenant une culture de bactéries probiotiques *Lactobacillus salivarius* (LS01) DSM 22775 déposées le 23.07.2009 à la Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH-DSMZ pour une utilisation dans le traitement préventif et/ou thérapeutique de pathologies choisies dans le groupe constitué des allergies, de l'atopie, de la rhinite allergique, de l'hypersensibilité alimentaire, de la dermatite atopique, de l'eczéma, de l'asthme ou des immunodéficiences.
2. Composition pour une utilisation selon la revendication 1, la composition comprenant une culture de bactéries probiotiques, dans laquelle la culture de bactéries probiotiques est la *Lactobacillus salivarius* (LS01) DSM 22775

déposée le 23.07.2009 à la Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH-DSMZ.

3. Composition pour une utilisation selon la revendication 1 ou 2 dans le traitement préventif et/ou thérapeutique de la dermatite atopique et/ou de l'eczéma.
4. Composition pour une utilisation selon la revendication 1 ou 2 dans le traitement préventif et/ou thérapeutique de la dermatite atopique.
5. Composition pour une utilisation selon l'une quelconque des revendications 1 à 4, dans laquelle lesdites cultures bactériennes sont présentes dans lesdites compositions sous la forme de bactéries vivantes, de bactéries mortes ou de leurs composants cellulaires, extraits cellulaires ou lysats.
6. Composition pour une utilisation selon l'une quelconque des revendications 1 à 5, dans laquelle ladite composition est sous forme solide pour une administration orale.
7. Composition pour une utilisation selon l'une quelconque des revendications 1 à 6, dans laquelle ladite culture de bactéries probiotiques est présente en concentration de 1×10^7 à 1×10^{11} CFU/g de composition, de préférence de 1×10^8 à 1×10^{10} CFU/g de composition.
8. Composition pour une utilisation selon l'une quelconque des revendications 1 à 7, dans laquelle ladite composition comprend en outre des fibres alimentaires dotées d'une activité prébiotique choisies parmi les fructooligosaccharides (FOS), l'inuline et la gomme guar partiellement hydrolysée (PHGG).
9. Composition pour une utilisation selon l'une quelconque des revendications 1 à 8, dans laquelle ladite-composition est une composition pharmaceutique, diététique, nutritionnelle ou nutraceutique.
10. Culture de bactéries probiotiques de *Lactobacillus salivarius* (LS01) DSM 22775 déposée le 23.07.2009 à la Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH-DSMZ pour une utilisation dans le traitement préventif et/ou thérapeutique de pathologies choisies dans le groupe constitué des allergies, de l'atopie, de la rhinite allergique, de l'hypersensibilité alimentaire, de la dermatite atopique, de l'eczéma, de l'asthme ou des immunodéficiences.
11. Culture pour une utilisation selon la revendication 10, dans laquelle ladite culture est destinée à une utilisation dans le traitement préventif et/ou thérapeutique de la dermatite atopique et/ou de l'eczéma.
12. Culture pour une utilisation selon la revendication 10, dans laquelle ladite culture est destinée à une utilisation dans le traitement préventif et/ou thérapeutique de la dermatite atopique.

FIGURE 1




Table 2

	Patients with probiotics (n= 31)			Placebo (n=15)			<i>p</i> *
	baseline	After 3 ms treatment	2 ms after suspension	baseline	After 3 ms treatment	2 ms after suspension	
SCORAD index	46.25 ± 3.70	29.45 ± 2.01	22.63 ± 2.81	35.00 ± 2.60	28.21 ± 1.53	27.43 ± 1.38	baseline vs. after 3 ms treatment: <i>p</i> < 0.001 baseline vs. 2 ms after suspension: <i>p</i> = 0.006
DLQ index	9.16 ± 0.80	6.58 ± 1.25	4.73 ± 0.75	6.52 ± 0.40	8.50 ± 1.27	5.29 ± 0.97	baseline vs. after 3 ms treatment: <i>p</i> < 0.024 baseline vs. 2 ms after suspension: <i>p</i> = 0.001

*significant differences in patients with probiotics

Table 9

	<i>L. salivarius</i> LS01		<i>B.breve</i> BR03	
	Recovery in faeces (% of patients treated)	Count interval (CFU/g)	Recovery in faeces (% of patients treated)	Count interval (CFU/g)
After 3 ms treatment	100%	$10^2 - 10^5$	90%	$10^5 - 10^8$
2 ms after suspension	25%	$10^2 - 10^4$	19%	$10^5 - 10^7$

Lactobacillus salivarius (LS01) DSM 22775 terápiás alkalmazásai

Szabadalmi igénypontok

1. A Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH-DSMZ intézménynél 2009. 07. 23-án letétbe helyezett *Lactobacillus salivarius* (LS01) DSM 22775 probiotikus baktériumtörzset tartalmazó készítmény allergiák, atopia, allergiás rhinitis, élelmiszer túlérzékenység, atopiás dermatitis, ekcéma, asztma vagy immunhiány közül kikerülő betegségek megelőző és/vagy gyógyító kezelésében történő alkalmazásra.
2. Készítmény az 1. igénypont szerinti alkalmazásra, ahol a készítmény egy probiotikus baktérium tenyészetét tartalmazza, ahol a probiotikus baktériumtenyészet a Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH-DSMZ intézménynél 2009. 07. 23-án letétbe helyezett *Lactobacillus salivarius* (LS01) DSM 22775.
3. Készítmény az 1. vagy 2. igénypont szerinti alkalmazásra, atopiás dermatitis és/vagy ekcéma megelőző és/vagy gyógyító kezelésében.
4. Készítmény az 1. vagy 2. igénypont szerinti alkalmazásra, atopiás dermatitis megelőző és/vagy gyógyító kezelésében.
5. Készítmény az 1-4. igénypontok bármelyike szerinti alkalmazásra, ahol a baktériumtenyészetek a készítményben élő baktérium, elölt baktérium vagy sejtkomponensek, sejtextraktumok vagy lizátumaik formájában vannak jelen.
6. Készítmény az 1-5. igénypontok bármelyike szerinti alkalmazásra, ahol a készítmény orális adagolásra szolgáló szilárd formában van jelen.
7. Készítmény az 1-6. igénypontok bármelyike szerinti alkalmazásra, ahol a probiotikus baktérium tenyészet a készítményre számítva 1×10^7 - 1×10^{11} CFU/g, előnyösen 1×10^8 - 1×10^{10} CFU/g koncentrációban van jelen.

8. Készítmény az 1-7. igénypontok bármelyike szerinti alkalmazásra, ahol a készítmény továbbá fruktooligoszacccharidok (FOS), inulin és részlegesen hidrolizált guar gumi (PHGG) közül választott prebiotikus hatással rendelkező étkezési rostokat tartalmaz.

9. Készítmény az 1-8. igénypontok bármelyike szerinti alkalmazásra, ahol a készítmény gyógyászati, étkezési, táplálkozási vagy gyógytáp készítmény.

10. A Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH-DSMZ intézménynél 2009. 07. 23-án letétbe helyezett *Lactobacillus salivarius* (LS01) DSM 22775 probiotikus baktérium tenyésztete allergiák, atopia, allergiás rhinitis, élelmiszer túlérzékenység, atopiás dermatitis, ekcéma, asztma vagy immunhiány közül kikerülő betegségek megelőző és/vagy gyógyító kezelésében történő alkalmazásra.

11. Tenyészet a 10. igénypont szerinti alkalmazásra, ahol a tenyészet atopiás dermatitis és/vagy ekcéma megelőző és/vagy gyógyító kezelésére szolgál.

12. Tenyészet a 10. igénypont szerinti alkalmazásra, ahol a tenyészet atopiás dermatitis megelőző és/vagy gyógyító kezelésére szolgál.

Szentorosi Zoltán
Szentorosi Zoltán ügyvéd
Szentorosi Zoltán Ügyvédi Iroda
H-1062 Budapest, Andrássy út 147.
Tel.: +36 1 460 1000, Fax: +36 1 460 1001
E-mail: zsoltan.szentorosi@szorosi.hu