



(22) Date de dépôt/Filing Date: 2007/01/22

(41) Mise à la disp. pub./Open to Public Insp.: 2007/07/26

(62) Demande originale/Original Application: 2 636 991

(30) Priorité/Priority: 2006/01/20 (US60/760,378)

(51) Cl.Int./Int.Cl. *A61K 38/46* (2006.01),
A61P 3/00 (2006.01)

(71) Demandeur/Applicant:
GENZYME CORPORATION, US

(72) Inventeurs/Inventors:
DODGE, JAMES, US;
PASSINI, MARCO, US;
SHIHABUDDIN, LAMYA, US;
CHENG, SENG, US

(74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : ADMINISTRATION D'ENZYME INTRAVENTRICULAIRE POUR DES MALADIES DE STOCKAGE DES
LYSOSOMES

(54) Title: INTRAVENTRICULAR ENZYME DELIVERY FOR LYSOSOMAL STORAGE DISEASES

(57) **Abrégé/Abstract:**

Lysosomal storage diseases can be successfully treated using intraventricular delivery of the enzyme which is etiologically deficient in the disease. The administration can be performed slowly to achieve maximum effect. Surprisingly, effects are seen on both sides of the blood-brain barrier, making this an ideal delivery means for lysosomal storage diseases which affect both brain and visceral organs.



ABSTRACT

Lysosomal storage diseases can be successfully treated using intraventricular delivery of the enzyme which is etiologically deficient in the disease. The administration can be performed slowly to achieve maximum effect. Surprisingly, effects are seen on both sides of the blood-brain barrier, making this an ideal delivery means for lysosomal storage diseases which affect both brain and visceral organs.

INTRAVENTRICULAR ENZYME DELIVERY FOR LYSOSOMAL STORAGE DISEASES

TECHNICAL FIELD OF THE INVENTION

[01] This invention is related to the area of lysosomal storage diseases. In particular, it relates to the treatment and/or prevention of these diseases by enzyme replacement therapy.

SUMMARY OF THE INVENTION

[02] A group of metabolic disorders known as lysosomal storage diseases (LSDs) includes over forty genetic disorders, many of which involve genetic defects in various lysosomal hydrolases. Representative lysosomal storage diseases and the associated defective enzymes are listed in Table I.

TABLE I

<u>Lysosomal storage disease</u>	<u>Defective enzyme</u>
Aspartylglucosaminuria	Aspartylglucosaminidase
Pabry	alpha.-Galactosidase A
Infantile Batten Disease* (CNL1)	Palmitoyl Protein Thioesterase
Classic Late Infantile batten Disease* (CNL2)	Tripeptidyl Peptidase
Juvenile Batten Disease* (CNL3)	Lysosomal Transmembrane Protein
Batten, other forms* (CNL4-CNL8)	Multiple gene products
Cystinosis	Cysteine transporter
Farber	Acid ceramidase
Fucosidosis	Acid .alpha.-L-fucosidase

Lysosomal storage disease

Galactosidosis
 Gaucher types 1, 2*, and 3* G.sub.M1
 gangliosidosis* Hunter*
 Hurler-Scheie*
 Kribs* alpha.-
 Mannosidosis* beta.-
 Mannosidosis*
 Maroteaux-Lamy
 Metachromatic
 leukodystrophy* Morquio
 A Morquio B

Mucopolysaccharidosis I/II*
 Niemann-Pick A*, B
 Niemann-Pick C*
 Poxnpe* Acid
 Sandhoff*
 Sanfilippo A*
 Sanfilippo B*

Sanfilippo C*
 Sanfilippo D*
 Schindler Disease* Schindler-
 Kanazaki
 Sialidosis
 Sly*
 Tay-Sachs*
 Wolman*

*CNS involvement

Defective enzyme

Protective protein/cathepsin A
 Acid .beta.-glucosidase, or
 Acid .beta.-galactosidase
 Iduronate-2-sulfatase
 alpha.-L-Iduronidase
 Galactocerebrosidase .
 Acid .alpha.-mannosidase
 Acid .beta.-mannosidase
 Arylsulfatase B
 Arylsulfatase A
 N-Acetylgalactosamine-6-sulfate
 Acid .beta.-galactosidase

N-Acetylglucosamine- 1-
 Acid sphingomyelinase
 NPC- 1
 .alpha.-glucosidase
 .beta.-Hexosaminidase B
 Heparan N-Sulfatase
 .alpha.-N-Acetylglucosaminidase

Acetyl-CoA: alpha.-glucosaminide
 N-Acetylglucosamine-6-sulfate
 .alpha.-N-Acetylglucosaminidase
 alpha.-N-Acetylglucosaminidase
 .alpha.-Neuraminidase .beta.-
 Glucuronidase .beta.-
 Hexosaminidase A
 Acid Lipase

- [03] The hallmark feature of LSDs is the abnormal accumulation of metabolites in the lysosomes which leads to the formation of large numbers of distended lysosomes in the perikaryon. A major challenge to treating LSDs (as opposed to treating an organ specific enzymopathy, *e.g.*, a liver-specific enzymopathy) is the need to reverse lysosomal storage pathology in multiple separate tissues. Some LSDs can be effectively treated by intravenous infusion of the missing enzyme, known as enzyme replacement therapy (ERT). For example, Gaucher type 1 patients have only visceral disease and respond favorably to ERT with recombinant glucocerebrosidase (Cerezyme, Genzyme Corp.). However, patients with metabolic disease that affects the CNS (*e.g.*: type 2 or 3 Gaucher disease) only respond partially to intravenous ERT because the replacement enzyme is prevented from entering the brain by the blood brain barrier (BBB). Furthermore, attempts to introduce a replacement enzyme into the brain by direct injection have been limited in part due to enzyme cytotoxicity at high local concentrations and limited parenchymal diffusion rates in the brain (Partridge, Peptide Drug Delivery to the Brain, Raven Press, 1991).
- [04] One exemplary LSD, is Niemann-Pick disease type A (NPA). According to UniProtKB/Swiss-Prot entry P17405, defects in the SMPDI gene, located on chromosome 11, (11p15.4-p15.1), are the cause of Niemann-Pick disease type A (NPA), also referred to as the classical infantile form of the disease. Niemann-Pick disease is a clinically and genetically heterogeneous recessive disorder. It is caused by the accumulation of sphingomyelin and other metabolically related lipids in the lysosomes, resulting in neurodegeneration starting from early life. Patients may show xanthomas, pigmentation, hepatosplenomegaly, lymphadenopathy and mental retardation. Niemann-Pick disease occurs more frequently among individuals of Ashkenazi Jewish ancestry than in the general population. NPA is characterized by very early onset in infancy and a rapidly progressive course leading to death by three years. The acid sphingomyelinase enzyme (ASM) that is defective in NPA converts sphingomyelin to ceramide. ASM also has phospholipase C activities toward 1,2-diacylglycerolphosphocholine and 1,2-diacylglycerolphosphoglycerol. The enzyme converts

Sphingomyelin + H₂O N-acylsphingosine + &ohne phosphate.

- [05] In accordance with the present invention, lysosomal storage diseases such as any of the diseases identified in Table 1 above, *e.g.*, Niemann-Pick disease type A or B, are treated and/or prevented using 'intraventricular delivery to the brain of the enzyme which is etiologically deficient in the disease. The administration can be performed slowly to achieve maximum effect. Effects are seen on both sides of the blood-brain barrier, making this a useful delivery means for lysosomal storage diseases which affect the brain and/or visceral organs. In a first aspect, the invention thus provides for a method of treating or preventing a lysosomal storage disease in patient, which disease is caused by an enzyme deficiency, the method comprising administering the enzyme to the patient via intraventricular delivery to the brain. In a related aspect, the invention provides for the use of an enzyme for the manufacture of a medicament for the treatment or prevention of a lysosomal storage disease in a patient, which disease is caused by a deficiency of the enzyme in the patient, wherein the treatment or prevention comprises the intraventricular administration of the enzyme to the brain. The enzyme deficiency may be caused by *e.g.*, a defect in the expression of the enzyme or by a mutation in the enzyme which leads to a reduced level of activity (*e.g.*, the enzyme being inactive) or an increased rate of clearance/breakdown of the enzyme in vivo. The deficiency may lead to an accumulation of an enzyme's substrate and the administration of the enzyme may lead to a reduction in the level of the substrate in the brain. The lysosomal storage disease may be any of the diseases identified in Table 1 above. The enzyme may be a lysosomal hydrolase.
- [06] According to one embodiment of the invention, a patient with Niemann-Pick A or B disease is treated. An acid sphingomyelinase is administered to the patient via intraventricular delivery to the brain in an amount sufficient to reduce sphingomyelin levels in said brain.

- [07] Another aspect of the invention is a kit for treating or preventing a lysosomal storage disease in patient, which disease is caused by an enzyme deficiency. The kit comprises the enzyme that is deficient, and a catheter and/or pump for delivery of the enzyme to one or more ventricles in the brain. The catheter and/or pump may be specifically designed and/or adapted for intraventricular delivery. According to one embodiment, the invention provides a kit for treating a patient with Niemann-Pick A or 13 disease. The kit comprises an acid sphingomyelinase, and a catheter for delivery of said acid sphingomyelinase to the patient's brain ventricles.
- [08] Yet another aspect of the invention is a kit for treating a patient with Niemann-Pick A or B disease. The kit comprises an acid sphingomyelinase and a pump for delivery of said acid sphingomyelinase to the patient's brain ventricles.
- [09] According to the invention, a patient can be treated who has a lysosomal storage disease which is caused by an enzyme deficiency which leads to accumulation of the enzyme's substrate. Such diseases include Gaucher disease, MPS I and II disease, Pompe disease, and Batten disease (CLN2), among others. The enzyme defective in the particular disease is administered to the patient via intraventricular delivery to the brain. It may be administered to the lateral ventricles and/or to the fourth ventricle. The rate of administration of the enzyme in accordance with the present invention is such that the administration of a single dose may be as a bolus or may take about 1-5 minutes, about 5-10 minutes, about 10-30 minutes, about 30-60 minutes, about 1-4 hours, or consumes more than four, five, six, seven, or eight hours. Substrate levels in said brain may thereby be reduced. The administration of a single dose may take more than 1 minute, more than 2 minutes, more than 5 minutes, more than 10 minutes, more than 20 minutes, more than 30 minutes, more than 1 hour, more than 2 hours, or more than 3 hours.
- [10] These and other embodiments which will be apparent to those of skill in the art upon reading the specification provide the art with methods and kits for the treatment and/or prevention of lysosomal storage diseases, in particular those involving both CNS and visceral pathologies.

BRIEF DESCRIPTION OF THE DRAWINGS

- [11] FIG. 1 shows diagram of sections of brain that were analyzed for sphingomyelin. S1 is at the front of brain and S5 is at the back of brain.
- [12] FIG. 2 shows that intraventricular administration of rhASM reduces SPM levels in the ASMKO mouse brain.
- [13] FIG. 3 shows intraventricular administration of rhASIVI reduces SPM levels in the ASMKO liver, spleen, and lung.
- [14] FIG. 4 shows hASM staining in the brain following intraventricular infusion.
- [15] FIG. 5 shows that intraventricular infusion of rhASM over a 6 h period reduces SPM levels in the ASMKO mouse brain.
- [16] FIG. 6 shows that intraventricular infusion of rhASM over a 6 h period reduces SPM levels in ASMKO liver, serum, and lung.
- [17] FIG. 7 shows documented hASM variants and their relationship to disease or enzyme activity.
- [18] FIG. 8 shows a cross section view of the brain with the ventricles indicated.
- [19] FIGS. 9A and 9B show lateral and superior views, respectively, of the ventricles.
- [20] FIG. 10 shows injection into the lateral ventricles.

DETAILED DESCRIPTION OF THE INVENTION

- [21] The practice of the present invention will employ, unless otherwise indicated, conventional techniques of immunology, molecular biology, microbiology, cell biology and recombinant DNA, which are within the skill of the art. See, e.g., Sambrook, Fritsch and Maniatis, *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd edition (1989); *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY* (F. M. Ausubel, *et al. eds.*, (1987)); the series *METHODS IN ENZYMOLOGY* (Academic Press, Inc.); *PCR 2: A PRACTICAL APPROACH* (M.J. MacPherson, B.D. Harries and G.R. Taylor eds. (1995)), Harlow and Lane, *eds.* (1988) *ANTIBODIES, A LABORATORY MANUAL*, and *ANIMAL CELL CULTURE* (R.I. Freshney, ed. (1987)).
- [22] As used in the specification and claims, the singular forms "a," "an" and "the" include plural references unless the context clearly dictates otherwise. For example, the term "a cell" includes a plurality of cells, including mixtures thereof.
- [23] As used herein, the term "comprising" is intended to mean that the compositions and methods include the recited elements, but not excluding others. "Consisting essentially of" when used to define compositions and methods, shall mean excluding other elements of any essential significance to the combination. Thus, a composition consisting essentially of the elements as defined herein would not exclude trace contaminants from the isolation and purification method and pharmaceutically acceptable carriers, such as phosphate buffered saline, preservatives, and the like. "Consisting of" shall mean excluding more than trace elements of other ingredients and excluding substantial method steps for administering the compositions or medicaments in accordance with this invention. Embodiments defined by each of these transition terms are within the scope of this invention.
- [23] All numerical designations, *e.g.*, pH, temperature, time, concentration, and molecular weight, including ranges, are approximations which are varied (+) or (-) by increments of 0.1. It is to be understood, although not always explicitly stated that all numerical designations are preceded by the term "about." It also is to be understood, although not

always explicitly stated, that the reagents described herein are merely exemplary and that equivalents of such that are known in the art may also be used.

- [25] The terms "therapeutic," "therapeutically effective amount," and their cognates refer to that amount of a substance, *e.g.*, enzyme or protein, that results in prevention or delay of onset, or amelioration, of one or more symptoms of a disease in a subject, or an attainment of a desired biological outcome, such as correction of neuropathology,. The term "therapeutic correction" refers to that degree of correction which results in prevention or delay of onset, or amelioration, of one or more symptoms in a subject. The effective amount can be determined by known empirical methods.
- [26] A "composition" or "medicament" is intended to encompass a combination of an active agent, *e.g.*, an enzyme, and a carrier or other material, *e.g.*, a compound or composition, which is inert (for example, a detectable agent or label) or active, such as an adjuvant, diluent, binder, stabilizer, buffer, salt, lipophilic solvent, preservative, adjuvant or the like, or a mixture of two or more of these substances- Carriers are preferably pharmaceutically acceptable. They may include pharmaceutical excipients and additives, proteins, peptides, amino acids, lipids, and carbohydrates (*e.g.*, sugars, including monosaccharides, di-, tri-, tetra-, and oligosaccharides; derivatized sugars such as alditols, aldonic acids, esterified sugars and the like; and polysaccharides or sugar polymers), which can be present singly or in combination, comprising alone or in combination 1-99.99% by weight or volume. Exemplary protein excipients include serum albumin such as human serum albumin (HA), recombinant human albumin (rHA), gelatin, casein, and the like. Representative amino acid/antibody components, which can also function in a buffering capacity, include alanine, glycine, arginine, betaine, histidine, glutamic acid, aspartic acid, cysteine, lysine, leucine, isoleucine, valine, methionine, phenylalanine, aspartame, and the like. Carbohydrate excipients are also intended within the scope of this invention, examples of which include but are not limited to monosaccharides such as fructose, maltose, galactose, glucose, D-mannose, sorbose, and the like; disaccharides, such as lactose, sucrose, trehalose, cellobiose, and the like; polysaccharides, such as

raffinose, melezitose, mattodextrins, dextrans, starches, and the like; and alditols, such as mannitol, xylitol, maltitol, lactitol, xylitol sorbitol (glucitol) and myoinositol.

- [27] The term carrier also includes a buffer or a pH adjusting agent or a composition containing the same; typically, the buffer is a salt prepared from an organic acid or base. Representative buffers include organic acid salts such as salts of citric acid, ascorbic acid, gluconic acid, carbonic acid, tartaric acid, succinic acid, acetic acid, or phthalic acid, Tris, tromethamine hydrochloride, or phosphate buffers. Additional carriers include polymeric excipients/additives such as polyvinylpyrrolidones, ficolls (a polymeric sugar), dextrates (e.g., cyclodextrins, such as 2-hydroxypropyl- α -cyclodextrin), polyethylene glycols, flavoring agents, antimicrobial agents, sweeteners, antioxidants, antistatic agents, surfactants (e.g., polysorbates such as "TWEEN 20" and "TWEEN 80"), lipids (e.g., phospholipids, fatty acids), steroids (e.g., cholesterol), and chelating agents (e.g., EDTA).
- [28] As used herein, the term "pharmaceutically acceptable carrier" encompasses any of the standard pharmaceutical carriers, such as a phosphate buffered saline solution, water, and emulsions, such as an oil/water or water/oil emulsion, and various types of wetting agents. The compositions and medicaments which are manufactured and/or used in accordance with the present invention and which include the particular enzyme whose deficiency is to be corrected can include stabilizers and preservatives and any of the above noted carriers with the additional proviso that they be acceptable for use in vivo. For example of carriers, stabilizers and adjuvants, see Martin REMINGTON'S PHARM. SCI., 15th Ed. (Mack Publ. Co., Easton (1975) and Williams & Williams, (1995), and in the "PHYSICIAN'S DESK REFERENCE", 52nd ed., Medical Economics, Montvale, N.J. (1998).
- [29] A "subject," "individual" or "patient" is used interchangeably herein, which refers to a vertebrate, preferably a mammal, more preferably a human. Mammals include, but are not limited to, mice, rats, monkeys, humans, farm animals, sport animals, and pets.

- [30] As used herein, the term "modulate" means to vary the amount or intensity of an effect or outcome, *e.g.*, to enhance, augment, diminish or reduce.
- [31] As used herein the term "ameliorate" is synonymous with "alleviate" and means to reduce or lighten. For example, one may ameliorate the symptoms of a disease or disorder by making them more bearable.
- [32] For identification of structures in the human brain, see, *e.g.*, The Human Brain: Surface, Three-Dimensional Sectional Anatomy With MR1, and Blood Supply, 2nd ed., eds. Deuteron et al., Springer Vela, 1999; Atlas of the Human Brain, eds. Mai et al., Academic Press; 1997; and Co-Planar Stereotaxic Atlas of the Human Brain: 3-Dimensional Proportional System: An Approach to Cerebral Imaging, eds. Tamarack et al., Thyme Medical Pub., 1988. For identification of structures in the mouse brain, see, *e.g.*, The Mouse Brain in Stereotaxic Coordinates, 2nd ed., Academic Press, 2000.
- [33] The inventors have discovered that intraventricular delivery to the brain of lysosomat hydrolase enzymes to patients who are deficient in the enzymes, leads to improved metabolic status of both the brain and the affected visceral (non-CNS) organs. This is particularly true when the delivery rate is slow, relative to a bolus delivery. Lysosomal storage diseases caused by a deficiency in a particular enzyme, such as those diseases identified in Table 1 above, may therefore be treated or prevented by the intraventricular administration of the respective enzyme. One particularly useful enzyme for treating Niemann-Pick A or 13 is acid sphingomyelinase (ASM), such as that shown in SEQ ID NO: 1.¹ One particularly useful enzyme for treating Gaucher disease is glucocerebrosidase. One particularly useful enzyme for treating MPS I disease is alpha-L-Iduronidase. One particularly useful enzyme for treating MPS II disease is iduronate-2-sulfatase. One particularly useful enzyme for treating Pompe disease, or glycogen storage disease type U (GSDII), also termed acid maltase deficiency (AMD) is acid alpha-glucosidase. One particularly useful enzyme for treating classic late infantile Batten

¹ Residues 1-46 constitute the signal sequence which is cleaved upon secretion.

disease (CLN2) is tripeptidyl peptidase. The enzymes that are used and/or administered in accordance with the present invention may be recombinant forms of the enzymes which are produced using methods that are well-known in the art. In one embodiment, the enzyme is a recombinant human enzyme.

- [34] Administration of lysosomal enzymes, and more particularly lysosomal hydrolase enzymes, to patients who are deficient in the enzymes may be performed into any one or more of the brain's ventricles, which are filled with cerebrospinal fluid (CSF). CSF is a clear fluid that fills the ventricles, is present in the subarachnoid space, and surrounds the brain and spinal cord. CSF is produced by the choroid plexuses and via the weeping or transmission of tissue fluid by the brain into the ventricles. The choroid plexus is a structure lining the floor of the lateral ventricle and the roof of the third and fourth ventricles. Certain studies have indicated that these structures are capable of producing 400-600 ccs of fluid per day consistent with an amount to fill the central nervous system spaces four times in a day. In adults, the volume of this fluid has been calculated to be from 125 to 150 ml (4-5 oz). The CSF is in continuous formation, circulation and absorption. Certain studies have indicated that approximately 430 to 450 ml (nearly 2 cups) of CSF may be produced every day. Certain calculations estimate that production equals approximately 0.35 ml per minute in adults and 0.15 per minute in infants. The choroid plexuses of the lateral ventricles produce the majority Of CSF. It flows through the foramina of Monro into the third ventricle where it is added to by production from the third ventricle and continues down through the aqueduct of Sylvius to the fourth ventricle. The fourth ventricle adds more CSF; the fluid then travels into the subarachnoid space through the foramina of Magendie and Luschka. It then circulates throughout the base of the brain, down around the spinal cord and, upward over the cerebral hemispheres. The CSF empties into the blood via the arachnid villi and intracranial vascular sinuses, thereby delivering the enzymes that are inflsed into the ventricles to not only the brain but also to the visceral organs that are known to be affected in LSDs.

- [35] Although a particular amino acid sequence is shown in SEQ ID NO: 1, variants of that sequence which retain activity, *e.g.*, normal variants in the human population, can be used as well. Typically these normal variants differ by just one or two residues from the sequence shown in SEQ ID NO: 1. The variants of SEQ ID NO: 1 that are to be used in accordance with the present invention, whether naturally occurring or not, should be at least 95%, 96%, 97%, 98%, or 99% identical to SEQ ID NO: 1. Variants of other enzymes may be used in accordance with the present invention. However, irrespective of the enzyme that is used, variants which are associated with disease or reduced activity should not be used. Typically the mature form of the enzyme will be delivered. In the case of SEQ ID NO: 1, the mature form will begin with residue 47 as shown in SEQ ID NO: 1. Variants which are associated with disease are shown in FIG. 7. In a similar manner, normal variants in the human population of such LSD enzymes as glucocerebrosidase, alpha-L-Iduronidase, iduronate-2-sulfatase, acid alpha- glucosidase, and tripeptidyl peptidase that which retain enzymatic activity can be used as well.
- [36] Kits according to the present invention are assemblages of separate components. While they can be packaged in a single container, they can be subpackaged separately. Even a single container can be divided into compartments. Typically a set of instructions will accompany the kit and provide instructions for delivering the enzymes, *e.g.*, the lysosomal hydrolase enzymes, intraventricularly. The instructions may be in printed form, in electronic form, as an instructional video or DVD, on a compact disc, on a floppy disc, on the Internet with an address provided in the package, or a combination of these means. Other components, such as diluents, buffers, solvents, tape, screws, and maintenance tools can be provided in addition to the enzyme, one or more cannulae or catheters, and/or a pump.
- [37] The populations treated by the methods of the invention include, but are not limited to, patients having, or who are at risk for developing, a neurometabolic disorder, *e.g.*, an LSD, such as the diseases listed in Table I, particularly if such a disease affects the CNS and visceral organs. In an illustrative embodiment, the disease is type A Niemann-Pick disease.

- [38] An ASM or other lysosomal hydrolase enzyme can be incorporated into a pharmaceutical composition useful to treat, *e.g.*, inhibit, attenuate, prevent, or ameliorate, a condition characterized by an insufficient level of a lysosomal hydrolase activity. The pharmaceutical composition will be administered to a subject suffering from a lysosomal hydrolase deficiency or someone who is at risk of developing said deficiency. The compositions should contain a therapeutic or prophylactic amount of the ASM or other lysosomal hydrolase enzyme, in a pharmaceutically-acceptable carrier. The pharmaceutical carrier can be any compatible, non-toxic substance suitable to deliver the polypeptides to the patient. Sterile water, alcohol, fats, and waxes may be used as the carrier. Pharmaceutically-acceptable adjuvants, buffering agents, dispersing agents, and the like, may also be incorporated into the pharmaceutical compositions. The carrier can be combined with the ASM or other lysosomal hydrolase enzyme in any form suitable for administration by intraventricular injection or infusion (which form is also possibly suitable for intravenous or intrathecal administration) or otherwise. Suitable carriers include, for example, physiological saline, bacteriostatic water, Cremophor EL.TM. (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS), other saline solutions, dextrose solutions, glycerol solutions, water and oils emulsions such as those made with oils of petroleum, animal, vegetable, or synthetic origin (peanut oil, soybean oil, mineral oil, or sesame oil). An artificial CSF can be used as a carrier. The carrier will preferably be sterile and free of pyrogens. The concentration of the ASM or other lysosomal hydrolase enzyme in the pharmaceutical composition can vary widely, *i.e.*, from at least about 0.01% by weight, to 0.1 % by weight, to about 1% weight, to as much as 20% by weight or more of the total composition.
- [39] For intraventricular administration of ASM or other lysosomal hydrolase enzyme, the composition must be sterile and should be fluid. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be

referable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in the composition.

- [40] Dosage of ASM or other lysosomal hydrolase enzyme may vary somewhat from individual to individual, depending on the particular enzyme and its specific in vivo activity, the route of administration, the medical condition, age, weight or sex of the patient, the patient's sensitivities to the ASM or other lysosomal hydrolase enzyme or components of the vehicle, and other factors which the attending physician will be capable of readily taking into account. While dosages may vary depending on the disease and the patient, the enzyme is generally administered to the patient in amounts of from about 0.1 to about 1000 milligrams per 50 kg of patient per month. In one embodiment, the enzyme is administered to the patient in amounts of about 1 to about 500 milligrams per 50 kg of patient per month. In other embodiments, the enzyme is administered to the patient in amounts of about 5 to about 300 milligrams per 50kg of patient per month, or about 10 to about 200 milligrams per 50kg of patient per month.
- [41] The rate of administration is such that the administration of a single dose may be administered as a bolus. A single dose may also be infused over about 1-5 minutes, about 5-10 minutes, about 10-30 minutes, about 30-60 minutes, about 1-4 hours, or consumes more than four, five, six, seven, or eight hours. It may take more than 1 minute, more than 2 minutes, more than 5 minutes, more than 10 minutes, more than 20 minutes, more than 30 minutes, more than 1 hour, more than 2 hours, or more than 3 hours. Applicants have observed that, while bolus intraventricular administration may be effective, slow infusion is very effective. While applicants do not wish to be bound by any particular theory of operation, it is believed that the slow infusion is more effective due to the turn-over of the cerebrospinal fluid (CSF). While estimates and calculations in the literature vary, the cerebrospinal fluid is believed to turn over within about 4, 5, 6, 7, or 8 hours in humans. In one embodiment, the slow infusion time of the invention should be metered so that it is . about equal to or greater than the turn-over time of the CSF. Turn-over time may depend on the species, size, and age of the subject but may be determined using methods known in the art. Infusion may also be continuous over a period of one or more days. The patient

may be treated once, twice, or three or more times a month, *e.g.*, weekly, *e.g.* every two weeks. Infusions may be repeated over the course of a subject's life as dictated by re-accumulation of the disease's substrate in the brain or visceral organs. Re-accumulation may be determined by any of the techniques that are well known in the art for the identification and quantization of the relevant substrate, which techniques may be performed on one or more samples taken from the brain and/or from one or more of the visceral organs. Such techniques include enzymatic assays and/or immunoassays, *e.g.* radioimmunoassays or ELISAs.

- [42] The CSF empties into the blood via the arachnid villi and intracranial vascular sinuses, thereby delivering the infused enzymes to the visceral organs that are known to be affected in LSDs. The visceral organs which are often affected in Niemann-Pick disease are the lungs, spleen, kidney, and liver. The slow intraventricular infusion provides diminished amounts of the substrate for an administered enzyme in at least the brain and potentially in visceral organs. The reduction in substrate accumulated in the brain, lungs, spleen, kidney, and/or liver may be dramatic. Reductions of greater than 10 %, 20 %, 30 %, 40 %, 50 %, 60 %, 70 %, 80 %, 90 % can be achieved. The reduction achieved is not necessarily uniform from patient to patient or even from organ to organ within a single patient. Reductions can be determined by any of the techniques that are well known in the art, *e.g.*, by enzymatic assays and/or immunoassay techniques, as discussed elsewhere herein.

- [43] If desired, the human brain structure can be correlated to similar structures in the brain of another mammal. For example, most mammals, including humans and rodents, show a similar topographical organization of the entorhinal-hippocampus projections, with neurons in the lateral part of both the lateral and medial entorhinal cortex projecting to the dorsal part or septal pole of the hippocampus, whereas the projection to the ventral hippocampus originates primarily from neurons in medial parts of the entorhinal cortex (Principles of Neural Science, 4th ed., eds Kandel et al., McGraw-Hill, 1991; The Rat Nervous System, 2nd ed., ed. Paxinos, Academic Press, 1995). Furthermore, layer II cells of the entorhinal cortex project to the dentate gyms, and they terminate in the outer

two-thirds of the molecular layer of the dentate gyrus. The axons from layer III cells project bilaterally to the cornu ammonis areas CA1 and CA3 of the hippocampus, terminating in the stratum lacunose molecular layer.

- [44] In an illustrative embodiment, the administration is accomplished by infusion of the LSD enzyme into one or both of the lateral ventricles of a subject or patient. By infusing into the lateral ventricles, the enzyme is delivered to the site in the brain in which the greatest amount of CSF is produced. The enzyme may also be infused into more than one ventricle of the brain. Treatment may consist of a single infusion per target site, or may be repeated. Multiple infusion/injection sites can be used. For example, the ventricles into which the enzyme is administered may include the lateral ventricles and the fourth ventricle. In some embodiments, in addition to the first administration site, a composition containing the LSD enzyme is administered to another site which can be contralateral or ipsilateral to the first administration site. Injections/infusions can be single or multiple, unilateral or bilateral.

- [45] To deliver the solution or other composition containing the enzyme specifically to a particular region of the central nervous system, such as to a particular ventricle, *e.g.*, to the lateral ventricles or to the fourth ventricle of the brain, it may be administered by stereotaxic microinjection. For example, on the day of surgery, patients will have the stereotaxic frame base fixed in place (screwed into the skull). The brain with stereotaxic frame base (MRI-compatible with fiducial markings) will be imaged using high resolution MRI. The MRI images will then be transferred to a computer that runs stereotaxic software. A series of coronal, sagittal and axial images will be used to determine the target site of vector injection, and trajectory. The software directly translates the trajectory into 3-dimensional coordinates appropriate for the stereotaxic frame. Burr holes are drilled above the entry site and the stereotaxic apparatus localized with the needle implanted at the given depth. The enzyme solution in a pharmaceutically acceptable carrier will then be injected. Additional routes of administration may be used, *e.g.*, superficial cortical application under direct visualization, or other non-stereotaxic application.

- [46] One way for delivering a slow infusion is to use a pump. Such pumps are commercially available, for example, from Alzet (Cupertino, CA) or Medtronic (Minneapolis, MN). The pump may be implantable. Another convenient way to administer the enzymes is to use a cannula or a catheter. The cannula or catheter may be used for multiple administrations separated in time. Cannulae and catheters can be implanted stereotactically. It is contemplated that multiple administrations will be used to treat the typical patient with a lysosomal storage disease. Catheters and pumps can be used separately or in combination.
- [47] The lysosomal storage diseases (LSD) includes over forty genetic disorders, many of which involve genetic defects in various lysosomal hydrolases. Representative lysosomal storage diseases and the associated defective enzymes are listed in Table 1.
- [48] Gaucher disease results as a consequence of an inherited deficiency of the lysosomal hydrolase glucocerebrosidase (GC), leading to the accumulation of its substrate, glucosylceramide (GL-1), in the lysosomes of histiocytes. The progressive accumulation of GL-1 in tissue macrophages (Gaucher cells) occurs in various tissues. The extent of the accumulation is dependent in part on the genotype. Clinically, three different Gaucher phenotypes are recognized, the non-neuropathic type 1, which is the most common with onset ranging from early childhood to adulthood, and the neuropathic types 2 and 3, presenting in infancy and early childhood, respectively. Any of these phenotypes may be treated in accordance with the present invention. The primary clinical manifestations common to all forms of Gaucher disease are hepatosplenomegaly, cytopenia, pathological bone fractures and, occasionally, pulmonary failure. A detailed discussion of Gaucher disease may be found in the Online Metabolic & Molecular Bases of Inherited Diseases, Part 16, Chapter 146 and 146.1 (2007). In patients with type 2 and type 3 Gaucher disease in whom there *is* significant central nervous system involvement, intraventricular delivery of the defective LSD enzyme leads to improved metabolic status of the brain and potentially the affected visceral (non-CNS) organs. Intraventricular delivery of the defective LSD enzyme in subjects with Gaucher type 1 disease leads to improved metabolic status of affected visceral (non-CNS) organs. There are animal models of

Gaucher disease, which have derived from mouse models created by targeted disruption of the corresponding mouse gene. For example, a Gaucher mouse model harboring the D409V mutation in the mouse GC locus exists (Xu, Y-H et al. (2003); *Ain. J. Pathol.* 163:2093-2101). The heterozygous mouse, *gba*D409V/null, exhibits~ 5% of normal GC activity in visceral tissues and develops lipid-engorged macrophages (Gaucher cells) in the liver, spleen, lung and bone marrow by 4 months of age. This model is a suitable system in which to evaluate the benefits of and to determine the conditions for the intraventricular delivery of the defective LSD enzyme in subjects with Gaucher disease.

- [49] Niemann-Pick disease (NPD) is a lysosomal storage disease and is an inherited neurometabolic disorder characterized by a genetic deficiency in acid sphingomyelinase (ASM; sphingomyelin cholinephosphohydrolase, EC 3.1.3.12). The lack of functional ASM protein results in the accumulation of sphingomyelin substrate within the lysosomes of neurons and glia throughout the brain. This leads to the formation of large numbers of distended lysosomes in the perikaryon, which are a hallmark feature and the primary cellular phenotype of type A NPD. The presence of distended lysosomes correlates with the loss of normal cellular function and a progressive neurodegenerative course that leads to death of the affected individual in early childhood (The Metabolic and Molecular Bases of Inherited Diseases, eds. Scriver *et al.*, McGraw-Hill, New York, 2001, pp. 35893610). Secondary cellular phenotypes (e.g. additional metabolic abnormalities) are also associated with this disease, notably the high level accumulation of cholesterol in the lysosomal compartment. Sphingomyelin has strong affinity for cholesterol, which results in the sequestering of large amounts of cholesterol in the lysosomes of ASMKO mice and human patients (Leventhal *et al.* (2001) *J. Biol. Chem.*, 276:44976-44983; Slotte (1997) *Subcell. Biochem.*, 28:277-293; and Viana *et al.* (1990) *J. Med. Genet.*, 27:499-504.) A detailed discussion of NPD disease may be found in the Online Metabolic & Molecular Base of Inherited Diseases, Part 16, Chapter 144 (2007). There are animal models of NPD: For example ASMKO mice are an accepted model of types A and B Niemann-Pick disease (Horinouchi et al. (1995) *Nat. Genetics*, 10:288-293; Jin et al. (2002) *J. Clin. Invest.*, 109:1183-1191; and Otterbach (1995) *Cell*, 81:1053-1061). Intraventricular

delivery of the defective LSD enzyme leads to improved metabolic status of the brain and the affected visceral (non-CNS) organs.

- [50] Mucopolysaccharidoses (MPS) are a group of lysosomal storage disorders caused by deficiencies of enzymes catalyzing the degradation of glycosaminoglycans (mucopolysaccharides). There are 11 known enzyme deficiencies that give rise to 7 distinct MPS, including MPS I (Hurler, Scheie, and Hurler-Scheie Syndromes) and MPS II (Hunter Syndrome). Any MPS can be treated in accordance with the present invention. A detailed discussion of MPS may be found in the Online Metabolic & Molecular Bases of Inherited Diseases, Part 16, Chapter 136 (2007). There are numerous animal models of MPS, which have derived from naturally occurring mutations in dogs, cats, rats, mice, and goats, as well as mouse models created by targeted disruption of the corresponding mouse gene. The biochemical and metabolic features of these animal models are generally quite similar to those found in humans; however, the clinical presentations may be milder. For example, accepted models for MPS I include a murine model [Clark, LA et al., Hum. Mol. Genet. (1997), 6:503] and a canine model [Menon, ICP et al., Genomics (1992), 14:763. For example, accepted models for MPS II include a mouse model [Muenzer, J. et al., (2002), Acta Paediatr. Suppl.;91(439):98-9]. In the MPS that have central nervous system involvement, such as is found in patients with MPS I and MPS intraventricular delivery of the defective LSD enzyme leads to improved metabolic status of the brain and potentially the affected visceral (non-CNS) organs.
- [51] Pompe disease, or glycogen storage disease type II (GSM), also termed acid maltase deficiency (AMD) is an inherited disorder of glycogen metabolism resulting from defects in activity of the lysosomal hydrolase acid alpha- glucosidase in all tissues of affected individuals. The enzyme deficiency results in intralysosomal accumulation of glycogen of normal structure in numerous tissues. The accumulation is most marked in cardiac and skeletal muscle and in hepatic tissues of infants with the generalized disorder. In late-onset GSDII, intralysosomal accumulation of glycogen is virtually limited to skeletal muscle and is of lesser magnitude. Electromyographic abnormalities suggestive of the diagnosis include pseudomyotonic discharges and irritability, but in juvenile- and adult-

onset patients, the abnormalities can vary in different muscles. CAT scans can reveal the site(s) of affected muscles. Most patients have elevated blood plasma levels of creatine kinase (CK) and elevations in hepatic enzymes, particularly in adult-onset patients, can be found. There are several naturally occurring animal models of the infantile- and late-onset disease. There is a knockout mouse model [Bijvoet AG et al., *Hum. Mol. Genet.* (1998); 7:53-62]. Ameliorative effects of enzyme therapy have been described in knockout mice [Raben, N et al., *Mol. Genet. Metab.* (2003); 80:159-69] and in a quail model. Intraventricular delivery of the defective LSD enzyme leads to improved metabolic status of the brain and potentially the affected visceral (non-CNS) organs.

- [52] The neuronal ceroid lipofuscinoses (NCL) are a group of neurodegenerative disorders distinguished from other neurodegenerative diseases by the accumulation of autofluorescent material ("aging pigment") in the brain and other tissues. The major clinical features include seizures, psychomotor deterioration, blindness, and premature death. Distinct subgroups of NCL have been recognized that differ in the age of onset of symptoms and the appearance of the storage material by electron microscopy. Three major groups—infantile (INCL), classical late infantile (LINCL), and juvenile (JNCL, also referred to as Batten disease)—are caused by autosomal recessive mutations in the CLN1, CLN2, and CLN3 genes, respectively. The protein products of the CLN1 (palmitoyl-protein thioesterase) and CLN2 (tripeptidyl peptidase or pepinase) genes are soluble lysosomal enzymes, whereas the CLN3 protein (battenin) is a lysosomal membrane protein, as is (tentatively) the CLN5 protein. The identification of mutations in genes encoding lysosomal proteins in several forms of NCL has led to the recognition of the lipofuscinoses as true lysosomal storage disorders. Any subgroup of NCL can be treated in accordance with the present invention. A detailed discussion of NCL disease may be found in the *Online Metabolic & Molecular Bases of Inherited Diseases*, Part 16, Chapter 154 (2007). Naturally occurring NCL disorders have been described in the sheep, dog, and mouse models have been derived by targeted disruption of a corresponding mouse gene [see e.g., Katz, ML et al., *J. Neurosci. Res.* (1999); 57:551-6; Cho, SK et al., *Glycobiology* (2005); 15:637-48.] Intraventricular delivery of the defective LSD enzyme

leads to improved metabolic status of the brain and possibly the affected visceral (non-CNS) organs.

- [53] A detailed discussion of additional lysosomal storage disorders disclosed in Table 1, in which intraventricular delivery of the defective LSD enzyme in the disease, may be found in the Online Metabolic & Molecular Bases of Inherited Diseases, Part 16 (2007).
- [54] The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific examples which are provided herein for purposes of illustration only, and are not intended to limit the scope of the invention.

EXAMPLE 1

Animal model

- [55] ASMKO mice are an accepted model of types A and B Niemann-Pick disease (Horinouchi et al. (1995) Nat. Genetics, 10:288-293; Jin et al. (2002) 3. Clin. Invest., 109:1183-1191; and Otterbach (1995) Cell, 81:1053-1061). Niemann-Pick disease (NPD) is classified as a lysosomal storage disease and is an inherited neurometabolic disorder characterized by a genetic deficiency in acid sphingomyelinase (ASM; sphingomyelin cholinephosphohydrolase, EC 3.1.3.12). The lack of functional ASM protein results in the accumulation of sphingomyelin substrate within the lysosomes of neurons and glia throughout the brain. This leads to the formation of large numbers of distended lysosomes in the perikaryon, which are a hallmark feature and the primary cellular phenotype of type A NPD. The presence of distended lysosomes correlates with the loss of normal cellular function and a progressive neurodegenerative course that leads to death of the affected individual in early childhood (The Metabolic and Molecular Bases of Inherited Diseases, eds. Scriver *et al.*, McGraw-Hill, New York, 2001, pp. 3589-3610). Secondary cellular phenotypes (*e.g.*, additional metabolic abnormalities) are also associated with this

disease, notably the high level accumulation of cholesterol in the lysosomal compartment. Sphingomyelin has strong affinity for cholesterol, which results in the sequestering of large amounts of cholesterol in the lysosomes of ASMKO mice and human patients (Leventhal *et al.* (2001) J. Biol. Chem., 276:44976-44983; Slotte (1997) Subcell. Biochem., 28:277-293; and Viana *et al.* (1990) J. Med. Genet., 27:499-504.)

EXAMPLE 2

"Continuous Intraventricular infusion of rhASM in the ASMKO mouse"

- [56] Goal: To determine what effect intraventricular infusion of recombinant human ASM (rhASM) has on storage pathology (*i.e.*, sphingomyelin and cholesterol storage) in the ASMKO mouse brain

- [57] Methods: ASMKO mice were stereotactically implanted with an indwelling guide cannula between 12 and 13 weeks of age. At 14 weeks of age mice were infused with .250 mg of hASM (n=5) over a 24 h period (~.01 mg/h) for four straight days (1 mg total was administered) using an infusion probe (fits inside the guide cannula) which is connected to a pump. Lyophilized hASM was dissolved in artificial cerebral spinal fluid (aCSF) prior to infusion. Mice were sacrificed 3 days post infusion. At sacrifice mice were overdosed with euthasol (>150 mg/kg) and then perfused with PBS or 4% paraformaldehyde. Brain, liver, lung and spleen were removed and analyzed for sphingomyelin (SPM) levels. Brain tissue was divided into 5 sections before SPM analysis (S1 = front of brain, S5 = back of the brain; see Fig. 1)

Table 2.

Group	Treatment	
ASMKO	.250 mg./24 h (1mg total)	5
ASMKO	None	4
WT	None	4

- [58] Results summary: Intraventricular infusion of hASM at .250mg/24 h for 4 continuous days (1 mg total) resulted in hASM staining and filipin (*i.e.*, cholesterol storage) clearance throughout the ASMKO brain. Biochemical analysis showed that intraventricular infusion of hASM also led to a global reduction in SPM levels throughout the brain. SPM levels were reduced to that of wild type (WT) levels. A significant reduction in SPM was also observed in the liver and spleen (a downward trend was seen in the lung).

EXAMPLE 3

"Intraventricular delivery of hASM in ASMKO Mice II"

- [59] Goal: to determine lowest efficacious dose over a 6 h infusion period.
- [60] Methods: ASMKO mice were stereotactically implanted with an indwelling guide cannula between 12 and 13 weeks of age. At 14 weeks of age mice were infused over a 6 hour period at one of the following doses of hASM: 10 mg/kg (.250 mg; n=12), 3 mg/kg (.075 mg; n=7), 1 mg/kg (.025 mg; n=7), .3 mg/kg (.0075 mg; n=7), or aCSF (artificial cerebral spinal fluid; n=7). Two mice from each dose level were perfused with 4% parformaldehyde immediately following the 6 h infusion to assess enzyme distribution in the brain (blood was also collected from these to determine serum hASM levels). The

remaining mice from each group were sacrificed 1 week post infusion. Brain, liver, and lung tissue from these mice was analyzed for SPM levels as in study 05-0208.

Table 3.

Group	Treatment	n
ASMKO	0.250 mg (10 mg/kg)	12
ASMKO	0.075 mg (3 mg/kg)	7
ASMKO	0.025 mg (1 mg/kg)	7
ASMKO	0.0075 mg(.3 mg/kg)	7
ASMKO	aCSF	7
WT	None	7

- [61] Results summary: Intraventricular hASM over a 6h period led to a significant reduction in SPM levels throughout the brain regardless of dose. Brain SPM levels in mice treated with doses $> .025$ mg were reduced to WT levels. Visceral organ SPM levels were also significantly reduced (but not to WT levels) in a dose dependent manner. In support of this finding hASM protein was also detected in the serum of ASMKO mice infused with hASM protein. Histological analysis showed that hASM protein was widely distributed throughout the brain (from S1 to S5) after intraventricular administration of hASM.

EXAMPLE 4

"Intraventricular infusion of rhASM in ASMKO mice III"

- [62] Goal: To determine (1) the time it takes for SPM to reaccumulate within the brain (and spinal cord) after a 6 h infusion of hASM (dose = .025 mg); (2) if there are sex differences in response to intraventricular hASM administration (previous experiments demonstrate that there are sex differences in substrate accumulation in the liver, whether or not this occurs in the brain is unknown).
- [63] Methods: ASMKO mice were stereotactically implanted with an indwelling guide cannula between 12 and 13 weeks of age. At 14 weeks of age mice were infused over a 6h period with .025 mg of hASM. After intraventricular delivery of hASM mice were sacrificed either at 1 week post infusion (n = 7 males, 7 females), or at 2 weeks post infusion (n = 7 males, 7 females) or at 3 weeks post infusion (n = 7 males, 7 females). At sacrifice the brain, spinal cord, liver and lung were removed for SPM analysis.

Table 4.

Group	Treatment	N	Sacrifice.
male ASMKO	.025 mg	7	1 week post infusion
Female ASMKO	.025 mg	7	1 week post infusion
male ASMKO	.025 mg	7	2 weeks post infusion
Female ASMKO	.025 mg	7	2 weeks post infusion
male ASMKO	.025 mg	7	3 weeks post infusion
Female ASMKO	.025 mg	7	3 weeks post infusion

male ASMKO	aCSF	7	1 week post infusion
FemaleASMKO	aCSF	7	1 week phst infusion
male WT	None	7	1 week post infusion
Female WT	None	7	1 week post infusion

[64] Tissue samples are prepared for SPM analysis.

EXAMPLE 5

"Effect of intraventricular infusion of rhASM on cognitive function in ASMKO mice"

[65] Goal: to determine if intraventricular infusion of rhASM alleviates diseased induced cognitive deficits in ASMKO mice.

[66] Methods: ASMKO mice are stereotactically implanted with an indwelling guide cannula between 9 and 10 weeks of age. At 13 weeks of age mice are infused over a 6h period with .025 mg of hASM. At 14 and 16 weeks of age mice undergo cognitive testing using the Barnes maze.

EXAMPLE 6

"hASM protein distribution within the ASMKO CNS after intraventricular infusion"

[67] Goal: to determine the distribution of hASM protein (as function of time) within the brain. and spinal cord of ASMKO mice after intraventricular infusion

- [68] Methods: ASMICO mice are stereotactically implanted with an indwelling guide cannula between 12 and 13 weeks of age. At 14 weeks of age mice are infused over a 6h period with :025 mg of hASM. Following infusion procedure mice are either sacrificed immediately or 1 week or 2 weeks or 3 weeks later.

Table 5. Indicates the infusion times that may be used with a particular enzyme for the treatment of the disease in which it is deficient as indicated in Table 1.

Enzyme for Infusion	INFUSION TIME										
	BOLUS	1-5 min	5 - 10 min	10 - 30 min	30 - 60 min	1-4 hrs	more than 4 hrs	more than 5 hrs	more than 6 hrs	more than 7 hrs	more than 8 hrs
Aspartylglucosaminidase .alpha.-Galactosidase A	X	X	X	X	X	X	X	X	X	X	X
Palmitoyl Protein Thioesterase	X	X	X	X	X	X	X	X	X	X	X
Tripeptidyl Peptidase	X	X	X	X	X	X	X			X	X
Lysosomal Transmembrane Protein	X	X	X	X	X	X	X	X	X	X	X
Cysteine transporter	X	X	X	X	X	X	X	X	X	X	X
Acid ceramidase	X	X	X	X	X	X	X	X	X		X
Acid .alpha.-L-fucosidase	X	X	X	X	X	X	X	X	X	X	X
Protective protein/cathepsin A	X	X	X	X	X	X	X	X	X	X	X
Acid.beta.-glucosidase, or Acid.beta.-galactosidase	X	X	X	X	X	X	X	X	X	X	X
Iduronate-2-sulfatase	X	X	X	X	X	X	X	X	X	X	X
alpha.L-Iduronidase	X	X	X	X	X	X	X	X	X	X	X
Galactocerebrosidase	X	X	X	X	X	X	X	X	X	X	X
Acid.alpha.-mannosidase	X	X	X	X	X	X	X	X	X	X	X
Acid.beta.-mannosidase	X	X	X	X	X	X	X	X	X	X	X
Arylsulfatase B	X	X	X	X	X	X	X	X	X	X	X
Arylsulfatase A	X	X	X	X	X	X	X	X	X	X	X
N-Acetylgalactosamine-6- sulfate	X	X	X	X	X	X	X	X	X	X	X
Acid.beta.-galactosidase	X	X	X	X	X	X	X	X	X	X	X
N-Acetylglucosamine-1	X	X	X	X	X	X	X	X	X	X	X
Acid sphingomyelinase	X	X	X	X	X	X	X	X	X	X	X

References

- 1) Belichenko PV, Dickson PI, Passage M, Jungles S, Mobley WC, Kakkis ED. Penetration, diffusion, and uptake of recombinant human alpha-l-iduronidase after intraventricular injection into the rat brain. *Mol Genet Metab.* 2005;86(1-2):141-9.
- 2) Kakkis E, McEntee M, Vogler C, Le S, Levy B, Belichenko P, Mobley W, Dickson P, Hanson S, Passage M. Intrathecal enzyme replacement therapy reduces lysosomal storage in the brain and meninges of the canine model of MPS I. *Mol Genet Metab.* 2004;83(1-2):163-74.
- 3) Bembi B, Ciana G, Zanatta M, et al. Cerebrospinal-fluid infusion of alglucerase in the treatment for acute neuronopathic Gaucher's disease. *Pediatr Res* 1995;38:A425.
- 4) Lonser RR, Walbridge S, Murray GJ, Aizenberg MR, Vortmeyer AO, Aerts JM, Brady RO, Oldfield EH. Convection perfusion of glucocerebrosidase for neuronopathic Gaucher's disease. *Ann Neurol.* 2005 Apr;57(4):542-8.

WE CLAIM:

1. Use of an enzyme for the manufacture of a medicament for prevention or treatment of a lysosomal storage disease in a patient, which disease is caused by a deficiency of the enzyme within the patient which leads to an accumulation of the enzyme's substrate,

wherein said enzyme is for intraventricular administration to the brain of the patient, wherein the administration of a single dose of the enzyme consumes more than six hours, and wherein the administration of the enzyme leads to a reduction in the level of the substrate in the brain and in one or more of the visceral organs and

wherein the enzyme is acid beta glucosidase, acid alpha glucosidase, alpha galactosidase A, alpha-L-iduronidase, iduronate-2-sulfatase, or tripeptidyl peptidase.

2. Use of an enzyme for the prevention or treatment of a lysosomal storage disease in a patient, which disease is caused by a deficiency of the enzyme within the patient which leads to an accumulation of the enzyme's substrate,

wherein said enzyme is for intraventricular administration to the brain of the patient, wherein the administration of a single dose of the enzyme consumes more than six hours, and wherein the administration of the enzyme leads to a reduction in the level of the substrate in the brain and in one or more of the visceral organs and

wherein the enzyme is acid beta glucosidase, acid alpha glucosidase, alpha galactosidase A, alpha-L-iduronidase, iduronate-2-sulfatase, or tripeptidyl peptidase.

3. The use of claim 1 or 2, wherein the administration of a single dose of the enzyme consumes more than seven hours.

4. The use of claim 1 or 2, wherein the administration of a single dose of the enzyme consumes more than eight hours.

5. The use of any one of claims 1 to 4, wherein said enzyme is for administration to the lateral ventricles and/or to the fourth ventricle of the brain.

6. The use of any one of claims 1 to 5, wherein said enzyme is for administration to one or both of the lateral ventricles of the brain.
7. The use of any one of claims 1 to 6, wherein the lysosomal storage disease is Gaucher disease and the enzyme is an acid beta glucosidase.
8. The use of any one of claims 1 to 6, wherein the lysosomal storage disease is Pompe disease and the enzyme is an acid alpha glucosidase.
9. The use of any one of claims 1 to 6, wherein the lysosomal storage disease is Fabry disease and the enzyme is an alpha galactosidase A.
10. The use of any one of claims 1 to 6, wherein the lysosomal storage disease is Mucopolysaccharidosis I syndrome and the enzyme is alpha-L-iduronidase.
11. The use of any one of claims 1 to 6, wherein the lysosomal storage disease is Mucopolysaccharidosis II syndrome and the enzyme is iduronate-2-sulfatase.
12. The use of any one of claims 1 to 6, wherein the lysosomal storage disease is classic late infantile Batten disease (CLN2) and the enzyme is tripeptidyl peptidase.
13. The use of any one of claims 1 to 12, wherein said prevention or treatment further comprises determining the re-accumulation of the substrate in the brain and/or in one or more of the visceral organs.
14. The use of any one of claims 1 to 13, wherein the enzyme is for administration using a pump and/or a catheter.
15. The use of any one of claims 1 to 14, wherein the enzyme is for administration using a plurality of infusions of the enzyme.

1/10

Fig. 1




FIG. 2.





FIG. 3.



4/10

FIG. 4.





FIG. 5






Fig. 6.

FIG. 7. Known variants

VARIANT	157	157	1	C -> R (seems to be less active).
VARIANT	242	242	1	G -> R (in NPB).
VARIANT	246	246	1	E -> Q (in NPA; 30% residual activity).
VARIANT	248	248	1	S -> R (in NPA).
VARIANT	302	302	1	L -> P (in NPA; in 23% of NPA Ashkenazi Jewish patients).
VARIANT	319	319	1	H -> Y (in NPA).
VARIANT	371	371	1	P -> S (in NPB).
VARIANT	382	382	1	M -> I (in NPA).
VARIANT	383	383	1	N -> S (in NPB).
VARIANT	389	389	1	N -> T (in NPA).
VARIANT	391	391	1	W -> G (in NPB; low sphingomyelin degradation rates).
VARIANT	421	421	1	H -> Y (in NPB).
VARIANT	436	436	1	S -> R (in NPB).
VARIANT	446	446	1	Y -> C (in NPA).
VARIANT	463	463	1	F -> S (in NPA).
VARIANT	475	475	1	P -> L (in NPA).
VARIANT	496	496	1	R -> L (in NPA; in 32% of NPA Ashkenazi Jewish patients).
VARIANT	537	537	1	Y -> H (in NPA).
VARIANT	577	577	1	G -> S (in NPA).
VARIANT	608	608	1	Missing (in NPB; prevalent among NPB patients from the North-African Maghreb region).

8/10






Fig. 8



Brain Ventricles: Lateral View



Brain Ventricles: Superior View

Figs. 9A and 9B.

10/10

Fig. 10.

